

REVIEW

Influence of open ocean biogeochemistry on aerosol and clouds: Recent findings and perspectives

Karine Sellegrì^{1*}, Rafel Simó^{2*}, Bingbing Wang^{3*}, Peter A. Alpert⁴, Katye Altieri⁵, Susannah Burrows⁶, Frances E. Hopkins⁷, Ilan Koren⁸, Isabel L. McCoy^{9,10}, Jurgita Ovadnevaite¹¹, Matthew Salter¹², and Julia Schmale¹³

Aerosols and clouds are key components of the marine atmosphere, impacting the Earth's radiative budget with a net cooling effect over the industrial era that counterbalances greenhouse gas warming, yet with an uncertain amplitude. Here we report recent advances in our understanding of how open ocean aerosol sources are modulated by ocean biogeochemistry and how they, in turn, shape cloud coverage and properties. We organize these findings in successive steps from ocean biogeochemical processes to particle formation by nucleation and sea spray emissions, further particle growth by condensation of gases, the potential to act as cloud condensation nuclei or ice nucleating particles, and finally, their effects on cloud formation, optical properties, and life cycle. We discuss how these processes may be impacted in a warming climate and the potential for ocean biogeochemistry—climate feedbacks through aerosols and clouds.

Keywords: Atmosphere, Ocean, Aerosol, Cloud, Biogeochemistry

1. Introduction

Clouds contribute about two-thirds of the Earth's albedo (Baker and Peter, 2008; Trenberth et al., 2009), but are still recognized as a component with a large uncertainty in future climate predictions (Bellouin et al., 2020; Zelinka et al., 2020; Ceppi and Nowack, 2021). Liquid clouds in the lower atmosphere, at the top of the marine boundary layer (MBL), are ubiquitous across the tropical and subtropical oceans. They form large marine stratocumulus decks over regions of upwelling in eastern ocean basins, which gradually evolve into trade cumulus clouds over the warmer central and western ocean basins (Wood, 2012). Liquid and mixed-phase (Danker et al., 2022) stratocumulus frequently occur in the midlatitudes as well, often in the cold sectors of midlatitude cyclones, before evolving to more broken clouds toward the equator (McCoy et al., 2017b; McCoy et al., 2023). At higher latitudes, the

presence of mixed phase and ice clouds further complicates our ability to predict cloud occurrences (D'Alessandro et al., 2019). Overall, marine clouds, which stretch from over the coastline to over remote marine environments, are shaped by seasonally variable natural processes. It is essential to understand the impact that naturally emitted aerosols have on cloud formation, and the sensitivity of remote aerosol–cloud–precipitation systems to climate change if we are to reduce the uncertainty in past and future climate predictions (Carslaw et al., 2013; Carslaw et al., 2017; Belloin et al., 2020; Regayre et al., 2020; McCoy et al., 2021; Myers et al., 2021a; Christensen et al., 2022).

Aerosols are coupled tightly to all cloud processes through the microphysical, chemical, and radiative pathways, initiating a number of feedbacks that contribute to the complexity of the climate system. Aerosols can serve

¹CNRS, LaMP, Université Clermont Auvergne, Aubière, France

²Institute of Marine Sciences, Consejo Superior de Investigaciones Científicas (CSIC), Barcelona, Spain

³State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen, China

⁴Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, Villigen PSI, Switzerland

⁵Department of Oceanography, University of Cape Town, Rondebosch, South Africa

⁶Atmospheric Science and Global Change Division, Pacific Northwest National Laboratory, Richland, WA, USA

⁷Plymouth Marine Laboratory, Plymouth, UK

⁸Department of Earth and Planetary Sciences, Weizmann Institute of Sciences, Rehovot, Israel

⁹Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

¹⁰NOAA Chemical Sciences Laboratory, Boulder, CO, USA

¹¹Centre for Climate and Air Pollution Studies, Ryan Institute, School of Natural Sciences, University of Galway, Ireland

¹²Bolin Centre for Climate Research, Stockholm, Sweden

¹³Extreme Environments Research Laboratory, École Polytechnique Fédérale de Lausanne (EPFL) Valais Wallis, Sion, Switzerland

* Corresponding authors:

Emails: karine.sellegrì@uca.fr; rsimo@icm.csic.es; bingbing.wang@xmu.edu.cn

as cloud condensation nuclei (CCN) or ice nucleating particles (INPs) (Heintzenberg and Charlson, 2009; Levin and Cotton, 2009) inducing changes in the cloud microphysics by affecting condensation/evaporation rates, latent heat release, collision rates and efficiency, and droplet mobility. These changes impact cloud macroscopic properties such as reflectance (albedo), lifetime, phase (liquid or ice), size, and precipitation (Altartatz et al., 2014; Rosenfeld et al., 2014). Reciprocally, marine clouds are wet chemistry reactors that may modify aerosol size and composition (Ervens et al., 2018), and potentially their droplet condensation (Shingler et al., 2016) and ice nucleation properties.

Challenges in establishing a well-defined preindustrial baseline (hence, understanding how the “natural” climate works) for aerosols and aerosol–cloud interactions lead to fundamental uncertainties in constraining climate sensitivity (i.e., the temperature increase corresponding to doubled CO₂) and simulating past and future climate (Carslaw et al., 2013). In particular, poorly understood natural aerosol emission processes contribute substantially to uncertainties in the prediction of aerosol forcing (Carslaw et al., 2017). Consequently, although the marine atmosphere constitutes a mixture of natural, locally produced components and species transported long distances from terrestrial sources, we will focus here on the relationships and interplay between open ocean biogeochemistry, natural marine aerosol, and clouds.

Marine aerosols can be emitted at the ocean surface from breaking waves, in the form of film drops from bubble film rupture, or jet drops from seawater droplets ejected from the air–seawater interface upon bubble burst. Droplets and drops that evaporate leaving particulate residues may subsist in the atmosphere as sea spray aerosols (SSA) and are defined as primary marine aerosols. Secondary marine aerosols are formed by nucleation or condensation of gas-phase precursors in the marine atmosphere. These gas-phase precursors may either be directly transferred to the atmosphere at the air–sea interface or be the result of chemical reactions in the atmosphere. Dimethylsulfide (DMS) is a biogenic trace gas that is the main source of sulfuric acid (H₂SO₄) and methane sulfonic acid (MSA) over the oceans, where they are major contributors to secondary aerosol production (de Leeuw et al., 2014). For this reason, DMS has been extensively investigated in the last 3 decades and, to date, it is the only marine biogenic aerosol precursor implemented in Earth System Models (Séférian et al., 2019; Bock et al., 2021). Yet, large uncertainties remain in the space and time resolved estimates of DMS concentrations and fluxes (Galí et al., 2018) as well as in the sensitivity of the climate system to DMS emissions (Fiddes et al., 2018).

Large research consortia and measurement campaigns aimed to integrate marine aerosol emissions and aerosol–cloud interactions have contributed new knowledge over the last 10 years: PEGASO (Plankton-derived Emission of trace Gases and Aerosols in the Southern Ocean; Dall’Osto et al., 2017), SOAP (Surface Ocean Aerosol Production; Law et al., 2017); NETCARE (Network on Climate and Aerosols: Addressing Key Uncertainties in Remote Canadian Environments; Abbatt et al., 2019); ACE-SPACE

(Antarctic Circumnavigation Expedition: Study of Preindustrial-like Aerosols and Their Climate Effects; Schmale et al., 2019); CSET (Cloud System Evolution in the Trades; Albrecht et al., 2019); MarParCloud (Marine biological production, organic aerosol Particles and marine Clouds: a process chain; van Pinxteren et al., 2020); PI-ICE (Ice–ocean–atmosphere interactions and their impacts on climate and ecosystems; Brean et al., 2021); Observations of Clouds, Aerosols, Precipitation, and Surface Radiation over the Southern Ocean (McFarquhar et al., 2021); ACE-ENA (Aerosol and Cloud Experiments in the Eastern North Atlantic; Wang et al., 2022); Sea2Cloud (“Are marine aerosols influencing clouds?”; Sellegri et al., 2023). However, these efforts also demonstrated that understanding the relationship between ocean emissions of primary and secondary aerosols and cloud properties still requires a more comprehensive knowledge of their marine sources and atmospheric interactions. One of the regions identified as particularly important for studying marine aerosol–cloud interactions is the Southern Ocean, which has received much attention during the last decade (Mallet et al., 2023). This region is little affected by anthropogenic aerosols (Hamilton et al., 2014), which allows for the study of the aerosol–cloud system under close to preindustrial conditions, that is, as natural as can be. Furthermore, it is in this region where our understanding of aerosol–cloud interactions is so poor that global models allow too much radiation to reach the surface, resulting in a large warm bias (Bodas-Salcedo et al., 2014; Cesana et al., 2022). Other regions where marine aerosol–cloud interactions are also important and that have been the focus of many recent studies are the high polar regions (see Willis et al., 2023).

In the present article, we provide a summary of the progress in the understanding of the successive steps linking ocean biogeochemistry and clouds over the last 10 years. The aim of this article is not to provide an extensive review of new knowledge but to shed light on certain key findings on critical processes and the identification of relevant unknowns. We focus on open ocean processes (and exclude coastal areas) with the aim to extract information most relevant for large-scale models and satellite retrievals. In Section 2, we address how gas-phase aerosol precursors are produced in seawater through biogeochemical processes before they are transferred to the atmosphere and involved in new particle formation (NPF), with a special focus on DMS. For more detailed chemical processes in the marine atmosphere, the reader is referred to Tinel et al. (2023). Section 3 is dedicated to SSA emission processes, and in particular to new findings about the influence of seawater biogeochemical properties on SSA fluxes. Once in the marine atmosphere, nucleated particles and SSA undergo further transformations that shape their size distribution and chemical composition. In Section 4, we address how new insights into the sources and chemical and physical transformations of marine aerosols are provided by field investigations that combine new chemical tracers with emerging analytical techniques (mass spectrometry, isotopes, etc.). In Sections 5 and 6,

we describe the latest advances in our understanding of the CCN and INP properties of marine aerosols, and of cloud properties in the remote marine atmosphere. In Section 7, we point to the impact that climate change may be having on ocean–aerosol–cloud interactions.

2. DMS and other potential precursors to particle nucleation

2.1. Nucleation and NPF occurrence in the open oceans

NPF is a chain of processes that starts with nucleation of highly condensable species into molecular clusters (approximately 1 nm in size), that grow by condensation and coagulation to a critical size which is thermodynamically stable (2–3 nm). Further rapid growth into the 10 nm size range is needed for particles to survive in the atmosphere, given their potential loss onto the preexisting particles by coagulation processes. Marine nucleation and early growth of newly formed particles have mostly been reported in coastal areas (see de Leeuw et al., 2014, Chapter 4 for a review), for understandable practical considerations, but also because of the scarcity of observed NPF by sizing instruments in the open ocean boundary layer. Here we focus on NPF events in the pristine open ocean, that is, large geographical areas that are most relevant to climate. Brean et al. (2021) reported some nucleation events over Antarctic waters, yet a very few of them were attributed to open ocean conditions. Baccarini et al. (2021) found no NPF events in marine air masses devoid of sea ice or coastal influence during 3 months of observations around Antarctica. The difficulty in elucidating the chemical mechanisms leading to nucleation and NPF in the remote open ocean atmosphere is largely due to this paucity of observations.

The paucity of NPF observations in the MBL, the lower atmosphere, led to the suggestion that most nucleation events occur in the upper troposphere, also termed free troposphere, with a subsequent reinjection into the MBL where newly formed particles would meet enough vapors to grow to a larger size (Quinn et al., 2017; Sanchez et al., 2018; Zheng et al., 2018; Sanchez et al., 2021). Williamson et al. (2019) showed that convective transport of air masses in the tropics results in NPF in the free troposphere, in line with past observations (Clarke et al., 1998). Recently, McCoy et al. (2021) added that the free troposphere of the Southern Ocean is the site of widespread and frequent NPF events

driven by cyclonic uplift, which directly impact CCN concentrations in the MBL.

The hypothesis of the free troposphere being the main contributor of new aerosols to the lower atmosphere has been questioned using satellite-retrieved CCN concentrations (Efraim et al., 2020). Also, Zheng et al. (2021) reported regular and frequent occurrence of NPF events in the *upper* remote MBL over the North Atlantic, facilitated in broken cloud fields. Additional indication that nucleation occurs in the MBL relies on a few chamber experiments in semi-controlled (“in the field”) conditions (Sellegrì et al., 2016) as well as observational evidence for open ocean NPF and growth in ambient air (O’Dowd et al., 2010); these authors pointed out that the newly formed particles grew at a slow rate of the order of 1 nm per hour, therefore preventing the detection of NPF events with the same criteria as over terrestrial areas. Along these lines, Peltola et al. (2022) recently reported that sub-10 nm particle concentrations in the MBL constitute a third of the total aerosol concentration, and that this pool of nanoparticles was not necessarily observed in air masses having traveled across the free troposphere. Therefore, they conclude that NPF events in marine air masses must be sought and detected with different criteria than those classically used for fast-growing particles in continental air masses. The detection of NPF in the open ocean needs to be revisited, with specificities relative to the low concentration of gaseous precursors that result in low cluster formation rates and slow growth. The identity of the different gaseous precursors contributing to NPF in the open ocean is another big unknown (**Figure 1**).

2.2. Potential nucleating chemicals in the open ocean atmosphere

2.2.1. Sulfur-containing species

The well-established role of H_2SO_4 as a ubiquitous nucleating species in most types of environments (Lee et al., 2019), together with its proven marine source through DMS emission and oxidation, point to this species as a natural candidate for NPF should this phenomenon ever occur in the MBL. However, DMS’s relatively long lifetime (around 1 day, Seinfeld and Pandis, 2006) implies that its oxidation to SO_2 and subsequently H_2SO_4 via photochemistry is unlikely to lead to atmospheric binary ($\text{H}_2\text{SO}_4\text{-H}_2\text{O}$) nucleation in the MBL. DMS would still take part in NPF if oxidants other than the OH radical make its oxidation




Figure 1. State of knowledge on nucleating species in the marine boundary layer of the open oceans.

significantly faster. Moreover, DMS-related nucleation is more likely to occur with the concurrent emission of volatile nitrogen compounds such as ammonia and amines. In this case, NPF may be favored through ternary nucleation of these bases with DMS-derived acids and water, as shown in experimental systems (Kirkby et al., 2011; Almeida et al., 2013). However, it is less likely to occur when multiphase chemistry dominates DMS fate, as it is expected to increase in-cloud MSA production at the expense of H_2SO_4 (Hoffmann et al., 2016). Jackson et al. (2020) suggested that DMS would contribute to the marine aerosol pool mostly via condensational growth on preexisting particles rather than through nucleation. In fact, direct measurements of sulfur acids and nitrogen bases in nucleating particles over the pristine ocean have mostly been collected in Antarctic air that is not completely free of coastal influences (Jokinen et al., 2018; Brean et al., 2021). The literature-based conclusion that DMS plays a role in NPF is mainly derived from observed covariations between chlorophyll-*a* (Chl-*a*) as a proxy for marine phytoplankton, MSA, and the number of marine aerosol particles. However, biologically rich waters can produce other chemical compounds than DMS, and MSA can condense on particles that have nucleated from other precursors, besides nucleating itself (Hodshire et al., 2019). Therefore, other chemical paths to particle nucleation precursors need further investigation. For instance, methanethiol (MeSH) is coproduced with DMS in seawater, it is more rapidly oxidized in the atmosphere than DMS and can contribute a large share of the oxidized sulfur burden in the marine atmosphere (Novak et al., 2022). MeSH fluxes have been recently reevaluated upward by a factor of 4 compared to earlier studies (Lawson et al., 2020; Kilgour et al., 2022). Hydroperoxymethyl thioformate (HPMTF, HOCH_2SCHO) has also been detected as another potential DMS oxidation product (Veres et al., 2020; Novak et al., 2021), and its ability to form new particles in the natural MBL should be investigated further. The atmospheric chemistry aspects related to HPMTF are detailed in Tinel et al. (2023).

2.2.2. Iodine

Atmospheric iodine has a unique marine source and is transferred to the MBL as molecular iodine (I_2) or volatile iodocarbons (mostly as CH_3I and CH_2I_2) (see Tinel et al., 2023 for a more complete review on iodine chemistry and chemical sources). Iodine oxides (HIO , HIO_3), formed from the oxidation of I_2 , are expected to be the most efficient nucleating species. Most marine nucleation studies pointing to iodine oxides have been conducted in coastal areas (Sipila et al., 2016; Beck et al., 2021) where macroalgae are the primary producers of iodine. However, a few observations indicate that plankton-derived iodine may be involved in nucleation, opening the way to open ocean NPF (Sellegri et al., 2016). Using chamber experiments, He et al. (2021) demonstrated that nucleation from iodic acid may be as efficient as that of H_2SO_4 in pristine environments. According to the flow tube experiments reported by Gómez Martín et al. (2022), iodine-containing clusters and particles are

formed by aggregation of I_2O_y and HIO_3 in ambient air where water vapor is available. Prados-Roman et al. (2015) estimated that about 75% of IO comes from inorganic precursors. However, a significant biogenic source was needed to explain the gap between measurements and modeled IO over the Southern Ocean (Inamdar et al., 2020). Huang et al. (2022) suggested that a fraction of iodine associated with organics condensing on iodine oxide clusters is recycled back into the gas phase, and this may lengthen iodine lifetime and represent an efficient transport mechanism to remote regions.

2.2.3. Terpenoids and other hydrocarbons

There is a large and diverse pool of dissolved volatile organic compound (VOC) in the surface ocean, which turns over rapidly, on the scale of hours to days (see Halsey and Giovannoni, 2023 for a full review). Isoprene (C_5H_8) is the most abundant marine terpenoid and is thought to be an efficient contributor to secondary aerosol (Yu et al., 2021 and references therein). Monoterpenes ($\text{C}_{10}\text{H}_{16}$) undergo oxidation to compounds of much lower volatility, highly oxidized molecules (HOM). In terrestrial environments, HOM were shown to contribute to the growth of newly formed particles (Ehn et al., 2014) as well as to nucleate by themselves and initiate NPF (Rose et al., 2018; Yan et al., 2018), but little is known of their role in the marine atmosphere. Marine monoterpenes, dominated by α -pinene, show mixing ratios ranging 5–200 ppt over the Southern Ocean. They are generally well correlated with isoprene and tracers of marine biological activity as shown by laboratory and field studies (Colomb et al., 2008; Shaw et al., 2010; Meskhidze et al., 2015; Kim et al., 2017). Other VOCs such as aromatic hydrocarbons (e.g., benzene, toluene, and xylene), previously thought to be exclusively associated with anthropogenic sources, may be produced by phytoplankton with concentrations similar to those of DMS (Rocco et al., 2021). Estimated oceanic emission fluxes of these aromatic compounds are enough to make them a potential significant source for secondary organic aerosol (SOA) in polar atmospheres (Wohl et al., 2023). However, any involvement in NPF events has not yet been investigated.

2.2.4. Other VOCs from surface photochemical reactions

In addition to biological VOCs directly emitted from oceans, recent works have shown that photochemical processes occurring at the very surface of the ocean lead to further production of VOCs with importance as aerosol precursors (Rossignol et al., 2016; Tinel et al., 2016; Brüggemann et al., 2018). Two main requirements for surface photochemical VOC formation and release are (a) sunlight to initiate photochemical reactions and (b) the development of an organic-rich sea surface microlayer at the air–water interface. Research continues to demonstrate that photochemical and subsequent reactions that involve iron, other ions common to seawater, and organic compounds (such as phytoplankton exudates), interact to form VOCs (Ciuraru et al., 2015; Fu et al., 2015; Huang et al., 2020; Stirchak et al., 2021;

Penezić et al., 2023). Upon reaction with the common oxidants like ozone and the OH radical, these VOCs may then be precursors to SOA formation and, potentially, even to nucleation in the MBL (Bernard et al., 2016; Alpert et al., 2017). This photochemical source of nucleation precursors is a rather new research topic compared to those mentioned in previous sections; it will need confirmation in the real ambient atmosphere by identifying tracers of surface photochemistry products. There is also a need for further laboratory work to figure out when photochemical production of VOCs may or may not be important (Rapf et al., 2018; Shrestha et al., 2018), and modeling studies to evaluate the global impact on total VOC emissions and their nucleation potential.

2.3. Biogeochemical sources of nucleating precursors

2.3.1. DMS production, spatial distribution, and temporal trends

In the surface ocean, DMS is produced mostly through the enzymatic breakdown of dimethylsulfoniopropionate (DMSP), a ubiquitous cellular component that serves multiple physiological functions, such as osmolyte, cryoprotectant, antioxidant, and overflow of excess reduced sulfur (Stefels et al., 2007). DMSP is synthesized mainly by phytoplankton, with intracellular concentrations that span orders of magnitude across taxa (McParland and Levine, 2019; Moran and Durham, 2019). It is also synthesized by bacteria (Curson et al., 2017), yet the quantitative significance of the bacterial production in the DMSP budget has not been evaluated. DMS can also be produced through enzymatic reduction of the other ubiquitous organosulfur compound, dimethylsulfoxide (DMSO; Spiess et al., 2009; Dixon et al., 2020), but the contribution of this process to pelagic gross DMS production remains uncertain due to the scarcity of measurements (Dixon et al., 2020; Herr et al., 2020). Recent advances in genetic approaches to understand the processes of DMSP metabolism and DMS production are making it possible to study how they evolved in marine microorganisms (Bullock et al., 2017), how they distribute in the ocean (Teng et al., 2021), and how they are part of the biogeochemistry of ocean sulfur (Moran and Durham, 2019).

The distribution patterns of DMS result from the interplay between the abundance and taxonomic patterns of DMSP-producing phytoplankton, their physiological status, and a complex network of DMS production and removal processes (Lizotte et al., 2012; Galí and Simó, 2015). On average, most of the produced DMS is consumed by biological (approximately 70%) and photochemical (approximately 20%) processes, allowing only approximately 10% to be vented to the atmosphere (Galí and Simó, 2015). Yet, these proportions may vary largely depending on irradiation, stratification, wind speed and mixing, and hence across regions, seasons, and even the diel cycle (Royer et al., 2016). Therefore, predicting the distribution of the DMS concentration and sea-to-air emission across space and time represents a formidable challenge. DMS concentration fields in the

surface ocean can be derived by producing climatologies from databases (Lana et al., 2011; Hulswar et al., 2022) or can be diagnosed semiempirically (statistically) from a set of environmental variables (Belviso et al., 2004). Pioneering statistical models included just a few diagnostic variables chosen from hypothetical phenomenological relationships to DMS (Belviso et al., 2004) and captured a limited share of the DMS variance (Halloran et al., 2010). Recently, more sophisticated machine-learning methods (like neural networks) with an increasing number of diagnostic environmental variables have notably improved DMS predictability at both regional and global scales (Wang et al., 2020; McNabb and Tortell, 2022; Mansour et al., 2023; McNabb and Tortell, 2023). However, a limitation of both the simple and the machine-learning diagnostic methods is that they rely on variables that cannot be synoptically diagnosed in near-real time, like the mixed layer depth or nitrate concentration. Consequently, they provide climatological maps but can hardly provide snapshots of the current DMS distribution, or interannual patterns to address the response of ocean DMS to global environmental change. Galí et al. (2018) proposed an algorithm to derive surface ocean DMSP and DMS concentrations from satellite-based variables such as Chl-*a* and solar radiation. Even though this algorithm does not work equally well everywhere over the global ocean, it represents an invaluable tool to produce regional synoptic maps and temporal trends of DMS (Galí et al., 2019). Remote sensing, however, still has the limitation of the incapacity to project into the future.

Alternatively, DMS fields can be predicted by prognostic modeling. Hitherto, the representation of DMS in prognostic biogeochemical models has barely paralleled advances in the knowledge of the processes in the DMS cycle. DMSP production and content in phytoplankton has been revealed to be less dependent on nutrients than thought as the DMSP quota of the strongest DMSP producers is less prone to regulation by nitrogen availability (McParland and Levine, 2019). Phytoplankton do not only act as the main source of DMSP, many also take up DMSP for unknown reasons (Vila-Costa et al., 2006; Spielmeyer et al., 2011). Consequently, they can represent a major sink that competes with bacterial degradation and has yet to be incorporated in DMS models. More than a decade ago, the first global prognostic model that included a sulfur module only represented 2 or 3 types of phytoplankton, each with its intracellular DMSP quota and mortality and exudation rates (see review in Bock et al., 2021). There was the appreciation that a better representation of phytoplankton composition and succession, bacterial processes, and solar radiation effects was required to improve DMS simulations (Le Clainche et al., 2010; Vogt et al., 2010). As an example, explicit incorporation of some of the strongest DMSP producers, *Phaeocystis*, in the Community Earth System Model attributed as much as 13% of global DMS emission to this phytoplankton genus (Wang et al., 2015). A process-based study using ship-borne mesocosm experiments found a positive relationship between the DMS flux and the nanophytoplankton cell abundance

(Rocco et al., 2023). On the other hand, the relative contributions of bacterial and algal DMSP-lysis enzymes to DMS production have not yet been fully assessed, nor has DMS production from DMSO, so that these processes are not yet represented in global prognostic models.

2.3.2. Biogenic production of other potential nucleation precursors

Seawater iodide originates from biological activity of microalgae and bacteria (Hepach et al., 2019; Hughes et al., 2021). Several phytoplanktonic groups consume iodate and produce iodide, with no functional-type specificity but with a larger yield for cultures that had reached a later senescent stage (Hepach et al., 2019). Like DMS, some other VOCs are produced as a regulated response to environmental stressors such as light, temperature, acidification, nutrient limitation, grazing, and viral-induced mortality (Halsey and Giovannoni, 2023). VOC production is commonly light dependent indicating a link to photosynthetic metabolism, but the diel cycles of reactive non-DMS VOCs are not described in the literature. Moore et al. (2020) suggest that VOCs are not only stress-induced metabolic end products but also intermediates in carbon metabolic pathways, which can diffuse out of cells. Furthermore, VOCs also serve as signal molecules in large aquatic organisms (Saha and Fink, 2022), but the same role in phytoplankton has been challenging to prove. Although phytoplankton are thought to be the primary source of most VOCs, other processes such as the degradation of dissolved organic matter (DOM) by heterotrophic bacteria also release VOCs. However, monoterpenes and other highly reactive VOC production by direct phytoplankton or bacterial production are poorly characterized.

Another suite of marine volatiles of great interest are ammonia (NH_3) and methylamines. The air–sea flux of NH_3 is sensitive to temperature, pH, and salinity (in decreasing order of importance) (Altieri et al., 2021). An interlink between the NH_3 and the DMS fluxes has been suggested (Johnson and Bell, 2008). This would arise from the complementation of the processes that give rise to DMS (and its precursor DMSP) and those that produce $\text{NH}_3/\text{NH}_4^+$ in the surface ocean. Recent work on NH_3 fluxes in the Southern Ocean suggests that the biogeochemistry that drives the seasonality in surface ocean NH_4^+ concentrations determines when the Southern Ocean

becomes a source of NH_3 to the atmosphere (Altieri et al., 2021). Regarding methylamines, concentrations of trimethylamine in the surface ocean are in the nM range and its sources are not well characterized. It's mainly attributed to the bacterial degradation of phytoplanktonic N-osmolytes such as glycine betaine, trimethylamine oxide, and choline (Carpenter et al., 2012; Mausz and Chen, 2019), which serve similar physiological functions as DMSP. Phytoplankton can be both a direct and indirect source of trimethylamine (Mausz and Chen, 2019), which can subsequently be transformed to dimethylamine and further to monomethylamine by marine bacteria (Sun et al., 2019).

With regard to VOCs produced through photochemical reactions at the sea surface microlayer, a dependence of fluxes on the concentration and composition of organic matter is to be expected. Therefore, biological processes that produce surfactants with the potential to accumulate in the surface microlayer will play a regulating role. For instance, lipids are released during the lysis of, or the predation on, phytoplankton cells, and subsequent enzymatic digestion leads to the formation of fatty acids (Budge et al., 2014; Cochran et al., 2016). In summary, **Figure 1** illustrates the recent investigations and the state of knowledge on the source, flux, and detection of particle nucleating species as discussed above.

3. Biogenic impact on SSA properties and emission flux

Despite the abundance of SSA flux parameterizations in the literature (see Russell et al., 2023 for a complete review), SSA flux estimations still suffer from large uncertainties. In recent global modeling exercises, the default sea spray emissions needed to be increased by around a factor of 3 to be consistent with measurements (Regayre et al., 2020), pointing to a lack of understanding of the processes involved in sea spray emissions. Using observational constraints, Bian et al. (2019) concluded that the negative bias in the aerosol optical depth simulated by a global model could partially be explained by missing sub-micron sea spray sources. A largely studied source of SSA is bubble bursting. When bubbles burst on the ocean surface, 2 types of droplets are formed, jet droplets leading to super-micron particles upon evaporation, and film droplets leading to sub-micron particles. **Figure 2** summarizes the recent investigations and the state of




Figure 2. State of knowledge on the dependence of sea spray sources on environmental variables.

knowledge on the dependence of sea spray properties and flux on environmental and biological variables which will be discussed as follows.

3.1. Factors influencing the SSA emission flux and parameterizations

Sea spray production is not only dependent on wind speed—surface temperature, salinity, and seawater composition have also been identified as important properties controlling the SSA emission flux (Fuentes et al., 2011; Hultin et al., 2011; Salter et al., 2015). The complexity of processes controlling SSA production via bubble bursting has resulted in many different formulations of the SSA source function that span over an order of magnitude for a given wind speed, especially in the smallest (<20 nm) and largest (>500 nm) size ranges (Ovadnevaite et al., 2014). This large spread may be due to different methods used to derive the size segregated sea spray flux. The SSA flux may be directly measured in ambient air using eddy correlation or gradient methods (Method A). A second method is the combination of transport models and observations of aerosol concentrations in a source–receptor approach (Method B). The last approach is laboratory-generated sea spray from seawater samples using bubble bursting devices that mimic the wave-breaking process (Method C). However, a large spread is also observed among sea spray flux formulations obtained with the same method. For Method A, wind speed, wave-state, or air entrainment in the SSA flux parameterizations reported in the literature are not always harmonized for a given seawater temperature, biological content, atmospheric washout state. For instance, since the SSA flux depends on both wind speed and temperature, the dependence of each variable should be investigated with the other variable kept constant. This is usually not done and for example regions of higher wind speed may correspond to regions of lower temperature implying a bias in the parameterizations relating SSA fluxes with wind speed.

Hartery et al. (2020) have provided a parameterization of a large SSA source function specific for the Southern Ocean that takes into account wet scavenging. Their results are substantially different from the traditionally used parameterizations such as in Jaeglé et al. (2011). The parameterization proposed by Hartery et al. improved the representation of the SSA flux in global climate models (Revell et al., 2019) as well as the correspondence of modeled CCN with Southern Ocean cloud observations (Kang et al., 2022). The different methods for providing a temperature dependence of the SSA flux seem to lead to different conclusions depending on the method used. Using Method B, Ovadnevaite et al. (2014), Grythe et al. (2014), and Liu et al. (2021) report a monotonic, positive relationship between the sea spray mass flux and the sea surface temperature (SST) over the whole temperature range investigated. Conversely, the temperature dependences of laboratory-generated sub-micron sea spray number flux (Method C) show opposite trends. For temperatures colder than 10°C, all studies report a strong increase of the sea spray number flux with decreasing temperature (Mårtensson et al., 2003; Salter et al.,

2014). This apparent contradiction in temperature dependences between Method B and Method C may be due to the size of SSA investigated. Salter et al. (2014) indeed reported different behaviors for sub- and super-micron SSA. Opposite behaviors between sub- and super-micron SSA are likely due to different production mechanisms as pointed out by Crocker et al. (2022).

Different temperature dependences of the SSA flux may also result from different influences from the biogeochemical properties of different seawater types (Forestieri et al., 2018; Christiansen et al., 2019). This area of research is largely understudied due to the difficulty in decoupling the chemical and biological effects from the temperature effect in studies of ambient-air SSA. Both types of variables (temperature and those derived from ocean biogeochemistry) have been shown to largely impact the SSA emission flux individually under controlled experimental conditions. Using seawater collected from 3 contrasting locations (Mediterranean Sea, Arctic coastal waters, and New Zealand), Sellegrì et al. (2021) reported that the artificially generated sea spray flux of particles larger than 100 nm, assimilated to CCN at 0.2% supersaturation, is linearly correlated to the seawater nanophytoplankton cell abundance for a given SST. They suggest that this effect results from biogenic surface-active components that modulate bubble lifetime and therefore the number SSA flux. A similar feature was found for SSA generated from sea ice samples, with a larger SSA number flux for the highest nanophytoplankton cell abundances (Dall'Osto et al., 2022). This feature was associated with the presence of humic-like substances. In Sellegrì et al. (2023), the SSA flux to SST relationship is quantified as a function of the seawater biogeochemical properties, with 30% decrease in the cyanobacterium *Synechococcus* leading to an average 4-fold increase in the SSA flux at 2°C SST compared to 8°C–12°C SST. This impact may be related to *Synechococcus*-derived DOM of specific properties, released via secretion, natural cell death, viral lysis, and predation (Fiore et al., 2015; Zhang et al., 2022).

3.2. Nascent SSA size distribution

SSA size distribution is important for atmospheric processing as it determines the surface area for condensation and reaction of nitrogen, sulfur, and organic species. It also determines the distances that particles can travel and transport the chemical species they are composed of (Neumann et al., 2016). Nascent SSA artificially generated during bubble bursting experiments consistently show a number size distribution with a sub-micron Aitken mode (i.e., around 40 nm) and small accumulation mode (i.e., around 100 nm) dominating the total size distribution, and frequently even a large nucleation mode contribution to the SSA number concentration (Schwier et al., 2015; Schwier et al., 2017 and references therein; Sellegrì et al., 2021). Due to the difficulty in segregating SSA from secondary aerosols in the ambient marine atmosphere, some models use the assumption that SSA size distribution can be approximated by a single broad mode encompassing particles from 100 nm to super-micron sizes, scaled from larger than 500 nm particles assimilated as

pure SSA (Modini et al., 2015). Using this methodology, the contribution of SSA number concentration to total aerosol concentrations in marine air masses may therefore be highly underestimated. This was recently shown by Xu et al. (2022), who used an hygroscopicity-derived ambient aerosol mixing state to confirm the large contribution of SSA to the Aitken and accumulation modes of natural marine ambient air. The effect of seawater biology on the SSA size distribution shape is not very clear and the organic content of nascent SSA has shown little impact on its size distribution shape (Sellegrì et al., 2021). Therefore, the stability of the ratio of super-micron to sub-micron aerosol generated in the laboratory could also be a valid method to infer a number concentration of sub-micron SSA from ambient air marine aerosol.

The detection of a nucleation mode in the ambient aerosol size distribution over remote areas does not necessarily point to secondary aerosol formation, as it was also detected during nascent sea spray experiments (Schwier et al., 2015; Schwier et al., 2017), especially in low SST conditions (Sellegrì et al., 2023). Conditions when these ultrafine SSA are produced are unclear. Surface-active compounds of the sea surface microlayer play a major role in the number of SSA emitted to the atmosphere (Modini et al., 2013), yet we don't know what biogeochemical or photochemical processes are at the origin of these surface-active components and their dependence on SST. Surface tension should be a measure to deploy in parallel to other seawater biogeochemical variables in the future and lab studies are needed to tackle this.

3.3. Nascent SSA chemical composition

Nascent SSA is a complex mixture of sea salt and organic matter, with organic matter comprising a substantial fraction of the total particle mass under certain conditions, especially for particles with diameters less than 1 μm (de Leeuw et al., 2014). The chemical composition of jet droplets is believed to reflect the composition of the bulk surface ocean (Burrows et al., 2014; Wang et al., 2017). The chemical composition of the film droplets is believed to reflect the composition of the film cap from which the droplets were born (Burrows et al., 2014). Given this, the chemical composition of these droplets is often assumed to be similar to that of the air–water interfaces present on bubble surfaces or at the ocean surface, that is, the sea surface microlayer. As such, these droplets are believed to be enriched in surface-active organic matter when the sea surface microlayer has a different composition than the underlying bulk seawater. It is therefore important to understand the factors driving sea surface microlayer enrichment in organic matter, which has received attention in the past decade but still lacks understanding both from the physical and chemical points of view (Tinnel et al., 2023).

The role that biological activity plays in determining the quantity and composition of organic matter present in nascent SSA has been an active area of research over the last 20 years and remains a large area of research (O'Dowd et al., 2015; Quinn et al., 2015; Ceburnis et al., 2016;

Gerard et al., 2016; Jayarathne et al., 2016; van Pinxteren et al., 2017; Beaupré et al., 2019; Frossard et al., 2019; Crocker et al., 2020; Radoman et al., 2022; Santander et al., 2022). The conclusions from the recent review by Bertram et al. (2018) can generally be characterized as falling into 2 groups. First, a series of studies have observed that the organic mass/ Na^+ ratios of nascent SSA (Facchini et al., 2008; O'Dowd et al., 2015) are highly dependent on the level of biological activity (e.g., the abundance of phytoplankton). Lower organic mass/ Na^+ ratios correspond to periods of low biological activity compared to periods of high biological activity, resulting in primary marine organic content being parameterized as a function of Chl-*a* in many modeling exercises (de Leeuw et al., 2014). In contrast, a second group of studies has observed little difference in the organic mass/ Na^+ ratio of nascent SSA when comparing oligotrophic and productive oceanic regions (Quinn et al., 2015) or during mesocosm phytoplankton experiments (Jayarathne et al., 2016; Schwier et al., 2017). Instead, some of these studies argue that a ubiquitous reservoir of organic mass in surface seawater is responsible for the organic mass enrichment of nascent SSA, overpowering any influence of local biological activity. These studies conclude that the organic mass fraction of nascent SSA can be approximated as constant across different ocean basins and seasons (Quinn et al., 2015; Saliba et al., 2019; Bates et al., 2020), a conclusion that is questioned in observation–modeling comparison exercises (Burrows et al., 2014).

Numerous studies have investigated the impact of ocean biology on the composition of the organic fraction of nascent SSA. For example, Gerard et al. (2016) observed a correlation between Chl-*a* and the concentration of anionic and cationic surfactants in ambient aerosol sampled over the Baltic Sea, and Frossard et al. (2019) found that surfactant concentrations in laboratory-generated SSA were correlated with the level of biological activity in seawater samples. However, transport of organic species to the atmosphere through SSA was not directly related to measures of primary productivity and was more likely related to more complex processes such as bacterial transformation and degradation of the primary products of plankton (Santander et al., 2022) and further linked to polysaccharides and fatty acids likely of bacterial origin (Glicker et al., 2022). Interestingly, Radoman et al. (2022) observed measurable changes in the composition of the organic mass present in laboratory-generated SSA during a phytoplankton bloom using high-resolution mass spectrometry, despite the organic mass fraction of the aerosol remaining unchanged during the sampling period, suggesting that the 2 parameters may be decoupled or have different dependency of the bloom stage (rise or demise). Overall, the transfer of organic carbon from seawater to aerosol appears to result from complex biological processes and finding direct universal relationships between a biological tracer (such as Chl-*a*) and the fraction of organic matter in nascent sea spray seems illusory.

A reconciliation between observations and modeling of the organic content of primary aerosol particles is proposed in the conceptual model OCEANFILMS (Organic

Compounds from Ecosystems to Aerosols: Natural Films and Interfaces via Langmuir Molecular Surfactants), which links sea spray chemistry with ocean biogeochemistry. This is done by mapping groups of organic compounds to the output of an ocean biogeochemistry model, and partitioning these organic groups into SSA on the basis of physico-chemical properties (absorptivity at the air–water interface, molecular weight, etc.) that are assigned to biological macromolecule types (proteins, polysaccharides, lipids, processed compounds, and humics) (Burrows et al., 2014; Burrows et al., 2022). The nature of the organic compounds determines the efficiency with which they are transferred to the atmosphere, resulting in a model where correlations between organics and Chl-*a* exist but are highly region- and season-dependent. These regional specific relationships further match recent seagoing campaigns (Bates et al., 2020) and seasonality observed at coastal stations such as Mace Head, Ireland.

4. Developments in the identification of marine aerosol sources from observations and modeling

Most marine aerosols are composed of salts (mainly sodium chloride) and sulfates, and a mixture of organic compounds. While sodium chloride is clearly primary, that is, a tracer for SSA, sulfates and organics may be primary or secondary. Source apportionment of marine aerosols depends strongly on our ability to chemically characterize particles across all sizes and independently of their primary or secondary nature. Several modeling studies (Gantt et al., 2011; Burrows et al., 2022) have used the water-insoluble organic carbon content of marine aerosol as a proxy for the primary organic source, based on observational studies (O'Dowd et al., 2004; van Pinxteren et al., 2017). However, the ratio of insoluble to soluble organic carbon in nascent SSA is variable (cf. Section 3; Freney et al., 2021) and therefore a potential source of uncertainty. The source apportionment between primary and secondary marine aerosol is further complicated by the fact that some of the molecular products of atmospheric photochemistry that contribute to marine SOA formation can also be produced within the seawater via either photochemical processes at the ocean surface or biological processing. For instance, Beaupré et al. (2019) observed that 19%–40% of the organic carbon present in nascent laboratory-generated SSA (less than 2 min of age) was chemically indistinguishable from highly aged SOA, and Freney et al. (2021) found that signatures of SOA, such as MSA, were already present in nascent sea spray. Enami et al. (2019) suggested that the oxidation of marine biogenic VOCs is facilitated by the presence of iodide in the surface microlayer, and their oxidation products are left in seawater or found in the marine primary organic aerosol.

A detailed concomitant analysis of the same chemical organic tracers in the atmosphere and ocean would be very valuable. For instance, a study by van Pinxteren et al. (2020) shows highly surface-active lipids being enriched in the sea surface microlayer. Looking for (and quantifying) these compounds in the aerosol phase would be helpful to assess their transfer function to the atmosphere. Using

X-ray spectro-microscopy, the organic matter associated with relatively freshly formed SSA particles was revealed to be dominated by acid and alcohol functionalities (Ault et al., 2013; Prather et al., 2013). Again, it would be valuable to apply a similar approach to seawater and surface microlayer samples. Recent advances in the measurement of carbon isotopes have also provided insight into marine organic matter sources. In the ocean, photosynthesizing plankton produce organic matter with lighter isotopes, indicated by a lower $^{13}\text{C}/^{12}\text{C}$ ratio. This particulate organic matter undergoes isotopic fractionation over the course of weeks to years as it is broken down and decomposed through trophic-level interactions, ultimately transitioning into isotopically heavier DOM, characterized by a higher $^{13}\text{C}/^{12}\text{C}$ ratio. Ejected SSA contain the C isotopic ratio of the associated marine DOM, which subsequently undergoes kinetic fractionation through SOA formation via photochemical and cloud-processing mechanisms. Consequently, the aerosol organic matter becomes isotopically heavier on a timescale of approximately 1 week. In this way, C isotopes are instrumental to trace the pathway of organic matter from the surface ocean to SSA and SOAs. For example, Ceburnis et al. (2016) found that most of the organic enrichment in marine aerosol over the southern Indian Ocean was attributable to fresh marine particulate organic matter. Consistent with these findings, Crocker et al. (2020) observed that the C isotopic composition of laboratory-generated SSA reflected changes in seawater composition during mesocosm phytoplankton blooms.

There is a severe lack of information on the sources and chemistry of marine Aitken mode particles (20–80 nm) due to measurement limitations. The Aitken mode is a persistent feature over the global oceans (Hoppel and Frick, 1990) and is critical because these particles are on their way to growing into the CCN size range, where they will influence cloud properties. Two main sources of Aitken mode particles are (i) primary sea spray emissions (see Section 3) and (ii) growth of nucleation mode particles by condensation of secondary compounds (e.g., Burkart et al., 2017; Zheng et al., 2020). The relative importance of the two sources remains unclear and is expected to vary by region. Significant advances have been made in the knowledge of sea spray sources of Aitken mode particles by analyzing laboratory-generated nascent SSA using single particle microscopic analyses (Prather et al., 2013) or hygroscopicity-derived chemical composition (Schwier et al., 2015; Schwier et al., 2017). These studies have clearly established that most small SSA contain organics at the point of emission. An early study by Oppo et al. (1999) proposed a theoretical model that explains this phenomenon as caused by adsorption of organic components on the surface of small saline droplets. In this model, small SSA have larger fractions of surface-adsorbed organic compounds due to the large surface-to-volume ratio of particles, while large SSA are composed primarily of salt.

Although the presence of high organic mass fractions in nascent SSA is well-established, it is more difficult to quantify the relative contributions of primary and secondary organic matter in the ambient Aitken mode particles.

The reason is that the small size and mass of Aitken particles makes it difficult to analyze their chemical composition. Traditional filter-based approaches require long sampling times which may introduce artifacts for the semi-volatile species and do not allow fine size discrimination. Online mass-spectrometric methods exist for the analysis of sub-50 nm particles (Lawler et al., 2021), but these systems are rare since they are custom-built, and difficult to operate. Therefore, the nature of the condensable gases responsible for the growth of nucleation mode particles into Aitken mode particles remains unclear. Instrumental development in several directions is needed: (a) a high-volume collection technique specific for Aitken mode particles, which can then be characterized by traditional filter-based chemical analyses; (b) offline organic and inorganic chemical analysis techniques with extremely low detection limits that can be applied with short sample collection times; and (c) making existing online mass spectrometry or Fourier Transform InfraRed spectroscopy techniques to target the sub-50 nm size range.

The processes that cause growth of particles from the Aitken to the accumulation mode are also still poorly understood. Growth can happen either through direct condensation of semi- or low-volatile gases issued from oxidation processes, through surface heterogeneous chemistry, or through aqueous phase processing. Indirect evidence of the contribution of oxygenic volatiles to dry formation of SOA can be derived from their detection in the gas phase. Organic acids, known precursors of SOA, have been detected in the Arctic MBL as well as in aerosol formed from photochemistry or DOM at the sea surface microlayer (Mungall et al., 2017). Once in the atmosphere, there is also the difficulty in segregating dry and wet formation of SOA. Cloud processing is often more efficient than the dry oxidation pathways, particularly in the case of MSA formation (Chen et al., 2018; Baccarini et al., 2021). To constrain the reaction pathways, isotopic analyses are again a powerful tool (e.g., Harris et al., 2014; Ishino et al., 2021; AuYang et al., 2022). Atmospheric aging hence poses a challenge to source attribution and the community needs to think of new tracers to be targeted. New developments, such as the Extractive Electrospray Ionization time-of-flight (EESI-TOF; Wang et al., 2021) for near-molecular aerosol speciation, the EESI-Ultrahigh-Resolution Mass Spectrometry (EESI-Orbitrap; Lee et al., 2020) allowing insights into compound structure, and the High-Resolution Time-of-Flight Chemical Ionization Mass Spectrometer with a Filter Inlet for Gases and AEROSols (FIGAERO-HRToF-CIMS; Siegel et al., 2021), providing molecular composition information, open up new routes to understand chemical reaction mechanisms, kinetics, and sources. So far, these emerging mass spectrometric techniques have only rarely been used in marine aerosol research, thereby representing a great underexplored potential.

An advantage of applying high-resolution spectrometric techniques on Aitken and accumulation mode particle (roughly 100–1,000 nm) characterization is that it allows the consolidation of results by statistical methods such as positive matrix factorization, which are established methods for source apportionment. Based

on the spectrometric spectra, fingerprints of recurring composition can be extracted and related to potential sources using chemical markers. Recent works in the Arctic (Moschos et al., 2022a, 2022b) and Atlantic oceans (Lawler et al., 2020; Huang et al., 2022), and the Sea of Japan (Loh et al., 2023) demonstrate the ability to distinguish among natural-terrestrial, natural-marine, and anthropogenic aerosol sources.

5. Marine CCN and INPs

5.1. Marine-related CCN

The properties of a CCN at a given water vapor supersaturation are reasonably well described by the Koehler theory that includes size and chemical components of the aerosol, as shown by several successful closures between modeled and measured CCN (Asmi et al., 2012 and references therein). A good estimation of the number of CCN at a given supersaturation could be obtained from the number of particles above a certain size (activation size), knowing the large chemical classes (organics, ammonium sulfate, sodium chloride) present in particles at this size. Fresh SSA CCN concentration could adequately be approximated from the number concentration of SSA larger than about 100 nm at 0.2% supersaturation (typical of marine low level clouds) (Sellegri et al., 2021). However, ambient SSA and marine secondary particles with high organic fractions tend to deviate from the theory due to the disconnection between the organic component hygroscopicity at subsaturated and supersaturated conditions and the organic surface tension effect (Ovadnevaite et al., 2011; Ovadnevaite et al., 2017). The global importance of this effect and deviation from the classical theory is still under debate, but Lowe et al. (2019) demonstrate that accounting for the surface phase becomes essential in clean environments such as the remote MBL under certain circumstances.

Many studies on marine CCN concentrations do not investigate what makes a marine particle a CCN, but rather what the contributions from different sources are (i.e., primary or secondary sources). At wind speed greater than 16 m s^{-1} , Fossum et al. (2018) estimated that SSA may contribute up to 100% of marine CCN-relevant aerosol particles present in the MBL in the Southern Ocean. On the other hand, non-sea-salt sulfate was argued to dominate CCN concentrations, either based on correlation with Aitken mode particles (Tatzelt et al., 2022) or from measurements of nonrefractory chemical composition in marine aerosol (by up to 60% during summer; Zheng et al., 2018; Zawadowicz et al., 2021). Sanchez et al. (2018) estimated “new sulfate” contributes about one-third of North Atlantic CCN during spring and autumn while sea spray dominates during winter. Biogenic organics were also found responsible for the growth of preexisting particles to CCN sizes, alone or by co-condensing with sulfates (Zheng et al., 2020). Secondary particles from DMS oxidation were also predicted to contribute to the MBL CCN burden in modeling exercises mostly during summer (Revell et al., 2019). McCoy et al. (2015) used top-down constraints from satellite observations of cloud droplet number

concentration to infer the relative contributions of DMS-derived sulfate aerosol, sea spray, and marine organic matter to the regional-scale seasonal variations in CCN. They found that DMS-related sources drive variations in the cloud droplet number concentration across the entire Southern Ocean, *via* sulfate aerosol at lower Southern Ocean latitudes (35°S–45°S) and impacts on SSA at higher latitudes (45°S–55°S). However, as discussed above, SSA can significantly contribute to the sub-micron particle mode, including Aitken mode particles (Lawler et al., 2021; Xu et al., 2022) and, thus, can potentially add more significantly to the CCN population than previously thought (Lauvset et al., 2017; Quinn et al., 2017).

5.2. Ice formation on marine-related particles

Atmospheric INP concentrations measured at marine observatories are often used to infer the ice nucleating abilities of marine aerosol particles (McCluskey et al., 2018a; Knopf et al., 2022). However, it is important to note that not all particles found over oceans or in the marine atmosphere are derived from marine sources. Ice nucleation studies on the natural atmospheric aerosol must deal with the complexity of measuring low INP concentration and determining INP composition from diverse sources. In general, great care is needed when implementing parameterizations of INP concentration for atmospheric ice production as INP overestimation can bias estimates of radiative balance, shown especially for the Southern Ocean (Vergara-Temprado, 2018). Using a combination of global modeling and observations, Burrows et al. (2013) suggested that the biogenic marine INP concentration dominated the total INP concentration in the Southern Ocean, but also pointed to a lack of observations in this area to confirm their conclusion. Zhao et al. (2021) showed that marine organic INP concentrations can be up to 1,000 times higher than dust INPs at 40°S. However, sources of INPs in models are still not well constrained as some parameterizations tend to underestimate the observed INP concentrations except for at temperatures representative of cirrus clouds (parameterization from McCluskey et al., 2018b) or tend to overestimate them except for temperatures representative of supercooled clouds (source functions from Wilson et al., 2015; Trueblood et al., 2021). It is therefore important to better constrain SSA INP sources, in particular their biogenic components.

5.2.1. Evidence of a biogenic source of marine INPs

A strong biogenic contribution to marine INP concentrations is suggested from the evidenced relationships between seawater and sea surface microlayer water particulate organic carbon and atmospheric INPs concentrations at warm temperatures representative of supercooled clouds (>20°C) (Wilson et al., 2015; Trueblood et al., 2021). In the North Atlantic, ice nucleation due to SSA at the warmest freezing temperatures always coincided with observed total phytoplankton cell concentrations in water on the order of 10^8 L^{-1} and when dominated by picoeukaryotes (Wilbourne et al., 2020). It was shown that INPs were relatively stable over the Southern Ocean and

increased due to coastal (either marine or additional terrestrial) sources with concentrations generally coinciding with fluorescent, hence potentially biogenic, particles (Landwehr et al., 2021; Moallemi et al., 2021). In the laboratory, early research showed that phytoplankton cultures contain INPs capable of forming ice as warm as -4°C (Schnell, 1975) through the liquid-to-ice phase transition known as immersion freezing. The first marine derived INPs were unambiguously identified no more than 4 decades later as the fragmented and intact phytoplankton cells of *Thalassiosira pseudonana*, *Nanochloris atomus*, and *Emiliana huxleyi* (Alpert et al., 2011a, 2011b). Uniquely, ice formation from both immersion freezing and deposition ice nucleation (the vapor-to-ice phase transition) was quantified. Ice nucleation experiments using cultures of diatoms, *Melosira arctica*, *Skeletonema marinoi*, and *T. pseudonana* as well as other phytoplankton and aerosolized sea surface microlayer water all show ice nucleation ability (Wilson et al., 2015; Ickes et al., 2020). Even after filtering cultures of marine phytoplankton and bacteria (0.1 μm filter), biogenic materials in the water retain freezing ability (Wilson et al., 2015; Ladino et al., 2016). This indicates that not only cell or cell fragments of phytoplankton are INPs, but also their exudates generated through their metabolism or lysis. SSA particles produced from cultures of *Prochlorococcus* from 3 different strains were INPs via deposition ice nucleation, with a link to polysaccharide, lipopolysaccharide, and protein content (Wolf et al., 2019).

Understanding of SSA particle production, composition, and ice nucleation was greatly enhanced through a series of laboratory breaking wave channel studies with actively growing mixed microorganism communities (Prather et al., 2013; DeMott et al., 2016). Through careful examination of seawater conditions that influence SSA production, the authors determined that SSA particles were INPs. However, they found higher INP concentrations prior to microbial growth and when most SSA particles were internal mixtures of sea salt and organic matter (Prather et al., 2013). This was contrary to expectations that increasing microbial growth would increase biogenic matter in aerosol particles and thereby, ice nucleation efficiency or INP concentrations. It was later confirmed that ice nucleation tended not to be enhanced as a function of microbial growth or biological markers, such as chlorophyll concentration, heterotrophic bacterial counts, and enzyme activity (DeMott et al., 2016; McCluskey et al., 2017). The complex balance between the numbers and efficiency of INPs produced, released, and consumed in water due to phytoplankton and bacteria metabolism prior to aerosolization has yet to be understood.

Single particle analysis of sub- and super-micron-sized SSA has aided their chemical and morphological identification, giving a unique look into the origin of why SSA nucleated ice at all. Recently, single SSA that formed ice was identified using X-ray spectro-microscopy and their chemical characteristics was compared against those SSA that did not form ice attempting to identify the chemical agents, peculiarities, or unique features responsible for ice nucleation (Alpert et al., 2022). It was directly found that

all SSA particles, INPs and non-INPs alike, were practically identical, and that the organic matter associated with them was entirely derived from phytoplankton and bacterial exudates (Alpert et al., 2022). Therefore, current analytical techniques failed to provide the relevant information on the feature that makes SSA a good INP at the temperatures considered. Further research with comprehensive techniques is needed to reveal the unique features of INPs, especially regarding warmer temperatures relevant for biological aerosol. Continuing to quantify marine INP concentrations is just as important to using that data for shaping theoretically based parameterizations. This may involve identifying water properties suspected to be important for ice nucleation, such as hydrogen bonding interaction between water and ice nucleating biogenic/organic surfaces with ions commonly present in seawater.

5.2.2. Parameterizing the sources of marine INPs

As a result of the recent wealth of research on ambient marine INPs, parameterizations of their emissions and ice nucleation efficacy have been proposed, some of which are suitable for application in atmospheric models. As previously mentioned, organic carbon has been shown to be a predictor for marine INPs in some studies and has been used to parameterize marine INP concentrations in the sea surface microlayer (Wilson et al., 2015) and in generated SSA (Trueblood et al., 2021). Other parameterizations include those based on the sea salt abundance in air as a reference to predict marine INP concentrations (Irish et al., 2019). INP concentrations normalized to the total sea spray surface area, n_s , measured either in the clean marine sector of ambient air or as laboratory generated SSA were parameterized as a function of air temperature (i.e., independently of biology) for use in modeling exercises (McCluskey et al., 2018b; Knopf et al., 2022). Nearly 2 years of ship-based measurements of INP concentrations were made from the Arctic to the Antarctic and showed that n_s has a similar slope as a function of temperature compared to over land but are typically less by about 2 orders of magnitude (Welti et al., 2020).

Ice nucleation rate coefficients, J_{het} , have also been parameterized from various field and laboratory studies, and they can be derived from nucleation theory and calculated

from observations as INP numbers normalized to surface area and time (Cornwell et al., 2021; Alpert et al., 2022; Knopf et al., 2022). A reliable link between thermodynamic control and heterogeneous ice nucleation kinetics is quantified only with knowledge of temperature and relative humidity (Knopf and Alpert, 2013). Alpert et al. (2022) developed a model that allows accurate prediction (within a factor of 10) of marine INPs using J_{het} from SSA size distributions, temperature, and relative humidity (i.e., a thermodynamic control) despite their various inorganic and biogenic components or microbial sources. Other studies previously mentioned (Ickes et al., 2020; Wagner et al., 2021) have been even more conservative, claiming the low level of understanding is insufficient for developing a generalized parameterization for ice nucleation due to SSA particles. This is partly due to the variations in ice nucleation abilities of phytoplankton cells, particle sources (i.e., sea surface microlayer or bulk water), aerosolization methods, INP measurement techniques, and the relative humidity particles are exposed to in experiments prior to ice formation which may promote the formation of solid salts. Much more research is needed to identify, quantify, and understand the fundamental processes for ice nucleation by marine aerosol particles due to various ice formation pathways and particle complexity.

We add that the degree of chemical or photochemical aging of SSA and any influence on ice nucleation ability apart from fresh SSA is far understudied by comparison. Some evidence suggests that ambient atmosphere aged SSA are as good INPs as nascent SSA particles (China et al., 2017; Lata et al., 2021; Knopf et al., 2022), but to the best of our knowledge, this has not been the subject of a targeted study. Furthermore, the degree of SSA aging due to condensation of secondary organic or inorganic material from the gas phase on the ice nucleation ability of ambient particles (whether of marine or terrestrial origin) is absent from scientific literature. **Figure 3** summarizes the recent investigations and the current understanding on properties of marine aerosols serving as CCN and INPs as discussed above.

6. Marine aerosol–cloud interactions, cloud properties, and radiative impact

Aerosol, cloud, and precipitation interactions develop through complex relationships between microphysical




Figure 3. State of knowledge on the cloud condensation nuclei (CCN) and ice nucleating particle (INP) properties of marine aerosol components.

(e.g., aerosol growth, droplet or ice crystal nucleation, time-evolution of droplet size distributions, and precipitation processes), dynamical and thermodynamical (e.g., system motions, mesoscale organization, phase changes and internal energy exchanges on the local scale), and large-scale, environmental processes (e.g., radiation, wind-driven surface fluxes and advection, and other meteorological controlling factors) (e.g., Wood, 2012; Boucher et al., 2013; Rosenfeld et al., 2014; Klein et al., 2017; Bellouin et al., 2020; Wall et al., 2022). These processes are further modulated by a network of positive and negative feedbacks spanning a range of scales (e.g., Stevens and Feingold, 2009; Wood, 2012; Boucher et al., 2013; Altaratz et al., 2014; Rosenfeld et al., 2014; Fan et al., 2016; Bellouin et al., 2020; Scott et al., 2020; Christensen et al., 2022).

Aerosols interact with cloud systems in 2 general ways. For a fixed liquid amount, clouds with increased CCN have more, smaller droplets and thus a higher albedo, proportional to the increased scattering surface area due to this aerosol perturbation (Twomey, 1977). Another response to this aerosol-driven microphysical change can also occur at the macrophysics scale, further modifying the cloud albedo (Bellouin et al., 2020). These adjustments encompass a broad range of mechanisms for liquid, mixed phase, and ice clouds, and their influence on the radiative properties of the cloud system varies by their cloud dynamics and environmental conditions (e.g., Bellouin et al., 2020; Wall et al., 2022; Zhang et al., 2022; Zhang and Feingold, 2023). Precipitation suppression (i.e., resulting in longer cloud lifetimes; Albrecht, 1989) and entrainment enhancement (i.e., reducing cloud liquid and thus brightness, e.g., Xue and Feingold, 2006; Bretherton et al., 2007) are 2 key adjustments to aerosol increases in liquid marine clouds.

Bubble bursting generates small concentrations of relatively large SSA (giant CCN with $r \geq 0.5 \mu\text{m}$) that are activated at the lowest supersaturations, modulate the maximum cloud supersaturation, and speed up drizzle formation (Jensen and Nugent, 2017; Prank et al., 2022). However, fine SSA and secondary aerosols (e.g., associated with marine biology) increase the number of potential CCN, delaying the onset of drizzle (e.g., Albrecht, 1989; Prank et al., 2022). Droplet growth consumes supersaturation by diffusion. The collective droplet surface area will thus determine the diffusion efficiency and latent heat release. Latent heat serves as a (cloud) internal fuel driving their dynamics and influencing their mesoscale organization (e.g., Bretherton and Blossey, 2017; Narenpitak et al., 2021; Janssens et al., 2023). Smaller droplets can be lifted higher up because of their higher mobility (Koren et al., 2015). Simultaneously, at the cloud's edges a larger number of small droplets imply more efficient evaporation and enhanced mixing. The competition between the cloud enhancing processes in the core and cloud suppressing processes in the margins implies that the final impact of larger aerosol number concentrations (for instance due to biologically richer conditions) on cloud properties and lifetime depends on the initial cloud size. Larger clouds with lower surface to volume ratio may benefit from the cloud enhancing processes while smaller clouds that

exhibit a relatively larger entrainment zone will experience suppression (Dagan et al., 2017).

In parallel, precipitation processes also depend heavily on the droplet size distribution, a function of total condensed mass, and aerosol number concentrations itself (Pruppacher and Klett, 1997). Clouds formed from higher aerosol concentrations tend to have smaller droplets with a narrower size distribution. This implies delay of the onset of efficient precipitation processes in biologically active cases, impacting cloud lifetime (Albrecht, 1989). This marine biology driven aerosol–cloud–precipitation interaction can occur in a complex, multistep process. For example, precipitating clouds reduce CCN and remove sub-cloud surface area available for condensation upon updraft, increasing peak supersaturation. In a biologically active case with large Aitken number concentrations, this increase in peak supersaturation increases activation and growth of Aitken mode particles into CCN, helping the cloud system to resist precipitation depletion and cloud break up (McCoy et al., 2021; Wyant et al., 2022). In contrast, in an aerosol-poor (low biology) case rain droplets can efficiently clean clouds and generate rain evaporation below cloud base. This can further influence cloud field organization, particularly through cold pool formation (e.g., Zuidema et al., 2018), which may further modulate the sensitivity of cloud systems to aerosol perturbations (e.g., Wall et al., 2022; Zhou and Feingold, 2023). This implies that aerosol has an important role in cloud field organization which can strongly affect the cloud amount, size, and lifetime and therefore their overall radiative effect (e.g., Eastman et al., 2022; Wall et al., 2022; Zhang et al., 2022; McCoy et al., 2023; Zhang and Feingold, 2023). The strong nonlinearities of these interactions are difficult to predict and warrant further examination.

Marine organic particles are an important consideration in modeling exercises on cloud impact (i.e., either added to the CCN and INP number concentrations or modulating the size and hygroscopicity of existing marine aerosols). Ignoring the fine SSA mode could largely diminish the indirect climate effect of SSA (Quinn et al., 2017). However, ambient measurements (Fossum et al., 2020) and modeling studies (Partanen et al., 2014) showed a positive (0.3 W m^{-2}) rather than negative indirect radiative forcing effect by SSA if sufficiently detailed aerosol microphysics and physics-based parameterizations of cloud activation are considered. This positive effect is ascribed to the tendency of SSA to suppress both the in-cloud supersaturation and the formation of CCN from non-sea-salt sulfate. On the other hand, addition of primary marine organics results in a small negative indirect effect (-0.07 W m^{-2}), showing the importance of knowing the sea spray composition (Partanen et al., 2014). Prank et al. (2022) noted that the sensitivity of clouds to variations in organic fraction of sea spray and hygroscopicity was relatively limited compared to differences between sea spray emission flux parameterizations. Another sensitivity study on marine organic particles, however, shows that a strong effect is observed in terms of significant changes in CCN and cloud droplet number only in the Southern Hemisphere over 40°S – 60°S , while the effect in terms of




Figure 4. State of knowledge on the biological influence impacting cloud properties.

significant changes in cloud ice mass and number concentrations is notable over 50° – 70° in both hemispheres (Zhao et al., 2021). None of these studies includes the effect that biogenic organics present in the sea water have on the SSA number emission flux, nor the potential impact of organics on the sea spray flux temperature dependence (Section 3.1). Including these processes could increase the sensitivity of cloud properties to marine biology in models.

Observational Southern Ocean cloud comparisons were successful when the OCEANFILMS model (cf. Section 3.2; McCoy et al., 2015) and improved sulfur chemistry and DMS oxidation parameterizations were implemented (Revell et al., 2019), resulting in a more realistic radiative impact (Mulcahy et al., 2018). Other global climate model simulations (i.e., in CAM6) corresponding to flight sampling were less successful as CCN and droplet number concentration were biased low, potentially due to underproduction of Aitken aerosols (McCoy et al., 2021). Neglecting biogenic NPF (cf. Section 2.2) in global climate models critically impacts the cloud-related radiative effect (Gordon et al., 2016), which is consistent with the difficulty of capturing remote marine environments in other recent climate models (McCoy et al., 2021). This further highlights the importance of global climate models incorporating more nuances of aerosol pathways and their interactions with the cloud-climate system (cf. Sections 2–5, 7, **Figure 2**).

There are also many uncertainties remaining on the impact of INPs and secondary ice formation processes on marine cloud properties (Bellouin et al., 2020). In the Southern Ocean, the underestimation of ice formation in CAM6 (D'Alessandro et al., 2019) results in too much cloud liquid and too little cloud ice in mixed-phase clouds. One reason could be that the cloud lifetime is reduced in models because their CCN/INP ratios are too low, because of either too low marine CCN or too high marine INPs concentrations (Vergara-Temprado et al., 2018). As noted by Field et al. (2017), another reason could be that most models misrepresent several important secondary ice formation mechanisms (e.g., rime splintering, ice–ice collision fragmentation, droplet shattering during freezing, and fragmentation during sublimation of ice bridges) which were found of major importance in the summertime Southern Ocean (Järvinen et al., 2022).

As for cloud measurements, which are rarely accessible in situ, a breakthrough is expected to come from satellite

remote sensing observations, which, in the last 5 years, have progressed significantly by increasing their spatial and temporal resolutions (e.g., TropOMI, GEMS), providing more detailed spectral information and developing new approaches for cloud inversion (Tzabari et al., 2021). Combining multiple satellite datasets resulted in better insights into complex processes such as detection and quantification of mixed-phase clouds in subpolar regions (Listowski et al., 2020), areas notoriously difficult to survey with a single sensor. Without being able to replace the ground-based evidence, the continued support and development of satellite observations, particularly of the aerosol, cloud, and climate system, is critical (National Academies of Sciences, Engineering, Medicine, 2018). Such satellite observing systems (e.g., ACCP, the Aerosol and Cloud, Convection and Precipitation Designated Observable, <https://science.nasa.gov/earth-science/decadal-surveys/decadal-accp/>) open new opportunities for marine science and complement the complex, expensive, and difficult to achieve in situ measurements in remote regions. **Figure 4** summarizes the recent investigations and the current understanding on the biological influence impacting cloud properties as discussed above.

7. Climate change impact and potential feedback loops

We presented recent knowledge on how marine biogeochemistry could impact climate via gas and aerosol emissions and their impact on clouds. Here we discuss how future climate may affect biogeochemical changes, closing feedback loops. The key feedback loop in which climate influences ocean biogeochemistry and in turn influences climate via DMS emissions and cloud formation is known as the CLAW hypothesis (Charlson et al., 1987). Even for DMS—the most studied aerosol-relevant biogenic compound—little progress has been made on its overall impact on climate due to uncertainties on fundamental mechanisms in its production and oxidation pathways (Ghahreman et al., 2017; Yu et al., 2021). While significant efforts have been undertaken to understand the effects of ocean biogenic emissions on climate, the opposite is less frequently investigated. If a changing climate (e.g., changes in temperature, ocean acidity and radiation via cloud cover) affects aerosol sources, there will likely be an additional aerosol-mediated cloud feedback (Gettelman and Sherwood, 2016; Gettelman et al., 2016). Climate controls on aerosol and cloud activation processes must be better

understood (Gordon et al., 2017; Tegen and Schepanski, 2018; Twohy et al., 2021).

Temperature—The average ocean surface temperature is expected to increase by 2.5°C by 2100, with consequences for marine biology (Bindoff et al., 2019). Increases in SST may have different impacts on SSA emissions. Struthers et al. (2013) predicted a general increase in sea salt aerosol number emissions due to both increasing surface wind speed (with nonuniform effects for different size ranges) and SST. However, taking into account the impact of biogeochemistry on sub-micron SSA emission may lead to different conclusions; for instance, ocean warming is expected to expand the distribution range and cell abundances of *Synechococcus* (Flombaum et al., 2013). Also, in a changing climate, the strong temperature and humidity dependence of ice nucleation rates may lead to less ice crystal formation (all other parameters equal). Due to the lack of understanding of the biological processes leading to more or less INPs, the impact of a redistribution of different phytoplankton assemblages on future INP concentrations is unknown.

CO₂—The oceanic uptake of anthropogenic CO₂ is leading to surface ocean acidification (Findlay and Turley, 2021). This acidification may result in climate feedbacks via changes to the net production and emission flux of short-lived climate-active gases, including DMS (Hopkins et al., 2020). The response of DMS to ocean acidification has been explored primarily by means of large-scale in situ mesocosm experiments, typically performed by exposing nutrient-induced blooms of phytoplankton to varying levels of CO₂ over a period of 4–6 weeks. In general, net DMS production decreases with increasing CO₂ concentrations (Hopkins et al., 2020). Inclusion of these results in Earth System Models has suggested a potential positive climate feedback driven by major reductions of DMS production in high-productivity areas, and consequent changes to aerosol and cloud properties that lead to surface warming (Six et al., 2013; Schwinger et al., 2017). In a first attempt to model the entire CLAW cycle, from ocean to clouds and back to ocean, Wang et al. (2018) concluded that truly coupled and dynamic marine ecosystem and sulfur models are needed if we are to assess the occurrence and strength of DMS-mediated climate feedback as presented in the CLAW hypothesis. Due to these limitations in model configuration, Earth System models diverge even in the sign of the simulated future trends of ocean DMS emission under global warming scenarios (Bock et al., 2021). Moreover, the effects of other stressors that act in concert with ocean acidification, such as warming, deoxygenation, and especially solar radiation, remain poorly explored although there has been some advances (Boyd et al., 2016; Husserr et al., 2017). Moving forward requires improved mechanistic representation of DMS-related processes in climate models, and the integration of experimental data which considers the influence of multiple climate stressors on DMS production.

Cloud coverage—Satellite records show that boundary layer cloud cover has decreased between the 1980s and the 2000s across the subtropics (Norris et al., 2016). This

resulted in a positive shortwave subtropical cloud feedback due to increasing SST (Myers and Norris, 2016; McCoy et al., 2017a). Positive subtropical feedback is expected under future climate warming while other regions and cloud types experience more varied, uncertain shortwave feedback (e.g., Myers et al., 2021b; Scott et al., 2020; Sherwood et al., 2020; Zelinka et al., 2020; Ceppi and Nowack, 2021; McCoy et al., 2023). Additional to this uncertainty on the current and future evolution of the cloud cover, there are large unknowns on how associated changes in solar radiation, and in particular ultraviolet light, influence the distribution and activity of phytoplankton assemblages, their grazers and the concurrent bacterioplankton across oceanic regions (Gao et al., 2007; Ruiz-González et al., 2013; Häder et al., 2015). These unknowns hamper our ability to anticipate how changes in exposure to solar radiation will result in changes in the planktonic release of cloud-forming substances, and what will be the climate feedbacks, although some progress has been made with a few trace gases (Galí et al., 2011; McLeod et al., 2021).

Future climate effects are foreseen to be most prominent at the poles, with many studies showing effects of reducing sea ice cover in the Arctic. Among other effects, the increase in open water surfaces would cause an increase in SSA production as well as changes in its composition (Kirpes et al., 2019; Burrows et al., 2022; Willis et al., 2023). In addition, an increase in open water in the polar oceans (in both hemispheres) allows for more marine gas-phase organic emissions, which can either contribute to NPF or to existing particle growth.

Finally, future studies should take advantage of increasing modeling capacities to run Earth system models at higher spatial resolution and/or with inclusion of more processes believed important such as those highlighted above. The main processes included in recent regional/global models are summarized in **Figure 5a** with an attempt to identify how are they expected to be impacted under climate change. To these, **Figure 5b** adds processes identified as phenomenologically important, but not yet well enough characterized to be included in regional/global models.

Overall, the recent observations mentioned across the sections of this article show the importance of conducting coordinated ocean–atmosphere studies embracing the diversity of the aerosol sources and understanding their biogeochemical drivers. The long-term measurements in remote locations should be co-localized in both the atmosphere and ocean, which is technically challenging but also missing due to a historical relative impermeability between the 2 research communities. Laboratory studies are essential for disentangling complexity and understanding processes. While results must be carefully considered, process-based studies linking ocean biogeochemistry and the atmospheric composition should be (i) exportable to the modeling community, hence guided by the processes identified as missing in the models, (ii) designed to include model-compatible variables, and (iii) quantitative. Lastly, utilization of long-term aerosol records to interpret past aerosol changes in terms of climate




Figure 5. The evolving view of marine ecosystem impacts on aerosols and climate. (a) Traditional view of the interconnections between marine ecosystems, aerosols, and clouds, and main processes that have been considered in regional/global model studies (but may not be included in all models), “+” signs indicate how future warming is expected to change total oceanic biomass, and how the resulting processes would be impacted under this high versus low biological activity. (b) Processes identified as important but not yet well characterized enough to be included in regional/global models (bold). The impact of climate change is not shown since different factors may result in opposing effects.

variability remains essential for robust future projections of aerosol distributions and feedbacks (Tegen and Schepanski, 2018).

Data accessibility statement

All data that were considered in this work are available in the cited literature.

Funding

This publication resulted in part from support from the U.S. National Science Foundation (Grant OCE-1840868) to the Scientific Committee on Oceanic Research (SCOR). K. Sellegrì and R. Simó received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreements nos. 771369 and 834162, respectively). B. Wang acknowledges the support from the National Science Foundation of China (grant nos. 41775133, 42075076, 42142017). S. Burrows was funded by the U.S. Department of Energy (DOE), Office of Science, Office of Biological and Environmental Research through the Early Career Research Program. J. Schmale holds the Ingvar Kamprad Chair for Extreme Environments Research sponsored by Ferring Pharmaceuticals. I. L. McCoy was

supported by the NOAA cooperative agreements NA17OAR4320101 and NA22OAR4320151.

Competing interests

The authors have declared that no competing interests exist.

Author contributions

Contributed to conception and design: KS, RS, BW.

Drafted and revised the article: All authors.

Approved the submitted version for publication: All authors.

References

Abbatt, JPD, Leaitch, WR, Aliabadi, AA, Bertram, AK, Blanchet, JP, Boivin-Rioux, A, Bozem, H, Burkart, J, Chang, RYW, Charette, J, Chaubey, JP, Christensen, RJ, Cirisan, A, Collins, DB, Croft, B, Dionne, J, Evans, GJ, Fletcher, CG, Galí, M, Ghahremaninezhad, R, Girard, E, Gong, W, Gosselin, M, Gourdal, M, Hanna, SJ, Hayashida, H, Herber, AB, Hesarakis, S, Hoor, P, Huang, L, Husscherr, R, Irish, VE, Keita, SA, Kodros, JK, Köllner, F, Kolonjari, F, Kunkel, D, Ladino, LA, Law, K, Levasseur,

- M, Libois, Q, Liggio, J, Lizotte, M, Macdonald, KM, Mahmood, R, Martin, RV, Mason, RH, Miller, LA, Moravek, A, Mortenson, E, Mungall, EL, Murphy, JG, Namazi, M, Norman, AL, O'Neill, NT, Pierce, JR, Russell, LM, Schneider, J, Schulz, H, Sharma, S, Si, M, Staebler, RM, Steiner, NS, Thomas, JL, von Salzen, K, Wentzell, JJB, Willis, MD, Wentworth, GR, Xu, JW, Yakobi-Hancock, JD. 2019. Overview paper: New insights into aerosol and climate in the Arctic. *Atmospheric Chemistry and Physics* **19**(4): 2527–2560. DOI: <http://dx.doi.org/10.5194/acp-19-2527-2019>.
- Albrecht, B, Ghatge, V, Mohrmann, J, Wood, R, Zuidema, P, Bretherton, C, Schwartz, C, Eloranta, E, Glienke, S, Donaher, S, Sarkar, M, McGibbon, J, Nugent, A, Shaw, RA, Fugal, J, Minnis, P, Pali-knoda, R, Lussier, L, Jensen, J, Vivekanandan, J, Ellis, S, Tsai, P, Rilling, R, Haggerty, J, Campos, T, Stell, M, Reeves, M, Beaton, S, Allison, J, Stossmeister, G, Hall, S, Schmidt, S. 2019. Cloud system evolution in the trades-CSET: Following the evolution of boundary layer cloud systems with the NSF/NCAR GV. *Bulletin of the American Meteorological Society* **100**(1): 93–121.
- Albrecht, BA. 1989. Aerosols, cloud microphysics, and fractional cloudiness. *Science* **245**(4923): 1227–1230.
- Almeida, J, Schobesberger, S, Kürten, A, Ortega, IK, Kupiainen-Mäkelä, O, Praplan, AP, Adamov, A, Amorim, A, Bianchi, F, Breitenlechner, M, David, A, Dommen, J, Donahue, NM, Downard, A, Dunne, E, Duplissy, J, Ehrhart, S, Flagan, RC, Franchin, A, Guida, R, Hakala, J, Hansel, A, Heinritzi, M, Henschel, H, Jokinen, T, Junninen, H, Kajos, M, Kangasluoma, J, Keskinen, H, Kupc, A, Kürten, T, Kvashin, AN, Laaksonen, A, Lehtipalo, K, Leiminger, M, Leppä, J, Loukonen, V, Makhmutov, V, Mathot, S, McGrath, MJ, Nieminen, T, Olenius, T, Onnela, A, Petäjä, T, Riccobono, F, Riipinen, I, Rissanen, M, Rondo, L, Ruuskanen, T, Santos, FD, Sarnela, N, Schallhart, S, Schnitzhofer, R, Seinfeld, JH, Simon, M, Sipilä, M, Stozhkov, Y, Stratmann, F, Tomé, A, Tröstl, J, Tsagkogeorgas, G, Vaattovaara, P, Viisanen, Y, Virtanen, A, Vrtala, A, Wagner, PE, Weingartner, E, Wex, H, Williamson, C, Wimmer, D, Ye, P, Yli-Juuti, T, Carslaw, KS, Kulmala, M, Curtius, J, Baltensperger, U, Worsnop, DR, Vehkamäki, H, Kirkby, J. 2013. Molecular understanding of sulphuric acid–amine particle nucleation in the atmosphere. *Nature* **502**(7472): 359–363. DOI: <https://doi.org/10.1038/nature12663>.
- Alpert, PA, Aller, JY, Knopf, DA. 2011a. Ice nucleation from aqueous NaCl droplets with and without marine diatoms. *Atmospheric Chemistry and Physics* **11**: 5539–5555. DOI: <http://dx.doi.org/10.5194/acp-11-5539-2011>.
- Alpert, PA, Aller, JY, Knopf, DA. 2011b. Initiation of the ice phase by marine biogenic surfaces in supersaturated gas and supercooled aqueous phases. *Physical Chemistry Chemical Physics* **13**(44): 19882–19894.
- Alpert, PA, Ciuraru, R, Rossignol, S, Passananti, M, Tinel, L, Perrier, S, Dupart, Y, Steimer, SS, Ammann, M, Donaldson, DJ, George, C. 2017. Fatty acid surfactant photochemistry results in new particle formation. *Scientific Reports* **7**(1): 12693. DOI: <http://dx.doi.org/10.1038/s41598-017-12601-2>.
- Alpert, PA, Kiltathau, WP, O'Brien, RE, Moffet, RC, Gilles, MK, Wang, B, Laskin, A, Aller, JY, Knopf, DA. 2022. Ice-nucleating agents in sea spray aerosol identified and quantified with a holistic multimodal freezing model. *Science Advances* **8**(44): eabq6842. DOI: <http://dx.doi.org/10.1126/sciadv.abq6842>.
- Altaratz, O, Koren, I, Remer, LA, Hirsch, E. 2014. Cloud invigoration by aerosols—Coupling between microphysics and dynamics. *Atmospheric Research* **140–141**: 38–60.
- Altieri, KE, Fawcett, SE, Hastings, MG. 2021. Reactive nitrogen cycling in the atmosphere and ocean. *Annual Review of Earth and Planetary Sciences* **49**: 523–550.
- Asmi, E, Freney, E, Hervo, M, Picard, D, Rose, C, Colomb, A, Sellegrì, K. 2012. Aerosol cloud activation in summer and winter at puy-de-Dôme high altitude site in France. *Atmospheric Chemistry and Physics* **12**(23): 11589–11607. DOI: <http://dx.doi.org/10.5194/acp-12-11589-2012>.
- Ault, AP, Moffet, RC, Baltrusaitis, J, Collins, DB, Ruppel, MJ, Cuadra-Rodriguez, LA, Zhao, D, Guasco, TL, Ebben, CJ, Geiger, FM, Bertram, TH, Prather, KA, Grassian, VH. 2013. Size-dependent changes in sea spray aerosol composition and properties with different seawater conditions. *Environmental Science & Technology* **47**(11): 5603–5612. DOI: <http://dx.doi.org/10.1021/es400416g>.
- AuYang, D, Chen, J, Zheng, W, Zhang, Y, Shi, G, Sonke, JE, Cartigny, P, Hongming, C, Yuan, W, Liu, L, Gai, P, Liu, C. 2022. South-hemispheric marine aerosol Hg and S isotope compositions reveal different oxidation pathways. *National Science Open* **1**(2): 20220014. DOI: <http://dx.doi.org/10.1360/nso/20220014>.
- Baccarini, A, Karlsson, L, Dommen, J, Duplessis, P, Vüllers, J, Brooks, IM, Saiz-Lopez, A, Salter, M, Tjernström, M, Baltensperger, U, Zieger, P, Schmale, J. 2021. Frequent new particle formation over the high Arctic pack ice by enhanced iodine emissions. *Nature Communications* **11**(1): 4924. DOI: <http://dx.doi.org/10.1038/s41467-020-18551-0>.
- Baker, MB, Peter, T. 2008. Small-scale cloud processes and climate. *Nature* **451**: 299–300. DOI: <http://dx.doi.org/10.1038/nature06594>.
- Bates, TS, Quinn, PK, Coffman, DJ, Johnson, JE, Upchurch, L, Saliba, G, Lewis, S, Graff, J, Russell, LM, Behrenfeld, MJ. 2020. Variability in marine plankton ecosystems are not observed in freshly emitted sea spray aerosol over the North Atlantic Ocean. *Geophysical Research Letters* **47**(1):

- e2019GL085938. DOI: <https://doi.org/10.1029/2019GL085938>.
- Beaupré, SR, Kieber, DJ, Keene, WC, Long, MS, Frossard, AA, Kinsey, JD, Duplessis, P, Chang, R, Maben, JR, Lu, X, Zhu, Y, Bisgrove, J.** 2019. Oceanic efflux of ancient marine dissolved organic carbon in primary marine aerosol. *Science Advances* **5**(10): eaax6535. DOI: <https://doi.org/10.1126/sciadv.aax6535>.
- Beck, LJ, Sarnela, N, Junninen, H, Hoppe, CJM, Garmash, O, Bianchi, F, Riva, M, Rose, C, Peräkylä, O, Wimmer, D, Kausiala, O, Jokinen, T, Ahonen, L, Mikkilä, J, Hakala, J, He, X-C, Kontkanen, J, Wolf, K, Cappelletti, D, Mazzola, M, Traversi, R, Petroselli, C, Viola, AP, Vitale, V, Lange, RD, Masling, A, Nøjgaard, JK, Krejci, R, Karlsson, L, Zieger, P, Jang, S, Lee, K, Vakkari, V, Lampilahti, J, Thakur, RC, Leino, K, Kangasluoma, J, Duplissy, E-M, Siivola, E, Marbouti, M, Jun, TY, Alfonso, S-L, Petäjä, T, Ehn, M, Worsnop, D, Skov, H, Kulmala, M, Kerminen, V-M, Sipilä, M.** 2021. Differing mechanisms of new particle formation at two Arctic sites. *Geophysical Research Letters* **48**(4): e2020GL091334. DOI: <http://dx.doi.org/10.1029/2020GL091334>.
- Bellouin, N, Quaas, J, Gryspeerdt, E, Kinne, S, Stier, P, Watson-Parris, D, Boucher, O, Carslaw, KS, Christensen, M, Daniau, AL, Dufresne, JL, Feingold, G, Fiedler, S, Forster, P, Gettelman, A, Haywood, JM, Lohmann, U, Malavelle, F, Mauritsen, T, McCoy, DT, Myhre, G, Mülmenstädt, J, Neubauer, D, Possner, A, Rugenstein, M, Sato, Y, Schulz, M, Schwartz, SE, Sourdeval, O, Stordelmo, T, Toll, V, Winker, D, Stevens, B.** 2020. Bounding global aerosol radiative forcing of climate change. *Reviews of Geophysics* **58**(1): e2019RG000660. DOI: <http://dx.doi.org/10.1029/2019RG000660>.
- Belviso, S, Bopp, L, Moulin, C, Orr, JC, Anderson, TR, Chu, S, Elliot, S, Maltrud, ME, Simó, R.** 2004. Comparison of global climatological maps of sea surface dimethyl sulfide. *Global Biogeochemical Cycles* **18**(3): GB3013. DOI: <http://dx.doi.org/10.1029/2003GB002193>.
- Bernard, F, Ciuraru, R, Boréave, A, George, C.** 2016. Photosensitized formation of secondary organic aerosols above the air/water interface. *Environmental Science & Technology* **50**(16): 8678–8686. DOI: <http://dx.doi.org/10.1021/acs.est.6b03520>.
- Bertram, TH, Cochran, RE, Grassian, VH, Stone, EA.** 2018. Sea spray aerosol chemical composition: Elemental and molecular mimics for laboratory studies of heterogeneous and multiphase reactions. *Chemical Society Reviews* **47**(7): 2374–2400. DOI: <https://doi.org/10.1039/C7CS00008A>.
- Bian, H, Froyd, K, Murphy, DM, Dibb, J, Darmenov, A, Chin, M, Colarco, PR, da Silva, A, Kucsera, TL, Schill, G, Yu, H, Bui, P, Dollner, M, Weinzierl, B, Smirnov, A.** 2019. Observationally constrained analysis of sea salt aerosol in the marine atmosphere. *Atmospheric Chemistry and Physics* **19**(16): 10773–10785. DOI: <http://dx.doi.org/10.5194/acp-19-10773-2019>.
- Bindoff, NL, Cheung, WWL, Kairo, JG, Arístegui, J, Guinder, VA, Hallberg, R, Hilmi, N, Jiao, N, Karim, MS, Levin, L, O'Donoghue, S, Purca Cui-capusa, SR, Rinkevich, B, Suga, T, Tagliabue, A, Williamson, P.** 2019. Changing ocean, marine ecosystems, and dependent communities, in Pörtner, H-O, Roberts, DC, Masson-Delmotte, V, Zhai, P, Tignor, M, Poloczanska, E, Mintenbeck, K, Alegria, A, Nicolai, M, Okem, A, Petzold, J, Rama, B eds., *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*.
- Bock, J, Michou, M, Nabat, P, Abe, M, Mulcahy, JP, Oivié, DJL, Schwinger, J, Suntharalingam, P, Tjiputra, J, van Hulten, M, Watanabe, M, Yool, A, Séférian, R.** 2021. Evaluation of ocean dimethylsulfide concentration and emission in CMIP6 models. *Biogeosciences* **18**(12): 3823–3860. DOI: <http://dx.doi.org/10.5194/bg-18-3823-2021>.
- Bodas-Salcedo, A, Williams, KD, Ringer, MA, Beau, I, Cole, JN, Dufresne, JL, Koshiro, T, Stevens, B, Wang, Z, Yokohata, T.** 2014. Origins of the solar radiation biases over the Southern Ocean in CFMIP2 models. *Journal of Climate* **27**: 41–56. DOI: <http://dx.doi.org/10.1175/JCLI-D-13-00169.1>.
- Boucher, O, Randall, D, Artaxo, P, Bretherton, C, Feingold, G, Forster, P, Kerminen, V-M, Kondo, Y, Liao, H, Lohmann, U, Rasch, P, Satheesh, SK, Sherwood, S, Stevens, B, Zhang, XY.** 2013. Clouds and aerosols in climate change 2013, in Stocker, TF, Qin, D, Plattner, G-K, Tignor, MMB, Allen, SK, Boschung, J, Nauels, A, Xia, Y, Bex, V, Midgley, PM eds., *The physical science basis Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK; New York, NY: Cambridge University Press: 571–658.
- Boyd, PW, Dillingham, P, McGraw, C, Armstrong, E, Cornwall, CE, Feng, Y, Hurd, C, Gault-Ringold, M, Roleda, MY, Timmins-Schiffman, E, Nunn, BL.** 2016. Physiological responses of a Southern Ocean diatom to complex future ocean conditions. *Nature Climate Change* **6**: 207–213. DOI: <http://dx.doi.org/10.1038/nclimate2811>.
- Brean, J, Dall'Osto, M, Simó, R, Shi, Z, Beddows, DCS, Harrison, RM.** 2021. Open ocean and coastal new particle formation from sulfuric acid and amines around the Antarctic Peninsula. *Nature Geoscience* **14**: 383–388. DOI: <http://dx.doi.org/10.1038/s41561-021-00751-y>.
- Bretherton, CS, Blossey, PN.** 2017. Understanding mesoscale aggregation of shallow cumulus convection using large-eddy simulation. *Journal of Advances in Modeling Earth Systems* **9**(8): 2798–2821. DOI: <http://dx.doi.org/10.1002/2017ms000981>.
- Bretherton, CS, Blossey, PN, Uchida, J.** 2007. Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo. *Geophysical*

- Research Letters* **34**(3). DOI: <https://doi.org/10.1029/2006GL027648>.
- Brüggemann, M, Hayeck, N, George, C.** 2018. Interfacial photochemistry at the ocean surface is a global source of organic vapors and aerosols. *Nature Communications* **9**(1): 2101. DOI: <http://dx.doi.org/10.1038/s41467-018-04528-7>.
- Budge, SM, Devred, E, Forget, M-H, Stuart, V, Trzcinski, MK, Sathyendranath, S, Platt, T.** 2014. Estimating concentrations of essential omega-3 fatty acids in the ocean: Supply and demand. *ICES Journal of Marine Science* **71**(7): 1885–1893.
- Bullock, HA, Luo, H, Whitman, WB.** 2017. Evolution of dimethylsulfoniopropionate metabolism in marine phytoplankton and bacteria. *Frontiers in Microbiology* **8**: 637. DOI: <http://dx.doi.org/10.3389/fmicb.2017.00637>.
- Burkart, J, Willis, MD, Bozem, H, Thomas, JL, Law, K, Hoor, P, Aliabadi, AA, Köllner, F, Schneider, J, Herber, A, Abbatt, JPD, Leaitch, WR.** 2017. Summertime observations of elevated levels of ultrafine particles in the high Arctic marine boundary layer. *Atmospheric Chemistry and Physics* **17**: 5515–5535. DOI: <https://doi.org/10.5194/acp-17-5515-2017>.
- Burrows, SM, Easter, RC, Liu, X, Ma, P-L, Wang, H, Elliott, SM, Singh, B, Zhang, K, Rasch, PJ.** 2022. OCEANFILMS (Organic Compounds from Ecosystems to Aerosols: Natural Films and Interfaces via Langmuir Molecular Surfactants) sea spray organic aerosol emissions—Implementation in a global climate model and impacts on clouds. *Atmospheric Chemistry and Physics* **22**: 5223–5251. DOI: <http://dx.doi.org/10.5194/acp-22-5223-2022>.
- Burrows, SM, Hoose, C, Pöschl, U, Lawrence, MG.** 2013. Ice nuclei in marine air: Biogenic particles or dust? *Atmospheric Chemistry and Physics* **13**(1): 245–267. DOI: <http://dx.doi.org/10.5194/acp-13-245-2013>.
- Burrows, SM, Ogunro, O, Frossard, AA, Russell, LM, Rasch, PJ, Elliott, SM.** 2014. A physically based framework for modeling the organic fractionation of sea spray aerosol from bubble film Langmuir equilibria. *Atmospheric Chemistry and Physics* **14**: 13601–13629. DOI: <http://dx.doi.org/10.5194/acp-14-13601-2014>.
- Carpenter, LJ, Archer, SD, Beale, R.** 2012. Ocean-atmosphere trace gas exchange. *Chemical Society Reviews* **41**: 6473–6506.
- Carslaw, KS, Gordon, H, Hamilton, DS, Johnson, JS, Regayre, LA, Yoshioka, M, Pringle, KJ.** 2017. Aerosols in the pre-industrial atmosphere. *Current Climate Change Reports* **3**(1): 1–15. DOI: <http://dx.doi.org/10.1007/s40641-017-0061-2>.
- Carslaw, KS, Lee, LA, Reddington, CL, Pringle, KJ, Rap, A, Forster, PM, Mann, GW, Spracklen, DV, Woodhouse, MT, Regayre, LA, Pierce, JR.** 2013. Large contribution of natural aerosols to uncertainty in indirect forcing. *Nature* **503**(7474): 67–71.
- Ceburnis, D, Rinaldi, M, Ovadnevaite, J, Martucci, G, Giulianelli, L, O'Dowd, CD.** 2016. Marine submicron aerosol gradients, sources and sinks. *Atmospheric Chemistry and Physics* **16**: 12425–12439. DOI: <https://doi.org/10.5194/acp-16-12425-2016>.
- Ceppi, P, Nowack, P.** 2021. Observational evidence that cloud feedback amplifies global warming. *Proceedings of the National Academy of Sciences of the United States of America* **118**(30): e2026290118. DOI: <http://dx.doi.org/10.1073/pnas.2026290118>.
- Cesana, GV, Khadir, T, Chepfer, H, Chiriaco, M.** 2022. Southern Ocean solar reflection biases in CMIP6 models linked to cloud phase and vertical structure representations. *Geophysical Research Letters* **49**: e2022GL099777. DOI: <http://dx.doi.org/10.1029/2022GL099777>.
- Charlson, R, Lovelock, J, Andreae, M, Warren, SG.** 1987. Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate. *Nature* **326**(6114): 655–661. DOI: <https://doi.org/10.1038/326655a0>.
- Chen, Q, Sherwen, T, Evans, M, Alexander, B.** 2018. DMS oxidation and sulfur aerosol formation in the marine troposphere: A focus on reactive halogen and multiphase chemistry. *Atmospheric Chemistry and Physics* **18**: 13617–13637. DOI: <https://doi.org/10.5194/acp-18-13617-2018>.
- China, S, Alpert, PA, Zhang, B, Schum, S, Dzepina, K, Wright, K, Owen, RC, Fialho, P, Mazzoleni, LR, Mazzoleni, C, Knopf, DA.** 2017. Ice cloud formation potential by free tropospheric particles from long-range transport over the Northern Atlantic Ocean. *Journal of Geophysical Research: Atmospheres* **122**(5): 3065–3079. DOI: <http://dx.doi.org/10.1002/2016JD025817>.
- Christensen, OB, Kjellström, E, Dieterich, C, Gröger, M, Meier, HEM.** 2022. Atmospheric regional climate projections for the Baltic Sea region until 2100. *Earth System Dynamics* **13**(1): 133–157. DOI: <http://dx.doi.org/10.5194/esd-13-133-2022>.
- Christiansen, S, Salter, ME, Gorokhova, E, Nguyen, QT, Bilde, M.** 2019. Sea spray aerosol formation: Laboratory results on the role of air entrainment, water temperature, and phytoplankton biomass. *Environmental Science & Technology* **53**(22): 13107–13116.
- Ciuraru, R, Fine, L, van Pinxteren, M, D'Anna, B, Herrmann, H, George, C.** 2015. Unravelling new processes at interfaces: Photochemical isoprene production at the sea surface. *Environmental Science & Technology* **49**(22): 13199–13205. DOI: <http://dx.doi.org/10.1021/acs.est.5b02388>.
- Clarke, AD, Varner, JL, Eisele, F, Mauldin, RL, Tanner, D, Litchy, M.** 1998. Particle production in the remote marine atmosphere: Cloud outflow and subsidence during ACE 1. *Journal of Geophysical Research: Atmospheres* **103**(D13): 16397–16409. DOI: <http://dx.doi.org/10.1029/97jd02987>.
- Cochran, RE, Laskina, O, Jayarathne, T, Laskin, A, Laskin, J, Lin, P, Sultana, C, Lee, C, Moore, KA, Cappa, CD, Bertram, TH, Prather, KA, Grassian, VH, Stone, EA.** 2016. Analysis of organic anionic surfactants in fine and coarse fractions of freshly emitted sea spray aerosol. *Environmental Science & Technology* **50**(5): 2477–2486.

- Colomb, A, Yassaa, N, Williams, J, Peeken, I, Lochte, K. 2008. Screening volatile organic compounds (VOCs) emissions from five marine phytoplankton species by head space gas chromatography/mass spectrometry (HS-GC/MS). *Journal of Environmental Monitoring* **10**: 325–330. DOI: <http://dx.doi.org/10.1039/b715312k>.
- Cornwell, GC, McCluskey, CS, DeMott, PJ, Prather, KA, Burrows, SM. 2021. Development of heterogeneous ice nucleation rate coefficient parameterizations from ambient measurements. *Geophysical Research Letters* **48**(23): e2021GL095359. DOI: <https://doi.org/10.1029/2021GL095359>.
- Crocker, DR, Hernandez, RE, Huang, HD, Pendergraft, MA, Cao, R, Dai, J, Morris, CK, Deane, GB, Prather, KA, Thiemens, MH. 2020. Biological influence on $\delta^{13}\text{C}$ and organic composition of nascent sea spray aerosol. *ACS Earth and Space Chemistry* **4**(9): 1686–1699. DOI: <https://doi.org/10.1021/acsearthspacechem.0c00072>.
- Crocker, DR, Kaluarachchi, CP, Cao, R, Dinasquet, J, Franklin, EB, Morris, CK, Amiri, S, Petras, D, Nguyen, T, Torres, RR, Martz, TR, Malfatti, F, Goldstein, AH, Tivanski, AV, Prather, KA, Thiemens, MH. 2022. Isotopic insights into organic composition differences between super-micron and sub-micron sea spray aerosol. *Environmental Science & Technology* **56**(14): 9947–9958.
- Curson, AR, Liu, J, Martinez, AB, Green, RT, Chan, Y, Carrion, O, Williams, BT, Zhang, SH, Yang, GP, Page, PCB, Zhang, XH, Todd, JD. 2017. Dimethylsulfoniopropionate biosynthesis in marine bacteria and identification of the key gene in this process. *Nature Microbiology* **2**: 17009. DOI: <http://dx.doi.org/10.1038/nmicrobiol.2017.9>.
- D'Alessandro, JJ, Diao, M, Wu, C, Liu, X, Jensen, JB, Stephens, BB. 2019. Cloud phase and relative humidity distributions over the Southern Ocean in austral summer based on in situ observations and CAM5 simulations. *Journal of Climate* **32**(10): 2781–2805. DOI: <http://dx.doi.org/10.1175/JCLI-D-18-0232.1>.
- Dagan, G, Koren, I, Altaratz, O, Heiblum, RH. 2017. Time-dependent, non-monotonic response of warm convective cloud fields to changes in aerosol loading. *Atmospheric Chemistry and Physics* **17**(12): 7435–7444.
- Dall'Osto, M, Ovadnevaite, J, Paglione, M, Beddows, DCS, Ceburnis, D, Cree, C, Cortés, P, Zamanillo, M, Nunes, SO, Pérez, GL, Ortega-Retuerta, E, Emelianov, M, Vaqué, D, Marrasé, C, Estrada, M, Sala, MM, Vidal, M, Fitzsimons, MF, Beale, R, Airs, R, Rinaldi, M, Decesari, S, Facchini, MC, Harrison, RM, O'Dowd, C, Simó, R. 2017. Antarctic sea ice region as a source of biogenic organic nitrogen in aerosols. *Scientific Reports* **7**: 6047. DOI: <https://doi.org/10.1038/s41598-017-06188-x>.
- Dall'Osto, M, Vaqué, D, Sotomayor-Garcia, A, Cabrera-Brufau, M, Estrada, M, Buchaca, T, Soler, M, Nunes, S, Zeppenfeld, S, van Pinxteren, M, Herrmann, H, Wex, H, Rinaldi, M, Paglione, M, Beddows, DCS, Harrison, RM, Berdalet, E. 2022. Sea ice microbiota in the Antarctic Peninsula modulates cloud-relevant sea spray aerosol production. *Frontiers in Marine Science* **9**: 827061. DOI: <http://dx.doi.org/10.3389/fmars.2022.827061>.
- Danker, J, Sourdeval, O, McCoy, IL, Wood, R, Possner A. 2022. Exploring relations between cloud morphology, cloud phase, and cloud radiative properties in Southern Ocean's stratocumulus clouds. *Atmospheric Chemistry and Physics* **22**(15): 10247–10265.
- de Leeuw, G, Guieu, C, Arneth, A, Bellouin, N, Bopp, L, Boyd, PW, Denier van der Gon, HAC, Desboeufs, KV, Dulac, F, Facchini, MC, Gantt, B, Langmann, B, Mahowald, NM, Maranon, E, O'Dowd, C, Olgun, N, Pulido-Villena, E, Rinaldi, M, Stephanou, EG, Wagener, T. 2014. Ocean–atmosphere interactions of particles, in Liss, PS, Johnson, MT eds., *Ocean-atmosphere interactions of gases and particles*. Berlin, Heidelberg, Germany: Springer Earth System Sciences: 171–246. DOI: http://dx.doi.org/10.1007/978-3-642-25643-1_4.
- DeMott, PJ, Hill, TCJ, McCluskey, CS, Prather, KA, Collins, DB, Sullivan, RC, Ruppel, MJ, Mason, RH, Irish, VE, Lee, T, Hwang, CY, Rhee, TS, Snider, JR, McMeeking, GR, Dhaniyala, S, Lewis, ER, Wentzell, JJB, Abbatt, J, Lee, C, Sultana, CM, Ault, AP, Axson, JL, Diaz Martinez, M, Venero, I, Santos-Figueroa, G, Stokes, MD, Deane, GB, Mayol-Bracero, OL, Grassian, VH, Bertram, TH, Bertram, AK, Moffett, BF, Franc, GD. 2016. Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust particles. *Proceedings of the National Academy of Sciences* **113**(21): 5797–5803. DOI: <https://doi.org/10.1073/pnas.1514034112>.
- Dixon, JL, Hopkins, FE, Stephens, JA, Schäfer, H. 2020. Seasonal changes in microbial dissolved organic sulfur transformations in coastal waters. *Microorganisms* **8**: 337. DOI: <http://dx.doi.org/10.3390/microorganisms8030337>.
- Eastman, R, McCoy, IL, Wood, R. 2022. Wind, rain, and the closed to open cell transition in subtropical marine stratocumulus. *Journal of Geophysical Research: Atmospheres* **127**(20): e2022JD036795. DOI: <https://doi.org/10.1029/2022JD036795>.
- Efraim, A, Rosenfeld, D, Schmale, J, Zhu, Y. 2020. Satellite retrieval of cloud condensation nuclei concentrations in marine stratocumulus by using clouds as CCN chambers. *Journal of Geophysical Research: Atmospheres* **125**: e2020JD032409. DOI: <http://dx.doi.org/10.1029/2020JD032409>.
- Ehn, M, Thornton, JA, Kleist, E, Sipilä, M, Junninen, H, Pullinen, I, Springer, M, Rubach, F, Tillmann, R, Lee, B, Lopez-Hilfiker, F, Andreas, S, Acir, IH, Rissanen, M, Jokinen, T, Schobesberger, S, Kangasluoma, J, Kontkanen, J, Nieminen, T, Kurtén, T, Nielsen, LB, Jørgensen, S, Kjaergaard, HG, Canagaratna, M, Maso, MD, Berndt, T, Petäjä, T, Wahner, A, Kerminen, VM, Kulmala, M,

- Worsnop, DR, Wildt, J, Mentel, TF.** 2014. A large source of low-volatility secondary organic aerosol. *Nature* **506**(7489): 476–479. DOI: <https://doi.org/10.1038/nature13032>.
- Enami, S, Hoffmann, MR, Colussi, AJ.** 2019. Iodide accelerates the processing of biogenic monoterpene emissions on marine aerosols. *ACS Omega* **4**: 7574–7580. DOI: <http://dx.doi.org/10.1021/acsomega.9b00024>.
- Ervens, B, Sorooshian, A, Aldhaif, AM, Shingler, T, Crosbie, E, Ziemba, L, Campuzano-Jost, P, Jimenez, JL, Wisthaler, A.** 2018. Is there an aerosol signature of chemical cloud processing? *Atmospheric Chemistry and Physics* **18**(21): 16099–16119. DOI: <http://dx.doi.org/10.5194/acp-18-16099-2018>.
- Facchini, MC, Rinaldi, M, Decesari, S, Carbone, C, Finessi, E, Mircea, M, Fuzzi, S, Ceburnis, D, Flanagan, R, Nilsson, ED, de Leeuw, G, Martino, M, Woeltjen, J, O'Dowd, CD.** 2008. Primary submicron marine aerosol dominated by insoluble organic colloids and aggregates. *Geophysical Research Letters* **35**(17): L17814. DOI: <https://doi.org/10.1029/2008GL034210>.
- Fan, J, Wang, Y, Rosenfeld, D, Liu, X.** 2016. Review of aerosol–cloud interactions: Mechanisms, significance, and challenges. *Journal of the Atmospheric Sciences* **73**(11): 4221–4252. DOI: <https://doi.org/10.1175/JAS-D-16-0037.1>.
- Fiddes, SL, Woodhouse, M, Nicholls, Z, Lane, TP, Schofield, R.** 2018. Cloud, precipitation and radiation responses to large perturbations in global dimethyl sulfide. *Atmospheric Chemistry and Physics* **18**: 10177–10198.
- Field, PR, Lawson, RP, Brown, PRA, Lloyd, G, Westbrook, C, Moiseev, D, Miltenberger, A, Nenes, A, Blyth, A, Choularton, T, Connolly, P, Buehl, J, Crosier, J, Cui, Z, Dearden, C, DeMott, P, Flossmann, A, Heymsfield, A, Huang, Y, Kalesse, H, Kanji, ZA, Korolev, A, Kirchgaessner, A, Lasher-Trapp, S, Leisner, T, McFarquhar, G, Phillips, V, Stith, J, Sullivan, S.** 2017. Secondary ice production: Current state of the science and recommendations for the future. *Meteorological Monographs* **58**: 7.1–7.20. DOI: <https://doi.org/10.1175/AMSMONOGRAPH-D-16-0014.1>.
- Findlay, HS, Turley, C.** 2021. Ocean acidification and climate change, in Letcher, TM ed., *Climate change*. Amsterdam, the Netherlands: Elsevier: 251–279.
- Fiore, CL, Longnecker, K, Kido Soule, MC, Kujawinski, EB.** 2015. Release of ecologically relevant metabolites by the cyanobacterium *Synechococcus elongatus* CCMP 1631. *Environmental Microbiology* **17**(10): 3949–3963. DOI: <http://dx.doi.org/10.1111/1462-2920.12899>.
- Flombaum, P, Gallegos, JL, Gordillo, RA, Rincón, J, Zabala, LL, Jiao, N, Karl, DM, Li, WK, Lomas, MW, Veneziano, D, Vera, CS, Vrugt, JA, Martiny, AC.** 2013. Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proceedings of the National Academy of Sciences of the United States of America* **110**(24): 9824–9829. DOI: <https://doi.org/10.1073/pnas.1307701110>.
- Forestieri, SD, Moore, KA, Martinez Borrero, R, Wang, A, Stokes, MD, Cappa, CD.** 2018. Temperature and composition dependence of sea spray aerosol production. *Geophysical Research Letters* **45**(14): 7218–7225.
- Fossum, KN, Ovadnevaite, J, Ceburnis, D, Dall'Osto, M, Marullo, S, Bellacicco, M, Simó, R, Liu, D, Flynn, M, Zuend, A, O'Dowd, C.** 2018. Summertime primary and secondary contributions to Southern Ocean cloud condensation nuclei. *Scientific Reports* **8**: 13844. DOI: <http://dx.doi.org/10.1038/s41598-018-32047-4>.
- Fossum, KN, Ovadnevaite, J, Ceburnis, D, Preißler, J, Snider, JR, Huang, R-J, Zuend, A, O'Dowd, C.** 2020. Sea-spray regulates sulfate cloud droplet activation over oceans. *npj Climate and Atmospheric Science* **3**: 14. DOI: <http://dx.doi.org/10.1038/s41612-020-0116-2>.
- Frenay, E, Sellegrì, K, Nicosia, A, Trueblood, JT, Rinaldi, M, Williams, LR, Prévôt, ASH, Thyssen, M, Grégori, G, Haëntjens, N, Dinasquet, J, Obernosterer, I, Van-Wambeke, F, Engel, A, Zäncker, B, Desboeufs, K, Asmi, E, Timmonen, H, Guieu, C.** 2021. Mediterranean nascent sea spray organic aerosol and relationships with seawater biogeochemistry. *Atmospheric Chemistry and Physics* **21**(13): 10625–10641. DOI: <http://dx.doi.org/10.5194/acp-21-10625-2021>.
- Frossard, AA, Long, MS, Keene, WC, Duplessis, P, Kinsey, JD, Maben, JR, Kieber, DJ, Chang, RY-W, Beaupré, SR, Cohen, RC, Lu, X, Bisgrove, J, Zhu, Y.** 2019. Marine aerosol production via detrainment of bubble plumes generated in natural seawater with a forced-air Venturi. *Journal of Geophysical Research: Atmospheres* **124**(20): 10931–10950. DOI: <https://doi.org/10.1029/2019JD030299>.
- Fu, H, Ciuraru, R, Dupart, Y, Passananti, M, Tinel, L, Rossignol, S, Perrier, S, Donaldson, DJ, Chen, J, George, C.** 2015. Photosensitized production of atmospherically reactive organic compounds at the air/aqueous interface. *Journal of the American Chemical Society* **137**(26): 8348–8351. DOI: <http://dx.doi.org/10.1021/jacs.5b04051>.
- Fuentes, E, Coe, H, Green, D, McFiggans, G.** 2011. On the impacts of phytoplankton-derived organic matter on the properties of the primary marine aerosol—Part 2: Composition, hygroscopicity and cloud condensation activity. *Atmospheric Chemistry and Physics* **11**: 2585–2602. DOI: <https://doi.org/10.5194/acp-11-2585-2011>.
- Galí, M, Devred, E, Babin, M, Levasseur, M.** 2019. Decadal increase in Arctic dimethylsulfide emission. *Proceedings of the National Academy of Sciences* **116**(39): 19311–19317. DOI: <https://doi.org/10.1073/pnas.1904378116>.
- Galí, M, Levasseur, M, Devred, E, Simó, R, Babin, M.** 2018. Sea-surface dimethylsulfide (DMS) concentration from

- satellite data at global and regional scales. *Biogeosciences* **15**: 3497–3519. DOI: <http://dx.doi.org/10.5194/BG-15-3497-2018>.
- Galí, M, Saló, V, Almeda, R, Calbet, A, Simó, R.** 2011. Stimulation of gross dimethylsulfide (DMS) production by solar radiation. *Geophysical Research Letters* **38**(15): L15612. DOI: <http://dx.doi.org/10.1029/2011GL048051>.
- Galí, M, Simó, R.** 2015. A meta-analysis of oceanic DMS and DMSP cycling processes: Disentangling the summer paradox. *Global Biogeochemical Cycle* **29**(4): 496–515. DOI: <http://dx.doi.org/10.1002/2014GB004940>.
- Gantt, B, Meskhidze, N, Facchini, MC, Rinaldi, M, Ceburnis, D, O'Dowd, CD.** 2011. Wind speed dependent size-resolved parameterization for the organic mass fraction of sea spray aerosol. *Atmospheric Chemistry and Physics* **11**(16): 8777–8790. DOI: <http://dx.doi.org/10.5194/acp-11-8777-2011>.
- Gao, K, Wu, Y, Li, G, Wu, H, Villafañe, VE, Helbling, EW.** 2007. Solar UV radiation drives CO₂ fixation in marine phytoplankton: A double-edged sword. *Plant Physiology* **144**(1): 54–59. DOI: <http://dx.doi.org/10.1104/pp.107.098491>.
- Gerard, V, Nozière, B, Baduel, C, Fine, L, Frossard, AA, Cohen, RC.** 2016. Anionic, cationic, and nonionic surfactants in atmospheric aerosols from the Baltic Coast at Askö, Sweden: Implications for cloud droplet activation. *Environmental Science & Technology* **50**(6): 2974–2982. DOI: <https://doi.org/10.1021/acs.est.5b05809>.
- Gettelman, A, Lin, L, Medeiros, B, Olson, J.** 2016. Climate feedback variance and the interaction of aerosol forcing and feedbacks. *Journal of Climate* **29**(18): 6659–6675. DOI: <https://doi.org/10.1175/JCLI-D-16-0151.1>.
- Gettelman, A, Sherwood, SC.** 2016. Processes responsible for cloud feedback. *Current Climate Change Reports* **2**: 179–189. DOI: <https://doi.org/10.1007/s40641-016-0052-8>.
- Ghahreman, R, Norman, A-L, Croft, B, Martin, RV, Pierce, JR, Burkart, J, Rempillo, O, Bozem, H, Kunkel, D, Thomas, JL, Aliabadi, AA, Wentworth, GR, Levasseur, M, Staebler, RM, Sharma, S, Leaitch, WR.** 2017. Boundary layer and free-tropospheric dimethyl sulfide in the Arctic spring and summer. *Atmospheric Chemistry and Physics* **17**(14): 8757–8770. DOI: <https://doi.org/10.5194/acp-17-8757-2017>.
- Glicker, HS, Lawler, MJ, Chee, S, Resch, J, Garofalo, LA, Mayer, KJ, Prather, K, Farmer, DK, Smith, JN.** 2022. Chemical composition of an ultrafine sea spray aerosol during the sea spray chemistry and particle evolution experiment. *ACS Earth and Space Chemistry* **6**(7): 1914–1923. DOI: <http://dx.doi.org/10.1021/acsearthspacechem.2c00127>.
- Gómez Martín, JC, Lewis, TR, James, AD, Saiz-Lopez, A, Plane, JMC.** 2022. Insights into the chemistry of iodine new particle formation: The role of iodine oxides and the source of iodic acid. *Journal of the American Chemical Society* **144**(21): 9240–9253.
- Gordon, H, Kirkby, J, Baltensperger, U, Bianchi, F, Breitenlechner, M, Curtius, J, Dias, A, Dommen, J, Donahue, NM, Dunne, EM, Duplissy, J, Ehrhart, S, Flagan, RC, Frege, C, Fuchs, C, Hansel, A, Hoyle, CR, Kulmala, M, Kürten, A, Lehtipalo, K, Makhmutov, V, Molteni, U, Rissanen, MP, Stozhkov, Y, Tröstl, J, Tsagkogeorgas, G, Wagner, R, Williamson, C, Wimmer, D, Winkler, PM, Yan, C, Carslaw, KS.** 2017. Causes and importance of new particle formation in the present-day and pre-industrial atmospheres. *Journal of Geophysical Research: Atmospheres* **122**(16): 8739–8760. DOI: <https://doi.org/10.1002/2017JD026844>.
- Gordon, H, Sengupta, K, Rap, A, Duplissy, J, Frege, C, Williamson, C, Heinritzi, M, Simon, M, Yan, C, Almeida, J, Tröstl, J, Nieminen, T, Ortega, IK, Wagner, R, Dunne, EM, Adamov, A, Amorim, A, Bernhammer, A-K, Bianchi, F, Breitenlechner, M, Brilke, S, Chen, X, Craven, JS, Dias, A, Ehrhart, S, Fischer, L, Flagan, RC, Franchin, A, Fuchs, C, Guida, R, Hakala, J, Hoyle, CR, Jokinen, T, Junninen, H, Kangasluoma, J, Kim, J, Kirkby, J, Krapf, M, Kürten, A, Laaksonen, A, Lehtipalo, K, Makhmutov, V, Mathot, S, Molteni, U, Monks, SA, Onnela, A, Peräkylä, O, Piel, F, Petäjä, T, Praplan, AP, Pringle, KJ, Richards, NAD, Rissanen, MP, Rondo, L, Sarnela, N, Schobesberger, S, Scott, CE, Seinfeld, JH, Sharma, S, Sipilä, M, Steiner, G, Stozhkov, Y, Stratmann, F, Tomé, A, Virtanen, A, Vogel, AL, Wagner, AC, Wagner, PE, Weingartner, E, Wimmer, D, Winkler, PM, Ye, P, Zhang, X, Hansel, A, Dommen, J, Donahue, NM, Worsnop, DR, Baltensperger, U, Kulmala, M, Curtius, J, Carslaw, KS.** 2016. Reduced anthropogenic aerosol radiative forcing caused by biogenic new particle formation. *Proceedings of the National Academy of Sciences* **113**(43): 12053–12058. DOI: <https://doi.org/10.1073/pnas.1602360113>.
- Grythe, H, Ström, J, Krejci, R, Quinn, P, Stohl, A.** 2014. A review of sea-spray aerosol source functions using a large global set of sea salt aerosol concentration measurements. *Atmospheric Chemistry and Physics* **14**(3): 1277–1297.
- Häder, DP, Williamson, CE, Wängberg, SA, Rautio, M, Rose, KC, Gao, K, Helbling, WE, Sinha, RP, Worrrest, R.** 2015. Effects of UV radiation on aquatic ecosystems and interactions with other environmental factors. *Photochemical and Photobiological Sciences* **14**: 108–126. DOI: <http://dx.doi.org/10.1039/C4PP90035A>.
- Halloran, P, Bell, T, Totterdell, I.** 2010. Can we trust empirical marine DMS parameterisations within projections of future climate? *Biogeosciences* **7**(5): 1645–1656.
- Halsey, KH, Giovannoni, SJ.** 2023. Biological controls on marine volatile organic compound emissions: A balancing act at the sea-air interface. *Earth-Science*

- Reviews* **240**: 104360. DOI: <http://dx.doi.org/10.1016/j.earscirev.2023.104360>.
- Hamilton, DS, Lee, LA, Pringle, KJ, Reddington, CL, Spracklen, DV, Carslaw, KS.** 2014. Occurrence of pristine aerosol environments on a polluted planet. *Proceedings of the National Academy of Sciences* **111**(52): 18466–18471. DOI: <https://doi.org/10.1073/pnas.1415440111>.
- Harris, E, Sinha, B, van Pinxteren, D, Schneider, J, Poulain, L, Collett, J, D'Anna, B, Fahlbusch, B, Foley, S, Fomba, KW, George, C, Gnauk, T, Henning, S, Lee, T, Mertes, S, Roth, A, Stratmann, F, Borrmann, S, Hoppe, P, Herrmann, H.** 2014. In-cloud sulfate addition to single particles resolved with sulfur isotope analysis during HCCT-2010. *Atmospheric Chemistry and Physics* **14**(8): 4219–4235. DOI: <https://doi.org/10.5194/acp-14-4219-2014>.
- Hartery, S, Toohey, D, Revell, L, Sellegri, K, Kuma, P, Harvey, M, McDonald, AJ.** 2020. Constraining the surface flux of sea spray particles from the Southern Ocean. *Journal of Geophysical Research: Atmospheres* **125**(4): e2019JD032026. DOI: <https://doi.org/10.1029/2019JD032026>.
- He, X-C, Tham, YJ, Dada, L, Wang, M, Finkenzeller, H, Stolzenburg, D, Iyer, S, Simon, M, Kürten, A, Shen, J, Rörup, B, Rissanen, M, Schobesberger, S, Baalbaki, R, Wang, DS, Koenig, TK, Jokinen, T, Sarnela, N, Beck, LJ, Almeida, J, Amanatidis, S, Amorim, A, Ataei, F, Baccarini, A, Bertozzi, B, Bianchi, F, Brilke, S, Caudillo, L, Chen, D, Chiu, R, Chu, B, Dias, A, Ding, A, Dommen, J, Duplissy, J, El Haddad, I, Gonzalez Carracedo, L, Granzin, M, Hansel, A, Heinritzi, M, Hofbauer, V, Junninen, H, Kangasluoma, J, Kemppainen, D, Kim, C, Kong, W, Krechmer, JE, Kvashin, A, Laitinen, T, Lamkaddam, H, Lee, CP, Lehtipalo, K, Leiminger, M, Li, Z, Makhmutov, V, Manninen, HE, Marie, G, Marten, R, Mathot, S, Mauldin, RL, Mentler, B, Möhler, O, Müller, T, Nie, W, Onnela, A, Petäjä, T, Pfeifer, J, Philippov, M, Ranjithkumar, A, Saiz-Lopez, A, Salma, I, Scholz, W, Schuchmann, S, Schulze, B, Steiner, G, Stozhkov, Y, Tauber, C, Tomé, A, Thakur, RC, Väisänen, O, Vazquez-Pufleau, M, Wagner, AC, Wang, Y, Weber, SK, Winkler, PM, Wu, Y, Xiao, M, Yan, C, Ye, Q, Ylisirniö, A, Zauner-Wieczorek, M, Zha, Q, Zhou, P, Flagan, RC, Curtius, J, Baltensperger, U, Kulmala, M, Kerminen, V-M, Kurtén, T, Donahue, NM, Volkamer, R, Kirkby, J, Worsnop, DR, Sipilä, M.** 2021. Role of iodine oxoacids in atmospheric aerosol nucleation. *Science* **371**(6529): 589–595.
- Heintzenberg, J, Charlson, RJ eds.** 2009. *Clouds in the perturbed climate system: Their relationship to energy balance, atmospheric dynamics, and precipitation*. Cambridge, MA: The MIT Press: 597.
- Hepach, H, Hughes, C, Hogg, K, Collings, S, Chance, R.** 2019. Common features of iodate to iodide reduction amongst a diverse range of marine phytoplankton. *Biogeosciences Discussions* 1–27. DOI: <http://dx.doi.org/10.5194/bg-2019-443>.
- Herr, AE, Dacey, JWH, Kiene, RP, McCulloch, RD, Schu-back, N, Tortell, PD.** 2020. Potential roles of dimethylsulfoxide in regional sulfur cycling and phytoplankton physiological ecology in the NE Subarctic Pacific. *Limnology and Oceanography* **66**(1): 76–94. DOI: <http://dx.doi.org/10.1002/lno.11589>.
- Hodshire, AL, Campuzano-Jost, P, Kodros, JK, Croft, B, Nault, BA, Schroder, JC, Jimenez, JL, Pierce, JR.** 2019. The potential role of methanesulfonic acid (MSA) in aerosol formation and growth and the associated radiative forcings. *Atmospheric Chemistry and Physics* **19**(5): 3137–3160. DOI: <http://dx.doi.org/10.5194/acp-19-3137-2019>.
- Hoffmann, EH, Tilgner, A, Schroedner, R, Bräuer, P, Wolke, R, Herrmann, H.** 2016. An advanced modeling study on the impacts and atmospheric implications of multiphase dimethyl sulfide chemistry. *Proceedings of the National Academy of Sciences of the United States of America* **113**: 11776–11781. DOI: <http://dx.doi.org/10.1073/pnas.1606320113>.
- Hopkins, FE, Suntharalingam, P, Gehlen, M, Andrews, O, Archer, SD, Bopp, L, Buitenhuis, E, Dadou, I, Duce, R, Goris, N, Jickells, T, Johnson, M, Keng, F, Law, CS, Lee, K, Liss, PS, Lizotte, M, Malin, G, Murrell, JC, Naik, H, Rees, AP, Schwinger, J, Williamson, P.** 2020. The impacts of ocean acidification on marine trace gases and the implications for atmospheric chemistry and climate. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **476**(2237): 201907692.
- Hoppel, WA, Frick, GM.** 1990. Submicron aerosol size distributions measured over the tropical and South Pacific. *Atmospheric Environment Part A General Topics* **24**(3): 645–659. DOI: [https://doi.org/10.1016/0960-1686\(90\)90020-N](https://doi.org/10.1016/0960-1686(90)90020-N).
- Huang, D, Wang, J, Xia, H, Zhang, Y, Bao, F, Li, M, Chen, C, Zhao, J.** 2020. Mechanistic insights on the photo-sensitized chemistry of a fatty acid at the air/water interface. *Environmental Science & Technology* **54**(21): 13448–13457. DOI: <https://doi.org/10.1021/acs.est.0c03793>.
- Huang, R-J, Hoffmann, T, Ovadnevaite, J, Laaksonen, A, Kokkola, H, Xu, W, Xu, W, Cerburnis, D, Zhang, R, Seinfeld, JH, O'Dowd, C.** 2022. Heterogeneous iodine-organic chemistry fast-tracks marine new particle formation. *Proceedings of the National Academy of Sciences of the United States of America* **119**(32): e2201729119. DOI: <http://dx.doi.org/10.1073/pnas.2201729119>.
- Hughes, C, Barton, E, Hepach, H, Chance, R, Pickering, M, Hogg, K, Pommerening-Röser, A, Wadley, MR, Stevens, DP, Jickells, TD.** 2021. Oxidation of iodide to iodate by cultures of marine ammonia-oxidising bacteria. *Marine Chemistry* **234**: 104000. DOI: <http://dx.doi.org/10.1016/j.marchem.2021.104000>.
- Hulswar, S, Simó, R, Galí, M, Bell, TG, Lana, A, Inamdar, S, Halloran, PR, Manville, G, Mahajan, AS.** 2022. Third revision of the global surface seawater

- dimethyl sulfide climatology (DMS-Rev3). *Earth System Science Data* **14**: 2963–2987. DOI: <http://dx.doi.org/10.5194/essd-14-2963-2022>.
- Hultin, KAH, Krejci, R, Pinhassi, J, Gomez-Consarnau, L, Mårtensson, EM, Hagström, A, Nilsson, ED. 2011. Aerosol and bacterial emissions from Baltic Seawater. *Atmospheric Research* **99**: 1–14. DOI: <http://dx.doi.org/10.1016/j.atmosres.2010.08.018>.
- Hussherr, R, Levasseur, M, Lizotte, M, Tremblay, J-É, Mol, J, Thomas, H, Gosselin, M, Starr, M, Miller, LA, Jarníková, T, Schuback, N, Mucci, A. 2017. Impact of ocean acidification on Arctic phytoplankton blooms and dimethyl sulfide concentration under simulated ice-free and under-ice conditions. *Biogeosciences* **14**: 2407–2427. DOI: <http://dx.doi.org/10.5194/bg-14-2407-2017>.
- Ickes, L, Porter, GCE, Wagner, R, Adams, MP, Bierbauer, S, Bertram, AK, Bilde, M, Christiansen, S, Ekman, AML, Gorokhova, E, Höhler, K, Kiselev, AA, Leck, C, Möhler, O, Murray, BJ, Schiebel, T, Ullrich, R, Salter, ME. 2020. The ice-nucleating activity of Arctic sea surface microlayer samples and marine algal cultures. *Atmospheric Chemistry and Physics* **20**: 11089–11117. DOI: <http://dx.doi.org/10.5194/acp-20-11089-2020>.
- Inamdar, S, Tinel, L, Chance, R, Carpenter, LJ, Sabu, P, Chacko, R, Tripathy, SC, Kerkar, AU, Sinha, AK, Bhaskar, PV, Sarkar, A, Roy, R, Sherwen, T, Cuevas, C, Saiz-Lopez, A, Ram, K, Mahajan, AS. 2020. Estimation of reactive inorganic iodine fluxes in the Indian and Southern Ocean marine boundary layer. *Atmospheric Chemistry and Physics* **20**: 12093–12114. DOI: <http://dx.doi.org/10.5194/acp-20-12093-2020>.
- Irish, VE, Hanna, SJ, Xi, Y, Boyer, M, Polishchuk, E, Ahmed, M, Chen, J, Abbatt, JPD, Gosselin, M, Chang, R, Miller, LA, Bertram, AK. 2019. Revisiting properties and concentrations of ice-nucleating particles in the sea surface microlayer and bulk seawater in the Canadian Arctic during summer. *Atmospheric Chemistry and Physics* **19**(11): 7775–7787. DOI: <http://dx.doi.org/10.5194/acp-19-7775-2019>.
- Ishino, S, Hattori, S, Legrand, M, Chen, Q, Alexander, B, Shao, J, Huang, J, Jaeglé, L, Jourdain, B, Preunkert, S, Yamada, A, Yoshida, N, Savarino, J. 2021. Regional characteristics of atmospheric sulfate formation in East Antarctica imprinted on ^{17}O -excess signature. *Journal of Geophysical Research: Atmospheres* **126**: e2020JD033583. DOI: <http://dx.doi.org/10.1029/2020JD033583>.
- Jackson, RL, Gabric, AJ, Woodhouse, MT, Swan, HB, Jones, GB, Cropp, R, Deschaseaux, ESM. 2020. Coral reef emissions of atmospheric dimethylsulfide and the influence on marine aerosols in the southern great barrier reef, Australia. *Journal of Geophysical Research: Atmospheres* **125**(7): e2019JD031837. DOI: <http://dx.doi.org/10.1029/2019JD031837>.
- Jaeglé, L, Quinn, PK, Bates, TS, Alexander, B, Lin, J-T. 2011. Global distribution of sea salt aerosols: New constraints from in situ and remote sensing observations. *Atmospheric Chemistry and Physics* **11**(7): 3137–3157. DOI: <http://dx.doi.org/10.5194/acp-11-3137-2011>.
- Janssens, M, de Arellano, JV-G, van Heerwaarden, CC, de Roode, SR, Siebesma, AP, Glassmeier, F. 2023. Nonprecipitating shallow cumulus convection is intrinsically unstable to length scale growth. *Journal of the Atmospheric Sciences* **80**: 849–870. DOI: <http://dx.doi.org/10.1175/JAS-D-22-0111.1>.
- Järvinen, E, McCluskey, CS, Waitz, F, Schnaiter, M, Bansemer, A, Bardeen, CG, Gettelman, A, Heymsfield, A, Stith, JL, Wu, W, D'Alessandro, JJ, McFarquhar, GM, Diao, M, Finlon, JA, Hill, TCJ, Levin, EJT, Moore, KA, DeMott, PJ. 2022. Evidence for secondary ice production in Southern Ocean maritime boundary layer clouds. *Journal of Geophysical Research: Atmospheres* **127**(16): e2021JD036411. DOI: <http://dx.doi.org/10.1029/2021JD036411>.
- Jayarathne, T, Sultana, CM, Lee, C, Malfatti, F, Cox, JL, Pendergraft, MA, Moore, KA, Azam, F, Tivanski, AV, Cappa, CD, Bertram, TH, Grassian, VH, Prather, KA, Stone, EA. 2016. Enrichment of saccharides and divalent cations in sea spray aerosol during two phytoplankton blooms. *Environmental Science & Technology* **50**(21): 11511–11520. DOI: <https://doi.org/10.1021/acs.est.6b02988>.
- Jensen, JB, Nugent, AD. 2017. Condensational growth of drops formed on giant sea-salt aerosol particles. *Journal of the Atmospheric Sciences* **74**(3): 679–697. DOI: <https://doi.org/10.1175/JAS-D-15-0370.1>.
- Johnson, MT, Bell, TG. 2008. Coupling between dimethylsulfide emissions and the ocean–atmosphere exchange of ammonia. *Environmental Chemistry* **5**(4): 259–267. DOI: <http://dx.doi.org/10.1071/en08030>.
- Jokinen, T, Sipilä, M, Kontkanen, J, Vakkari, V, Tisler, P, Duplissy, E-M, Junninen, H, Kangasluoma, J, Manninen, HE, Petäjä, T, Kulmala, M, Worsnop, DR, Kirkby, J, Virkkula, A, Kerminen, V-M. 2018. Ion-induced sulfuric acid–ammonia nucleation drives particle formation in coastal Antarctica. *Science Advances* **4**(4): eaat9744. DOI: <https://doi.org/10.1126/sciadv.aat9744>.
- Kang, L, Marchand, RT, Wood, R, McCoy, IL. 2022. Coalescence scavenging drives droplet number concentration in Southern Ocean low clouds. *Geophysical Research Letters* **49**(7): e2022GL097819.
- Kilgour, DB, Novak, GA, Sauer, JS, Moore, AN, Dinisquet, J, Amiri, S, Franklin, EB, Mayer, K, Winter, M, Morris, CK, Price, T, Malfatti, F, Crocker, DR, Lee, C, Cappa, CD, Goldstein, AH, Prather, KA, Bertram, TH. 2022. Marine gas-phase sulfur emissions during an induced phytoplankton bloom. *Atmospheric Chemistry and Physics* **22**(3): 1601–1613. DOI: <https://doi.org/10.5194/acp-22-1601-2022>.
- Kim, MJ, Novak, GA, Zoerb, MC, Yang, M, Blomquist, BW, Huebert, BJ, Cappa, CD, Bertram, TH. 2017. Air-sea exchange of biogenic volatile organic

- compounds and the impact on aerosol particle size distributions. *Geophysical Research Letters* **44**(8): 3887–3896. DOI: <http://dx.doi.org/10.1002/2017GL072975>.
- Kirkby, J, Curtius, J, Almeida, J, Dunne, E, Duplissy, J, Ehrhart, S, Franchin, A, Gagné, S, Ickes, L, Kürten, A, Kupc, A, Metzger, A, Riccobono, F, Rondo, L, Schobesberger, S, Tsagkogeorgas, G, Wimmer, D, Amorim, A, Bianchi, F, Breitenlechner, M, David, A, Dommen, J, Downard, A, Ehn, M, Flanagan, RC, Haider, S, Hansel, A, Hauser, D, Jud, W, Junninen, H, Kreissl, F, Kvashin, A, Laaksonen, A, Lehtipalo, K, Lima, J, Lovejoy, ER, Makhmutov, V, Mathot, S, Mikkilä, J, Minginette, P, Mogo, S, Nieminen, T, Onnela, A, Pereira, P, Petäjä, T, Schnitzhofer, R, Seinfeld, JH, Sipilä, M, Stozhkov, Y, Stratmann, F, Tomé, A, Vanhanen, J, Viisanen, Y, Vrtala, A, Wagner, PE, Walther, H, Weingartner, E, Wex, H, Winkler, PM, Carslaw, KS, Worsnop, DR, Baltensperger, U, Kulmala, M.** 2011. Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation. *Nature* **476**(7359): 429–433. DOI: <https://doi.org/10.1038/nature10343>.
- Kirpes, RM, Bonanno, D, May, NW, Fraund, M, Barget, AJ, Moffet, RC, Ault, AP, Pratt, KA.** 2019. Winter-time Arctic sea spray aerosol composition controlled by sea ice lead microbiology. *ACS Central Science* **5**: 1760–1767. DOI: <http://dx.doi.org/10.1021/acscentsci.9b00541>.
- Klein, SA, Hall, A, Norris, JR, Pincus, R.** 2017. Low-cloud feedbacks from cloud-controlling factors: A review. *Surveys in Geophysics* **38**(6): 1307–1329. DOI: <https://doi.org/10.1007/s10712-017-9433-3>.
- Knopf, DA, Alpert, PA.** 2013. A water activity based model of heterogeneous ice nucleation kinetics for freezing of water and aqueous solution droplets. *Faraday Discussions* **165**: 513–534.
- Knopf, DA, Charnawskas, JC, Wang, P, Wong, B, Tomlin, JM, Jankowski, KA, Fraund, M, Veghte, DP, China, S, Laskin, A, Moffet, RC, Gilles, MK, Aller, JY, Marcus, MA, Raveh-Rubin, S, Wang, J.** 2022. Micro-spectroscopic and freezing characterization of ice-nucleating particles collected in the marine boundary layer in the eastern North Atlantic. *Atmospheric Chemistry and Physics* **22**(8): 5377–5398. DOI: <http://dx.doi.org/10.5194/acp-22-5377-2022>.
- Koren, I, Altaratz, O, Dagan, G.** 2015. Aerosol effect on the mobility of cloud droplets. *Environmental Research Letters* **10**(10): 104011.
- Ladino, LA, Yakobi-Hancock, JD, Kilthau, WP, Mason, RH, Si, M, Li, J, Miller, LA, Schiller, CL, Huffman, A, Aller, JY, Knopf, DA, Bertram, AK, Abbatt, JPD.** 2016. Addressing the ice nucleating abilities of marine aerosol: A combination of deposition mode laboratory and field measurements. *Atmospheric Environment* **132**: 1–10. DOI: <http://dx.doi.org/10.1016/j.atmosenv.2016.02.028>.
- Lana, A, Bell, TG, Simó, R, Vallina, SM, Ballabrera-Poy, J, Kettle, AJ, Dachs, J, Bopp, L, Saltzman, ES, Stefels, J, Johnson, JE, Liss, PS.** 2011. An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean. *Global Biogeochemical Cycles* **25**: BG1004. DOI: <http://dx.doi.org/10.1029/2010GB003850>.
- Landwehr, S, Volpi, M, Haumann, FA, Robinson, CM, Thurnherr, I, Ferracci, V, Baccarini, A, Thomas, J, Gorodetskaya, I, Tatzelt, C, Henning, S, Modini, RL, Forrer, HJ, Lin, Y, Cassar, N, Simó, R, Hassler, C, Moallemi, A, Fawcett, SE, Harris, N, Airs, R, Derkani, MH, Alberello, A, Toffoli, A, Chen, G, Rodríguez-Ros, P, Zamanillo, M, Cortés-Greus, P, Xue, L, Bolas, CG, Leonard, KC, Perez-Cruz, F, Walton, D, Schmale, J.** 2021. Exploring the coupled ocean and atmosphere system with a data science approach applied to observations from the Antarctic Circumnavigation Expedition. *Earth System Dynamics* **12**(4): 1295–1369. DOI: <http://dx.doi.org/10.5194/esd-12-1295-2021>.
- Lata, NN, Zhang, B, Schum, S, Mazzoleni, L, Brimberry, R, Marcus, MA, Cantrell, WH, Fialho, P, Mazzoleni, C, China, S.** 2021. Aerosol composition, mixing state, and phase state of free tropospheric particles and their role in ice cloud formation. *ACS Earth and Space Chemistry* **5**(12): 3499–3510. DOI: <http://dx.doi.org/10.1021/acsearthspacechem.1c00315>.
- Lauvset, SK, Tjiputra, J, Muri, H.** 2017. Climate engineering and the ocean: Effects on biogeochemistry and primary production. *Biogeosciences* **14**(24): 5675–5691. DOI: <http://dx.doi.org/10.5194/bg-14-5675-2017>.
- Law, CS, Smith, MJ, Harvey, MJ, Bell, TG, Cravigan, LT, Elliott, FC, Lawson, SJ, Lizotte, M, Marriner, A, McGregor, J, Ristovski, Z, Safi, KA, Saltzman, ES, Vaattovaara, P, Walker, CF.** 2017. Overview and preliminary results of the Surface Ocean Aerosol Production (SOAP) campaign. *Atmospheric Chemistry and Physics* **17**(22): 13645–13667. DOI: <http://dx.doi.org/10.5194/acp-17-13645-2017>.
- Lawler, MJ, Lewis, SL, Russell, LM, Quinn, PK, Bates, TS, Coffman, DJ, Upchurch, LM, Saltzman, ES.** 2020. North Atlantic marine organic aerosol characterized by novel offline thermal desorption mass spectrometry: Polysaccharides, recalcitrant material, and secondary organics. *Atmospheric Chemistry and Physics* **20**(24): 16007–16022. DOI: <https://doi.org/10.5194/acp-20-16007-2020>.
- Lawler, MJ, Saltzman, ES, Karlsson, L, Zieger, P, Salter, M, Baccarini, A, Schmale, J, Leck, C.** 2021. New insights into the composition and origins of ultra-fine aerosol in the summertime high Arctic. *Geophysical Research Letters* **48**(21): e2021GL094395. DOI: <http://dx.doi.org/10.1029/2021GL094395>.
- Lawson, SJ, Law, CS, Harvey, MJ, Bell, TG, Walker, CF, de Bruyn, WJ, Saltzman, ES.** 2020. Methanethiol, dimethyl sulfide and acetone over biologically productive waters in the southwest Pacific Ocean. *Atmospheric Chemistry and Physics* **20**(5): 3061–3078. DOI: <http://dx.doi.org/10.5194/acp-20-3061-2020>.

- Le Clainche, Y, Vézina, A, Levasseur, M, Cropp, RA, Gunson, JR, Vallina, SM, Vogt, M, Lancelot, C, Allen, JI, Archer, SD, Bopp, L, Deal, C, Elliott, S, Jin, M, Malin, G, Schoemann, V, Simó, R, Six, KD, Stefels, J. 2010. A first appraisal of prognostic ocean DMS models and prospects for their use in climate models. *Global Biogeochemical Cycles* **24**(3): GB3021. DOI: <http://dx.doi.org/10.1029/2009GB003721>.
- Lee, CP, Riva, M, Wang, D, Tomaz, S, Li, D, Perrier, S, Slowik, JG, Bourgain, F, Schmale, J, Prevot, ASH, Baltensperger, U, George, C, El Haddad, I. 2020. Online aerosol chemical characterization by extractive electrospray ionization—Ultrahigh-Resolution Mass Spectrometry (EESI-Orbitrap). *Environmental Science & Technology* **54**(7): 3871–3880. DOI: <https://doi.org/10.1021/acs.est.9b07090>.
- Lee, S-H, Gordon, H, Yu, H, Lehtipalo, K, Haley, R, Li, Y, Zhang, R. 2019. New particle formation in the atmosphere: From molecular clusters to global climate. *Journal of Geophysical Research: Atmospheres* **124**(13): 7098–7146. DOI: <https://doi.org/10.1029/2018JD029356>.
- Levin, Z, Cotton, WR eds. 2009. *Aerosol pollution impact on precipitation: A scientific review*. Dordrecht, the Netherlands: Springer.
- Listowski, C, Rojo, M, Claud, C, Delanoë, J, Rysman, J-F, Cazenave, Q, Noer, G. 2020. New insights into the vertical structure of clouds in polar lows, using radar-lidar satellite observations. *Geophysical Research Letters* **47**(17): e2020GL088785. DOI: <https://doi.org/10.1029/2020GL088785>.
- Liu, S, Liu, CC, Froyd, KD, Schill, GP, Murphy, DM, Bui, TP, Dean-Day, JM, Weinzierl, B, Dollner, M, Diskin, GS, Chen, G, Gao, R-S. 2021. Sea spray aerosol concentration modulated by sea surface temperature. *Proceedings of the National Academy of Sciences of the United States of America* **118**(9): e2020583118. DOI: <http://dx.doi.org/10.1073/pnas.2020583118>.
- Lizotte, M, Levasseur, M, Michaud, S, Scarratt, MG, Merzouk, A, Gosselin, M, Pommier, J, Rivkin, RB, Kiene, RP. 2012. Macroscale patterns of the biological cycling of dimethylsulfoniopropionate (DMSP) and dimethylsulfide (DMS) in the Northwest Atlantic. *Biogeochemistry* **110**: 183–200. DOI: <http://dx.doi.org/10.1007/s10533-011-9698-4>.
- Loh, A, Kim, D, An, JG, Choi, N, Yim, UH. 2023. Chemical characterization of sub-micron aerosols over the East Sea (Sea of Japan). *Science of The Total Environment* **856**(Part 1): 159173. DOI: <https://doi.org/10.1016/j.scitotenv.2022.159173>.
- Lowe, SJ, Partridge, DG, Davies, JF, Wilson, KR, Topping, D, Riipinen, I. 2019. Key drivers of cloud response to surface-active organics. *Nature Communications* **10**: 5214. DOI: <http://dx.doi.org/10.1038/s41467-019-12982-0>.
- Mallet, MD, Humphries, RS, Fiddes, SL, Alexander, SP, Altieri, K, Angot, H, Anilkumar, N, Bartels-Rausch, T, Creamean, J, Dall'Osto, M, Dommergue, A, Frey, M, Henning, S, Lannuzel, D, Lapere, R, Mace, GG, Mahajan, AS, McFarquhar, GM, Meiners, KM, Miljevic, B, Peeken, I, Protat, A, Schmale, J, Steiner, N, Sellegri, K, Simó, R, Thomas, JL, Willis, MD, Winton, VHL, Woodhouse, MT. 2023. Untangling the influence of Antarctic and southern Ocean life on clouds. *Elementa: Science of the Anthropocene* **11**(1): 00130. DOI: <http://dx.doi.org/10.1525/elementa.2022.00130>.
- Mansour, K, Decesari, S, Ceburnis, D, Ovadnevaite, J, Rinaldi, M. 2023. Machine learning for prediction of daily sea surface dimethylsulfide concentration and emission flux over the north Atlantic Ocean (1998–2021). *Science of the Total Environment* **871**: 162123. DOI: <http://dx.doi.org/10.1016/j.scitotenv.2023.162123>.
- Mårtensson, EM, Nilsson, ED, de Leeuw, G, Cohen, LH, Hansson, H-C. 2003. Laboratory simulations and parameterization of the primary marine aerosol production. *Journal of Geophysical Research: Atmospheres* **108**(D9): 4297. DOI: <https://doi.org/10.1029/2002JD002263>.
- Mausz, MA, Chen, Y. 2019. Microbiology and ecology of methylated amine metabolism in marine ecosystems. *Current Issues in Molecular Biology* **33**: 133–148. DOI: <https://doi.org/10.21775/cimb.033.133>.
- McCluskey, CS, Hill, TCJ, Humphries, RS, Rauker, AM, Moreau, S, Strutton, PG, Chambers, SD, Williams, AG, McRobert, I, Ward, J, Keywood, MD, Harnwell, J, Ponsonby, W, Loh, ZM, Krummel, PB, Protat, A, Kreidenweis, SM, DeMott, PJ. 2018a. Observations of ice nucleating particles over Southern Ocean waters. *Geophysical Research Letters* **45**(21). DOI: <https://doi.org/10.1029/2018GL079981>.
- McCluskey, CS, Hill, TCJ, Malfatti, F, Sultana, CM, Lee, C, Santander, MV, Beall, CM, Moore, KA, Cornwall, GC, Collins, DB, Prather, KA, Jayarathne, T, Stone, EA, Azam, F, Kreidenweis, SM, DeMott, PJ. 2017. A dynamic link between ice nucleating particles released in nascent sea spray aerosol and oceanic biological activity during two mesocosm experiments. *Journal of the Atmospheric Sciences* **74**(1): 151–166. DOI: <https://doi.org/10.1175/JAS-D-16-00871>.
- McCluskey, CS, Ovadnevaite, J, Rinaldi, M, Atkinson, J, Belosi, F, Ceburnis, D, Marullo, S, Hill, TCJ, Lohmann, U, Kanji, ZA, O'Dowd, C, Kreidenweis, SM, DeMott, PJ. 2018b. Marine and terrestrial organic ice-nucleating particles in pristine marine to continentally influenced Northeast Atlantic air masses. *Journal of Geophysical Research: Atmospheres* **123**(11): 6196–6212. DOI: <http://dx.doi.org/10.1029/2017JD028033>.
- McCoy, DT, Burrows, SM, Wood, R, Grosvenor, DP, Elliott, SM, Ma, PL, Rasch, PJ, Hartmann, DL. 2015. Natural aerosols explain seasonal and spatial patterns of Southern Ocean cloud albedo. *Science Advances* **1**(6): e1500157. DOI: <http://dx.doi.org/10.1126/sciadv.1500157>.

- McCoy, DT, Eastman, R, Hartmann, DL, Wood, R.** 2017a. The change in low cloud cover in a warmed climate inferred from AIRS, MODIS, and ERA-Interim. *Journal of Climate* **30**(10): 3609–3620. DOI: <http://dx.doi.org/10.1175/JCLI-D-15-0734.1>.
- McCoy, IL, Bretherton, CS, Wood, R, Twohy, CH, Gettelman, A, Bardeen, CG, Toohey, DW.** 2021. Influences of recent particle formation on Southern Ocean aerosol variability and low cloud properties. *Journal of Geophysical Research: Atmospheres* **126**(8): e2020JD033529. DOI: <http://dx.doi.org/10.1029/2020JD033529>.
- McCoy, IL, McCoy, DT, Wood, R, Zuidema, P, Bender, FAM.** 2023. The role of mesoscale cloud morphology in the shortwave cloud feedback. *Geophysical Research Letters* **50**(2): e2022GL101042. DOI: <http://dx.doi.org/10.1029/2022GL101042>.
- McCoy, IL, Wood, R, Fletcher, JK.** 2017b. Identifying meteorological controls on open and closed mesoscale cellular convection associated with marine cold air outbreaks. *Journal of Geophysical Research: Atmospheres* **122**(21): 11678–11702.
- McFarquhar, GM, Bretherton, CS, Marchand, R, Protat, A, DeMott, PJ, Alexander, SP, Roberts, GC, Twohy, CH, Toohey, D, Siems, S, Huang, Y, Wood, R, Rauber, RM, Lasher-Trapp, S, Jensen, J, Stith, JL, Mace, J, Um, J, Järvinen, E, Schnaiter, M, Gettelman, A, Sanchez, KJ, McCluskey, CS, Russell, LM, McCoy, IL, Atlas, RL, Bardeen, CG, Moore, KA, Hill, TCJ, Humphries, RS, Keywood, MD, Ristovski, Z, Cravigan, L, Schofield, R, Fairall, C, Mallet, MD, Kreidenweis, SM, Rainwater, B, D'Alessandro, J, Wang, Y, Wu, W, Saliba, G, Levin, EJT, Ding, S, Lang, F, Truong, SCH, Wolff, C, Haggerty, J, Harvey, MJ, Klekociuk, AR, McDonald, A.** 2021. Observations of clouds, aerosols, precipitation, and surface radiation over the Southern Ocean: An overview of CAPRICORN, MARCUS, MICRE, and SOCRATES. *Bulletin of the American Meteorological Society* **102**(4): E894–E928. DOI: <http://dx.doi.org/10.1175/BAMS-D-20-0132.1>.
- McLeod, AR, Brand, T, Campbell, CN, Davidson, K, Hatton, AD.** 2021. Ultraviolet radiation drives emission of climate-relevant gases from marine phytoplankton. *Journal of Geophysical Research: Biogeosciences* **126**(9): e2021JG006345.
- McNabb, BJ, Tortell, PD.** 2022. Improved prediction of dimethyl sulfide (DMS) distributions in the northeast subarctic Pacific using machine-learning algorithms. *Biogeosciences* **19**(6): 1705–1721. DOI: <http://dx.doi.org/10.5194/bg-19-1705-2022>.
- McNabb, BJ, Tortell, PD.** 2023. Oceanographic controls on Southern Ocean dimethyl sulfide distributions revealed by machine learning algorithms. *Limnology and Oceanography* **68**(3): 616–630. DOI: <http://dx.doi.org/10.1002/lno.12298>.
- McParland, EL, Levine, NM.** 2019. The role of differential DMSP production and community composition in predicting variability of global surface DMSP concentrations. *Limnology and Oceanography* **64**(2): 757–773.
- Meskhidze, N, Sabolis, A, Reed, R, Kamykowski, D.** 2015. Quantifying environmental stress-induced emissions of algal isoprene and monoterpenes using laboratory measurements. *Biogeosciences* **12**(3): 637–651. DOI: <http://dx.doi.org/10.5194/bg-12-637-2015>.
- Moallemi, A, Landwehr, S, Robinson, C, Simó, R, Zamanillo, M, Chen, G, Baccarini, A, Schnaiter, M, Henning, S, Modini, RL, Gysel-Beer, M, Schmale, J.** 2021. Sources, occurrence and characteristics of fluorescent biological aerosol particles measured over the pristine Southern Ocean. *Journal of Geophysical Research: Atmospheres* **126**(11): e2021JD034811. DOI: <http://dx.doi.org/10.1029/2021JD034811>.
- Modini, RL, Frossard, AA, Ahlm, L, Russell, LM, Corrigan, CE, Roberts, GC, Hawkins, LN, Schroder, JC, Bertram, AK, Zhao, R, Lee, AKY, Abbatt, JPD, Lin, J, Nenes, A, Wang, Z, Wonaschütz, A, Sorooshian, A, Noone, KJ, Jonsson, H, Seinfeld, JH, Toom, D, Macdonald, AM, Leaitch, WR.** 2015. Primary marine aerosol-cloud interactions off the coast of California. *Journal of Geophysical Research: Atmospheres* **120**(9): 4282–4303. DOI: <http://dx.doi.org/10.1002/2014JD022963>.
- Modini, RL, Russell, LM, Deane, GB, Stokes, MD.** 2013. Effect of soluble surfactant on bubble persistence and bubble-produced aerosol particles. *Journal of Geophysical Research: Atmospheres* **118**(3): 1388–1400. DOI: <http://dx.doi.org/10.1002/jgrd.50186>.
- Moore, ER, Davie-Martin, CL, Giovannoni, SJ, Halsey, KH.** 2020. Pelagibacter metabolism of diatom-derived volatile organic compounds imposes an energetic tax on photosynthetic carbon fixation. *Environmental Microbiology* **22**(5): 1720–1733.
- Moran, MA, Durham, BP.** 2019. Sulfur metabolites in the pelagic ocean. *Nature Reviews Microbiology* **17**: 665–678. DOI: <http://dx.doi.org/10.1038/s41579-019-0250-1>.
- Moschos, V, Dzepina, K, Bhattu, D, Lamkaddam, H, Casotto, R, Daellenbach, KR, Canonaco, F, Rai, P, Aas, W, Becagli, S, Calzolari, G, Eleftheriadis, K, Moffett, CE, Schnelle-Kreis, J, Severi, M, Sharma, S, Skov, H, Vestenius, M, Zhang, W, Hakola, H, Hellén, H, Huang, L, Jaffrezo, J-L, Massling, A, Nøjgaard, JK, Petäjä, T, Popovicheva, O, Sheesley, RJ, Traversi, R, Yttri, KE, Schmale, J, Prévôt, ASH, Baltensperger, U, El Haddad, I.** 2022a. Equal abundance of summertime natural and wintertime anthropogenic Arctic organic aerosols. *Nature Geoscience* **15**(3): 196–202. DOI: <https://doi.org/10.1038/s41561-021-00891-1>.
- Moschos, V, Schmale, J, Aas, W, Becagli, S, Calzolari, G, Eleftheriadis, K, Moffett, CE, Schnelle-Kreis, J, Severi, M, Sharma, S, Skov, H, Vestenius, M, Zhang, W, Hakola, H, Hellén, H, Huang, L,**

- Jaffrezo, J-L, Massling, A, Nøjgaard, JK, Petäjä, T, Popovicheva, O, Sheesley, RJ, Traversi, R, Yttri, KE, Prévôt, ASH, Baltensperger, U, El Haddad, I. 2022b. Elucidating the present-day chemical composition, seasonality and source regions of climate-relevant aerosols across the Arctic land surface. *Environmental Research Letters* **17**(3): 034032. DOI: <https://doi.org/10.1088/1748-9326/ac444b>.
- Mulcahy, JP, Jones, C, Sellar, A, Johnson, B, Boutle, IA, Jones, A, Andrews, T, Rumbold, ST, Mollard, J, Bellouin, N, Johnson, CE, Williams, KD, Grosvenor, DP, McCoy, DT. 2018. Improved aerosol processes and effective radiative forcing in HadGEM3 and UKESM1. *Journal of Advances in Modeling Earth Systems* **10**(11): 2786–2805. DOI: <https://doi.org/10.1029/2018MS001464>.
- Mungall, EL, Abbatt, JPD, Wentzell, JJB, Lee, AKY, Thomas, JL, Blais, M, Gosselin, M, Miller, LA, Papakyriakou, T, Willis, MD, Liggio, J. 2017. Microlayer source of oxygenated volatile organic compounds in the summertime marine Arctic boundary layer. *Proceedings of the National Academy of Sciences of the United States of America* **114**(24): 6203–6208. DOI: <http://dx.doi.org/10.1073/pnas.1620571114>.
- Myers, KF, Doran, PT, Cook, J, Kotcher, JE, Myers, TA. 2021a. Consensus revisited: Quantifying scientific agreement on climate change and climate expertise among Earth scientists 10 years later. *Environmental Research Letters* **16**(10): 104030.
- Myers, TA, Norris, JR. 2016. Reducing the uncertainty in subtropical cloud feedback. *Geophysical Research Letters* **43**(5): 2144–2148 DOI: <http://dx.doi.org/10.1002/2015GL067416>.
- Myers, TA, Scott, RC, Zelinka, MD, Klein, SA, Norris, JR, Caldwell, PM. 2021b. Observational constraints on low cloud feedback reduce uncertainty of climate sensitivity. *Nature Climate Change* **11**: 501–507. DOI: <https://doi.org/10.1038/s41558-021-01039-0>.
- Narenpitak, P, Kazil, J, Yamaguchi, T, Quinn, P, Feingold, G. 2021. From sugar to flowers: A transition of shallow cumulus organization during ATOMIC. *Journal of Advances in Modeling Earth Systems* **13**(10): e2021MS002619. DOI: <http://dx.doi.org/10.1029/2021ms002619>.
- National Academies of Sciences, Engineering, Medicine. 2018. *Thriving on our changing planet: A decadal strategy for earth observation from space*. Washington, DC: The National Academies Press. DOI: <http://dx.doi.org/10.17226/24938>.
- Neumann, D, Matthias, V, Bieser, J, Aulinger, A, Quante, M. 2016. A comparison of sea salt emission parameterizations in northwestern Europe using a chemistry transport model setup. *Atmospheric Chemistry and Physics* **16**(15): 9905–9933. DOI: <http://dx.doi.org/10.5194/acp-16-9905-2016>.
- Norris, JR, Allen, RJ, Evan, AT, Zelinka, MD, O'Dell, CW, Klein, SA. 2016. Evidence for climate change in the satellite cloud record. *Nature* **536**: 72–75. DOI: <http://dx.doi.org/10.1038/nature18273>.
- Novak, GA, Fite, CH, Holmes, CD, Veres, PR, Neuman, JA, Faloona, I, Thornton, JA, Wolfe, GM, Vermeuel, MP, Jernigan, CM, Peischl, J, Ryerson, TB, Thompson, CR, Bourgeois, I, Warneke, C, Gkatzelis, GI, Coggon, MM, Sekimoto, K, Bui, TP, Dean-Day, J, Diskin, GS, DiGangi, JP, Nowak, JB, Moore, RH, Wiggins, EB, Winstead, EL, Robinson, C, Thornhill, KL, Sanchez, KJ, Hall, SR, Ullmann, K, Dollner, M, Weinzierl, B, Blake, DR, Bertram, TH. 2021. Rapid cloud removal of dimethyl sulfide oxidation products limits SO₂ and cloud condensation nuclei production in the marine atmosphere. *Proceedings of the National Academy of Sciences of the United States of America* **118**(42): e2110472118. DOI: <http://dx.doi.org/10.1073/pnas.2110472118>.
- Novak, GA, Kilgour, DB, Jernigan, CM, Vermeuel, MP, Bertram, TH. 2022. Oceanic emissions of dimethyl sulfide and methanethiol and their contribution to sulfur dioxide production in the marine atmosphere. *Atmospheric Chemistry and Physics* **22**(9): 6309–6325. DOI: <http://dx.doi.org/10.5194/acp-22-6309-2022>.
- O'Dowd, C, Ceburnis, D, Ovadnevaite, J, Bialek, J, Stengel, DB, Zacharias, M, Nitschke, U, Connan, S, Rinaldi, M, Fuzzi, S, Decesari, S, Facchini, MC, Marullo, S, Santolieri, R, Dell'Anno, A, Corinaldesi, C, Tangherlini, M, Danovaro, R. 2015. Connecting marine productivity to sea-spray via nanoscale biological processes: Phytoplankton Dance or Death Disco? *Scientific Reports* **5**: 14883. DOI: <https://doi.org/10.1038/srep14883>.
- O'Dowd, C, Monahan, C, Dall'Osto, M. 2010. On the occurrence of open ocean particle production and growth events. *Geophysical Research Letters* **37**(19): L19805.
- O'Dowd, CD, Facchini, MC, Cavalli, F, Ceburnis, D, Mircea, M, Decesari, S, Fuzzi, S, Yoon, YJ, Putaud, J-P. 2004. Biogenically driven organic contribution to marine aerosol. *Nature* **431**(7009): 676–680. DOI: <https://doi.org/10.1038/nature02959>.
- Oppo, C, Bellandi, S, Degli Innocenti, N, Stortini, AM, Loglio, G, Schiavuta, E, Cini, R. 1999. Surfactant components of marine organic matter as agents for biogeochemical fractionation and pollutant transport via marine aerosols. *Marine Chemistry* **63**(3–4): 235–253. DOI: [http://dx.doi.org/10.1016/S0304-4203\(98\)00065-6](http://dx.doi.org/10.1016/S0304-4203(98)00065-6).
- Ovadnevaite, J, Ceburnis, D, Martucci, G, Bialek, J, Monahan, C, Rinaldi, M, Facchini, MC, Berresheim, H, Worsnop, DR, O'Dowd, C. 2011. Primary marine organic aerosol: A dichotomy of low hygroscopicity and high CCN activity. *Geophysical Research Letters* **38**(21): L21806. DOI: <http://dx.doi.org/10.1029/2011gl048869>.
- Ovadnevaite, J, Manders, A, de Leeuw, G, Ceburnis, D, Monahan, C, Partanen, AI, Korhonen, H, O'Dowd, CD. 2014. A sea spray aerosol flux parameterization encapsulating wave state. *Atmospheric*

- Chemistry and Physics* **14**(4): 1837–1852. DOI: <http://dx.doi.org/10.5194/acp-14-1837-2014>.
- Ovadnevaite, J, Zuend, A, Laaksonen, A, Sanchez, KJ, Roberts, G, Ceburnis, D, Decesari, S, Rinaldi, M, Hodas, N, Facchini, MC, Seinfeld, JH, O'Dowd, C.** 2017. Surface tension prevails over solute effect in organic-influenced cloud droplet activation. *Nature* **546**: 637–641. DOI: <http://dx.doi.org/10.1038/nature22806>.
- Partanen, AI, Dunne, EM, Bergman, T, Laakso, A, Kokkola, H, Ovadnevaite, J, Sogacheva, L, Baisnée, D, Sciare, J, Manders, A, O'Dowd, C, de Leeuw, G, Korhonen, H.** 2014. Global modelling of direct and indirect effects of sea spray aerosol using a source function encapsulating wave state. *Atmospheric Chemistry and Physics* **14**(21): 11731–11752. DOI: <http://dx.doi.org/10.5194/acp-14-11731-2014>.
- Peltola, M, Rose, C, Trueblood, JV, Gray, S, Harvey, M, Sellegri, K.** 2022. New particle formation in coastal New Zealand with a focus on open-ocean air masses. *Atmospheric Chemistry and Physics* **22**(9): 6231–6254. DOI: <http://dx.doi.org/10.5194/acp-22-6231-2022>.
- Penezić, A, Wang, X, Perrier, S, George, C, Frka, S.** 2023. Interfacial photochemistry of marine diatom lipids: Abiotic production of volatile organic compounds and new particle formation. *Chemosphere* **313**: 137510. DOI: <https://doi.org/10.1016/j.chemosphere.2022.137510>.
- Prados-Roman, C, Cuevas, CA, Hay, T, Fernandez, RP, Mahajan, AS, Royer, SJ, Galí, M, Simó, R, Dachs, J, Großmann, K, Kinnison, DE, Lamarque, JF, Saiz-Lopez, A.** 2015. Iodine oxide in the global marine boundary layer. *Atmospheric Chemistry and Physics* **15**(2): 583–593. DOI: <http://dx.doi.org/10.5194/acp-15-583-2015>.
- Prank, M, Tonttila, J, Ahola, J, Kokkola, H, Kühn, T, Romakkaniemi, S, Raatikainen, T.** 2022. Impacts of marine organic emissions on low-level stratiform clouds—A large eddy simulator study. *Atmospheric Chemistry and Physics* **22**(16): 10971–10992. DOI: <http://dx.doi.org/10.5194/acp-22-10971-2022>.
- Prather, KA, Bertram, TH, Grassian, VH, Deane, GB, Dale Stokes, M, DeMott, PJ, Aluwihare, LI, Palenik, BP, Azam, F, Seinfeld, JH, Moffet, RC, Molina, MJ, Cappa, CD, Geiger, FM, Roberts, GC, Russell, LM, Ault, AP, Baltrusaitis, J, Collins, DB, Corrigan, CE, Cuadra-Rodriguez, LA, Ebben, CJ, Forestieri, SD, Guasco, TL, Hersey, SP, Kim, MJ, Lambert, WF, Modini, RL, Mui, W, Pedler, BE, Ruppel, MJ, Ryder, OS, Schoepp, NG, Sullivan, RC, Zhao, D.** 2013. Bringing the ocean into the laboratory to probe the chemical complexity of sea spray aerosol. *Proceedings of the National Academy of Sciences* **110**(19): 7550–7555. DOI: <https://doi.org/10.1073/pnas.1300262110>.
- Pruppacher, HR, Klett, JD.** 1997. *Microphysics of clouds and precipitation* (2nd ed.). Dordrecht, the Netherlands: Kluwer Academic: 954.
- Quinn, PK, Coffman, DJ, Johnson, JE, Upchurch, LM, Bates, TS.** 2017. Small fraction of marine cloud condensation nuclei made up of sea spray aerosol. *Nature Geoscience* **10**: 674–679. DOI: <http://dx.doi.org/10.1038/ngeo3003>.
- Quinn, PK, Collins, DB, Grassian, VH, Prather, KA, Bates, TS.** 2015. Chemistry and related properties of freshly emitted sea spray aerosol. *Chemical Reviews* **115**(10): 4383–4399. DOI: <https://doi.org/10.1021/cr500713g>.
- Radoman, N, Christiansen, S, Johansson, JH, Hawkes, JA, Bilde, M, Cousins, IT, Salter, ME.** 2022. Probing the impact of a phytoplankton bloom on the chemistry of nascent sea spray aerosol using high-resolution mass spectrometry. *Environmental Science: Atmospheres* **2**: 1152–1169. DOI: <https://doi.org/10.1039/D2EA00028H>.
- Rapf, RJ, Perkins, RJ, Dooley, MR, Kroll, JA, Carpenter, BK, Vaida, V.** 2018. Environmental processing of lipids driven by aqueous photochemistry of α -keto acids. *ACS Central Science* **4**(5): 624–630. DOI: <http://dx.doi.org/10.1021/acscentsci.8b00124>.
- Regayre, LA, Schmale, J, Johnson, JS, Tatzelt, C, Baccarini, A, Henning, S, Yoshioka, M, Stratmann, F, Gysel-Beer, M, Grosvenor, DP, Carslaw, KS.** 2020. The value of remote marine aerosol measurements for constraining radiative forcing uncertainty. *Atmospheric Chemistry and Physics* **20**(16): 10063–10072. DOI: <http://dx.doi.org/10.5194/acp-20-10063-2020>.
- Revell, LE, Kremser, S, Hartery, S, Harvey, M, Mulcahy, JP, Williams, J, Morgenstern, O, McDonald, AJ, Varma, V, Bird, L, Schuddeboom, A.** 2019. The sensitivity of Southern Ocean aerosols and cloud microphysics to sea spray and sulfate aerosol production in the HadGEM3-GA7.1 chemistry–climate model. *Atmospheric Chemistry and Physics* **19**(24): 15447–15466. DOI: <http://dx.doi.org/10.5194/acp-19-15447-2019>.
- Rocco, M, Dunne, E, Peltola, M, Barr, N, Williams, J, Colomb, A, Safi, K, Saint-Macary, A, Marriner, A, Deppeler, S, Harnwell, J, Law, C, Sellegri, K.** 2021. Oceanic phytoplankton are a potentially important source of benzenoids to the remote marine atmosphere. *Nature Communications Earth & Environment* **2**(1): 175.
- Rocco, M, Dunne, E, Saint-Macary, A, Peltola, M, Barthelmeß, T, Barr, N, Safi, K, Marriner, A, Deppler, S, Harnwell, J, Engel, A, Colomb, A, Saiz-Lopez, A, Harvey, M, Law, CS, Sellegri, K.** 2023. Air-sea fluxes of dimethyl sulphide and methanethiol in the South-West Pacific. *EGU sphere* [preprint]. DOI: <http://dx.doi.org/10.5194/egusphere-2023-516>.
- Rose, C, Zha, Q, Dada, L, Yan, C, Lehtipalo, K, Junninen, H, Buenrostro Mazon, S, Jokinen, T, Sarnela, N, Sipilä, M, Petäjä, T, Kerminen, V-M, Bianchi, F, Kulmala, M.** 2018. Observations of biogenic ion-induced cluster formation in the atmosphere.

- Science Advances* **4**(4): eaar5218. DOI: <https://doi.org/10.1126/sciadv.aar5218>.
- Rosenfeld, D, Andreae, MO, Asmi, A, Chin, M, de Leeuw, G, Donovan, DP, Kahn, R, Kivekas, N, Kulmala, M, Lau, W, Schmidt, KS, Suni, T, Wagner, T, Wild, M, Quaas, J. 2014. Global observations of aerosol-cloud-precipitation-climate interactions. *Reviews of Geophysics* **52**(4): 750–808.
- Rossignol, S, Tinel, L, Bianco, A, Passananti, M, Briggante, M, Donaldson, DJ, George, C. 2016. Atmospheric photochemistry at a fatty acid-coated air-water interface. *Science* **353**(6300): 699–702. DOI: <http://dx.doi.org/10.1126/science.aaf3617>.
- Royer, SJ, Galí, M, Mahajan, AS, Ross, ON, Pérez, GL, Saltzman, ES, Simó, R. 2016. A high-resolution time-depth view of dimethylsulphide cycling in the surface sea. *Scientific Reports* **6**: 32325. DOI: <http://dx.doi.org/10.1038/srep32325>.
- Ruiz-González, C, Simó, R, Sommaruga, R, Gasol, JM. 2013. Away from darkness: A review on the effects of solar radiation on heterotrophic bacterioplankton activity. *Frontiers in Microbiology*. DOI: <http://dx.doi.org/10.3389/fmicb.2013.00131>.
- Russell, LM, Moore, RH, Burrows, SM, Quinn, PK. 2023. Ocean flux of salt, sulfate, and organic components to atmospheric aerosol. *Earth-Science Reviews* **239**(4): 104364. DOI: <http://dx.doi.org/10.1016/j.earscirev.2023.104364>.
- Saha, M, Fink, P. 2022. Algal volatiles—The overlooked chemical language of aquatic primary producers. *Biological Reviews* **97**(6): 2162–2173.
- Saliba, G, Chen, CL, Lewis, S, Russell, LM, Rivellini, LH, Lee, AKY, Quinn, PK, Bates, TS, Haëntjens, N, Boss, ES, Karp-Boss, L, Baetge, N, Carlson, CA, Behrenfeld, MJ. 2019. Factors driving the seasonal and hourly variability of sea-spray aerosol number in the North Atlantic. *Marine Sciences Faculty Scholarship* 223. DOI: https://digitalcommons.library.umaine.edu/sms_facpub/223.
- Salter, ME, Nilsson, ED, Butcher, A, Bilde, M. 2014. On the seawater temperature dependence of the sea spray aerosol generated by a continuous plunging jet. *Journal of Geophysical Research: Atmospheres* **119**(14): 9052–9072. DOI: <https://doi.org/10.1002/2013JD021376>.
- Salter, ME, Zieger, P, Acosta Navarro, JC, Grythe, H, Kirkevåg, A, Rosati, B, Riipinen, I, Nilsson, ED. 2015. An empirically derived inorganic sea spray source function incorporating sea surface temperature. *Atmospheric Chemistry and Physics* **15**: 11047–11066. DOI: <https://doi.org/10.5194/acp-15-11047-2015>.
- Sanchez, KJ, Chen, CL, Russell, LM, Betha, R, Liu, J, Price, DJ, Massoli, P, Ziemba, LD, Crosbie, EC, Moore, RH, Müller, M, Schiller, SA, Wisthaler, A, Lee, AKY, Quinn, PK, Bates, TS, Porter, J, Bell, TG, Saltzman, ES, Vaillancourt, RD, Behrenfeld, MJ. 2018. Substantial seasonal contribution of observed biogenic sulfate particles to cloud condensation nuclei. *Scientific Reports* **8**: 3235. DOI: <http://dx.doi.org/10.1038/s41598-018-21590-9>.
- Sanchez, KJ, Roberts, GC, Saliba, G, Russell, LM, Twohy, C, Reeves, MJ, Humphries, RS, Keywood, MD, Ward, JP, McRobert, LM. 2021. Measurement report: Cloud processes and the transport of biological emissions affect Southern Ocean particle and cloud condensation nuclei concentrations. *Atmospheric Chemistry and Physics* **21**(5): 3427–3446.
- Santander, MV, Schiffer, JM, Lee, C, Axson, JL, Tauber, MJ, Prather, KA. 2022. Factors controlling the transfer of biogenic organic species from seawater to sea spray aerosol. *Scientific Reports* **12**(1): 1–10. DOI: <https://doi.org/10.1038/s41598-022-07335-9>.
- Schmale, J, Baccarini, A, Thurnherr, I, Henning, S, Efraim, A, Regayre, L, Bolas, C, Hartmann, M, Welti, A, Lehtipalo, K, Aemisegger, F, Tatzelt, C, Landwehr, S, Modini, R, Tummon, F, Johnson, JS, Harri, N, Schnaiterale, M, Toffoli, S, Derkani, M, Bukowiecki, N, Stratmann, F, Dommen, J, Baltensperger, U, Wernli, H, Rosenfeld, D, Gysel-Beer, M, Carslaw, KS. 2019. Overview of the Antarctic Circumnavigation Expedition: Study of Preindustrial-like Aerosols and Their Climate Effects (ACE-SPACE). *Bulletin of the American Meteorological Society* **100**(11): 2260–2283. DOI: <http://dx.doi.org/10.1175/BAMS-D-18-0187.1>.
- Schnell, RC. 1975. Ice nuclei produced by laboratory cultured marine phytoplankton. *Geophysical Research Letters* **2**(11): 500–502. DOI: <http://dx.doi.org/10.1029/GL002i011p00500>.
- Schwier, AN, Rose, C, Asmi, E, Ebling, AM, Landing, WM, Marro, S, Pedrotti, LM, Sallon, A, Iuculano, F, Agusti, S, Tsiola, A, Pitta, P, Louis, J, Guieu, C, Gazeau, F, Sellegri, K. 2015. Primary marine aerosol emissions from the Mediterranean Sea during pre-bloom and oligotrophic conditions: Correlations to seawater chlorophyll a from a mesocosm study. *Atmospheric Chemistry and Physics* **15**(14): 7961–7976. DOI: <http://dx.doi.org/10.5194/acp-15-7961-2015>.
- Schwier, AN, Sellegri, K, Mas, S, Charrière, B, Pey, J, Rose, C, Temime-Roussel, B, Parin, D, Jaffrezo, JL, Picard, D, Sempéré, R, Marchand, N, D'Anna, B. 2017. Primary marine aerosol physical flux and chemical composition during a nutrient enrichment experiment in mesocosms in the Mediterranean Sea. *Atmospheric Chemistry and Physics* **17**(23): 14645–14660. DOI: <http://dx.doi.org/10.5194/acp-17-14645-2017>.
- Schwinger, J, Tjiputra, J, Goris, N, Six, KD, Kirkevåg, A, Seland, O, Heinze, C, Ilyina, T. 2017. Amplification of global warming through pH dependence of DMS production simulated with a fully coupled earth system model. *Biogeosciences* **14**(15): 3633–3648.
- Scott, RC, Myers, TA, Norris, JR, Zelinka, MD, Klein, SA, Sun, M, Doelling, DR. 2020. Observed sensitivity of low-cloud radiative effects to meteorological perturbations over the global oceans. *Journal of Climate*

- 33(17): 7717–7734. DOI: <https://doi.org/10.1175/JCLI-D-19-1028.1>.
- Séférian, R, Nabat, P, Michou, M, Saint-Martin, D, Voltaire, A, Colin, J, Decharme, B, Delire, C, Berthet, S, Chevallier, M, Sénési, S, Franchisteguy, L, Vial, J, Mallet, M, Joetzjer, E, Geoffroy, O, Guérémy, JF, Moine, MP, Msadek, R, Ribes, A, Rocher, M, Roehrig, R, Salas-y-Méla, D, Sanchez, E, Terray, L, Valcke, S, Waldman, R, Aumont, O, Bopp, L, Deshayes, J, Éthé, C, Madec, G. 2019. Evaluation of CNRM earth system model, CNRM-ESM2-1: Role of earth system processes in present-day and future climate. *Journal of Advances in Modeling Earth Systems* **11**(12): 4182–4227. DOI: <http://dx.doi.org/10.1029/2019MS001791>.
- Seinfeld, JH, Pandis, SN. 2006. *Atmospheric chemistry and physics: From air pollution to climate change* (2nd ed.). New York, NY: John Wiley & Sons.
- Sellegri, K, Barthelmeß, T, Trueblood, J, Cristi, A, Freney, E, Rose, C, Barr, N, Harvey, M, Safi, K, Deppler, S, Thompson, K, Dillon, W, Engel, A, Law, C. 2023. Quantified effect of seawater biogeochemistry on the temperature dependence of sea spray aerosol fluxes. *Atmospheric Chemistry and Physics* [preprint]. DOI: <http://dx.doi.org/10.5194/acp-2022-790>.
- Sellegri, K, Nicosia, A, Freney, E, Uitz, J, Thyssen, M, Grégori, G, Engel, A, Zäncker, B, Haëntjens, N, Mas, S, Picard, D, Saint-Macary, A, Peltola, M, Rose, C, Trueblood, J, Lefevre, D, D'Anna, B, Desboeuf, K, Meskhidze, N, Guieu, C, Law, CS. 2021. Surface ocean microbiota determine cloud precursors. *Scientific Reports* **11**: 281. DOI: <http://dx.doi.org/10.1038/s41598-020-78097-5>.
- Sellegri, K, Pey, J, Rose, C, Culot, A, DeWitt, HL, Mas, S, Schwier, AN, Temime-Roussel, B, Charriere, B, Saiz-Lopez, A, Mahajan, AS, Parin, D, Kukui, A, Sempere, R, D'anna, B, Marchand, N. 2016. Evidence of atmospheric nanoparticle formation from emissions of marine microorganisms. *Geophysical Research Letters* **43**(12): 6596–6603. DOI: <http://dx.doi.org/10.1002/2016GL069389>.
- Shaw, SL, Gantt, B, Meskhidze, N. 2010. Production and emissions of marine isoprene and monoterpenes: A review. *Advances in Meteorology* **2010**: 1–24. DOI: <http://dx.doi.org/10.1155/2010/408696>.
- Sherwood, SC, Webb, MJ, Annan, JD, Armour, KC, Forster, PM, Hargreaves, JC, Hegerl, G, Klein, SA, Marvel, KD, Rohling, EJ, Watanabe, M, Andrews, T, Braconnot, P, Bretherton, CS, Foster, GL, Hausfather, Z, von der Heydt, AS, Knutti, R, Mauritsen, T, Norris, JR, Proistosescu, C, Rugenstein, M, Schmidt, GA, Tokarska, KB, Zelinka, MD. 2020. An assessment of Earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics* **58**(4): e2019RG000678. DOI: <https://doi.org/10.1029/2019RG000678>.
- Shingler, T, Crosbie, E, Ortega, A, Shiraiwa, M, Zuend, A, Beyersdorf, A, Ziemba, L, Anderson, B, Thornhill, L, Perring, AE, Schwarz, JP, Campazano-Jost, P, Day, DA, Jimenez, JL, Hair, JW, Mikoviny, T, Wisthaler, A, Sorooshian, A. 2016. Airborne characterization of subsaturated aerosol hygroscopicity and dry refractive index from the surface to 6.5 km during the SEAC⁴RS campaign. *Journal of Geophysical Research: Atmospheres* **121**(8): 4188–4210. DOI: <http://dx.doi.org/10.1002/2015JD024498>.
- Shrestha, M, Luo, M, Li, Y, Xiang, B, Xiong, W, Grassian, VH. 2018. Let there be light: Stability of palmitic acid monolayers at the air/salt water interface in the presence and absence of simulated solar light and a photosensitizer. *Chemical Science* **9**(26): 5716–5723.
- Siegel, K, Karlsson, L, Zieger, P, Baccarini, A, Schmale, J, Lawler, M, Salter, M, Leck, C, Ekman, AML, Riipinen, I, Mohr, C. 2021. Insights into the molecular composition of semi-volatile aerosols in the summertime central Arctic Ocean using FIGAERO-CIMS. *Environmental Science: Atmospheres* **1**: 161–175. DOI: <https://doi.org/10.1039/D0EA00023J>.
- Sipilä, M, Sarnela, N, Jokinen, T, Henschel, H, Junninen, H, Kontkanen, J, Richters, S, Kangasluoma, J, Franchin, A, Peräkylä, O, Rissanen, MP, Ehn, M, Vehkamäki, H, Kurten, T, Berndt, T, Petäjä, T, Worsnop, D, Ceburnis, D, Kerminen, VM, Kulmala, M, O'Dowd, C. 2016. Molecular-scale evidence of aerosol particle formation via sequential addition of HIO₃. *Nature* **537**(7621): 532–534. DOI: <http://dx.doi.org/10.1038/nature19314>.
- Six, KD, Kloster, S, Ilyina, T, Archer, SD, Zhang, K, Maier-Reimer, E. 2013. Global warming amplified by reduced Sulphur fluxes as a result of ocean acidification. *Nature Climate Change* **3**: 975–978.
- Spielmeyer, A, Gebser, B, Pohnert, G. 2011. Investigations of the uptake of dimethylsulfoniopropionate by phytoplankton. *Chembiochem: A European Journal of Chemical Biology* **12**(15): 2276–2279.
- Spiese, CE, Kieber, DJ, Nomura, CT, Kiene, RP. 2009. Reduction of dimethylsulfoxide to dimethylsulfide by marine phytoplankton. *Limnology and Oceanography* **54**(2): 560–570.
- Stefels, J, Steinke, M, Turner, S, Malin, G, Belviso, S. 2007. Environmental constraints on the production and removal of the climatically active gas dimethylsulphide (DMS) and implications for ecosystem modelling. *Biogeochemistry* **83**: 245–275. DOI: <https://doi.org/10.1007/s10533-007-9091-5>.
- Stevens, B, Feingold, G. 2009. Untangling aerosol effects on clouds and precipitation in a buffered system. *Nature* **461**: 607–613. DOI: <http://dx.doi.org/10.1038/nature08281>.
- Stirchak, LT, Abis, L, Kalalian, C, George, C, Donaldson, DJ. 2021. Differences in photosensitized release of VOCs from illuminated seawater versus freshwater surfaces. *ACS Earth and Space Chemistry* **5**(9): 2233–2242. DOI: <http://dx.doi.org/10.1021/acsearthspacechem.1c00063>.
- Struthers, H, Ekman, AML, Glantz, P, Iversen, T, Kirkevåg, A, Seland, Ø, Mårtensson, EM, Noone, K, Nilsson, ED. 2013. Climate-induced changes in sea salt aerosol number emissions: 1870 to 2100.

- Journal of Geophysical Research: Atmospheres* **118**(2): 670–682. DOI: <http://dx.doi.org/10.1002/jgrd.50129>.
- Sun, J, Mausz, MA, Chen, Y, Giovannoni, SJ. 2019. Microbial trimethylamine metabolism in marine environments. *Environmental Microbiology* **21**(2). DOI: <https://doi.org/10.1111/1462-2920.14461>.
- Tatzelt, C, Henning, S, Welti, A, Baccarini, A, Hartmann, M, Gysel-Beer, M, van Pinxteren, M, Modini, RL, Schmale, J, Stratmann, F. 2022. Circum-Antarctic abundance and properties of CCN and INPs. *Atmospheric Chemistry and Physics* **22**(14): 9721–9745. DOI: <http://dx.doi.org/10.5194/acp-22-9721-2022>.
- Tegen, I, Schepanski, K. 2018. Climate feedback on aerosol emission and atmospheric concentrations. *Current Climate Change Reports* **4**: 1–10 DOI: <http://dx.doi.org/10.1007/s40641-018-0086-1>.
- Teng, ZJ, Qin, QL, Zhang, W, Li, J, Fu, HH, Wang, P, Lan, M, Luo, G, He, J, McMin, A, Wang, M, Chen, XL, Zhang, YZ, Chen, Y, Li, CY. 2021. Biogeographic traits of dimethyl sulfide and dimethylsulfoniopropionate cycling in polar oceans. *Microbiome* **9**: 207. DOI: <http://dx.doi.org/10.1186/s40168-021-01153-3>.
- Tinel, L, Abbatt, JMD, Saltzman, E, Engel, A, Fernandez, R, Li, Q, Mahajan, AS, Nicewonger, M, Novak, G, Saiz-Lopez, A, Schneider, S, Wang, S. 2023. Impacts of ocean biogeochemistry on atmospheric chemistry. *Elementa: Science of the Anthropocene* **11**(1): 00032.
- Tinel, L, Rossignol, S, Bianco, A, Passananti, M, Perrier, S, Wang, X, Brigante, M, Donaldson, DJ, George, C. 2016. Mechanistic insights on the photosensitized chemistry of a fatty acid at the air/water interface. *Environmental Science & Technology* **50**(20): 11041–11048. DOI: <http://dx.doi.org/10.1021/acs.est.6b03165>.
- Trenberth, KE, Fasullo, JT, Kiehl, J. 2009. Earth's global energy budget. *Bulletin of the American Meteorological Society* **90**(3): 311–323.
- Trueblood, JV, Nicosia, A, Engel, A, Zäncker, B, Rinaldi, M, Freney, E, Thyssen, M, Obernosterer, I, Dinassquet, J, Belosi, F, Tovar-Sánchez, A, Rodriguez-Romero, A, Santachiara, G, Guieu, C, Sellegri, K. 2021. A two-component parameterization of marine ice-nucleating particles based on seawater biology and sea spray aerosol measurements in the Mediterranean Sea. *Atmospheric Chemistry and Physics* **21**(6): 4659–4676. DOI: <https://doi.org/10.5194/acp-21-4659-2021>.
- Twohy, CH, DeMott, PJ, Russell, LM, Toohey, DW, Rainwater, B, Geiss, R, Sanchez, KJ, Lewis, S, Roberts, GC, Humphries, RS, McCluskey, CS, Moore, KA, Selleck, PW, Keywood, MD, Ward, JP, McRobert, IM. 2021. Cloud-nucleating particles over the Southern Ocean in a changing climate. *Earth's Future* **9**(3): e2020EF001673. DOI: <https://doi.org/10.1029/2020EF001673>.
- Twomey, S. 1977. The influence of pollution on the short-wave albedo of clouds. *Journal of the Atmospheric Sciences* **34**(7): 1149–1152.
- Tzabari, M, Holodovsky, V, Shubi, O, Eytan, E, Altaratz, O, Koren, I, Aumann, A, Schilling, K, Schechner, YY. 2021. CloudCT 3D volumetric tomography: Considerations for imager preference, comparing visible light, short-wave infrared, and polarized imagers, in Kupinski, MK, Shaw, JA, Snik, F eds., *Polarization science and remote sensing X* (vol. **11833**). San Diego, CA: SPIE: 19–26.
- van Pinxteren, M, Barthel, S, Fomba, KW, Müller, K, von Tümpling, W, Herrmann, H. 2017. The influence of environmental drivers on the enrichment of organic carbon in the sea surface microlayer and in submicron aerosol particles—Measurements from the Atlantic Ocean. *Elementa: Science of the Anthropocene* **5**: 35. DOI: <https://doi.org/10.1525/elementa.225>.
- van Pinxteren, M, Fomba, KW, Triesch, N, Stolle, C, Wurl, O, Bahlmann, E, Gong, X, Voigtländer, J, Wex, H, Robinson, TB, Barthel, S, Zeppenfeld, S, Hoffmann, EH, Roveretto, M, Li, C, Grosselin, B, Daële, V, Senf, F, van Pinxteren, D, Manzi, M, Zabalegui, N, Frka, S, Gašparović, B, Pereira, R, Li, T, Wen, L, Li, J, Zhu, C, Chen, H, Chen, J, Fiedler, B, von Tümpling, W, Read, KA, Punjabi, S, Lewis, AC, Hopkins, JR, Carpenter, LJ, Peeken, I, Rixen, T, Schulz-Bull, D, Monge, ME, Mellouki, A, George, C, Stratmann, F, Herrmann, H. 2020. Marine organic matter in the remote environment of the Cape Verde islands—An introduction and overview to the MarParCloud campaign. *Atmospheric Chemistry and Physics* **20**(11): 6921–6951. DOI: <http://dx.doi.org/10.5194/acp-20-6921-2020>.
- Veres, PR, Neuman, JA, Bertram, TH, Assaf, E, Wolfe, GM, Williamson, CJ, Weinzierl, B, Tilmes, S, Thompson, CR, Thames, AB, Schroder, JC, Saiz-Lopez, A, Rollins, AW, Roberts, JM, Price, D, Peischl, J, Nault, BA, Möller, KH, Miller, DO, Meinardi, S, Li, Q, Lamarque, J-F, Kupc, A, Kjaergaard, HG, Kinnison, D, Jimenez, JL, Jernigan, CM, Hornbrook, RS, Hills, A, Dollner, M, Day, DA, Cuevas, CA, Campuzano-Jost, P, Burkholder, J, Bui, TP, Brune, WH, Brown, SS, Brock, CA, Bourgeois, I, Blake, DR, Apel, EC, Ryerson, TB. 2020. Global airborne sampling reveals a previously unobserved dimethyl sulfide oxidation mechanism in the marine atmosphere. *Proceedings of the National Academy of Sciences of the United States of America* **117**(9): 4505–4510. DOI: <https://doi.org/10.1073/pnas.1919344117>.
- Vergara-Temprado, J, Miltenberger, AK, Furtado, K, Grosvenor, DP, Shipway, BJ, Hill, AA, Wilkinson, JM, Field, PR, Murray, BJ, Carslaw, KS. 2018. Strong control of Southern Ocean cloud reflectivity by ice-nucleating particles. *Proceedings of the National Academy of Sciences of the United States of America* **115**(11): 2687–2692. DOI: <http://dx.doi.org/10.1073/pnas.1721627115>.

- Vila-Costa, M, Simó, R, Harada, H, Gasol, JM, Slezak, D, Kiene, RP. 2006. Dimethylsulfoniopropionate uptake by marine phytoplankton. *Science* **314**(5799): 652–654.
- Vogt, M, Vallina, SM, Buitenhuis, ET, Bopp, L, Le Quéré, C. 2010. Simulating dimethylsulphide seasonality with the dynamic green ocean model PlankTOM5. *Journal of Geophysical Research: Oceans* **115**(C6): C06021. DOI: <http://dx.doi.org/10.1029/2009JC005529>.
- Wagner, R, Ickes, L, Bertram, AK, Els, N, Gorokhova, E, Möhler, O, Murray, BJ, Umo, NS, Salter, ME. 2021. Heterogeneous ice nucleation ability of aerosol particles generated from Arctic sea surface microlayer and surface seawater samples at cirrus temperatures. *Atmospheric Chemistry and Physics* **21**(18): 13903–13930. DOI: <https://doi.org/10.5194/acp-21-13903-2021>.
- Wall, CJ, Norris, JR, Possner, A, McCoy, DT, McCoy, IL, Lutsko, NJ. 2022. Assessing effective radiative forcing from aerosol–cloud interactions over the global ocean. *Proceedings of the National Academy of Sciences* **119**(46): e2210481119. DOI: <https://doi.org/10.1073/pnas.2210481119>.
- Wang, DS, Lee, CP, Krechmer, JE, Majluf, F, Tong, Y, Canagaratna, MR, Schmale, J, Prévôt, ASH, Baltensperger, U, Dommen, J, El Haddad, I, Slowik, JG, Bell, DM. 2021. Constraining the response factors of an extractive electrospray ionization mass spectrometer for near-molecular aerosol speciation. *Atmospheric Measurement Techniques* **14**(11): 6955–6972. DOI: <https://doi.org/10.5194/amt-14-6955-2021>.
- Wang, J, Wood, R, Jensen, MP, Chiu, JC, Liu, Y, Lamer, K, Desai, N, Giangrande, SE, Knopf, DA, Kollias, P, Laskin, A, Liu, X, Lu, C, Mechem, D, Mei, F, Starzec, M, Tomlinson, J, Wang, Y, Yum, SS, Zheng, G, Aiken, AC, Azevedo, EB, Blanchard, Y, China, S, Dong, X, Gallo, F, Gao, S, Ghate, VP, Glienke, S, Goldberger, L, Hardin, JC, Kuang, C, Luke, EP, Matthews, AA, Miller, MA, Moffet, R, Pekour, M, Schmid, B, Sedlacek, AJ, Shaw, RA, Shilling, JE, Sullivan, A, Suski, K, Veghte, DP, Weber, R, Wyant, M, Yeom, J, Zawadowicz, M, Zhang, Z. 2022. Aerosol and Cloud Experiments in the Eastern North Atlantic (ACE-ENA). *Bulletin of the American Meteorological Society* **103**(2): E619–E641.
- Wang, S, Elliott, S, Maltrud, M, Cameron-Smith, P. 2015. Influence of explicit phaeocystis parameterizations on the global distribution of marine dimethyl sulfide. *Journal of Geophysical Research: Biogeosciences* **120**(11): 2158–2177.
- Wang, S, Maltrud, M, Elliott, S, Cameron-Smith, P, Jonko, A. 2018. Influence of dimethyl sulfide on the carbon cycle and biological production. *Biogeochemistry* **138**: 49–68. DOI: <http://dx.doi.org/10.1007/s10533-018-0430-5>.
- Wang, WL, Song, G, Primeau, F, Saltzman, ES, Bell, TG, Moore, JK. 2020. Global ocean dimethyl sulfide climatology estimated from observations and an artificial neural network. *Biogeosciences* **17**(21): 5335–5354. DOI: <http://dx.doi.org/10.5194/bg-17-5335-2020>.
- Wang, X, Deane, GB, Moore, KA, Ryder, OS, Stokes, MD, Beall, CM, Collins, DB, Santander, MV, Burrows, SM, Sultana, CM, Prather, KA. 2017. The role of jet and film drops in controlling the mixing state of submicron sea spray aerosol particles. *Proceedings of the National Academy of Sciences* **114**(27): 6978–6983. DOI: <https://doi.org/10.1073/pnas.1702420114>.
- Welti, A, Bigg, EK, DeMott, PJ, Gong, X, Hartmann, M, Harvey, M, Henning, S, Herenz, P, Hill, TCJ, Hornblow, B, Leck, C, Löffler, M, McCluskey, CS, Rauber, AM, Schmale, J, Tatzelt, C, van Pinxteren, M, Stratmann, F. 2020. Ship-based measurements of ice nuclei concentrations over the Arctic, Atlantic, Pacific and Southern Oceans. *Atmospheric Chemistry and Physics* **20**(23): 15191–15206. DOI: <http://dx.doi.org/10.5194/acp-20-15191-2020>.
- Wilbourn, EK, Thornton, DCO, Ott, C, Graff, J, Quinn, PK, Bates, TS, Betha, R, Russell, LM, Behrenfeld, MJ, Brooks, SD. 2020. Ice nucleation by marine aerosols over the north Atlantic Ocean in late spring. *Journal of Geophysical Research: Atmospheres* **125**: e2019JD030913. DOI: <http://dx.doi.org/10.1029/2019JD030913>.
- Williamson, CJ, Kupc, A, Axisa, D, Bilsback, KR, Bui, T, Campuzano-Jost, P, Dollner, M, Froyd, KD, Hodshire, AL, Jimenez, JL, Kodros, JK, Luo, G, Murphy, DM, Nault, BA, Ray, EA, Weinzierl, B, Wilson, JC, Yu, F, Yu, P, Pierce, JR, Brock, CA. 2019. A large source of cloud condensation nuclei from new particle formation in the tropics. *Nature* **574**: 399–403. DOI: <http://dx.doi.org/10.1038/s41586-019-1638-9>.
- Willis, M, Lannuzel, D, Else, B, Angot, H, Campbell, K, Crabeck, O, Delille, B, Hayashida, H, Lizotte, M, Loose, B, Meiners, KM, Miller, L, Moreau, S, Nomura, D, Prytherch, J, Schmale, J, Steiner, N, Tedesco, L, Thomas, J. 2023. Polar oceans and sea ice in a changing climate. *Elementa: Science of the Anthropocene* **11**(1): 00056. DOI: <https://doi.org/10.1525/elementa.2023.00056>.
- Wilson, TW, Ladino, LA, Alpert, PA, Breckels, MN, Brooks, IM, Browse, J, Burrows, SM, Carslaw, KS, Huffman, JA, Judd, C, Kilhau, WP, Mason, RH, McFiggans, G, Miller, LA, Nájera, JJ, Polishchuk, E, Rae, S, Schiller, CL, Si, M, Temprado, JV, Whale, TF, Wong, JPS, Wurl, O, Yakobi-Hancock, JD, Abbatt, JPD, Aller, JY, Bertram, AK, Knopf, DA, Murray, BJ. 2015. A marine biogenic source of atmospheric ice-nucleating particles. *Nature* **525**(7568): 234–238.
- Wohl, C, Li, Q, Cuevas, CA, Fernandez, RP, Yang, M, Saiz-Lopez, A, Simó, R. 2023. Marine biogenic emissions of benzene and toluene and their contribution to secondary organic aerosols over the polar oceans. *Science Advances* **9**: eadd9031. DOI: <https://doi.org/10.1126/sciadv.add9031>.

- Wolf, MJ, Coe, A, Dove, LA, Zawadowicz, MA, Dooley, K, Biller, SJ, Zhang, Y, Chisholm, SW, Cziczo, DJ.** 2019. Investigating the heterogeneous ice nucleation of sea spray aerosols using *Prochlorococcus* as a model source of marine organic matter. *Environmental Science & Technology* **53**(3): 1139–1149. DOI: <http://dx.doi.org/10.1021/acs.est.8b05150>.
- Wood, R.** 2012. Stratocumulus clouds. *Monthly Weather Review* **140**(8): 2373–2423.
- Wyant, MC, Bretherton, CS, Wood, R, Blossey, PN, McCoy, IL.** 2022. High free-tropospheric Aitken-mode aerosol concentrations buffer cloud droplet concentrations in large-eddy simulations of precipitating stratocumulus. *Journal of Advances in Modeling Earth Systems* **14**(6): e2021MS002930. DOI: <https://doi.org/10.1029/2021MS002930>.
- Xu, W, Ovadnevaite, J, Fossum, KN, Lin, C, Huang, RJ, Ceburnis, D, O'Dowd, C.** 2022. Sea spray as an obscured source for marine cloud nuclei. *Nature Geoscience* **15**(4): 282–286. DOI: <http://dx.doi.org/10.1038/s41561-022-00917-2>.
- Xue, H, Feingold, G.** 2006. Large-eddy simulations of trade wind cumuli: Investigation of aerosol indirect effects. *Journal of the Atmospheric Sciences* **63**(6): 1605–1622. DOI: <https://doi.org/10.1175/JAS3706.1>.
- Yan, C, Dada, L, Rose, C, Jokinen, T, Nie, W, Schobesberger, S, Junninen, H, Lehtipalo, K, Sarnela, N, Makkonen, U, Garmash, O, Wang, Y, Zha, Q, Paasonen, P, Bianchi, F, Sipilä, M, Ehn, M, Petäjä, T, Kerminen, V-M, Worsnop, DR, Kulmala, M.** 2018. The role of H₂SO₄-NH₃ anion clusters in ion-induced aerosol nucleation mechanisms in the boreal forest. *Atmospheric Chemistry and Physics* **18**(18): 13231–13243. DOI: <https://doi.org/10.5194/acp-18-13231-2018>.
- Yu, Z, Li, Y.** 2021. Marine volatile organic compounds and their impacts on marine aerosol—A review. *Science of the Total Environment* **768**: 145054.
- Zawadowicz, MA, Suski, K, Liu, J, Pekour, M, Fast, J, Mei, F, Sedlacek, AJ, Springston, S, Wang, Y, Zaveri, RA, Wood, R, Wang, J, Shilling, JE.** 2021. Aircraft measurements of aerosol and trace gas chemistry in the eastern North Atlantic. *Atmospheric Chemistry and Physics* **21**(10): 7983–8002. DOI: <http://dx.doi.org/10.5194/acp-21-7983-2021>.
- Zelinka, MD, Myers, TA, McCoy, DT, Po-Chedley, S, Caldwell, PM, Ceppi, P, Klein, SA, Taylor, KE.** 2020. Causes of higher climate sensitivity in CMIP6 models. *Geophysical Research Letters* **47**(1): e2019GL085782.
- Zhang, J, Feingold, G.** 2023. Distinct regional meteorological influences on low-cloud albedo susceptibility over global marine stratocumulus regions. *Atmospheric Chemistry and Physics* **23**(2): 1073–1090. DOI: <https://doi.org/10.5194/acp-23-1073-2023>.
- Zhang, J, Liu, J, Liu, D, Chen, X, Shi, Q, He, C, Li, G.** 2022. Temperature rise increases the bioavailability of marine *Synechococcus*-derived dissolved organic matter. *Frontiers in Microbiology*. DOI: <http://dx.doi.org/10.3389/fmicb.2022.838707>.
- Zhao, X, Liu, X, Burrows, SM, Shi, Y.** 2021. Effects of marine organic aerosols as sources of immersion-mode ice-nucleating particles on high-latitude mixed-phase clouds. *Atmospheric Chemistry and Physics* **21**(4): 2305–2327. DOI: <http://dx.doi.org/10.5194/acp-21-2305-2021>.
- Zheng, G, Kuang, C, Uin, J, Watson, T, Wang, J.** 2020. Large contribution of organics to condensational growth and formation of Cloud Condensation Nuclei (CCN) in the remote marine boundary layer. *Atmospheric Chemistry and Physics* **20**(21): 12515–12525. DOI: <http://dx.doi.org/10.5194/acp-20-12515-2020>.
- Zheng, G, Wang, Y, Aiken, AC, Gallo, F, Jensen, MP, Kollias, P, Kuang, C, Luke, E, Springston, S, Uin, J, Wood, R, Wang, J.** 2018. Marine boundary layer aerosol in the eastern North Atlantic: Seasonal variations and key controlling processes. *Atmospheric Chemistry and Physics* **18**(23): 17615–17635.
- Zheng, G, Wang, Y, Wood, R, Jensen, MP, Kuang, C, McCoy, IL, Matthews, A, Mei, F, Tomlinson, JM, Shilling, JE, Zawadowicz, MA, Crosbie, E, Moore, R, Ziemba, L, Andreae, MO, Wang, J.** 2021. New particle formation in the remote marine boundary layer. *Nature Communications* **12**: 1–10. DOI: <http://dx.doi.org/10.1038/s41467-020-20773-1>.
- Zhou, X, Feingold, G.** 2023. Impacts of mesoscale cloud organization on aerosol-induced cloud water adjustment and cloud brightness. *Geophysical Research Letters* **50**(13): e2023GL103417. DOI: <http://dx.doi.org/10.1029/2023GL103417>.
- Zuidema, P, Alvarado, M, Chiu, C, DeSzoeko, S, Fairall, C, Feingold, G, Freedman, A, Ghan, S, Haywood, J, Kollias, P, Lewis, E, McFarquhar, G, McComiskey, A, Mechem, D, Onasch, T, Redemann, J, Romps, D, Turner, D, Wang, H, Wood, R, Yuter, S, Zhu, P.** 2018. Layered Atlantic Smoke Interactions with Clouds (LASIC) Field Campaign Report [Staford, R ed.]. ARM Climate Research Facility Report DOE/SC-ARM-18-018, U.S. Department of Energy, Office of Science.

How to cite this article: Sellegrì, K, Simó, R, Wang, B, Alpert, PA, Altieri, K, Burrows, S, Hopkins, FE, Koren, I, McCoy, IL, Ovadnevaite, J, Salter, M, Schmale, J. 2024. Influence of open ocean biogeochemistry on aerosol and clouds: Recent findings and perspectives. *Elementa: Science of the Anthropocene* 12(1). DOI: <https://doi.org/10.1525/elementa.2023.00058>

Domain Editor-in-Chief: Detlev Helmig, Boulder AIR LLC, Boulder, CO, USA

Knowledge Domain: Atmospheric Science

Part of an Elementa Special Feature: Boundary Shift: The Air-Sea Interface in a Changing Climate

Published: July 29, 2024 **Accepted:** May 09, 2024 **Submitted:** April 24, 2023

Copyright: © 2024 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.



Elem Sci Anth is a peer-reviewed open access journal published by University of California Press.

OPEN ACCESS 