



Geometry & Topology

Volume 27 (2023)

Contact three-manifolds with exactly two simple Reeb orbits

DANIEL CRISTOFARO-GARDINER

UMBERTO HRYNIEWICZ

MICHAEL HUTCHINGS

HUI LIU



Contact three-manifolds with exactly two simple Reeb orbits

DANIEL CRISTOFARO-GARDINER

UMBERTO HRYNIEWICZ

MICHAEL HUTCHINGS

HUI LIU

It is known that every contact form on a closed three-manifold has at least two simple Reeb orbits, and a generic contact form has infinitely many. We show that if there are exactly two simple Reeb orbits, then the contact form is nondegenerate. Combined with a previous result, this implies that the three-manifold is diffeomorphic to the three-sphere or a lens space, and the two simple Reeb orbits are the core circles of a genus-one Heegaard splitting. We also obtain further information about the Reeb dynamics and the contact structure. For example, the Reeb flow has a disk-like global surface of section and so its dynamics are described by a pseudorotation, the contact structure is universally tight, and in the case of the three-sphere the contact volume and the periods and rotation numbers of the simple Reeb orbits satisfy the same relations as for an irrational ellipsoid.

37J99, 53E50; 53D42

1 Introduction

1.1 Statement of results

Let Y be a closed oriented three-manifold. Recall that a *contact form* on Y is a 1-form λ on Y such that $\lambda \wedge d\lambda > 0$. A contact form λ has an associated *Reeb vector field* R defined by the equations

$$d\lambda(R, \cdot) = 0, \quad \lambda(R) = 1.$$

A *Reeb orbit* is a periodic orbit of R , ie a map

$$\gamma: \mathbb{R}/T\mathbb{Z} \rightarrow Y, \quad \gamma'(t) = R(\gamma(t)),$$

for some $T > 0$, modulo reparametrization of the domain by translations. The number T is the period, also called the *symplectic action*, of γ . We say that the Reeb orbit γ is

simple if the map γ is an embedding. Every Reeb orbit is the k -fold cover of a simple Reeb orbit for some positive integer k .

The three-dimensional case of the Weinstein conjecture, which was proved in full generality by Taubes [42], asserts that a contact form on a closed three-manifold has at least one Reeb orbit; see [28] for a survey. It was further shown in [11] that a contact form on a closed three-manifold has at least two simple Reeb orbits. This lower bound is the best possible without further hypotheses:

Example 1.1 Recall that if Y is a compact hypersurface in $\mathbb{R}^4 = \mathbb{C}^2$ which is “star-shaped” (transverse to the radial vector field), then the standard Liouville form

$$(1-1) \quad \lambda = \frac{1}{2} \sum_{i=1}^2 (x_i dy_i - y_i dx_i)$$

restricts to a contact form on Y . If Y is the three-dimensional ellipsoid

$$\partial E(a, b) = \left\{ z \in \mathbb{C}^2 \mid \frac{\pi |z_1|^2}{a} + \frac{\pi |z_2|^2}{b} = 1 \right\}$$

and if a/b is irrational, then there are exactly two simple Reeb orbits, corresponding to the circles in Y where $z_2 = 0$ and $z_1 = 0$, with periods a and b , respectively.

One can also take quotients of the above irrational ellipsoids by finite cyclic group actions to obtain contact forms on lens spaces with exactly two simple Reeb orbits.

It is conjectured that, in fact, every contact form on a closed connected three-manifold has either two or infinitely many simple Reeb orbits. This was proved by Colin, Dehornoy and Rechtman [8] for contact forms that are nondegenerate (see the definition below), extending a result of [12]. It was also shown by Irie [35] that, for a C^∞ -generic contact form on a closed three-manifold, there are infinitely many simple Reeb orbits, and moreover their images are dense in the three-manifold.

The goal of this paper is to give detailed information about the “exceptional” case of contact forms on a closed three-manifold with exactly two simple Reeb orbits.

To state the first result, let $\xi = \text{Ker}(\lambda)$ denote the contact structure determined by λ . This is a rank-2 vector bundle with a linear symplectic form $d\lambda$. If $\gamma: \mathbb{R}/T\mathbb{Z} \rightarrow Y$ is a Reeb orbit, then the derivative of the time T flow of R restricts to a symplectic linear map

$$(1-2) \quad P_\gamma: (\xi_{\gamma(0)}, d\lambda) \rightarrow (\xi_{\gamma(0)}, d\lambda),$$

which we call the *linearized return map*. We say that γ is *nondegenerate* if 1 is not an eigenvalue of P_γ ; this condition is invariant under reparametrization of γ . We say

that the contact form λ is nondegenerate if all Reeb orbits (including nonsimple ones) are nondegenerate. The set of nondegenerate contact forms is residual in the set of all contact forms with the C^∞ -topology. The Reeb orbit γ is called *hyperbolic* if P_γ has eigenvalues in $\mathbb{R} \setminus \{\pm 1\}$. The Reeb orbit γ is called *elliptic* if the eigenvalues of P_γ are of the form $e^{\pm 2\pi i \phi}$, and *irrationally elliptic* if moreover ϕ is irrational. If γ is irrationally elliptic, then γ and all of its covers are nondegenerate, because the linearized return map for the k -fold cover of γ has eigenvalues $e^{\pm 2\pi i k \phi}$.

Many results about Reeb dynamics and related questions assume some kind of nondegeneracy hypothesis or allow only certain kinds of degeneracies. One of the main points of the present work is that we can derive our results without making any such assumption.

Theorem 1.2 *Let Y be a closed three-manifold, and let λ be a contact form on Y with exactly two simple Reeb orbits. Then λ is nondegenerate and, moreover, both simple Reeb orbits are irrationally elliptic.*

Theorem 1.2 might seem surprising in view of known results about critical points of real-valued functions on finite-dimensional manifolds. For example, on the two-torus the minimal number of critical points is three, and when there are only three critical points they cannot all be nondegenerate. We refer the reader to Remark 1.12 for related discussion.

As a corollary of Theorem 1.2, we obtain the following topological constraint:

Corollary 1.3 *Let Y be a closed three-manifold, and let λ be a contact form on Y with exactly two simple Reeb orbits. Then Y is diffeomorphic to a lens space.¹ Moreover, the two simple Reeb orbits are the core circles of a genus-one Heegaard splitting of Y .*

Proof This was shown in [33, Theorem 1.3 and Section 4.8] under the additional hypothesis that λ is nondegenerate. By Theorem 1.2, this nondegeneracy automatically holds. $\square$

Remark 1.4 A special case of Theorem 1.2, where Y is a compact convex hypersurface in $\mathbb{R}^4$ with the restriction of the standard Liouville form (1-1), was previously shown by Wang, Hu and Long [45, Theorem 1.4].

¹Here and below our convention is that S^3 is a lens space, but $S^1 \times S^2$ is not.

We also obtain additional dynamical information. To state the result, recall that the *contact volume* of (Y, λ) is defined by

$$\text{vol}(Y, \lambda) := \int_Y \lambda \wedge d\lambda.$$

Theorem 1.5 *Let Y be a lens space and let λ be a contact form on Y with exactly two simple Reeb orbits, γ_1 and γ_2 . Then:*

- (a) *Let $p = |\pi_1(Y)| < \infty$, let $T_i \in \mathbb{R}$ denote the period of γ_i , and let $\phi_i \in \mathbb{R}$ denote the “Seifert rotation number” of γ_i ; see Definition 4.3. Then*

$$\text{vol}(Y, \lambda) = pT_1T_2 = T_1^2/\phi_1 = T_2^2/\phi_2.$$

- (b) *λ is dynamically convex, and the contact structure $\xi = \text{Ker}(\lambda)$ is universally tight.*²

Example 1.6 For the ellipsoid in Example 1.1, we have $T_1 = a$, $T_2 = b$, $\phi_1 = a/b$, $\phi_2 = b/a$, $p = 1$, and $\text{vol} = ab$. Thus Theorem 1.5(a) implies that if $Y = S^3$, then the periods T_i , the rotation numbers ϕ_i , and the contact volume satisfy the same relations as for an ellipsoid. For $Y = S^3$, under the additional assumptions that λ is nondegenerate and ξ is the standard contact structure, it was previously shown by Bourgeois, Cieliebak and Ekholm [5] and Gürel [19] that “action–index relations” hold, implying that the periods T_i and rotation numbers ϕ_i satisfy the same relations as for an ellipsoid. The equation $\text{vol} = T_1T_2$ that we prove in this case answers [4, Question 2].

Remark 1.7 There exist contact forms on S^3 with exactly two simple Reeb orbits which are not strictly contactomorphic to ellipsoids. One way to see this is to start from Katok’s construction [37] of Finsler metrics on S^2 with exactly two closed geodesics, such that the Liouville measure on the unit tangent bundle is ergodic for the geodesic flow. Such a geodesic flow can then be lifted to a Reeb flow on the standard contact 3–sphere with the same properties. Another way to see this is by Albers, Geiges and Zehmisch [1], who showed that the pseudorotations from Fayad and Katok [15] can be realized as the return map on a disk-like global surface of section for a Reeb flow on the standard contact 3–sphere with precisely two periodic orbits; see Section 1.3 below. On the other hand, Helmut Hofer has suggested to the authors in private

²Recall that a contact form on a three-manifold Y with $c_1(\xi)|_{\pi_2(Y)} = 0$ is called *dynamically convex* if $\text{CZ}(\gamma) \geq 3$ for every contractible Reeb orbit γ , where CZ denotes the Conley–Zehnder index (see Section 2.2) computed with respect to a trivialization which extends over a disc bounded by γ . A contact structure on Y is *universally tight* if its pullback to the universal cover of Y is tight.

correspondence (2021) that perhaps imposing the additional condition that the rotation numbers of the two Reeb orbits are Diophantine forces the contact form to be strictly contactomorphic to an ellipsoid; see Fayad and Krikorian [16, Question 6].

Remark 1.8 As shown by Honda [23, Proposition 5.1] (see Cornwell [10, page 17] for more explanation), each lens space has either one or two universally tight contact structures up to isotopy, and when there are two they are contactomorphic (and one is obtained from the other by reversing its orientation). Consequently, in Theorem 1.5(b), the contact structure is contactomorphic to a “standard” contact structure on the lens space obtained as in Example 1.1. In particular, universally tight contact structures on lens spaces are precisely the ones that admit contact forms with exactly two simple Reeb orbits. Some other results obtaining information about contact structures from Reeb dynamics can be found in work by Etnyre and Ghrist [14], Hofer, Wysocki and Zehnder [20; 22], and [24].

Remark 1.9 We also obtain information about the knot types of the simple Reeb orbits γ_1 and γ_2 . It follows from the Heegaard splitting in Corollary 1.3 that these are p –unknotted. We further show in Section 5 that their self-linking number is -1 when $p = 1$; similar arguments show that, for general p , their rational self-linking number, as defined by Baker and Etnyre [2], equals $-1/p$.

1.2 Outline of the proofs

We now briefly describe the proofs of Theorems 1.2 and 1.5.

A key ingredient in these proofs, as well as in the related papers [11; 12], is the “volume property” in embedded contact homology, which was proved in [13]. The embedded contact homology (ECH) of (Y, λ) is the homology of a chain complex which is built out of Reeb orbits, and whose differential counts (mostly) embedded pseudoholomorphic curves in $\mathbb{R} \times Y$; see the lecture notes [30] and the review in Section 2. The version of the volume property that we will use here asserts that if Y is a closed connected 3–manifold with a contact form λ , then

$$\lim_{k \rightarrow \infty} \frac{c_{\sigma_k}(Y, \lambda)^2}{k} = 2 \operatorname{vol}(Y, \lambda).$$

Here $\{\sigma_k\}$ is a “ U –sequence” in ECH, and c_{σ_k} is a “spectral invariant” associated to σ_k , which is the total symplectic action of a certain finite set of Reeb orbits determined by σ_k ; these notions are reviewed in Section 2.

The outline of the proof of Theorem 1.2 is as follows. Let γ_1 and γ_2 denote the two simple Reeb orbits, and let T_1 and T_2 denote their periods. Simple applications of the volume property from [11; 12] (just using the $k^{1/2}$ growth rate of the spectral invariants and not the exact relation with contact volume) show that the homology classes $[\gamma_i] \in H_1(Y)$ are torsion, and the ratio T_1/T_2 is irrational. A more precise use of the volume property then gives the relations

$$(1-3) \quad \phi_i = \frac{T_i^2}{\text{vol}(Y, \lambda)}, \quad \ell(\gamma_1, \gamma_2) = \frac{T_1 T_2}{\text{vol}(Y, \lambda)},$$

where $\phi_i \in \mathbb{R}$ is the Seifert rotation number that appears in Theorem 1.5(a), while $\ell(\gamma_1, \gamma_2) \in \mathbb{Q}$ is the linking number of γ_1 and γ_2 ; see Definition 4.2. The proof of (1-3) also depends on a new estimate for the behavior of the ECH index (the grading on the ECH chain complex) under perturbations of possibly degenerate contact forms, which is proved in Section 3.

The equations (1-3) imply the relations

$$(1-4) \quad \phi_1 = \ell(\gamma_1, \gamma_2) \frac{T_1}{T_2}, \quad \phi_2 = \ell(\gamma_1, \gamma_2) \frac{T_2}{T_1}.$$

Since $\ell(\gamma_1, \gamma_2)$ is rational and T_1/T_2 is irrational, it follows that ϕ_1 and ϕ_2 are irrational. This implies that γ_1 and γ_2 are rationally elliptic (see Section 4.4), which completes the proof of Theorem 1.2.

The Heegaard decomposition in Corollary 1.3 implies that $\ell(\gamma_1, \gamma_2) = 1/p$, and combined with (1-4) this proves Theorem 1.5(a). The proof of Theorem 1.5(b) uses additional calculations in Section 5 to deduce dynamical convexity and universal tightness from information about the numbers ϕ_i .

1.3 Pseudorotations

The contact forms studied here are analogous to “pseudorotations”, defined in various ways as maps in some class with the minimum number of periodic orbits. For example, according to Ginzburg and Gürel [18] a *Hamiltonian pseudorotation* of $\mathbb{C}P^n$ is defined to be a Hamiltonian symplectomorphism of $\mathbb{C}P^n$ with $n + 1$ fixed points and no other periodic points; see eg Le Roux and Seyfaddini [39], Shelukhin [41] and Çineli, Ginzburg and Gürel [7] for generalizations to other symplectic manifolds. More classically, we consider here pseudorotations of the open or closed disk defined as area-preserving homeomorphisms with one fixed point and no other periodic points; see eg Bramham [6] and Fayad and Katok [15].

In fact, there is a direct connection between the contact forms considered here and pseudorotations of the closed disk. Recall that given a closed three-manifold Y with a contact form λ , a *disk-like global surface of section* for the Reeb flow is an immersed disk, with boundary on a Reeb orbit, embedded and transverse to the Reeb flow in the interior, such that the Reeb flow starting at any point in Y hits the disk both forwards and backwards in time.

Corollary 1.10 *Let Y be a closed three-manifold, and let λ be a contact form on Y with exactly two simple Reeb orbits. Then both orbits bound disk-like global surfaces of section whose associated return maps define smooth pseudorotations of the open disk.*

Proof By Theorem 1.2, Corollary 1.3 and Theorem 1.5, Y is a lens space and λ is nondegenerate and dynamically convex. As explained in Remark 1.9, both orbits are p -unknotted, with self-linking number $-1/p$. Hence, the result follows from [24, Theorem 1.12]. $\square$

Remark 1.11 Conversely, as mentioned above, at least some pseudorotations of the closed disk can be “suspended” to contact forms on S^3 with exactly two simple Reeb orbits; see Albers, Geiges and Zehmisch [1].

Remark 1.12 It is shown by Collier, Kerman, Reiniger, Turmunkh and Zimmer [9] — see also Franks [17] — that, for a Hamiltonian pseudorotation of $\mathbb{C}P^1$, each fixed point is strongly nondegenerate, meaning that the linearized return map and its higher powers are nondegenerate, and moreover the fixed points are irrationally elliptic, similarly to Theorem 1.2. It is an open question whether every pseudorotation of $\mathbb{C}P^n$ for $n > 1$ is strongly nondegenerate, and one can ask analogous questions for pseudorotations of more general symplectic manifolds.

Remark 1.13 For pseudorotations of D^2 , in many cases, Joly [36] and Pirnapasov [40] proved identities related to Theorem 1.5(a).

Remark 1.14 One can arrange in the statement of Corollary 1.10 that the first return maps on the obtained disk-like global surfaces of section extend smoothly to the boundary and preserve a smooth 2-form that defines an area form in the interior. Moreover, one can conjugate such a return map by a homeomorphism to obtain a pseudorotation of the closed disk which is smooth in the interior.

Acknowledgments Cristofaro-Gardiner was partially supported by NSF grants DMS-1711976 and DMS-2105471. Hryniewicz was partially supported by the DFG SFB/TRR

191 *Symplectic structures in geometry, algebra and dynamics*, 281071066–TRR 191. Hutchings was partially supported by NSF grant DMS–2005437. Liu was partially supported by NSFC (Nos. 12022111 and 11771341) and the Fundamental Research Funds for the Central Universities (No. 2042021kf1059).

Key discussions for this project took place when Cristofaro-Gardiner and Liu were visiting Peking University for the *Workshop on gauge theory and Floer homology* in December 2019. We thank Peking University for their hospitality during this visit. This research was completed while Cristofaro-Gardiner was on a von Neumann fellowship at the Institute for Advanced Study and he thanks the institute for their wonderful support. We thank Viktor Ginzburg and Helmut Hofer for helpful comments.

We thank the referee for careful reading of the paper.

2 Preliminaries

In this section we review the material about embedded contact homology that is needed for the proofs of Theorems 1.2 and 1.5. We include a new, slight extension of the definition of the ECH index to degenerate contact forms.

Throughout this section fix a closed oriented three-manifold Y and a contact form λ on Y , and let $\xi = \text{Ker}(\lambda)$ denote the associated contact structure.

2.1 Topological preliminaries

We now recall some topological notions we will need, following the treatment in [27]. These were originally introduced in a slightly different context in [26].

Definition 2.1 An *orbit set* is a finite set of pairs $\alpha = \{(\alpha_i, m_i)\}$ where the α_i are distinct simple Reeb orbits, and the m_i are positive integers. We define the homology class of the orbit set α by

$$[\alpha] = \sum_i m_i [\alpha_i] \in H_1(Y).$$

Definition 2.2 If $\alpha = \{(\alpha_i, m_i)\}$ and $\beta = \{(\beta_j, n_j)\}$ are orbit sets with $[\alpha] = [\beta]$, define $H_2(Y, \alpha, \beta)$ to be the set of 2-chains Z in Y with $\partial Z = \sum_i m_i \alpha_i - \sum_j n_j \beta_j$, modulo boundaries of 3-chains. The set $H_2(Y, \alpha, \beta)$ is an affine space over $H_2(Y)$.

Given orbit sets α and β as above, let $Z \in H_2(Y, \alpha, \beta)$, and let τ be a homotopy class of symplectic trivialization of the contact structure ξ over the Reeb orbits α_i and β_j .

Definition 2.3 (see [27, Section 2.5]) Define the *relative first Chern class*

$$c_\tau(\alpha, \beta, Z) \in \mathbb{Z}$$

as follows. Let S be a compact oriented surface with boundary and let $f: S \rightarrow Y$ be a smooth map representing the class Z . Let ψ be a section of $f^*\xi$ which, on each boundary component, is nonvanishing and constant with respect to τ . Define $c_\tau(\alpha, \beta, Z)$ to be the algebraic count of zeroes of ψ .

Definition 2.4 [27, Section 2.7] An *admissible representative* of $Z \in H_2(Y, \alpha, \beta)$ is a smooth map $f: S \rightarrow [-1, 1] \times Y$ where S is a compact oriented surface with boundary, the restriction of f to ∂S consists of positively oriented covers of $\{1\} \times \alpha_i$ with total multiplicity m_i and negatively oriented covers of $\{-1\} \times \beta_j$ with total multiplicity n_j , the composition of f with the projection $[-1, 1] \times Y \rightarrow Y$ represents the class Z , the restriction of f to the interior of S is an embedding, and f is transverse to $\{-1, 1\} \times Y$.

Definition 2.5 [27, Section 2.7] If $Z, Z' \in H_2(Y, \alpha, \beta)$, define the *relative intersection pairing*

$$Q_\tau(Z, Z') \in \mathbb{Z}$$

as follows. Let S and S' be admissible representatives of Z and Z' , respectively, whose interiors are transverse and do not intersect near the boundary. Define

$$(2-1) \quad Q_\tau(Z, Z') = \#(\text{int}(S) \cap \text{int}(S')) - \sum_i \ell_\tau(\xi_i^+, \xi_i^{+'}) + \sum_j \ell_\tau(\xi_j^-, \xi_j^{-'}).$$

Here $\#$ denotes the signed count of intersections, while the remaining terms are linking numbers defined as follows. For $\epsilon > 0$ small, the intersection of S with $\{1 - \epsilon\} \times Y$ consists of the union over i of a braid ξ_i^+ in a neighborhood of α_i (see Section 3.1), while the intersection of S with $\{-1 + \epsilon\} \times Y$ consists of the union over j of a braid ξ_j^- in a neighborhood of β_j . Likewise, S' determines braids $\xi_i^{+'}$ and $\xi_j^{-'}$. The notation ℓ_τ indicates the linking number in a neighborhood of α_i or β_j computed using the trivialization τ ; see [27, Section 2.6] for details and sign conventions.

When $Z = Z'$, we write³

$$Q_\tau(\alpha, \beta, Z) = Q_\tau(Z, Z).$$

³An alternative equivalent definition of $Q_\tau(\alpha, \beta, Z)$ is given in [30, Section 3.3], which does not include the linking number terms in (2-1). There the admissible representatives S and S' are required to satisfy additional conditions which force these linking number terms to be zero.

As explained in [27], the relative first Chern class $c_\tau(\alpha, \beta, Z)$ and the relative self-intersection number $Q_\tau(\alpha, \beta, Z)$ depend only on α, β, Z and τ . Moreover, if we change Z by adding $A \in H_2(Y)$, then

$$(2-2) \quad c_\tau(\alpha, \beta, Z + A) - c_\tau(\alpha, \beta, Z) = \langle c_1(\xi), A \rangle,$$

$$(2-3) \quad Q_\tau(\alpha, \beta, Z + A) - Q_\tau(\alpha, \beta, Z) = 2[\alpha] \cdot A.$$

Remark 2.6 If γ is a third orbit set, if τ is a trivialization of ξ over the Reeb orbits in α, β and γ , and if $W \in H_2(Y, \beta, \gamma)$, then we have the additivity properties

$$c_\tau(\alpha, \beta, Z) + c_\tau(\beta, \gamma, W) = c_\tau(\alpha, \gamma, Z + W),$$

$$Q_\tau(\alpha, \beta, Z) + Q_\tau(\beta, \gamma, W) = Q_\tau(\alpha, \gamma, Z + W).$$

Note also that the definition of c_τ makes sense more generally if the α_i and β_j are transverse knots. Likewise the definition of Q_τ makes sense if the α_i and β_j are knots and τ is an oriented trivialization of their normal bundles.

2.2 The ECH index

Let $\gamma: \mathbb{R}/T\mathbb{Z} \rightarrow Y$ be a Reeb orbit and let τ be a symplectic trivialization of $\gamma^*\xi$. The derivative of the time t Reeb flow from $\xi_{\gamma(0)}$ to $\xi_{\gamma(t)}$, with respect to τ , is a 2×2 symplectic matrix $\Phi(t)$. The family of symplectic matrices $\{\Phi(t)\}_{t \in [0, T]}$ induces a family of diffeomorphisms of S^1 in the universal cover of $\text{Diff}(S^1)$, which has a dynamical rotation number, here denoted by $\theta_\tau(\gamma) \in \mathbb{R}$. We call this real number the *rotation number* of γ with respect to τ and denote it by $\theta_\tau(\gamma) \in \mathbb{R}$; it depends only on γ and the homotopy class of τ . When $\theta_\tau(\gamma) \notin \frac{1}{2}\mathbb{Z}$, the eigenvalues of the linearized return map (1-2) are $e^{\pm 2\pi i \theta_\tau(\gamma)}$.

Definition 2.7 Define the *Conley–Zehnder index*

$$(2-4) \quad \text{CZ}_\tau(\gamma) = \lfloor \theta_\tau(\gamma) \rfloor + \lceil \theta_\tau(\gamma) \rceil \in \mathbb{Z}.$$

Remark 2.8 The above definition agrees with the usual Conley–Zehnder index when γ is nondegenerate. When γ is degenerate, it is common to give a different definition of the Conley–Zehnder index, as the minimum of the Conley–Zehnder indices of nondegenerate perturbations of γ , and this will sometimes differ from our definition by 1. For our purposes, especially to obtain an estimate as in Proposition 3.1 below (possibly with a different constant), it does not matter which of these definitions of the Conley–Zehnder index we use for degenerate Reeb orbits.

Notation 2.9 If $\alpha = \{(\alpha_i, m_i)\}$ is an orbit set and if τ is a trivialization of ξ over all of the Reeb orbits α_i , define

$$(2-5) \quad \text{CZ}_\tau^I(\alpha) = \sum_i \sum_{k=1}^{m_i} \text{CZ}_\tau(\alpha_i^k).$$

Here γ^k denotes the k^{th} iterate of γ .

Definition 2.10 Let α and β be orbit sets with $[\alpha] = [\beta] \in H_1(Y)$, and $Z \in H_2(Y, \alpha, \beta)$. Define the *ECH index*

$$(2-6) \quad I(\alpha, \beta, Z) = c_\tau(\alpha, \beta, Z) + Q_\tau(\alpha, \beta, Z) + \text{CZ}_\tau^I(\alpha) - \text{CZ}_\tau^I(\beta) \in \mathbb{Z}.$$

The above agrees with the usual definition of the ECH index — see eg [30, Section 3.4] — when the contact form is nondegenerate. It is explained, for example in [27, Section 2.8], why $I(\alpha, \beta, Z)$ depends only on α , β and Z , and not on τ . Moreover, it follows from (2-2) and (2-3) that, if we change Z by adding $A \in H_2(Y)$, then

$$(2-7) \quad I(\alpha, \beta, Z + A) - I(\alpha, \beta, Z) = \langle c_1(\xi) + 2 \text{PD}(\Gamma), A \rangle,$$

where $\Gamma = [\alpha] = [\beta] \in H_1(Y)$ and PD denotes the Poincaré dual. By Remark 2.6,

$$(2-8) \quad I(\alpha, \beta, Z) + I(\beta, \gamma, W) = I(\alpha, \gamma, Z + W).$$

2.3 Embedded contact homology

In this subsection assume that the contact form λ is nondegenerate. Let $\Gamma \in H_1(Y)$. We now review how to define the embedded contact homology $\text{ECH}_*(Y, \xi, \Gamma)$. More details may be found in [30].

Definition 2.11 An *ECH generator* is an orbit set $\alpha = \{(\alpha_i, m_i)\}$ such that $m_i = 1$ whenever α_i is hyperbolic.

Definition 2.12 Define $\text{ECC}_*(Y, \lambda, \Gamma)$ to be the vector space⁴ over $\mathbb{Z}/2$ generated by ECH generators α with $[\alpha] = \Gamma$. This vector space has a relative $\mathbb{Z}/d$ -grading, where d denotes the divisibility of $c_1(\xi) + 2 \text{PD}(\Gamma) \in H^2(Y; \mathbb{Z})$; if α and β are two generators, then their grading difference is $I(\alpha, \beta, Z) \bmod d$ for any $Z \in H_2(Y, \alpha, \beta)$. This makes sense by (2-7) and (2-8).

⁴It is also possible to use $\mathbb{Z}$ coefficients, as explained in [32, Section 9], but this has not been necessary for the applications of ECH so far.

Remark 2.13 In the special case where $c_1(\xi) \in H^2(Y; \mathbb{Z})$ is torsion and $\Gamma = 0$, the chain complex $\text{ECC}_*(Y, \lambda, 0)$ has a canonical absolute $\mathbb{Z}$ -grading defined by

$$I(\alpha) = I(\alpha, \emptyset, Z) \in \mathbb{Z}$$

for any $Z \in H_2(Y, \alpha, \emptyset)$. This is well defined by (2-7).

Definition 2.14 An almost complex structure J on $\mathbb{R} \times Y$ is λ -compatible if $J\partial_s = R$, where s denotes the $\mathbb{R}$ coordinate, J is invariant under the $\mathbb{R}$ action on $\mathbb{R} \times Y$ by translation of s , and $J(\xi) = \xi$, rotating positively with respect to $d\lambda$.

If J is a generic λ -compatible almost complex structure, one defines a differential

$$\partial_J : \text{ECC}_*(Y, \lambda, \Gamma) \rightarrow \text{ECC}_{*-1}(Y, \lambda, \Gamma)$$

whose coefficient from α to β is a count of “ J -holomorphic currents” that represent classes $Z \in H_2(Y, \alpha, \beta)$ with ECH index $I(\alpha, \beta, Z) = 1$; see [30, Section 3] for details. It is shown in [31] that $\partial_J^2 = 0$. The *embedded contact homology* $\text{ECH}_*(Y, \lambda, \Gamma, J)$ is defined to be the homology of the chain complex $(\text{ECC}_*(Y, \lambda, \Gamma), \partial_J)$. A theorem of Taubes [43], tensored with $\mathbb{Z}/2$, asserts that there is a canonical isomorphism

$$(2-9) \quad \text{ECH}_*(Y, \lambda, \Gamma, J) = \widehat{HM}^{-*}(Y, \mathfrak{s}_\xi + \text{PD}(\Gamma)) \otimes \mathbb{Z}/2,$$

where the right-hand side is a version of Seiberg–Witten Floer cohomology as defined by Kronheimer and Mrowka [38], and $\mathfrak{s}_\xi$ is a spin-c structure on Y determined by ξ . In particular, ECH depends only⁵ on the triple (Y, ξ, Γ) , and so we can denote it by $\text{ECH}_*(Y, \xi, \Gamma)$.

When Y is connected, there is also a well-defined “ U -map”

$$(2-10) \quad U : \text{ECH}_*(Y, \xi, \Gamma) \rightarrow \text{ECH}_{*-2}(Y, \xi, \Gamma).$$

This is induced by a chain map

$$U_{J,z} : (\text{ECC}_*(Y, \lambda, \Gamma), \partial_J) \rightarrow (\text{ECC}_{*-2}(Y, \xi, \Gamma), \partial_J)$$

which counts J -holomorphic currents with ECH index 2 passing through a generic basepoint $z \in \mathbb{R} \times Y$. The assumption that Y is connected implies that the induced map on homology does not depend on the choice of basepoint z ; see [33, Section 2.5] for details. Taubes showed in [44, Theorem 1.1] that under the isomorphism (2-9), the map on homology induced by $U_{J,z}$ agrees with a corresponding map on Seiberg–Witten Floer cohomology. We thus obtain a well-defined U -map (2-10).

⁵In a sense, ECH does not depend on the contact structure either; see [30, Remark 1.7] for explanation.

Definition 2.15 A U -sequence for Γ is a sequence $\{\sigma_k\}_{k \geq 1}$ where each σ_k is a nonzero homogeneous class in $\text{ECH}_*(Y, \xi, \Gamma)$, and $U\sigma_{k+1} = \sigma_k$ for each $k \geq 1$.

We will need the following nontriviality result for the U -map, which is proved by combining Taubes' isomorphism (2-9) with results from Kronheimer and Mrowka [38]:

Proposition 2.16 [12, Proposition 2.3] *If $c_1(\xi) + 2\text{PD}(\Gamma) \in H^2(Y; \mathbb{Z})$ is torsion, then a U -sequence for Γ exists.*

2.4 Spectral invariants

If $\alpha = \{(\alpha_i, m_i)\}$ is an orbit set, define its *symplectic action* by

$$\mathcal{A}(\alpha) = \sum_i m_i \int_{\alpha_i} \lambda.$$

Note here that $\int_{\alpha_i} \lambda$ agrees with the period of α_i , because $\lambda(R) = 1$.

Assume now that λ is nondegenerate. For $L \in \mathbb{R}$, define $\text{ECC}_*^L(Y, \lambda, \Gamma)$ to be the subspace of $\text{ECC}_*(Y, \lambda, \Gamma)$ spanned by ECH generators α with symplectic action $\mathcal{A}(\alpha) < L$. It follows from the definition of “ λ -compatible almost complex structure” that ∂_J maps ECC_*^L to itself; see [30, Section 1.4]. We define the *filtered ECH* to be the homology of this subcomplex, which we denote by $\text{ECH}_*^L(Y, \lambda, \Gamma)$. The inclusion of chain complexes induces a map

$$\iota_L : \text{ECH}_*^L(Y, \lambda, \Gamma) \rightarrow \text{ECH}_*(Y, \xi, \Gamma).$$

It is shown in [34, Theorem 1.3] that the filtered homology $\text{ECH}_*^L(Y, \lambda, \Gamma)$ and the map ι_L do not depend on the choice of J . However, unlike the usual ECH, filtered ECH does depend on the contact form λ and not just on the contact structure ξ .

Definition 2.17 [29] If $0 \neq \sigma \in \text{ECH}_*(Y, \xi, \Gamma)$, define the *spectral invariant*

$$c_\sigma(Y, \lambda) = \inf\{L \mid \sigma \in \text{Im}(\iota_L)\} \in \mathbb{R}.$$

An equivalent definition is that $c_\sigma(Y, \lambda)$ is the minimum L such that the class σ can be represented by a cycle in the chain complex $(\text{ECC}_*(Y, \lambda, \Gamma), \partial_J)$ which is a sum of ECH generators each having symplectic action $\leq L$. In particular, by definition, $c_\sigma(Y, \lambda) = \mathcal{A}(\alpha)$ for some ECH generator α with $[\alpha] = \Gamma$.

We can change the contact form λ , without changing the contact structure ξ , by multiplying λ by a smooth function $f: Y \rightarrow \mathbb{R}^{>0}$. As explained in [11, Section 2.5], it turns out that even when λ is degenerate, one can still define $c_\sigma(Y, \lambda)$ as a limit of spectral invariants $c_\sigma(Y, f_n \lambda)$ where $f_n \lambda$ is nondegenerate and $f_n \rightarrow 1$ in C^0 .

These spectral invariants have the following important properties:

Proposition 2.18 *Let Y be a closed connected three-manifold, and let λ be a (possibly degenerate) contact form on Y . Then:*

(a) *If $0 \neq \sigma \in \text{ECH}_*(Y, \xi, \Gamma)$, then*

$$c_\sigma(Y, \lambda) = \mathcal{A}(\alpha)$$

for some orbit set α with $[\alpha] = \Gamma$.

(b) *If $\sigma \in \text{ECH}_*(Y, \xi, \Gamma)$ and $U\sigma \neq 0$, then*

$$(2-11) \quad c_{U\sigma}(Y, \lambda) \leq c_\sigma(Y, \lambda).$$

If there are only finitely many simple Reeb orbits, then the inequality (2-11) is strict.

(c) **Volume property** *If $c_1(\xi) + 2 \text{PD}(\Gamma) \in H^2(Y; \mathbb{Z})$ is torsion, and if $\{\sigma_k\}_{k \geq 1}$ is a U -sequence for Γ , then*

$$\lim_{k \rightarrow \infty} \frac{c_{\sigma_k}(Y, \lambda)^2}{k} = 2 \text{vol}(Y, \lambda).$$

Proof As noted above, part (a) holds by definition when λ is nondegenerate, and in the degenerate case it follows from a compactness argument for Reeb orbits; see [11, Lemma 3.1(a)].

If λ is nondegenerate, then since the chain map $U_{J,z}$ counts J -holomorphic curves it decreases symplectic action like the differential, so strict inequality in (2-11) holds. The not necessarily strict inequality (2-11) in the degenerate case follows by a limiting argument. The fact that (2-11) is strict for degenerate contact forms with only finitely many simple Reeb orbits⁶ is proved by a more subtle compactness argument for holomorphic curves in [11, Lemma 3.1(b)].

Part (c), the most nontrivial part, is a special case of [13, Theorem 1.3]. □

⁶The equality $c_{U\sigma}(Y, \lambda) = c_\sigma(Y, \lambda)$ is possible for degenerate contact forms with infinitely many simple Reeb orbits. This happens, for example, for some classes σ when Y is an ellipsoid $\partial E(a, b)$ with a/b rational.

3 The ECH index and perturbations

The goal of this section is to prove Proposition 3.1 below, which gives an upper bound on how much the ECH index can change when one perturbs the contact form. This is an important ingredient in the proof of Theorems 1.2 and 1.5.

To state the proposition, let λ be a contact form on a closed three-manifold Y , and let $\lambda_n = f_n \lambda$ be a sequence of contact forms with $f_n \rightarrow 1$ in C^2 . In the case of interest, λ will be degenerate, while each of the contact forms λ_n will be nondegenerate.

Fix an orbit set $\alpha = \{(\alpha_i, m_i)\}$ for λ , and let N be a disjoint union of tubular neighborhoods N_i of the simple Reeb orbits α_i . Consider a sequence of orbit sets $\alpha(n)$ for λ_n that converges to α as currents. In particular this implies that if n is sufficiently large, and if we write $\alpha' = \alpha(n)$, then α' is contained in N , and its intersection with N_i is homologous in N_i to $m_i \alpha_i$. There is then a unique $W_\alpha \in H_2(Y, \alpha', \alpha)$ that is contained in N .

Likewise fix an orbit set $\beta = \{(\beta_j, n_j)\}$ for λ along with disjoint tubular neighborhoods of the simple Reeb orbits β_j , and consider a sequence of orbit sets $\beta(n)$ for λ_n that converges to β as currents. Then, for k sufficiently large, writing $\beta' = \beta(n)$, we obtain a distinguished $W_\beta \in H_2(Y, \beta', \beta)$.

For fixed large n there is now a bijection

$$H_2(Y, \alpha, \beta) \simeq H_2(Y, \alpha', \beta')$$

sending $Z \in H_2(Y, \alpha, \beta)$ to

$$Z' = Z + W_\alpha - W_\beta \in H_2(Y, \alpha', \beta').$$

Proposition 3.1 *With the notation as above, for fixed orbit sets α and β , if n is sufficiently large, then*

$$|I(\alpha, \beta, Z) - I(\alpha', \beta', Z')| \leq 2 \left(\sum_i m_i + \sum_j n_j \right).$$

Here $I(\alpha, \beta, Z)$ denotes the ECH index for λ , and $I(\alpha', \beta', Z')$ denotes the ECH index for λ_n .

3.1 Reduction to a local statement

We now reduce Proposition 3.1 to a local statement, Proposition 3.3, below.

Let γ be an oriented knot in Y , and let N be a tubular neighborhood of γ with an identification $N \simeq S^1 \times D^2$. By a “braid in N with d strands”, we mean an oriented

knot in N which is positively transverse to the D^2 fibers and which intersects each fiber d times.

Definition 3.2 • A *weighted braid* in N with m strands is a finite set of pairs $\zeta = \{(\zeta_i, m_i)\}$ where the ζ_i are disjoint braids in N with d_i strands, the m_i are positive integers and $\sum_i m_i d_i = m$.

• If τ is an oriented trivialization of the normal bundle of γ , then for $i \neq j$ there is a well-defined linking number $\ell_\tau(\zeta_i, \zeta_j) \in \mathbb{Z}$, as discussed in Section 2.1. Similarly, for each i there is a well-defined writhe $w_\tau(\gamma_i) \in \mathbb{Z}$; see [27, Section 2.6]. Define the *writhe* of the weighted braid ζ by

$$(3-1) \quad w_\tau(\zeta) = \sum_i m_i^2 w_\tau(\zeta_i) + \sum_{i \neq j} m_i m_j \ell_\tau(\zeta_i, \zeta_j).$$

Suppose now that γ is a simple Reeb orbit for λ , and that the normal bundle identification $N \simeq S^1 \times D^2$ above is chosen so that the Reeb vector field for λ is transverse to the D^2 fibers. If $\lambda' = f\lambda$ with f sufficiently C^2 close to 1, then the Reeb vector field for λ' in N is also transverse to the D^2 fibers. Suppose that this is the case.

Let $\gamma' = \{(\gamma'_k, m_k)\}$ be an orbit set for λ' which is contained in N . We can regard γ' as a weighted braid with m strands for some positive integer m . Also note that a trivialization τ of $\gamma^*\xi$ extends to a trivialization of ξ over the entire tubular neighborhood N , and thus canonically induces a homotopy class of trivialization τ' of ξ over the Reeb orbits γ'_k . We can now state:

Proposition 3.3 *With the notation as above, if λ' is sufficiently C^2 close to λ and if γ' is sufficiently close to $m\gamma$ as a current, then*

$$(3-2) \quad \left| -w_\tau(\gamma') - \text{CZ}_{\tau'}^I(\gamma') + \sum_{l=1}^m \text{CZ}_\tau(\gamma^l) \right| \leq 2m.$$

Proof of Proposition 3.1 assuming Proposition 3.3 By shrinking the tubular neighborhoods, we can assume without loss of generality that the chosen tubular neighborhood of each orbit α_i or β_j has an identification with $S^1 \times D^2$, in which the Reeb flow of λ is transverse to the D^2 fibers.

In the orbit set α' , each pair (α_i, m_i) gets replaced by an orbit set α'_i which represents a weighted braid with m_i strands in the tubular neighborhood of α_i . Likewise, each pair (β_j, n_j) gets replaced by an orbit set β'_j which represents a weighted braid with n_j strands in the tubular neighborhood of β_j . Let τ be a homotopy class of symplectic

trivializations of ξ over the Reeb orbits α_i and β_j . As in Proposition 3.3, this canonically induces a homotopy class of symplectic trivializations τ' over the Reeb orbits in the orbit sets α'_i and β'_j .

Because τ and τ' extend to a trivialization of ξ over the tubular neighborhoods containing W_α and W_β , it follows from the definition of the relative first Chern class that

$$(3-3) \quad c_{\tau'}(\alpha', \beta', Z') = c_\tau(\alpha, \beta, Z).$$

By Proposition 3.3, if n is sufficiently large, then

$$(3-4) \quad \begin{aligned} \left| -w_\tau(\alpha'_i) - CZ_{\tau'}^I(\alpha'_i) + \sum_{k=1}^{m_i} CZ_\tau(\alpha_i^k) \right| &\leq 2m_i, \\ \left| -w_\tau(\beta'_j) - CZ_{\tau'}^I(\beta'_j) + \sum_{l=1}^{n_j} CZ_\tau(\beta_j^l) \right| &\leq 2n_j. \end{aligned}$$

By (2-6), (3-3) and (3-4), to complete the proof of Proposition 3.1 it is enough to show

$$(3-5) \quad Q_{\tau'}(\alpha', \beta', Z') = Q_\tau(\alpha, \beta, Z) + \sum_i w_\tau(\alpha'_i) - \sum_j w_\tau(\beta'_j).$$

To prove (3-5), by Remark 2.6 it is enough to show

$$Q_\tau(\alpha', \alpha, W_\alpha) = \sum_i w_\tau(\alpha'_i), \quad Q_\tau(\beta', \beta, W_\beta) = \sum_j w_\tau(\beta'_j).$$

Since the chosen tubular neighborhoods of the Reeb orbits of α_i are disjoint, and the chosen tubular neighborhoods of the Reeb orbits of β_j are disjoint, the above equations follow from Lemma 3.4. $\square$

Lemma 3.4 *Let $\zeta = \{(\zeta_i, m_i)\}$ be a weighted braid with m strands as in Definition 3.2. Let W be the unique relative homology class in $H_2(N, \zeta, (\gamma, m))$. Then*

$$(3-6) \quad Q_\tau(\zeta, (\gamma, m), W) = w_\tau(\zeta).$$

Here τ defines a trivialization of the vertical tangent bundle of $N \rightarrow \gamma$ which then induces a trivialization of the normal bundle of each braid ζ_i .

Proof We can make an admissible representative S for W — see Definition 2.4 — whose intersection with $\{1 - \epsilon\} \times N$ consists of m_i parallel (with respect to τ) copies of each ζ_i , and which shrinks radially towards γ as the $[-1, 1]$ coordinate on $[-1, 1] \times N$ goes down to -1 . We can make another such admissible representative S' , disjoint from S , whose intersection with $\{1 - \epsilon\} \times N$ is parallel to the first and which likewise shrinks radially towards γ . Then in (2-1), the intersection number term vanishes. The

first linking number term in (2-1) also vanishes, as it is a sum of linking numbers of braids in neighborhoods of the ξ_i ; for each i , the braid from S and the braid from S' , with respect to τ , are trivial and parallel, and thus have linking number zero. The second linking number term in (2-1) is a linking number in a neighborhood of γ and equals $w_\tau(\zeta)$. $\square$

3.2 The structure of the braids

To prove Proposition 3.3, let γ be a simple Reeb orbit of λ , let N be a tubular neighborhood of γ as in Definition 3.2, and let τ be a trivialization of $\gamma^*\xi$. Let θ denote the rotation number $\theta_\tau(\gamma) \in \mathbb{R}$.

Suppose first that θ is irrational. Then the Reeb orbit γ and all of its covers are nondegenerate. Consequently, when λ' is sufficiently C^2 close to λ , there is a unique Reeb orbit γ'_0 for λ' close (as a current) to γ , and for n large the only possibility for the orbit set γ' is that it is the singleton set $\gamma' = \{(\gamma'_0, m)\}$. In this case Proposition 3.3 holds because $w_\tau(\gamma') = 0$ and the left-hand side of (3-2) is zero.

The nontrivial case of Proposition 3.3 is when the rotation number θ is rational. In this case we need to investigate the braids that can arise in γ' . The idea in what follows is to first analyze the case where the rotation number is an integer, and then reduce the general case to this one by taking an appropriate cover of a neighborhood of γ .

We start with the case where the rotation number is an integer. Here the picture is simple: each braid has just one strand, and the linking number of any two braids is given by the rotation number. More precisely:

Lemma 3.5 *With the above notation, suppose that the rotation number θ is an integer a . Let $\lambda_n = f_n\lambda$ with $f_n \rightarrow 1$ in C^2 . Then:*

- (a) *For a fixed positive integer d , if $\{\alpha_n\}$ is a sequence where each α_n is a simple Reeb orbit for λ_n in N which is a braid with d strands, with α_n converging as currents to $d\gamma$ as $n \rightarrow \infty$, then $d = 1$, and in particular the writhe $w_\tau(\alpha_n)$ equals 0 for n large enough.*
- (b) *Given two sequences of simple Reeb orbits $\{\alpha_n\}$ and $\{\beta_n\}$ as in (a) with $\alpha_n \neq \beta_n$ for each n , if n is sufficiently large, then the linking number $\ell_\tau(\alpha_n, \beta_n)$ equals a .*

This lemma is proved in Section 3.3 below. We now consider the case where the rotation number is a rational number a/b that is not an integer. Here there is a similarly nice

picture: each new simple Reeb orbit that can appear can be treated, for our purposes, like an (a, b) torus braid; see also Remark 3.7. More precisely:

Lemma 3.6 *With the above notation, suppose that the rotation number is $\theta = a/b$, where a and b are relatively prime integers with $b > 1$. Let $\lambda_n = f_n \lambda$ with $f_n \rightarrow 1$ in C^2 . Then:*

- (a) *For n sufficiently large, there is a unique simple Reeb orbit γ'_0 for λ_n that is close to γ as a current.*
- (b) *For a fixed integer $d > 1$, if $\{\alpha_n\}$ is a sequence where each α_n is a simple Reeb orbit for λ_n in N which is a braid with d strands, with α_n converging as currents to $d\gamma$ as $n \rightarrow \infty$, then $d = b$, the writhe $w_\tau(\alpha_n)$ equals $a(b-1)$, and the linking number $\ell_\tau(\gamma'_0, \alpha_n)$ equals a .*
- (c) *Given two sequences of Reeb orbits $\{\alpha_n\}$ and $\{\beta_n\}$ as in (b) with $\alpha_n \neq \beta_n$ for each n , if n is sufficiently large, then the linking number $\ell_\tau(\alpha_n, \beta_n)$ equals ab .*

Remark 3.7 In Lemma 3.6(b), we expect that one can further show that if n is sufficiently large then α_n is an (a, b) torus braid around γ'_0 ; however, we do not need this.

Proof of Lemma 3.6 assuming Lemma 3.5 Part (a) holds because the Reeb orbit γ is nondegenerate.

To prove part (b), we first note that, by the same argument as for (a), we must have that $d \geq b$, because for $0 < d < b$ the d^{th} iterate of γ has rotation number $da/b \notin \mathbb{Z}$.

Now let $\tilde{N}$ denote the b -fold cyclic cover of the tubular neighborhood N , with the pullback of the contact form λ_n . There is a unique simple Reeb orbit $\tilde{\gamma}'_0$ in $\tilde{N}$ whose projection to N is a b -fold cover of γ'_0 . In addition, by lifting the Reeb orbit α_n to a Reeb trajectory in $\tilde{N}$ and extending it by the Reeb flow if needed, we obtain a simple Reeb orbit $\tilde{\alpha}_n$ in $\tilde{N}$ whose projection to N is a cover of α_n . By Lemma 3.5(a), if n is sufficiently large, then $\tilde{\alpha}_n$ is a braid with one strand in $\tilde{N}$, hence α_n has at most b strands. Thus, $d = b$. By Lemma 3.5(b) we have $\ell_\tau(\tilde{\gamma}'_0, \tilde{\alpha}_n) = a$ in $\tilde{N}$, and it follows that $\ell_\tau(\gamma'_0, \alpha_n) = a$.

We now compute the writhe $w_\tau(\alpha_n)$. There are b possibilities for the Reeb orbit $\tilde{\alpha}_n$ in the previous paragraph, which we denote by η_l for $l \in \mathbb{Z}/b$, ordered so that the $\mathbb{Z}/b$ action on $\tilde{N}$ by deck transformations sends η_l to η_{l+1} . The writhe $w_\tau(\alpha_n)$ is a signed count of crossings of two strands of α_n . Each such crossing corresponds to a crossing of some η_l with some $\eta_{l'}$ for $l \neq l'$, as well as crossings of η_{l+p} with $\eta_{l'+p}$

for $p = 1, \dots, b-1$, obtained from the first crossing by deck transformations. On the other hand, the linking number of η_l with $\eta_{l'}$ is one half the signed count of crossings of η_l with $\eta_{l'}$. Thus we obtain

$$w_\tau(\alpha_n) = \frac{1}{b} \sum_{l \neq l'} \ell_\tau(\eta_l, \eta_{l'}) = \frac{1}{b} b(b-1)a = a(b-1).$$

Here we are using Lemma 3.5(b) to get that $\ell_\tau(\eta_l, \eta_{l'}) = a$ when $l \neq l'$.

We now prove (c). Similarly to the previous calculation, each crossing counted by the linking number $\ell_\tau(\alpha_n, \beta_n)$ corresponds to b crossings of some lift of α_n (extended to a simple Reeb orbit) with some lift of β_n (extended to a simple Reeb orbit). Thus the linking number we want is $1/b$ times the sum of the linking number of each of the b extended lifts of α_n with each of the b extended lifts of β_n , which is $(1/b)b^2a = ab$. $\square$

Proof of Proposition 3.3 As explained above, we can assume that $\theta = a/b$, where a and b are relatively prime integers with $b > 0$. When $a/b \notin \mathbb{Z}$, the orbit set γ' consists of the orbit γ'_0 from Lemma 3.6(a) with multiplicity m_0 for some $m_0 \geq 0$, together with orbits γ'_k for $k \neq 0$ with multiplicities $m_k > 0$. When $a/b \in \mathbb{Z}$ the same is true except that we do not necessarily have a unique γ'_0 and we can take $m_0 = 0$. Since each γ'_k for $k \neq 0$ is close to a b -fold cover of γ , we have

$$(3-7) \quad m_0 + b \sum_{k \neq 0} m_k = m.$$

By (3-1) and Lemmas 3.5 and 3.6, if λ' is sufficiently C^2 close to λ and if γ' is sufficiently close to $m\gamma$ as a current, then

$$(3-8) \quad w_\tau(\gamma') = a(b-1) \sum_{k \neq 0} m_k^2 + 2am_0 \sum_{k \neq 0} m_k + ab \sum_{0 \neq k \neq k' \neq 0} m_k m'_{k'}.$$

Now we consider Conley–Zehnder indices. By (2-5),

$$(3-9) \quad \text{CZ}_{\tau'}^{I'}(\gamma') = \sum_{l=1}^{m_0} \text{CZ}_{\tau'}((\gamma'_0)^l) + \sum_{k \neq 0} \sum_{l=1}^{m_k} \text{CZ}_{\tau'}((\gamma'_k)^l).$$

For a positive integer $l \leq m$, if λ' is sufficiently close to λ , then with respect to τ the Reeb orbit $(\gamma'_0)^l$ has rotation number close to $(a/b)l$, and each Reeb orbit $(\gamma'_k)^l$ for $k \neq 0$ has rotation number close to al . Then, by (2-4) and (3-9),

$$\left| \text{CZ}_{\tau'}^{I'}(\gamma') - \sum_{l=1}^{m_0} \frac{2al}{b} - \sum_{k \neq 0} \sum_{l=1}^{m_k} 2al \right| \leq m_0 + \sum_{k \neq 0} m_k.$$

It follows from this and (3-7) that

$$(3-10) \quad \left| \text{CZ}_{\tau'}^I(\gamma') - \frac{a}{b}(m_0^2 + m_0) - a \sum_{k \neq 0} (m_k^2 + m_k) \right| \leq m.$$

Finally, by (2-4),

$$\left| \sum_{l=1}^m \text{CZ}_{\tau}(\gamma^l) - \frac{a}{b}(m^2 + m) \right| \leq m.$$

Then, by (3-7),

$$(3-11) \quad \left| \sum_{l=1}^m \text{CZ}_{\tau}(\gamma^l) - \frac{a}{b}(m_0^2 + m_0) - a(2m_0 + 1) \sum_{k \neq 0} m_k - ab \left(\sum_{k \neq 0} m_k \right)^2 \right| \leq m.$$

Combining (3-8), (3-10) and (3-11) gives the desired estimate (3-2). $\square$

3.3 Perturbations of degenerate flows

To conclude the proof of Proposition 3.3 we now prove Lemma 3.5.

As in the statement of the lemma, let γ be a simple Reeb orbit of λ of period T , and let $\lambda_n = f_n \lambda$, where $f_n \rightarrow 1$ in C^2 . Let ϕ^t and ϕ_n^t denote the time t flows of the Reeb vector fields for λ and λ_n , respectively. Let $p \in \gamma$, and let $P_{\gamma}: \xi_p \rightarrow \xi_p$ denote the linearized return map (1-2).

Lemma 3.8 *Let $\{(p_n, T_n)\}_{n=1, \dots}$ be a sequence in $Y \times (0, \infty)$ satisfying:*

- (c1) $\phi_n^{T_n}(p_n) = p_n \rightarrow p$.
- (c2) $\phi_n^{T_n/j}(p_n) \neq p_n$ for all integers $j \geq 2$ and all n .
- (c3) $T_n \rightarrow T_{\infty} \in [0, \infty)$.

Then one of the following alternatives holds:

- (a1) $T_{\infty} = T$.
- (a2) $T_{\infty} = Td$ for some integer $d \geq 2$, and the eigenvalues of P_{γ} that are roots of unity of degree d generate multiplicatively all roots of unity of order d .

Proof This is a special case of a result of Bangert [3, Proposition 1] for C^1 flows. $\square$

In the situation of Lemma 3.5, more can be said:

Corollary 3.9 *Suppose that the eigenvalues of P_{γ} are real and positive. Let $\{(p_n, T_n)\}$ be a sequence satisfying conditions (c1), (c2) and (c3) of Lemma 3.8. Then alternative (a2) does not hold.*

Proof The only root of unity that can be an eigenvalue of P_γ is 1, hence the set of eigenvalues of P_γ does not generate multiplicatively the group of roots of unity of order d when $d \geq 2$. $\square$

Proof of Lemma 3.5 Part (a) follows from Corollary 3.9.

To prove part (b), fix a diffeomorphism Φ from the tubular neighborhood N of γ to $(\mathbb{R}/T\mathbb{Z}) \times \mathbb{C}$ such that γ corresponds to $(\mathbb{R}/T\mathbb{Z}) \times \{0\}$, the Reeb vector field R_n of λ_n is transverse to the $\mathbb{C}$ fibers for n sufficiently large (assume that n is this large below), and the derivative of Φ in the normal direction along γ agrees with the trivialization τ . We omit the diffeomorphism Φ from the notation below and write points in N using the coordinates $(t, z) \in (\mathbb{R}/T\mathbb{Z}) \times \mathbb{C}$.

By part (a), by taking n large enough we can assume that α_n and β_n have the same period as γ . After reparametrization, the Reeb orbit α_n is given by a map

$$\mathbb{R}/T\mathbb{Z} \rightarrow (\mathbb{R}/T\mathbb{Z}) \times \mathbb{C}, \quad t \mapsto (t, \hat{\alpha}_n(t)),$$

where $\hat{\alpha}_n: \mathbb{R}/T\mathbb{Z} \rightarrow \mathbb{C}$. Likewise the Reeb orbit β_n is given by a map $\hat{\beta}_n: \mathbb{R}/T\mathbb{Z} \rightarrow \mathbb{C}$. We have

$$(3-12) \quad \ell_\tau(\alpha_n, \beta_n) = \text{wind}(\hat{\alpha}_n - \hat{\beta}_n),$$

where the right-hand side denotes the winding number of the loop

$$\hat{\alpha}_n - \hat{\beta}_n: \mathbb{R}/T\mathbb{Z} \rightarrow \mathbb{C}^*.$$

We now compute the right-hand side of (3-12). There is a convex neighborhood U of 0 in $\mathbb{C}$ such that, if n is sufficiently large (which we assume below), then the following two conditions hold: First, $\hat{\alpha}_n(0), \hat{\beta}_n(0) \in U$. Second, for each $t \in [0, T]$ there is a well-defined map $\psi_n^t: U \rightarrow \mathbb{C}$ such that, for $z \in U$, the flow of the Reeb vector field R_n starting at $(0, z)$ first hits $\{t\} \times \mathbb{C}$ at the point $(t, \psi_n^t(z))$. In particular, it follows from the definition that

$$(3-13) \quad \hat{\alpha}_n(t) = \psi_n^t(\hat{\alpha}_n(0)), \quad \hat{\beta}_n(t) = \psi_n^t(\hat{\beta}_n(0)).$$

Now consider the derivative of ψ_n^t , which we denote by

$$D\psi_n^t: U \times \mathbb{C} \rightarrow \mathbb{C}.$$

By (3-13), we may apply the fundamental theorem of calculus to the function

$$s \mapsto \psi_n^t(s\hat{\alpha}_n(0) + (1-s)\hat{\beta}_n(0))$$

to obtain

$$(3-14) \quad \hat{\alpha}_n(t) - \hat{\beta}_n(t) = \int_0^1 D\psi_n^t(s\hat{\alpha}_n(0) + (1-s)\hat{\beta}_n(0), \hat{\alpha}_n(0) - \hat{\beta}_n(0)) ds.$$

By the convergence of λ_n , if n is sufficiently large (which we assume below), then the amount that $D\psi_n^t(s\hat{\alpha}_n(0) + (1-s)\hat{\beta}_n(0), \cdot)$ rotates any vector as compared to $D\psi_n^t(0, \cdot)$ can be made arbitrarily small. It follows that the integrand in (3-14), and hence $\hat{\alpha}_n(t) - \hat{\beta}_n(t)$, has positive inner product with $D\psi_n^t(0, \hat{\alpha}_n(0) - \hat{\beta}_n(0))$. Thus, the right-hand side of (3-12) differs by less than $\frac{1}{4}$ from the rotation number (the change in argument divided by 2π) of the path

$$(3-15) \quad [0, T] \rightarrow \mathbb{C}^*, \quad t \mapsto D\psi_n^t(0, \hat{\alpha}_n(0) - \hat{\beta}_n(0)).$$

The rotation number of the linearized Reeb flow along γ differs from the rotation number of any individual vector by less than $\frac{1}{2}$. Hence, by again applying convergence of the λ_n as above, if n is sufficiently large then the rotation number of the path (3-15) differs by less than $\frac{1}{2}$ from a . Since the right-hand side of (3-12) is an integer which differs by less than $\frac{3}{4}$ from a , it must equal a . $\square$

4 Two simple Reeb orbits implies nondegenerate

We now prove Theorem 1.2. Throughout this section assume that Y is a closed connected three-manifold and λ is a contact form on Y with exactly two simple Reeb orbits γ_1 and γ_2 of periods T_1 and T_2 , respectively.

4.1 The homology classes of the Reeb orbits

Lemma 4.1 *The classes $[\gamma_i] \in H_1(Y)$ and $c_1(\xi) \in H^2(Y; \mathbb{Z})$ are torsion.*

Proof We use a similar argument to the proof of [12, Theorem 1.7].

Since every oriented three-manifold is spin, we can choose $\Gamma \in H_1(Y)$ such that $c_1(\xi) + 2\text{PD}(\Gamma) = 0 \in H^2(Y; \mathbb{Z})$. By Proposition 2.16, there exists a U -sequence $\{\sigma_k\}_{\geq 1}$ for Γ . Write $c_k = c_{\sigma_k}(Y, \lambda) \in \mathbb{R}$.

By Proposition 2.18(a),

$$c_k = m_{1,k}T_1 + m_{2,k}T_2$$

for some nonnegative integers $m_{1,k}$ and $m_{2,k}$, and furthermore

$$(4-1) \quad m_{1,k}[\gamma_1] + m_{2,k}[\gamma_2] = \Gamma \in H_1(Y).$$

By Proposition 2.18(b), the sequence $\{c_k\}$ is strictly increasing. It then follows from (4-1) that there are infinitely many integral linear combinations of $[\gamma_1]$ and $[\gamma_2]$ that have the same value in $H_1(Y)$. Thus the kernel of the map

$$(4-2) \quad \mathbb{Z}^2 \rightarrow H_1(Y), \quad (m_1, m_2) \mapsto m_1[\gamma_1] + m_2[\gamma_2],$$

has rank at least 1.

In fact, the kernel of the map (4-2) must have rank at least 2; otherwise c_k would grow at least linearly in k , contradicting the sublinear growth in the volume property in Proposition 2.18(c). It follows that $[\gamma_1]$ and $[\gamma_2]$ are torsion. Since $c_1(\xi) + 2 \text{PD}(\Gamma) = 0$, we deduce that $c_1(\xi)$ is also torsion. $\square$

4.2 Computing the ECH index

If m_1 and m_2 are nonnegative integers, we use the notation $\gamma_1^{m_1} \gamma_2^{m_2}$ to indicate the orbit set $\{(\gamma_1, m_1), (\gamma_2, m_2)\}$, with the element (γ_i, m_i) omitted when $m_i = 0$. Write $\alpha = \gamma_1^{m_1} \gamma_2^{m_2}$. If $[\alpha] = 0$, then it follows from Remark 2.13 and Lemma 4.1 that $I(\alpha) \in \mathbb{Z}$ is defined. We now give an explicit computation of $I(\alpha)$, following [33, Section 4.7].

Definition 4.2 Define the *linking number*

$$(4-3) \quad \ell(\gamma_1, \gamma_2) := \frac{\ell(\gamma_1^{l_1}, \gamma_2^{l_2})}{l_1 l_2} \in \mathbb{Q},$$

where l_1 and l_2 are positive integers such that $l_i[\gamma_i] = 0 \in H_1(Y)$, and on the right-hand side ℓ denotes the usual integer-valued linking number of disjoint nullhomologous loops.

Definition 4.3 For $i = 1, 2$, define the *Seifert rotation number* $\phi_i \in \mathbb{R}$ as follows. Let τ be a trivialization of ξ over γ_i . Let $\theta_{i,\tau} = \theta_\tau(\gamma_i) \in \mathbb{R}$ denote the rotation number of γ_i with respect to τ . Let l_i be a positive integer such that $l_i[\gamma_i] = 0$. Define

$$(4-4) \quad Q_{i,\tau} := \frac{Q_\tau(\gamma_i^{l_i})}{l_i^2} \in \mathbb{Q},$$

where $Q_\tau(\gamma_i^{l_i})$ is shorthand for $Q_\tau(\gamma_i^{l_i}, \emptyset, Z)$ for any $Z \in H_2(Y, \gamma_i^{l_i}, \emptyset)$. Note that $Q_{i,\tau}$ does not depend on Z by (2-3), and it does not depend on l_i either because Q_τ is quadratic in the relative homology class. Finally, define

$$(4-5) \quad \phi_i := Q_{i,\tau} + \theta_{i,\tau} \in \mathbb{R}.$$

The number ϕ_i does not depend on the choice of trivialization τ by the change of trivialization formulas in [27, Section 2].

Remark 4.4 When γ_i is nullhomologous, one can alternatively describe ϕ_i as follows. Let Σ be a Seifert surface spanned by γ_i . There is a distinguished homotopy class of trivialization τ' of ξ over γ_i , the “Seifert framing”, for which the normal vector to Σ has winding number zero around γ_i . We have $Q_{i,\tau'} = 0$ by [27, Lemma 3.10]. It then follows that $\phi_i = \theta_{\tau'}(\gamma_i)$. In the general case when γ_i is rationally nullhomologous, one can similarly describe ϕ_i as the rotation number with respect to a rational framing of γ_i determined by a rational Seifert surface.

Lemma 4.5 If $m_1[\gamma_1] + m_2[\gamma_2] = 0 \in H_1(Y)$, then

$$(4-6) \quad I(\gamma_1^{m_1} \gamma_2^{m_2}) = \phi_1 m_1^2 + \phi_2 m_2^2 + 2\ell(\gamma_1, \gamma_2) m_1 m_2 + O(m_1 + m_2).$$

Proof Let l_i be a positive integer with $l_i[\gamma_i] = 0$. Similarly to (4-4), define

$$c_{i,\tau} := \frac{c_\tau(\gamma_i^{l_i})}{l_i} \in \mathbb{Q},$$

where $c_\tau(\gamma_i^{l_i})$ is shorthand for $c_\tau(\gamma_i^{l_i}, \emptyset, Z)$ for any $Z \in H_2(Y, \gamma_i^{l_i}, \emptyset)$. Then $c_{i,\tau}$ does not depend on Z by (2-2) since $c_1(\xi)$ is torsion, and it is independent of the choice of l_i because c_τ is linear in the relative homology class Z .

It follows from the definition of the ECH index and the facts that c_τ and Q_τ are linear and quadratic in the relative homology class (see [33, Section 4.2]) that

$$I(\gamma_1^{m_1} \gamma_2^{m_2}) = \sum_{i=1}^2 (m_i c_{i,\tau} + m_i^2 Q_{i,\tau}) + 2m_1 m_2 \ell(\gamma_1, \gamma_2) + \sum_{i=1}^2 \sum_{k=1}^{m_i} ([k\theta_{i,\tau}] + [k\theta_{i,\tau}]),$$

where $Q_{i,\tau}$ and $\theta_{i,\tau}$ are as in (4-5). Plugging in the approximation

$$\sum_{i=1}^2 \sum_{k=1}^{m_i} ([k\theta_{i,\tau}] + [k\theta_{i,\tau}]) = \sum_{i=1}^2 m_i^2 \theta_{i,\tau} + O(m_1 + m_2)$$

then gives (4-6). □

4.3 Using the volume property

Lemma 4.6 The Seifert rotation numbers and linking number are given by

$$\phi_i = \frac{T_i^2}{\text{vol}(Y, \lambda)}, \quad \ell(\gamma_1, \gamma_2) = \frac{T_1 T_2}{\text{vol}(Y, \lambda)}.$$

Proof Both sides of the above equations are invariant under scaling the contact form by a positive constant, so we may assume without loss of generality that $\text{vol}(Y, \lambda) = 1$.

By Proposition 2.16 and Lemma 4.1, we can choose a U -sequence $\{\sigma_k\}_{k \geq 1}$ for $\Gamma = 0$. Since the U map has degree -2 , there is a constant $C \in \mathbb{Z}$ such that, for each positive integer k , the class σ_k has grading $C + 2k$. By Proposition 2.18(a), for each positive integer k there are nonnegative integers $m_{1,k}$ and $m_{2,k}$ such that

$$(4-7) \quad c_{\sigma_k}(Y, \lambda) = m_{1,k}T_1 + m_{2,k}T_2.$$

By the volume property of Proposition 2.18(c),

$$(4-8) \quad 2k = (m_{1,k}T_1 + m_{2,k}T_2)^2 + o(k).$$

Fix k and write $\alpha_k = \gamma_1^{m_{1,k}} \gamma_2^{m_{2,k}}$. If λ' is a sufficiently C^2 close nondegenerate perturbation of λ , then by the same compactness argument that proves Proposition 2.18(a) there is an orbit set α'_k close to α_k as a current such that $I(\alpha'_k) = C + 2k$ (and also $\int_{\alpha'_k} \lambda'$ is close to $c_{\sigma_k}(Y, \lambda)$, although we do not need this). By Proposition 3.1,

$$C + 2k = I(\alpha_k) + O(m_{1,k} + m_{2,k}).$$

Combining this with Lemma 4.5, we get

$$(4-9) \quad 2k = \phi_1 m_{1,k}^2 + \phi_2 m_{2,k}^2 + 2\ell(\gamma_1, \gamma_2)m_{1,k}m_{2,k} + O(m_{1,k} + m_{2,k}).$$

Putting together (4-8) and (4-9),

$$\begin{aligned} (\phi_1 - T_1^2)m_{1,k}^2 + (\phi_2 - T_2^2)m_{2,k}^2 + 2(\ell(\gamma_1, \gamma_2) - T_1T_2)m_{1,k}m_{2,k} \\ = O(m_{1,k} + m_{2,k}) + o(k). \end{aligned}$$

Consequently, if the sequence $(m_{2,k}/m_{1,k})_{k \geq 1}$ has an accumulation point $S \in [0, \infty]$, then the line in the (x, y) -plane of slope S through the origin is in the null space of the quadratic form

$$f(x, y) = (\phi_1 - T_1^2)x^2 + (\phi_2 - T_2^2)y^2 + 2(\ell(\gamma_1, \gamma_2) - T_1T_2)xy.$$

To complete the proof of the lemma, it now suffices to show that the sequence $(m_{2,k}/m_{1,k})_{k \geq 1}$ has at least three accumulation points, as then the quadratic form f must vanish identically. We claim that in fact this sequence has infinitely many accumulation points.

If the sequence has only finitely many accumulation points $S_1, \dots, S_n$, then for every $\epsilon > 0$ there exists $R > 0$ such that every point $(m_{1,k}, m_{2,k})$ is contained in the union of the disk $x^2 + y^2 \leq R^2$ and the cones around the lines of slope $S_1, \dots, S_n$ with angular width ϵ .

Since $\lim_{k \rightarrow \infty} c_{\sigma_k}^2/k = 2$, and since the points $(m_{1,k}, m_{2,k})$ are pairwise distinct by Proposition 2.18(b), by (4-7) it follows that, for large L , the number of points $(m_{1,k}, m_{2,k})$ contained in the triangle $T_1x + T_2y \leq L$ for $x \geq 0$ and $y \geq 0$ is approximately $\frac{1}{2}L^2$. As a result, there exists $\delta > 0$ such that, for all L sufficiently large, the fraction of lattice points in the above triangle that are contained in the sequence $(m_{1,k}, m_{2,k})_{k \geq 1}$ is at least δ . This gives a contradiction if ϵ in the previous paragraph is chosen sufficiently small. $\square$

4.4 Completing the proof of nondegeneracy

Proof of Theorem 1.2 The ratio T_1/T_2 is irrational⁷ by [11, Theorem 1.3]. Also, $\ell(\gamma_1, \gamma_2)$ is rational by the definition (4-3). It then follows from Lemma 4.6 that ϕ_1 and ϕ_2 are irrational.

By (4-5), since $Q_{i,\tau}$ is rational, it follows that the rotation number $\theta_{i,\tau}$ is irrational. Then P_{γ_i} has eigenvalues $e^{\pm 2\pi i \theta_{i,\tau}}$, so the Reeb orbits γ_i are irrationally elliptic. As explained in Section 1.1, it follows that all covers of γ_i are nondegenerate, so λ is nondegenerate. $\square$

5 Additional dynamical information

To finish up, we now prove Theorem 1.5.

To prepare for the proof, recall that if Y is a closed oriented three-manifold, if ξ is a contact structure on Y with $c_1(\xi) = 0 \in H^2(Y; \mathbb{Z})$, and if γ is a nullhomologous transverse knot, then the *self-linking number* $\text{sl}(\gamma) \in \mathbb{Z}$ is defined to be the difference between the Seifert framing (see Remark 4.4) and the framing given by a global trivialization of ξ . In the notation of Section 4.2,

$$(5-1) \quad \text{sl}(\gamma) = Q_\tau(\gamma) - c_\tau(\gamma),$$

where τ is any trivialization of $\xi|_\gamma$.

Now suppose that γ above is a simple Reeb orbit. Let $\phi(\gamma) \in \mathbb{R}$ denote the rotation number of γ with respect to the Seifert framing as in Section 4.2, and let $\theta(\gamma) \in \mathbb{R}$ denote the rotation number of γ with respect to a global trivialization of ξ . Also, let

$$\text{CZ}(\gamma) = \lfloor \theta(\gamma) \rfloor + \lceil \theta(\gamma) \rceil \in \mathbb{Z}$$

⁷The proof is simple: if T_1/T_2 is rational, T_1 and T_2 are both integer multiples of a single number, so Proposition 2.18(b) implies that the spectral invariants associated to a U -sequence grow at least linearly, contradicting the volume property.

denote the Conley–Zehnder index of γ with respect to a global trivialization. It follows from (5-1) that

$$(5-2) \quad \phi(\gamma) = \theta(\gamma) + \text{sl}(\gamma).$$

Proof of Theorem 1.5 By Corollary 1.3, γ_1 and γ_2 are the core circles of a genus-one Heegaard splitting of Y . It follows from this topological description that $\ell(\gamma_1, \gamma_2) = 1/p$. Part (a) of the theorem then follows from Lemma 4.6.

To prove part (b), suppose first that $Y = S^3$. We know from Theorem 1.2 that λ is nondegenerate and there are no hyperbolic Reeb orbits. Then ξ is tight, because otherwise [21, Theorem 1.4] would give a hyperbolic Reeb orbit. Moreover, it follows from [25, Theorem 1.3], combined with [21, Theorem 1.4] and the fact that there are no Reeb orbits with $\text{CZ} = 2$ (since Reeb orbits with even Conley–Zehnder index have integer rotation number and thus are hyperbolic), that one of the simple Reeb orbits, say γ_1 , satisfies $\text{sl}(\gamma_1) = -1$ and $\text{CZ}(\gamma_1) = 3$, and is the binding of an open book decomposition with pages that are disk-like global surfaces of section for the Reeb flow. The return map on a page preserves an area form with finite total area, and hence it has a fixed point by Brouwer’s translation theorem. This fixed point corresponds to the simple Reeb orbit γ_2 , which is transverse to the pages of the open book. Since, on $S^3 \setminus \gamma_1$, the tangent spaces of the pages define a distribution that is isotopic to ξ keeping transversality with the Reeb direction, $\text{sl}(\gamma_2) = -1$. Since $\text{CZ}(\gamma_1) = 3$, we have $\theta(\gamma_1) \in (1, 2)$, so, by (5-2), $\phi_1 \in (0, 1)$. By Lemma 4.6 as used in (1-4), we have $\phi_1\phi_2 = 1$, so $\phi_2 > 1$. By (5-2), again, $\theta(\gamma_2) > 2$. It follows that all iterates of γ_1 and γ_2 have $\theta > 1$, so λ is dynamically convex.

To prove part (b) in the general case, let $\tilde{\lambda}$ denote the pullback of the contact form λ to the universal cover S^3 of Y . It follows from the Heegaard decomposition that γ_1 and γ_2 each have order p in $\pi_1(Y)$. Consequently $\tilde{\lambda}$ has exactly two simple Reeb orbits $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$, which project to γ_1 and γ_2 as p -fold coverings. By the previous paragraph, $(S^3, \tilde{\lambda})$ is dynamically convex and tight, and it follows that (Y, λ) is dynamically convex and universally tight. $\square$

References

- [1] **P Albers, H Geiges, K Zehmisch**, *Pseudorotations of the 2-disc and Reeb flows on the 3-sphere*, Ergodic Theory Dynam. Systems 42 (2022) 402–436 MR Zbl

- [2] **K Baker, J Etnyre**, *Rational linking and contact geometry*, from “Perspectives in analysis, geometry, and topology” (I Itenberg, B Jöricke, M Passare, editors), Progr. Math. 296, Springer (2012) 19–37 MR Zbl
- [3] **V Bangert**, *On the lengths of closed geodesics on almost round spheres*, Math. Z. 191 (1986) 549–558 MR Zbl
- [4] **D Bechara, Sr, U L Hryniewicz, P A S Salomão**, *On the relation between action and linking*, J. Mod. Dyn. 17 (2021) 319–336 MR Zbl
- [5] **F Bourgeois, K Cieliebak, T Ekholm**, *A note on Reeb dynamics on the tight 3-sphere*, J. Mod. Dyn. 1 (2007) 597–613 MR Zbl
- [6] **B Bramham**, *Periodic approximations of irrational pseudo-rotations using pseudoholomorphic curves*, Ann. of Math. 181 (2015) 1033–1086 MR Zbl
- [7] **E Çinel, V L Ginzburg, B Z Gürel**, *Pseudo-rotations and holomorphic curves*, Selecta Math. 26 (2020) art. id. 78 MR Zbl
- [8] **V Colin, P Dehornoy, A Rechtman**, *On the existence of supporting broken book decompositions for contact forms in dimension 3*, Invent. Math. 231 (2023) 1489–1539 MR Zbl
- [9] **B Collier, E Kerman, B M Reiniger, B Turmunkh, A Zimmer**, *A symplectic proof of a theorem of Franks*, Compos. Math. 148 (2012) 1969–1984 MR Zbl
- [10] **C R Cornwell**, *Berge duals and universally tight contact structures*, Topology Appl. 236 (2018) 26–43 MR Zbl
- [11] **D Cristofaro-Gardiner, M Hutchings**, *From one Reeb orbit to two*, J. Differential Geom. 102 (2016) 25–36 MR Zbl
- [12] **D Cristofaro-Gardiner, M Hutchings, D Pomerleano**, *Torsion contact forms in three dimensions have two or infinitely many Reeb orbits*, Geom. Topol. 23 (2019) 3601–3645 MR Zbl
- [13] **D Cristofaro-Gardiner, M Hutchings, V G B Ramos**, *The asymptotics of ECH capacities*, Invent. Math. 199 (2015) 187–214 MR Zbl
- [14] **J Etnyre, R Ghrist**, *Tight contact structures via dynamics*, Proc. Amer. Math. Soc. 127 (1999) 3697–3706 MR Zbl
- [15] **B Fayad, A Katok**, *Constructions in elliptic dynamics*, Ergodic Theory Dynam. Systems 24 (2004) 1477–1520 MR Zbl
- [16] **B Fayad, R Krikorian**, *Some questions around quasi-periodic dynamics*, from “Proceedings of the International Congress of Mathematicians” (B Sirakov, P N de Souza, M Viana, editors), volume 3, World Sci., Hackensack, NJ (2018) 1927–1950 MR Zbl
- [17] **J Franks**, *Geodesics on S^2 and periodic points of annulus homeomorphisms*, Invent. Math. 108 (1992) 403–418 MR Zbl
- [18] **V L Ginzburg, B Z Gürel**, *Hamiltonian pseudo-rotations of projective spaces*, Invent. Math. 214 (2018) 1081–1130 MR Zbl

- [19] **B Z Gürel**, *Perfect Reeb flows and action-index relations*, *Geom. Dedicata* 174 (2015) 105–120 MR Zbl
- [20] **H Hofer, K Wysocki, E Zehnder**, *A characterisation of the tight three-sphere*, *Duke Math. J.* 81 (1995) 159–226 MR Zbl
- [21] **H Hofer, K Wysocki, E Zehnder**, *Unknotted periodic orbits for Reeb flows on the three-sphere*, *Topol. Methods Nonlinear Anal.* 7 (1996) 219–244 MR Zbl
- [22] **H Hofer, K Wysocki, E Zehnder**, *A characterization of the tight 3-sphere, II*, *Comm. Pure Appl. Math.* 52 (1999) 1139–1177 MR Zbl
- [23] **K Honda**, *On the classification of tight contact structures, I*, *Geom. Topol.* 4 (2000) 309–368 MR Zbl
- [24] **U L Hryniewicz, J E Licata, P A S Salomão**, *A dynamical characterization of universally tight lens spaces*, *Proc. Lond. Math. Soc.* 110 (2015) 213–269 MR Zbl
- [25] **U Hryniewicz, P A S Salomão**, *On the existence of disk-like global sections for Reeb flows on the tight 3-sphere*, *Duke Math. J.* 160 (2011) 415–465 MR Zbl
- [26] **M Hutchings**, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, *J. Eur. Math. Soc.* 4 (2002) 313–361 MR Zbl
- [27] **M Hutchings**, *The embedded contact homology index revisited*, from “New perspectives and challenges in symplectic field theory” (M Abreu, F Lalonde, L Polterovich, editors), CRM Proc. Lecture Notes 49, Amer. Math. Soc., Providence, RI (2009) 263–297 MR Zbl
- [28] **M Hutchings**, *Taubes’s proof of the Weinstein conjecture in dimension three*, *Bull. Amer. Math. Soc.* 47 (2010) 73–125 MR Zbl
- [29] **M Hutchings**, *Quantitative embedded contact homology*, *J. Differential Geom.* 88 (2011) 231–266 MR Zbl
- [30] **M Hutchings**, *Lecture notes on embedded contact homology*, from “Contact and symplectic topology” (F Bourgeois, V Colin, A Stipsicz, editors), Bolyai Soc. Math. Stud. 26, János Bolyai Math. Soc., Budapest (2014) 389–484 MR Zbl
- [31] **M Hutchings, C H Taubes**, *Gluing pseudoholomorphic curves along branched covered cylinders, I*, *J. Symplectic Geom.* 5 (2007) 43–137 MR Zbl
- [32] **M Hutchings, C H Taubes**, *Gluing pseudoholomorphic curves along branched covered cylinders, II*, *J. Symplectic Geom.* 7 (2009) 29–133 MR Zbl
- [33] **M Hutchings, C H Taubes**, *The Weinstein conjecture for stable Hamiltonian structures*, *Geom. Topol.* 13 (2009) 901–941 MR Zbl
- [34] **M Hutchings, C H Taubes**, *Proof of the Arnold chord conjecture in three dimensions, II*, *Geom. Topol.* 17 (2013) 2601–2688 MR Zbl
- [35] **K Irie**, *Dense existence of periodic Reeb orbits and ECH spectral invariants*, *J. Mod. Dyn.* 9 (2015) 357–363 MR Zbl

- [36] **B Joly**, *The Calabi invariant for Hamiltonian diffeomorphisms of the unit disk*, preprint (2021) arXiv 2102.09352
- [37] **A B Katok**, *Ergodic perturbations of degenerate integrable Hamiltonian systems*, *Izv. Akad. Nauk SSSR Ser. Mat.* 37 (1973) 539–576 MR Zbl In Russian; translated in *Math. USSR Izv.* 7 (1973) 535–571
- [38] **P Kronheimer, T Mrowka**, *Monopoles and three-manifolds*, *New Math. Monogr.* 10, Cambridge Univ. Press (2007) MR Zbl
- [39] **F Le Roux, S Seyfaddini**, *The Anosov–Katok method and pseudo-rotations in symplectic dynamics*, *J. Fixed Point Theory Appl.* 24 (2022) art. id. 36 MR Zbl
- [40] **A Pirnapasov**, *Hutchings’ inequality for the Calabi invariant revisited with an application to pseudo-rotations*, preprint (2021) arXiv 2102.09533
- [41] **E Shelukhin**, *Pseudo-rotations and Steenrod squares*, *J. Mod. Dyn.* 16 (2020) 289–304 MR Zbl
- [42] **C H Taubes**, *The Seiberg–Witten equations and the Weinstein conjecture*, *Geom. Topol.* 11 (2007) 2117–2202 MR Zbl
- [43] **C H Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, I*, *Geom. Topol.* 14 (2010) 2497–2581 MR Zbl
- [44] **C H Taubes**, *Embedded contact homology and Seiberg–Witten Floer cohomology, V*, *Geom. Topol.* 14 (2010) 2961–3000 MR Zbl
- [45] **W Wang, X Hu, Y Long**, *Resonance identity, stability, and multiplicity of closed characteristics on compact convex hypersurfaces*, *Duke Math. J.* 139 (2007) 411–462 MR Zbl

DCG: *Department of Mathematics, University of California, Santa Cruz
Santa Cruz, CA, United States*

DCG: *School of Mathematics, Institute for Advanced Study
Princeton, NJ, United States*

Current address for DCG: *Department of Mathematics, University of Maryland
College Park, MD, United States*

UH: *RWTH Aachen
Aachen, Germany*

MH: *Department of Mathematics, University of California, Berkeley
Berkeley, CA, United States*

HL: *School of Mathematics and Statistics, Wuhan University
Wuhan, China*

dcrstof@umd.edu, hryniewicz@mathga.rwth-aachen.de,
hutchings@berkeley.edu, huiliu00031514@whu.edu.cn

Proposed: András I Stipsicz

Received: 29 September 2021

Seconded: Ciprian Manolescu, Leonid Polterovich

Revised: 24 January 2022

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITOR

András I Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Dan Abramovich	Brown University dan_abramovich@brown.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Frances Kirwan	University of Oxford frances.kirwan@balliol.oxford.ac.uk
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	John Lott	University of California, Berkeley lott@math.berkeley.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Haynes Miller	Massachusetts Institute of Technology hmm@math.mit.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Tomasz Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Walter Neumann	Columbia University neumann@math.columbia.edu
Benson Farb	University of Chicago farb@math.uchicago.edu	Jean-Pierre Otal	Université Paul Sabatier, Toulouse otal@math.univ-toulouse.fr
Steve Ferry	Rutgers University sferry@math.rutgers.edu	Peter Ozsváth	Princeton University petero@math.princeton.edu
David M Fisher	Rice University davidfisher@rice.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Mike Freedman	Microsoft Research michaelf@microsoft.com	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David Gabai	Princeton University gabai@princeton.edu	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Peter Teichner	Max Planck Institut für Mathematik teichner@mac.com
Cameron Gordon	University of Texas gordon@math.utexas.edu	Richard P Thomas	Imperial College, London richard.thomas@imperial.ac.uk
Lothar Götsche	Abdus Salam Int. Centre for Th. Physics gotsche@ictp.trieste.it	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Mark Gross	University of Cambridge mgross@dpmms.cam.ac.uk	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2023 is US \$740/year for the electronic version, and \$1030/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY



mathematical sciences publishers
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 27 Issue 9 (pages 3387–3831) 2023

Hyperbolic groups acting improperly	3387
DANIEL GROVES and JASON FOX MANNING	
Cyclic homology, S^1 -equivariant Floer cohomology and Calabi–Yau structures	3461
SHEEL GANATRA	
Congruences on K-theoretic Gromov–Witten invariants	3585
JÉRÉMY GUÉRE	
Moduli of spherical tori with one conical point	3619
ALEXANDRE EREMENKO, GABRIELE MONDELLO and DMITRI PANOV	
The derivative map for diffeomorphism of disks: an example	3699
DIARMUID CROWLEY, THOMAS SCHICK and WOLFGANG STEIMLE	
On self-shrinkers of medium entropy in $\mathbb{R}^4$	3715
ALEXANDER MRAMOR	
The Gromov–Hausdorff distance between spheres	3733
SUNHYUK LIM, FACUNDO MÉMOLI and ZANE SMITH	
Contact three-manifolds with exactly two simple Reeb orbits	3801
DANIEL CRISTOFARO-GARDINER, UMBERTO HRYNIEWICZ, MICHAEL HUTCHINGS and HUI LIU	