

GEOGRAPHY OF SYMPLECTIC LEFSCHETZ FIBRATIONS AND RATIONAL BLOWDOWNS

R. İNANÇ BAYKUR, MUSTAFA KORKMAZ, AND JONATHAN SIMONE

ABSTRACT. We produce simply-connected, minimal, symplectic Lefschetz fibrations realizing all the lattice points in the symplectic geography plane below the Noether line. This provides a *symplectic* extension of the classical works populating the complex geography plane with holomorphic Lefschetz fibrations. Our examples are obtained by rationally blowing down Lefschetz fibrations with clustered nodal fibers, the total spaces of which are potentially new homotopy elliptic surfaces. Similarly, clustering nodal fibers on higher genera Lefschetz fibrations on standard rational surfaces, we get rational blowdown configurations that yield new constructions of small symplectic exotic 4-manifolds. We present an example of a construction of a minimal symplectic exotic $\mathbb{C}P^2 \# 5\overline{\mathbb{C}P^2}$ through this procedure applied to a genus-3 fibration.

1. INTRODUCTION

The *symplectic geography problem*, inspired by the study of compact complex algebraic surfaces by Persson et al. [38, 39, 50], asks which pairs of integers (a, b) can be realized as the holomorphic Euler characteristic $\chi_h = a$ and the first Chern number $c_1^2 = b$ of a closed minimal symplectic 4-manifold [22, 28]. It is well-known that these invariants depend only on the underlying homotopy type of the 4-manifold X , satisfying the identities $\chi_h = \frac{1}{4}(\chi + \sigma)$ and $c_1^2 = 2\chi + 3\sigma$, where χ and σ are the Euler characteristic and the signature of X . Both coordinates are positive for minimal simply-connected 4-manifolds of general type, which are usually the focus of the geography problem.

In the case of compact complex algebraic surfaces, the geography plane was populated almost exclusively by surfaces that are the total spaces of singular fibrations over complex curves [8, 9, 38–40, 43, 50], where most lattice points in the region $8a \geq b \geq 2a - 6$ are realized by minimal holomorphic Lefschetz fibrations; see e.g. [50]. The associated invariants of a compact complex surface of general type satisfy both the *Bogomolov-Miyaoka-Yau inequality* $9\chi_h \geq c_1^2$ and the *Noether inequality* $c_1^2 \geq 2\chi_h - 6$.

Perhaps the most striking difference between the complex and symplectic geography is that the Noether inequality fails for symplectic 4-manifolds [1, 14, 22, 41]. However, in this case, there are only sporadic examples realizing lattice points in the region $2a - 6 \geq b > 0$ as Lefschetz fibrations, which were suggested by Fintushel and Stern in [16]; see Remark 9. The families of examples we produce in our first theorem will in particular contain minimal simply-connected Lefschetz fibrations

Received by the editors February 3, 2022, and, in revised form, February 6, 2023.
 2020 *Mathematics Subject Classification.* Primary 57K43, 57K41, 57R55, 57M05.
 The first author was supported by the NSF grant DMS-2005327.




FIGURE 1. The lattice points in the shaded region $\mathcal{R}$ are populated by our minimal simply-connected symplectic Lefschetz fibrations.

populating this entire region. The interested reader can refer to Remark 12 for a further discussion on the geography of symplectic Lefschetz fibrations.

Before we formulate the theorem, let us recall the rational blowdown operation. The *rational blowdown* is performed by cutting out a regular neighborhood of a configuration of spheres, called a plumbing, embedded in an ambient 4-manifold and replacing it with a rational homology 4-ball. Since the rational blowdown was introduced by Fintushel-Stern in [15] and generalized by Park in [34], it has been used to construct many exotic 4-manifolds; e.g. [33, 35–37, 45, 52], most of which are symplectic due to Symington [47], [48]. The appeal of this construction, along with other similar symplectic cut-and-paste operations (e.g. [23, 42]), is that the fundamental group and Seiberg-Witten invariant calculations are relatively routine. The first step in this process is locating a suitable configuration of spheres in some 4-manifold. Partially because there is a well-known classification of singular fibers on elliptic fibrations, elliptic fibrations have been the most popular starting place to search for plumbings that can be symplectically rationally blown down (e.g. most of the above referenced articles start with elliptic fibrations). More recently, Akhmedov and Sakallı utilized the classification of singular fibers on certain holomorphic genus-2 Lefschetz fibrations for rational blowdowns [3].

In this article, our starting point will be genus $g > 1$ Lefschetz fibrations on rational surfaces and homotopy elliptic surfaces, corresponding to positive factorizations in the mapping class group $\text{Mod}(\Sigma_g)$. The particular positive factorizations we use allow us to cluster nodal singularities and obtain some extremal configurations for our rational blowdown procedures. Thus, our first set of results will be on clustering nodal singularities in genus- g Lefschetz fibrations on $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}^2}$; see Lemmas 4 and 6. Coupling these ideas with twisted fiber sums, for each $g \geq 2$, we construct minimal symplectic genus- g Lefschetz fibrations (Z_{g+1}, f_{g+1}) , where Z_{g+1} is a homotopy elliptic surface $E(g+1)$. These will contain $2g+2$ disjoint embedded symplectic (-4) -spheres on the fibers we can then rationally blow down

to get our first result. Consider the region

$$\mathcal{R} = \{(a, b) \in \mathbb{Z}^2 \mid a \geq 3 \text{ and } 0 < b \leq 2a\}$$

shown in Figure 1. We have:

Theorem 1. *For each point $(a, b) \in \mathcal{R}$, there exists a minimal simply-connected non-spin symplectic genus $g = a - 1$ Lefschetz fibration $(Z_{a,b}, f_{a,b})$ satisfying $\chi_h(Z_{a,b}) = a$ and $c_1^2(Z_{a,b}) = b$ obtained from (Z_a, f_a) by rationally blowing down b many (-4) -spheres contained in the fibers.*

Similar to the regular blowdown, each time we rationally blow down a (-4) -sphere, c_1^2 increases by one whereas χ_h does not change. So, we see that the portion of the geography plane highlighted in Figure 1 is populated by $Z_{a,b}$ as we vary $(a, b) \in \mathcal{R}$.

Curiously, the homotopy elliptic surfaces Z_{g+1} we build are not diffeomorphic to well-known homotopy elliptic surfaces obtained from $E(g+1)$ by logarithmic transforms or knot surgery on an elliptic fiber for any $g > 3$; we show this in Proposition 8. The symplectic 4-manifolds $Z_{a,b}$ with $0 < b < a - 3$ are moreover interesting in connection to a conjecture of Fintushel and Stern on the number of basic classes recently proven by Feehan and Leness [13]; see Remark 11.

Although there are known exotic symplectic 4-manifolds in the homeomorphism classes of $\mathbb{CP}^2 \# m\overline{\mathbb{CP}^2}$ for as small as $m = 2$ [2] (also see [20]), to date, the smallest symplectic exotic 4-manifold produced using the rational blowdown operation is an exotic $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}^2}$ [32, 33] (also see [19, 37, 52] for examples that are not known to be symplectic). Despite the first breakthroughs in constructions of small exotic rational surfaces all being via rational blowdowns [19, 35, 37, 45], and a massive amount of literature on applications of rational blowdowns and their generalizations over two decades, Question 2 is still open:

Question 2. *Is there an exotic $\mathbb{CP}^2 \# m\overline{\mathbb{CP}^2}$ with $m < 5$ that can be obtained from a standard rational surface via rational blowdowns? If so, what is the smallest such m ?*

In principle, by expanding our view to higher genus Lefschetz fibrations, we open the door to finding larger configurations of spheres that can be rationally blown down, leading to possibly smaller exotic 4-manifolds. Adapting this approach, we tackled this problem and we discovered many new constructions of examples right on the border. To illustrate, we will present a relatively simple construction, where we will start with a particular genus-3 Lefschetz fibration on $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}^2}$. We construct this Lefschetz fibration by refactoring the monodromy associated to the hyperelliptic Lefschetz fibration on $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}^2}$ by clustering nodes; see Lemma 6. After some blow-ups and one blow-down, we find a fairly simple configuration of spheres embedded in $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}^2}$ that can be rationally blown down. We then prove the following.

Theorem 3. *There exists a minimal symplectic exotic $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}^2}$ obtained by rationally blowing down a blow-up of a genus-3 Lefschetz fibration on a standard rational surface.*

Peculiarly, yet another reason for us to choose this example is that only a slight potential improvement of the positive factorization we employed in the proof of this theorem would result in an exotic $\mathbb{CP}^2 \# 4\overline{\mathbb{CP}^2}$; see Remark 17. In fact, we

expect the answer to the existence part of Question 2 to be yes, despite our own efforts thus far falling short, noting that our investigation has mostly focused on configurations in low genera fibrations.

Our paper is organized as follows. In Section 2, we construct positive factorizations for Lefschetz fibrations with clusters of nodes; see Lemmas 4 and 6. We use these factorizations to then prove Theorem 1 and Proposition 8 in Section 3, and Theorem 3 in Section 4. Throughout the article, any 4-manifold we consider will be compact, connected, smooth, oriented and without boundary, unless explicitly stated otherwise.

2. CLUSTERING NODES IN HIGHER GENERA LEFSCHETZ FIBRATIONS

Let Σ_g^m denote a compact connected orientable surface of genus g with m boundary components and let $\text{Mod}(\Sigma_g^m)$ be its *mapping class group*, the group of isotopy classes of orientation-preserving diffeomorphisms of Σ_g^m that restrict to the identity on the boundary $\partial\Sigma_g^m$. We write Σ_g for Σ_g^0 . For a simple closed curve c on Σ_g^m , we denote the positive (right-handed) Dehn twist along c by t_c . The conjugation of a group element ψ by ϕ , namely $\phi\psi\phi^{-1}$, will be written as ψ^ϕ . A conjugate of a positive Dehn twist satisfies $t_c^\phi = t_{\phi(c)}$, so it is also a positive Dehn twist, for any $\phi \in \text{Mod}(\Sigma_g^m)$. The reader should not confuse the power t_c^k of a Dehn twist with the conjugate element t_c^ϕ , which can be differentiated by k always denoting an integer and ϕ a mapping class.

Any product of positive of Dehn twists P that is equal to the identity in $\text{Mod}(\Sigma_g)$ gives rise to a (symplectic) Lefschetz fibration with generic fiber Σ_g . A lifting of $P = 1$ to $\text{Mod}(\Sigma_g^m)$ specifies m sections together with their self intersections. If a subfactor of P contains Dehn twists along disjoint curves, then, after a small perturbation, we get an associated singular fiber with *clustered nodes*. Topologically, clustering nodes in certain ways gives rise to many 2-spheres embedded in the singular fibers (e.g. a string of (-2) -spheres intersecting transversely). Our strategy is to manipulate particular positive factorizations in order to cluster nodes; in doing so, we are able to construct Lefschetz fibrations containing desirable configurations of spheres that can be further manipulated into configurations that can be rationally blown down. Often most of the rational blowdown configurations we seek for will be contained in a just a few singular fibers with clustered nodes.

Let $c_1, \dots, c_{2g+1}$ denote the curves on Σ_g depicted in Figure 2 and let us set $t_i = t_{c_i}$ for a short-hand notation. The element


$$h = t_1 t_2 \cdots t_{2g} t_{2g+1} t_{2g+1} t_{2g} \cdots t_2 t_1$$

is the hyperelliptic involution on Σ_g which fixes each c_i setwise. We thus obtain a positive factorization of a genus- g Lefschetz fibration from $h^2 = 1$, which is known to be a genus- g hyperelliptic Lefschetz fibration on $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}^2}$.

Lemma 4. *Let $g \geq 1$. For any integers $p, q \geq 0$ satisfying $p + q = 4g + 4$, there are positive factorizations in $\text{Mod}(\Sigma_g)$ of the form*

$$t_1^p \cdot t_3^q \cdot D_{p,q,g} = 1,$$

where $D_{p,q,g}$ is a product of $4g$ positive Dehn twists, and $t_i = t_{c_i}$ for c_i as in Figure 2.

FIGURE 2. Curves on Σ_g .

Proof. The statement will follow from a sequence of manipulations in $\text{Mod}(\Sigma_g)$:

$$\begin{aligned}
 1 &= h^2 \\
 &= h \cdot (t_1 t_2 \cdots t_{2g+1} t_{2g+1} \cdots t_2 t_1) \\
 &= t_1 t_2 \cdots t_{2g+1} \cdot h \cdot t_{2g+1} \cdots t_2 t_1 \\
 &= t_1 t_2 \cdots t_{2g+1} \cdot (t_1 t_2 \cdots t_{2g+1} t_{2g+1} \cdots t_2 t_1) \cdot t_{2g+1} \cdots t_2 t_1,
 \end{aligned}$$

which follows from h fixing every c_i and therefore commuting with every t_{c_i} . Since t_1 commutes with t_j for all $j > 2$, the two t_1 elements on the ends of the factorization $(t_1 t_2 \cdots t_{2g+1} t_{2g+1} \cdots t_2 t_1)$ in the middle of the last equation above can be moved outward to the third and third-to-last positions, obtaining the equation

$$= t_1 (t_2 t_1) t_3 \cdots t_{2g+1} \cdot (t_2 \cdots t_{2g+1} t_{2g+1} \cdots t_2) \cdot t_{2g+1} \cdots t_3 (t_1 t_2) t_1.$$

Iterating the same step for each t_i , for $i = 1, \dots, 2g$, yields the next equality:

$$= t_1 (t_2 t_1) (t_3 t_2) \cdots (t_{2g+1} t_{2g}) t_{2g+1} \cdot t_{2g+1} (t_{2g} t_{2g+1}) \cdots (t_2 t_3) (t_1 t_2) t_1.$$

After a cyclic permutation we can bring t_1 on the left to the far right, recalling that the whole product is equal to the identity. Applying a sequence of braid relations $t_i t_{i+1} t_i = t_{i+1} t_i t_{i+1}$ we can then carry each t_{2g+1} in the middle of the factorization all the way to the right to derive

$$\begin{aligned}
 &= (t_2 t_1) (t_3 t_2) \cdots (t_{2g+1} t_{2g}) t_{2g+1} \cdot t_{2g+1} (t_{2g} t_{2g+1}) \cdots (t_2 t_3) (t_1 t_2) \cdot t_1^2 \\
 &= (t_2 t_1) (t_3 t_2) \cdots (t_{2g+1} t_{2g}) \cdot (t_{2g} t_{2g+1}) \cdots (t_2 t_3) (t_1 t_2) \cdot t_1^4.
 \end{aligned}$$

Now, conjugating t_1 with t_2 , we obtain

$$= t_1^{t_2} t_2 (t_3 t_2) \cdots (t_{2g+1} t_{2g}) \cdot (t_{2g} t_{2g+1}) \cdots (t_2 t_3) (t_1 t_2) \cdot t_1^4.$$

Using braid relations again, we then carry this t_2 first to the very center as t_{2g+1} and then to far right as t_1 as follows

$$\begin{aligned}
 &= t_1^{t_2} (t_3 t_2) \cdots (t_{2g+1} t_{2g}) \cdot t_{2g+1} \cdot (t_{2g} t_{2g+1}) \cdots (t_2 t_3) (t_1 t_2) \cdot t_1^4 \\
 &= t_1^{t_2} (t_3 t_2) \cdots (t_{2g+1} t_{2g}) \cdot (t_{2g} t_{2g+1}) \cdots (t_2 t_3) (t_1 t_2) \cdot t_1^5.
 \end{aligned}$$

Repeating the same steps this time for the leftmost t_3 we obtain

$$= t_1^{t_2} t_2^{t_3} (t_4 t_3) \cdots (t_{2g+1} t_{2g}) \cdot (t_{2g} t_{2g+1}) \cdots (t_2 t_3) (t_1 t_2) \cdot t_1^6.$$

Iterate the same sequence of modifications for each t_{i+1} that appears in a pair $(t_{i+1}t_i)$ (each grouped in separate parentheses) on the left half of the factorization, for $i = 1, 2, \dots, 2g + 1$, in this order. Then repeat for each t_i that appear in pairs (t_it_{i+1}) on the right half of the factorization, for $i = 2g, \dots, 2, 1$, in this order. We obtain

$$= t_1^{t_2} t_2^{t_3} t_3^{t_4} \dots t_{2g}^{t_{2g+1}} \cdot t_{2g+1}^{t_{2g}} \dots t_3^{t_2} t_2^{t_1} \cdot t_1^{4g+4}.$$

Finally we observe that $(t_3^{t_2} t_2^{t_1}) t_1 = t_3 (t_3^{t_2} t_2^{t_1})$ by braid relations. For $p + q = 4g + 4$, repeating this q times we get

$$\begin{aligned} &= t_1^{t_2} t_2^{t_3} t_3^{t_4} \dots t_{2g}^{t_{2g+1}} \cdot t_{2g+1}^{t_{2g}} \dots t_4^{t_3} \cdot t_3^q \cdot (t_3^{t_2} t_2^{t_1}) \cdot t_1^p \\ &= t_1^{t_2} t_2^{t_3} t_3^{t_4} \dots t_{2g}^{t_{2g+1}} \cdot t_{2g+1}^{t_{2g}} \dots t_4^{t_3} \cdot t_3^{t_2} t_3^q t_2^{t_1} t_3^q \cdot t_3^q \cdot t_1^p \\ &= t_1^p \cdot t_3^q \cdot D_{p,q,g}, \end{aligned}$$

where the very last equation is obtained by cyclic permutation and also that t_1^p and t_3^q commute with each other. $\square$

Remark 5. We expect that the positive factorizations $t_1^{4g+4} D = 1$ in $\text{Mod}(\Sigma_g)$ we obtained above to be optimal for any $g \geq 2$, that is, one cannot perturb the given Lefschetz fibration on $X := \mathbb{CP}^2 \# (4g + 5) \overline{\mathbb{CP}^2}$ to cluster any more identical nodes (i.e. with isotopic vanishing cycles) on a single singular fiber. In fact, a cluster corresponding to the t_1^k factor yields a chain of $k - 1$ (-2) -spheres which span a negative-definite subspace of V of $H_2(X; \mathbb{R})$ for the intersection form Q_X . Moreover, for the symplectic regular fiber F , we have $[F] \neq 0$ in $H_2(X; \mathbb{R})$ and $[F] \in V^\perp$, the orthogonal complement of V with respect to Q_X . Because $Q_X|_{V^\perp}$ is nondegenerate, there is an additional class in $V^\perp$ with negative square. Hence $4g + 5 = b^-(X) \geq k$, which demonstrates that we are at most one off from clustering the maximal number of nodes. Generally, we conjecture that for a fiber sum indecomposable (see e.g. [4] for the definition) genus $g \geq 2$ Lefschetz fibration, the maximal number of nodes one can cluster like this is $4g + 4$, realized by the fibrations we get on the rational surface $\mathbb{CP}^2 \# (4g + 5) \overline{\mathbb{CP}^2}$. Note that this number is $4g + 5$ when $g = 1$.

We will now construct a particular positive factorization for a genus-3 Lefschetz fibration. In this case, for the rational blowdown configurations we desire to get, it will be essential to identify some sections as well. We will thus produce a positive factorization of the boundary multi-twist $t_{\delta_1} t_{\delta_2}$ in $\text{Mod}(\Sigma_3^2)$. (See e.g. [6] for how boundary twists yield sections.)




FIGURE 3. Curves on Σ_3^2 .

Lemma 6. *There is a positive factorization in $\text{Mod}(\Sigma_3^2)$*

$$(t_1^{14} t_a t_b t_4 t_6) \cdot (t_1 t_3 t_5 t_7)^{t_2^{-1} t_5 t_6^{-1}} \cdot D_6 = t_{\delta_1} t_{\delta_2},$$

where D_6 is a product of 6 positive Dehn twists, $t_i = t_{c_i}$, and c_i, a, b, c, δ_j are as in Figure 3.

Proof. The statement will follow from a sequence of manipulations this time in $\text{Mod}(\Sigma_3^2)$. We start with the well-known 7-chain relation:

$$\begin{aligned} t_{\delta_1} t_{\delta_2} &= (t_7 \cdots t_2 t_1)^8 \\ &= (t_7 \cdots t_2 t_1)(t_7 \cdots t_2 t_1)(t_7 \cdots t_2 t_1)(t_7 \cdots t_2 t_1) \\ &\quad \cdot (t_7 \cdots t_2 t_1)(t_7 \cdots t_2 t_1)(t_7 \cdots t_2 t_1)(t_7 \cdots t_2 t_1). \end{aligned}$$

Note that $t_i(t_7 \cdots t_2 t_1) = (t_7 \cdots t_2 t_1)t_{i+1}$ for each $i = 1, \dots, 6$. We can thus move the t_1 in the first parentheses over the next six $(t_7 \cdots t_2 t_1)$ factors, so it becomes t_7 . We then move the t_1 in the second parentheses over the next five $(t_7 \cdots t_2 t_1)$ factors, so it becomes t_6 . Repeating this for each t_1 in the first six parentheses, we get:

$$= (t_7 \cdots t_3 t_2)^7 \cdot (t_1 t_2 \cdots t_6 t_7 \cdot t_7 t_6 \cdots t_2 t_1).$$

By the same method, we can now move t_2 factors to arrive at the factorization

$$= (t_7 t_6 t_5 t_4 t_3)^6 \cdot (t_2 t_3 \cdots t_6 t_7 \cdot t_7 t_6 \cdots t_3 t_2) \cdot (t_1 t_2 \cdots t_6 t_7 \cdot t_7 t_6 \cdots t_2 t_1).$$

Using the 5-chain relation to substitute $(t_7 t_6 t_5 t_4 t_3)^6$ with $t_a t_b$, and braid relations to rewrite the rest of the factorization, we obtain

$$= (t_a t_b) \cdot (t_1 t_2 \cdots t_6 t_7 \cdot t_1 t_2 \cdots t_5 t_6 \cdot t_6 t_5 \cdots t_2 t_1 \cdot t_7 t_6 \cdots t_2 t_1).$$

We can then use commutativity relations to rearrange the factorization into

$$= (t_a t_b t_1) \cdot (t_2 t_1)(t_3 t_2)(t_4 t_3)(t_5 t_4)(t_6 t_5)(t_7 t_6) \cdot (t_6 t_7)(t_5 t_6)(t_4 t_5)(t_3 t_4)(t_2 t_3)(t_1 t_2) t_1.$$

Now, employing the same argument we had in the proof of Lemma 4, through a sequence of conjugations and braid relations, we can cluster t_1 s on the right we obtain

$$= (t_a t_b t_1) \cdot t_1^{t_2} \cdot t_2^{t_3} \cdot t_3^{t_4} \cdot t_4^{t_5} \cdot t_5^{t_6} \cdot t_6^{t_7} \cdot t_7^{t_6} \cdot t_6^{t_5} \cdot t_5^{t_4} \cdot t_4^{t_3} \cdot t_3^{t_2} \cdot t_2^{t_1} \cdot t_1^{13}.$$

We then bring all the t_1 factors together by cyclic permutation of t_1^{13} and the commutativity of the remaining t_1 factor with $t_a t_b$, which yields

$$= (t_1^{14} t_a t_b) \cdot t_1^{t_2} \cdot t_2^{t_3} \cdot t_3^{t_4} \cdot t_4^{t_5} \cdot t_5^{t_6} \cdot t_6^{t_7} \cdot t_7^{t_6} \cdot t_6^{t_5} \cdot t_5^{t_4} \cdot t_4^{t_3} \cdot t_3^{t_2} \cdot t_2^{t_1}.$$

Next carry the $t_5^{t_4}$ factor all the way to the right (note that it commutes with $t_2^{t_1}$) to obtain

$$= (t_1^{14} t_a t_b) \cdot t_1^{t_2} \cdot t_2^{t_3} \cdot t_3^{t_4} \cdot t_4^{t_5} \cdot t_5^{t_6} \cdot t_6^{t_7} \cdot t_7^{t_6} \cdot t_6^{t_5} \cdot (t_4^{t_3} t_3^{t_2})^{t_5^{t_4}} \cdot t_2^{t_1} \cdot t_1^{t_4}.$$

By cyclic permutation and commutativity of $t_5^{t_4}$ with $t_1^{14} t_a t_b$, we can then carry the $t_5^{t_4}$ factor and place it in the first cluster to obtain

$$= (t_1^{14} t_a t_b t_5^{t_4}) \cdot t_1^{t_2} \cdot t_2^{t_3} \cdot t_3^{t_4} \cdot t_4^{t_5} \cdot t_5^{t_6} \cdot t_6^{t_7} \cdot t_7^{t_6} \cdot t_6^{t_5} \cdot (t_4^{t_3} t_3^{t_2})^{t_5^{t_4}} \cdot t_2^{t_1}.$$

By moving away the Dehn twists in between by conjugations, we create a second cluster as follows

$$= (t_1^{14} t_a t_b t_5^{t_4}) \cdot (t_2^{t_3} t_5^{t_6} t_6^{t_7} t_6^{t_5} t_2^{t_1}) \cdot D'_6,$$

where D'_6 denotes the product of the remaining six positive Dehn twists. Here $t_2^{t_3}$ and $t_5^{t_6}$ commute, so we can move the latter into the first cluster to get

$$= (t_1^{14} t_a t_b t_5^{t_4} t_5^{t_6}) \cdot (t_2^{t_1} t_2^{t_3} t_6^{t_5} t_6^{t_7}) \cdot D'_6.$$

In the next step, we repeatedly make use of the following observation: if x and y intersect at one point, then the braid relation $t_x t_y t_x = t_y t_x t_y$ implies that $t_y^{-1} t_x t_y = t_x t_y t_x^{-1}$, and in turn, we have $t_x^{t_y^{-1}} = t_y^{t_x}$. So we can rewrite the last factorization as

$$= (t_1^{14} t_a t_b t_4^{t_5^{-1}} t_6^{t_5^{-1}}) \cdot (t_1^{t_2^{-1}} t_3^{t_2^{-1}} t_5^{t_6^{-1}} t_7^{t_6^{-1}}) \cdot D'_6.$$

Since $t_1^{14} t_a t_b$ commutes with t_5^{-1} , $t_1 t_3$ commutes with t_6^{-1} , and $t_5 t_7$ commutes with t_2^{-1} , we obtain

$$= (t_1^{14} t_a t_b t_4 t_6)^{t_5^{-1}} \cdot (t_1 t_3 t_5 t_7)^{t_2^{-1} t_6^{-1}} \cdot D'_6.$$

We can conjugate this expression with t_5 to obtain:

$$= (t_1^{14} t_a t_b t_4 t_6) \cdot (t_1 t_3 t_5 t_7)^{t_5 t_2^{-1} t_6^{-1}} \cdot D'_6.$$

Finally, noting that t_5 commutes with t_2^{-1} , the factorization in the statement of the lemma follows. $\square$

Remark 7. Halfway through the proof of Lemma 6, we arrive at

$$t_{\delta_1} t_{\delta_2} = (t_a t_b) \cdot (t_1 t_2 \cdots t_6 t_7 \cdot t_1 t_2 \cdots t_5 t_6 \cdot t_6 t_5 \cdots t_2 t_1 \cdot t_7 t_6 \cdots t_2 t_1)$$

via the 5-chain and 7-chain relations. Notice that if the two boundary components Σ_3^2 are capped off, we get $t_a = t_b = t_1$ so that this relation reduces to a relation that is Hurwitz equivalent to the relation in Lemma 4 for $g = 3$, $p = 16$, $q = 0$. Since the remaining manipulations in the proof of Lemma 6 involve commutativity, conjugation, and braid relations, the total space of the Leschetz fibration with monodromy factorization given in the statement of Lemma 6 is $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}}^2$.

3. GEOGRAPHY OF MINIMAL SYMPLECTIC LEFSCHETZ FIBRATIONS

Consider the genus $g \geq 2$ Lefschetz fibration on $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}}^2$ corresponding to the positive factorization $t_1^{2g+2} \cdot t_3^{2g+2} \cdot D_g = 1$ in $\text{Mod}(\Sigma_g)$ derived in Lemma 4, where $D_g := D_{2g+2, 2g+2, g}$. Performing an untwisted fiber sum of this fibration with itself is known to yield the complex surface $E(g+1)$. (To see it note that this fibration on $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}}^2$, after a small perturbation, is isomorphic to a holomorphic fibration and the untwisted fiber sum of two copies can be realized as a double branched cover ramified along two copies of the regular fiber.)

Instead, we are interested in the result of a twisted fiber sum, which yields a homotopy $E(g+1)$. After a global conjugation, the monodromy factorization of the above fibration can be written as

$$A_g \cdot t_{a_1}^{2g+2} \cdot t_{a_3}^{2g+2} = 1, \text{ and also as } t_{b_1}^{2g+2} \cdot t_{b_3}^{2g+2} \cdot B_g = 1 \text{ in } \text{Mod}(\Sigma_g),$$

where a_1, a_3, b_1, b_3 are the curves shown in Figure 4 and A_g and B_g are each products of $4g$ positive Dehn twists. Then, let (Z_{g+1}, f_{g+1}) denote the Lefschetz fibration with monodromy factorization

$$A_g \cdot t_{a_1}^{2g+2} \cdot t_{a_3}^{2g+2} \cdot t_{b_1}^{2g+2} \cdot t_{b_3}^{2g+2} \cdot B_g = 1 \text{ in } \text{Mod}(\Sigma_g),$$

which is a *twisted* fiber sum of two copies of the fibration we had on $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}}^2$.

We claim that Z_{g+1} is simply-connected. One way to see this is the following: our Lefschetz fibration on $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}^2}$, after a small perturbation, becomes isomorphic to the standard holomorphic Lefschetz fibration, which has many sections; see e.g. [49]. Therefore, by Seifert Van-Kampen, the complement of the regular fiber in the simply-connected 4-manifold $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}^2}$ is necessarily simply-connected, and in turn, the twisted fiber sum Z_{g+1} obtained by gluing the two complements is also simply-connected.

Easy Euler characteristic and signature calculations show that we have $\chi(Z_{g+1}) = 12g + 12 = \chi(E(g+1))$ and $\sigma(Z_{g+1}) = -8g - 8 = \sigma(E(g+1))$, so that $\chi_h(Z_{g+1}) = g + 1$ and $c_1^2(Z_{g+1}) = 0$. Moreover, the fiber sum Z_{g+1} is minimal by [51] (also see [4]). When g is even, Z_{g+1} is certainly not spin, for instance by Rokhlin's theorem. On the other hand, when g is odd, with some extra care, one can show that Z_{g+1} is spin using the spin criterion for monodromy factorizations of Lefschetz fibrations in [5, 46]. (Here the twisting matters; one may for instance take the second positive factorization in the fiber sum as the image of the first factorization under the obvious hyperelliptic involution, followed by a cyclic permutation.) Hence by Freedman [21], Z_{g+1} is homeomorphic to $E(g+1)$ for all $g \geq 2$.




FIGURE 4. Lantern curves in the twisted fiber sum (Z_{g+1}, f_{g+1}) .

We will next show that Z_{g+1} is not diffeomorphic to $E(g+1)$. A natural followup question is whether it is diffeomorphic to a well-known homotopy elliptic surface in the literature. Recall that a homotopy $E(n)$ obtained by logarithmic transformations for relatively prime $p, q \geq 1$ is denoted by $E(n)_{p,q}$, and the one obtained by knot surgery along a generic elliptic fiber is denoted by $E(n)_L$. While any $E(n)_{p,q}$ is symplectic, $E(n)_L$ is symplectic if and only if L is fibered, and when that is the case, $E(n)_L$ admits a genus $2l + (n-1)$ Lefschetz fibration [15, 17, 18], where l is the genus of L .

Proposition 8. *If $g > 3$, then Z_{g+1} is not diffeomorphic to $E(g+1)_{p,q}$ for any $p, q \geq 1$ or $E(g+1)_L$ for any knot L . In particular, Z_{g+1} is not diffeomorphic to $E(g+1)$.*

Proof. Let K denote the canonical class of Z_{g+1} and let S denote an embedded symplectic sphere with square -4 . By the adjunction equality, $\langle K, S \rangle = -\chi(S) - [S]^2 = 2$. Suppose Z_{g+1} is diffeomorphic to $E(g+1)_{p,q}$. Then $K = ((g+1)pq - p - q)f$, where f is a primitive class such that $pqf = F$ is the elliptic fiber class in $E(g+1)$. Thus $2 = \langle K, S \rangle = ((g+1)pq - p - q)f \cdot S$. Thus $(g+1)pq - p - q \in \{1, 2\}$. Now since $pq - p - q \geq -1$ and $pq \geq 1$ for all integers $p, q \geq 1$, we necessarily have that $g \leq 3$.

Next suppose $Z_{g+1} = E(g+1)_L$. If L is not a fibered knot, then $E(g+1)_L$ does not admit a symplectic structure [17], so Z_{g+1} cannot be diffeomorphic to it. Assume L is a fibered knot of genus $l \geq 0$. We first claim that the canonical class must be of the form $K = (2l + g - 1)F$, where F is the elliptic fiber class of $E(g+1)$ and l is the genus of L . Let Σ denote a generic fiber of the symplectic Lefschetz fibration on $E(g+1)_L$, which has genus $2l + g$ and square 0. From the construction, we know that F is a bisection of this fibration and $F \cdot \Sigma = 2$. By the adjunction equality, $\langle K, \Sigma \rangle = 2(2l + g) - 2 = 4l + 2g - 2$. By [17], $\mathcal{SW}_{E(g+1)_L} = \mathcal{SW}_{E(g+1)} \cdot \Delta_L(t^2)$, where $t = \exp(F)$ and Δ_L is the symmetrized Alexander polynomial of L ; consequently, the canonical class K must be of the form mF for some integer m . Thus we have $4l + 2g - 2 = m\langle F, \Sigma \rangle = 2m$ and so $m = 2l + g - 1$, as claimed. Finally, $2 = \langle K, S \rangle = (2l + g - 1)F \cdot S$, which implies $g \leq 3$.

If we take $p = q = 1$ for $E(g+1)_{p,q}$ or L the unknot for $E(g+1)_L$, we get back the standard $E(g+1)$. So the arguments above show in particular that Z_{g+1} is not diffeomorphic to $E(g+1)$. $\square$

The monodromy factorization of (Z_{g+1}, f_{g+1}) can be rewritten as

$$(1) \quad A_g \cdot (t_{a_1} t_{a_3} t_{b_1} t_{b_3})^{2g+2} \cdot B_g = 1 \text{ in } \text{Mod}(\Sigma_g),$$

where a_1, a_3, b_1, b_3 can be seen to create a lantern configuration; see Figure 4. Equivalently, clustering the corresponding four nodes on the same singular fiber we get a symplectic (-4) -sphere contained in the regular fiber. By [11], lantern substitution along this configuration will provide a new positive factorization for a symplectic Lefschetz fibration that would be obtained from the former by rationally blowing down the corresponding (-4) -sphere. As the above monodromy factorization shows, there are $2g + 2$ of these (-4) -spheres in Z_{g+1} .

Set $a = g - 1$. For each pair of integers (a, b) with $a \geq 3$ and $0 < b \leq 2a$, let $(Z_{a,b}, f_{a,b})$ denote the symplectic Lefschetz fibration obtained by rationally blowing down b of the above (-4) -spheres in (Z_a, f_a) . All $Z_{a,b}$ are minimal by [10] and we have $\chi_h(Z_{a,b}) = \chi_h(Z_a) = a$ and $c_1^2(Z_{a,b}) = c_1^2(Z_a) + b = b$. Thus the collection

$$\{Z_{a,b} \mid a \geq 3 \text{ and } 0 < b \leq 2a\}$$

of minimal symplectic 4-manifolds fills the region of the geography plane $\mathcal{R}$ in Figure 1.

The claim that $Z_{a,b}$ is non-spin follows from the following observation: Each time we apply a lantern substitution along a $(t_{a_1} t_{a_3} t_{b_1} t_{b_3})$ subfactor in the monodromy factorization, we get a new vanishing cycle that separates the pairs $\{a_1, b_1\}$ and $\{a_3, b_3\}$. This is a separating curve, and after a small perturbation of the fibration, we get a reducible fiber that contains a genus-1 (and a genus $g-1$) fiber component with self-intersection -1 . Hence, the intersection form of the ambient manifold has to be odd.

It remains to show that $Z_{a,b}$ are all simply-connected. This is fairly easy when $b < 2a$. Recall that if $\{v_j\}$ are the vanishing cycles of a genus- g Lefschetz fibration (X, f) with a section, the fundamental group of X is equal to $\pi_1(\Sigma_g)/N$ where N is the subgroup of $\pi_1(\Sigma_g)$ normally generated by $\{v_j\}$. Therefore, the vanishing cycles of the Lefschetz fibration we started with on the simply-connected 4-manifold $\mathbb{CP}^2 \# (4g+5)\overline{\mathbb{CP}}^2$ should normally generate all of $\pi_1(\Sigma_g)$. In turn, the vanishing cycles in the factor $A_g t_{a_1} t_{a_3}$ (or in $t_{b_1} t_{b_3} B_g$) would alone normally generate all of $\pi_1(\Sigma_g)$. But observe that until we make substitutions along all $t_{a_1} t_{a_3} t_{b_1} t_{b_3}$

factors, the original collection of the vanishing cycles for the Lefschetz fibration on $\mathbb{CP}^2 \# (4g + 5)\overline{\mathbb{CP}^2}$ (obviously after a global conjugation) are still part of the monodromy, which implies that $Z_{a,b}$ we get for $b < 2a$ is simply-connected. For the remaining $b = 2a$ case, one can similarly check that there are still an abundant collection of vanishing cycles in $Z_{a,2a}$ that normally generate the fundamental group of the fiber. (In fact, it suffices to observe that if we remove a_1 and a_3 from the collection, but add c_3 , which is one of the curves we get after the lantern substitution, the vanishing cycles in hand still kill the entire fundamental group of the fiber.)

This concludes the proof of Theorem 1.

Remark 9. There is at least one more example we can quickly cook up in the region $\mathcal{R}$ below the Noether line. Consider the genus-3 Lefschetz fibration on $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}^2}$ with monodromy factorization:

$$(t_1 t_2 t_3 t_4 t_5 t_6 t_7^2 t_6 t_5 t_4 t_3 t_2 t_1)^2 = 1 \text{ in } \text{Mod}(\Sigma_3),$$

where $t_i = t_{c_i}$ and c_i are as shown in Figure 2. This factorization can be re-written as

$$(t_1 t_3 t_5 t_7)^4 D = 1 \text{ in } \text{Mod}(\Sigma_3),$$

where D is a product of positive Dehn twists. Applying one lantern substitution along the factor $t_1 t_3 t_5 t_7$ results in a genus-3 Lefschetz fibration, the total space of which can be shown to be $\mathbb{CP}^2 \# 16\overline{\mathbb{CP}^2}$. If we now take the fiber sum of these two fibrations, we get a genus-3 Lefschetz fibration whose total space is a minimal [4, 51] symplectic 4-manifold with $\chi_h = 4$ and $c_1^2 = 1$, violating the Noether inequality.

The only other examples of relatively minimal Lefschetz fibrations in the literature we know of which realize some lattice points in the region below the Noether line (and above $c_1^2 = 0$) are the ones described by Fintushel and Stern in [16]. By taking fiber sums of two different holomorphic genus- g Lefschetz fibrations on rational surfaces, they get examples on the line $c_1^2 = \chi_h - 3$ for any $\chi_h > 3$ with $\chi_h \not\equiv 0 \pmod{3}$. Provided one determines the topological invariants that are only implicitly expressed in [16], one can obtain more examples following their construction scheme, but still with several limitations in the way of populating a large region; for example, one needs to have $g = \frac{1}{2}(p-1)(q-2) = \frac{1}{2}(p'-1)(q'-2)$ for pairs of relatively prime positive integers p, q and p', q' with $\{p, q\} \neq \{p', q'\}$. In particular, for every prime g , there is only one lattice point with $\chi_h = g + 1$ realized by this construction.

Moreover, a common aspect of all these sporadic examples is that they are fiber sums. In contrast, we suspect that most (perhaps all) of our examples $(Z_{a,b}, f_{a,b})$ with $a > 4$ and $b > 0$ are fiber sum indecomposable, and not diffeomorphic to any of these other examples.

Remark 10. Recall that Z_{g+1} is a homotopy $E(g+1)$ for all $a \geq 3$; in particular, Z_{g+1} is spin when $g+1$ is even. However, we can populate all lattice points on the line $c_1^2 = 0$ (with $\chi_h \geq 3$) with *non-spin* minimal simply-connected symplectic genus- g Lefschetz fibrations as well. This can be achieved by using a different twisted fiber sum than the one used to form Z_{g+1} . Let (Y_{g+1}, h_{g+1}) denote the Lefschetz fibration with monodromy factorization

$$A_g \cdot t_{a_1}^{2g+2} \cdot t_{a_3}^{2g+2} \cdot t_1^{2g+2} \cdot t_3^{2g+2} \cdot D_g = 1 \text{ in } \text{Mod}(\Sigma_g),$$

where $A_g \cdot t_{a_1}^{2g+2} \cdot t_{a_3}^{2g+2} = 1$ and $t_1^{2g+2} \cdot t_3^{2g+2} \cdot D_g = 1$ are the monodromy factorizations described previously. Rewriting this monodromy as $(t_{a_1} t_{a_3} t_3) \cdot D = 1$, where D is a

product of positive Dehn twists, it is clear that there is a fiber containing a sphere with self-intersection -3 , implying that the intersection form of Y_{g+1} is odd.

One may wonder if we can as well generate *spin* symplectic Lefschetz fibrations in the region $\mathcal{R}$, and this too is possible by varying our construction and using the *spin substitution* techniques developed in [5]. In this case, we would instead start with a twisted fiber sum (Y'_{g+1}, f'_{g+1}) on the $c_1^2 = 0$ line with a monodromy factorization

$$t_{a_5}^{4g+4} \cdot t_{b_5}^{4g+4} \cdot D'_g = 1 \text{ in } \text{Mod}(\Sigma_g),$$

for $g \geq 3$ and odd, where $\{a_5, b_5\}$ is the next pair of curves one can imagine in Figure 4, which cobound a subsurface Σ_2^2 of Σ_g . One can then perform 5-chain substitutions along each $t_{a_5} t_{b_5}$ factor to create the desired examples; cf. the proof of Theorem A in [5].

Remark 11. Let X be a simply-connected, almost complex 4-manifold of Seiberg-Witten simple type. Fintushel and Stern conjectured that if the characteristic numbers of X satisfy $0 \leq c_1^2 \leq \chi_h - 3$ then X has at least $\chi_h - c_1^2 - 2$ Seiberg-Witten basic classes. This conjecture was first confirmed for 4-manifolds of *superconformal simple type* in a physics paper [27], and was later completely proved by Feehan and Leness in [13]. Note that any symplectic X is of Seiberg-Witten simple type by work of Taubes. Moreover, in [12] Feehan and Leness, building on the examples in [14] with $c_1^2 = \chi_h - 3$, observed that there are *non-minimal* symplectic 4-manifolds at all the lattice points with $0 \leq c_1^2 < \chi_h - 3$ for which the bound on the number of basic classes is sharp. While it is beyond the scope of our work here, it would be interesting to determine if the same can be true for minimal examples, and in particular seeing if the number of basic classes of $Z_{a,b}$ with $0 \leq b < a - 3$ is $a - b - 2$.

Remark 12. Although there is an extensive literature on the geography of (semi-stable) holomorphic fibrations on compact complex surfaces, the situation in the symplectic case has not been understood well, at least until very recently; see e.g. [30, 44]. The recent works of [7, 29] showed the existence of simply-connected Lefschetz fibrations violating Xiao's famous slope inequality, which is always satisfied by holomorphic fibrations. Most recently, it was established in [5] that there are simply-connected Lefschetz fibrations with positive signatures—whereas it is still not known if such examples exist in the holomorphic case. Our examples in this article demonstrate further contrast between the geography of complex and symplectic Lefschetz fibrations.

4. A MINIMAL SYMPLECTIC EXOTIC $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$

Consider the Lefschetz fibration with monodromy factorization

$$(t_1^{14} t_a t_b t_4 t_6) \cdot (t_1 t_3 t_5 t_7)^{t_2^{-1} t_5 t_6^{-1}} \cdot D_6 = 1 \text{ in } \text{Mod}(\Sigma_3)$$

given by Lemma 6. As mentioned in Remark 7, the total space is $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}}^2$. Moreover, this fibration includes two singular fibers given by the clusters $t_1^{14} t_a t_b t_4 t_6$ and $(t_1 t_3 t_5 t_7)^{t_2^{-1} t_5 t_6^{-1}}$, along with the two (-1) -sections corresponding to the boundary components δ_1 and δ_2 in Figure 3. The first singular fiber F_1 is comprised of a string of 15 transversely intersecting embedded (-2) -spheres and an immersed (-2) -sphere that transversely intersects the first and last (-2) -spheres of the string. The second singular fiber F_2 is comprised of two (-4) -spheres intersecting each other transversely four times. The (-1) -sections intersect F_1 in the

first and fifteenth (-2) -spheres and intersect F_2 in each of the (-4) -spheres. This configuration is depicted schematically in Figure 5a.

Starting with this configuration of spheres, we will perform 16 blowups and a single blowdown to locate a plumbing P symplectically embedded $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2$ that can be symplectically rationally blown down to a symplectic exotic $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$. To prove that the blown down manifold X is homeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$, we will apply Freedman's Theorem. To prove X is not diffeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$, we will consider its symplectic Kodaira dimension. To this end, we first need to understand the homology classes of the fibers and sections of our Lefschetz fibration on $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}}^2$.





FIGURE 5.

Proposition 13. Let $\{h, e_1, \dots, e_{17}\}$ denote the standard basis of $H_2(\mathbb{CP}^2 \# 17\overline{\mathbb{CP}}^2)$. The homology classes of the configuration of spheres in Figure 5a are the homology classes shown in Figure 5b.

Proof. To determine these homology classes, we first symplectically blow down the configuration of spheres in $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}}^2$ 16 times. Upon doing so, the ambient 4-manifold is either $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ or $S^2 \times S^2$. We first show that it is the latter. Starting with the configuration shown in the first diagram of Figure 6, blow down the two (-1) -sections to obtain the next diagram in the figure. Starting with either green (-1) -sphere, sequentially blow down seven times. Then repeat starting with the other green (-1) -sphere. The result will be the configuration of spheres A , B , C , and D depicted in the last diagram. With abuse of notation, let A , B , C , and D denote both the spheres and their homology classes. It is clear that $A^2 = 12$, $B^2 = 0$, $C^2 = D^2 = 4$, $B \cdot C = B \cdot D = 1$, and $C \cdot D = 4$. Note that when we blew down the green (-1) -spheres in the second diagram, we introduced triple points. We then sequentially blow down at these triple points six times each. Thus we have that $A \cdot C = A \cdot D = 7$.

Suppose this configuration lives in $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ and let $\{h, e\}$ be the standard basis for $H_2(\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2)$. Then its canonical class is $K = \text{PD}(-3h + e)$. Let $B = xh + ye$. Then $B^2 = x^2 - y^2 = 0$ and $\langle -K, B \rangle = 3x + y$. Moreover, by the adjunction

FIGURE 6. Blowing down to $S^2 \times S^2$.

equality, $\langle -K, B \rangle = B^2 + 2 - 2g(B) = 2$. Combining these equations, we find that $x = 1, y = -1$ and so $B = h - e$. A similar computation shows that $C = 2h$. This implies that $C \cdot B = 2$, which is not the case. Thus the configuration of spheres must be in $S^2 \times S^2$.

Let s and f denote the standard section and fiber generators of $H_2(S^2 \times S^2)$. The canonical class is $K = \text{PD}(-2s - 2f)$. Let $A = xs + yf$. Then $A^2 = 2xy = 12$ and $\langle -K, A \rangle = 2x + 2y$. Moreover, since the homology class of A can be represented by a genus-2 surface, the adjunction equality gives us $\langle -K, A \rangle = 10$. Combining these equations, we easily see that $(x, y) = (2, 3)$ or $(3, 2)$. We may assume the former so that $A = 2s + 3f$. Now let $B = xs + yf$. A similar argument shows that $(x, y) = (0, 1)$ or $(1, 0)$. Since $A \cdot B = 2$, we must have the former and so $B = f$. Continuing in this way, we see that $C = D = s + 2f$.

Now that we know the homology classes of A, B, C , and D , we will reverse the blowdown process via blowups to recover the original configuration of spheres in $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}^2}$ along with their homology classes. This blowup process is shown in Figure 7. The first diagram shows the configuration of spheres A, B, C, D in $S^2 \times S^2$. Consider the intersection between A, B , and C , which is marked by a red point. We blow up this point seven times and call the exceptional spheres $d_1, \dots, d_7$. The resulting configuration is shown in the second diagram of Figure 7. Next, blow up seven times in the same way starting at the red point in the second diagram to obtain the third diagram and call the new exceptional spheres $d_9, \dots, d_{15}$. Finally, blow up at the final two red points to obtain the fourth diagram, which is the original configuration of spheres provided by the Lefschetz fibration. Call these last two exceptional spheres d_8 and d_{16} . The homology classes of the spheres in this final configuration is shown in the fourth diagram of Figure 7.




FIGURE 7. Blowing up $S^2 \times S^2$ to recover the homology classes of the original Lefschetz fibration configuration.

Let $\{h, e_1, \dots, e_{17}\}$ be the standard basis for $H_2(\mathbb{CP}^2 \# 17\overline{\mathbb{CP}^2})$. By performing the change of basis $f \mapsto h - e_1$, $s \mapsto h - e_2$, $d_1 \mapsto h - e_1 - e_2$, and $d_i \mapsto e_{i+1}$ for all $2 \leq i \leq 16$, we obtain the homology classes shown in Figure 5b. $\square$

Let P be the linear plumbing with weights

$$(-2, \dots, -2, -3, \underbrace{-2, \dots, -2}_3, -20, \underbrace{-2, \dots, -2}_{15}, -6, -7).$$

By symplectically blowing up the Lefschetz fibration on $\mathbb{CP}^2 \# 17\overline{\mathbb{CP}^2}$ sixteen times, and blowing down once, we can find P symplectically embedded in $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}^2}$. This process is shown in Figure 8. Let $u_1, \dots, u_{27}$ denote the homology classes of the spheres of P , as shown in Figure 9a. These homology classes are found explicitly in Figure 8, but for simplicity, they are recorded in the table in Figure 9b.

The boundary of P is the lens space $L(585^2, 291914)$, which bounds a rational homology 4-ball B , by Lisca [26]. Moreover, by Park [34] and Symington [48], P can be symplectically rationally blown down. Let $Z = \mathbb{CP}^2 \# 32\overline{\mathbb{CP}^2} \setminus P$ and let $X = Z \cup B$ be the result of the symplectic rational blowdown.

Proposition 14. X is homeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}^2}$.

Proof. We first show that X is simply connected. By the Seifert Van-Kampen Theorem, $\pi_1(X) = \pi_1(Z) *_{\pi_1(\partial P)} \pi_1(B)$. Since $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}^2}$ is simply connected, the map $\pi_1(\partial P) \rightarrow \pi_1(Z)$ induced by inclusion is surjective. Moreover, the map $\pi_1(\partial P) \cong \mathbb{Z}_{585^2} \rightarrow \pi_1(B) \cong \mathbb{Z}_{585}$ induced by inclusion is surjective ([34]). Thus, it suffices to show that Z is simply connected. Now since $\pi_1(\partial P)$ is abelian, so is $\pi_1(Z)$; consequently, $\pi_1(Z) \cong H_1(Z)$ and so it suffices to show that $H_1(Z)$ is trivial.





FIGURE 8. Sixteen blowups and one blowdown.

For $1 \leq i \leq 27$, let $\mu_i \in H_1(\partial P)$ denote the homology class of the meridian of the i th surgery curve in the obvious surgery diagram for ∂P . Let $\overline{\mu}_i$ denote the image of μ_i in $H_1(Z)$. Then $H_1(\partial P) \cong \mathbb{Z}_{585^2}$ is generated by μ_1 and the elements $\mu_1, \dots, \mu_{10}$ satisfy the relations

$$2\mu_1 = \mu_2, \quad 2\mu_i = \mu_{i-1} + \mu_{i+1} \text{ for } 2 \leq i \leq 5 \text{ and } 6 \leq i \leq 9, \text{ and } \quad 3\mu_6 = \mu_5 + \mu_7.$$

Combining these relations, we have that $\mu_6 = 6\mu_1$ and $\mu_{10} = 34\mu_1$. Since $H_1(\partial P)$ surjects onto $H_1(Z)$, $H_1(Z)$ is generated by $\overline{\mu}_1$ and satisfies the relations $\overline{\mu}_6 = 6\overline{\mu}_1$ and $\overline{\mu}_{10} = 34\overline{\mu}_1$.

In Z , $\overline{\mu}_6$ can be represented by the equator of the exceptional sphere e_{30} ; thus $\overline{\mu}_6 = 0$ and so $6\overline{\mu}_1 = \overline{\mu}_6 = 0$. Similarly, $\overline{\mu}_1$ and $\overline{\mu}_{10}$ can be represented by circles on the exceptional sphere e_{29} which cobound an annulus; hence $\overline{\mu}_1 = -\overline{\mu}_{10}$ and so


 FIGURE 9. The plumbing P and its homology classes.

$\overline{\mu}_1 = \overline{\mu}_{10} = 34\overline{\mu}_1$, or $35\overline{\mu}_1 = 0$. Since 6 and 35 are relatively prime, we readily obtain $\overline{\mu}_1 = 0$, proving that $H_1(Z)$ is trivial. Thus X is simply connected.

Now, $\sigma(X) = \sigma(\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2) - \sigma(P) + \sigma(B) = -4 = \sigma(\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2)$ and $\chi(X) = \chi(\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2) - \chi(P) + \chi(B) = \chi(\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2)$. Since the signatures of X and $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$ are not divisible by 16, both manifolds are odd. Thus, by Freedman's theorem [21], X is homeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$. $\square$

Proposition 15. X is not diffeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$.

Proof. Let $K_{\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2}$ denote the canonical class of $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$. By Theorem D in [25], $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$ has a unique symplectic structure up to diffeomorphism and deformation. Consequently, since the standard symplectic structure ν on $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$ satisfies $(K_{\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2}) \cdot \nu < 0$, $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$ does not admit a symplectic structure ν satisfying $(K_{\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2}) \cdot \nu > 0$. Let K_X denote the canonical class of X and let ω_X denote the symplectic class of X . We will show that $K_X \cdot \omega_X > 0$, proving that X is not diffeomorphic to $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$.

The canonical class of $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2$ is given by $K = PD(-3h + \sum_{i=1, i \neq 9}^{33} e_i)$. It is well-known that $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2$ admits a symplectic structure compatible with K and whose cohomology class can be represented by $\omega = PD(ah - \sum_{i=1, i \neq 9}^{33} b_i e_i)$, where

$a > b_0 > \cdots > b_{33}$ and $a > \sum_{\substack{i=1 \\ i \neq 9}}^{33} b_i$ (see, for example, Lemma 5.4 in [23]). Note that $K \cdot \omega = -3a + \sum_{\substack{i=1 \\ i \neq 9}}^{32} b_i$.

Give $H_2(P)$ the basis $\{u_1, \dots, u_{27}\}$ and give $H^2(P)$ the hom-dual basis $\{\gamma_1, \dots, \gamma_{27}\}$. Then $K|_P = \sum_{i=1}^{27} (K \cdot u_i) \gamma_i = \gamma_6 + 18\gamma_{10} + 4\gamma_{26} + 5\gamma_{27}$ and

$$\begin{aligned} \omega|_P &= \sum_{i=1}^{27} (\omega \cdot u_i) \gamma_i \\ &= (b_{27} - b_{28})\gamma_1 + (b_{26} - b_{27})\gamma_2 + \cdots + (b_{23} - b_{24})\gamma_5 \\ &\quad + (b_{22} - b_{23} - b_{30})\gamma_6 + (b_{21} - b_{22})\gamma_7 + \cdots + (b_{19} - b_{20})\gamma_9 \\ &\quad + (4a - 2b_1 - \sum_{i=2}^8 b_i - \sum_{i=10}^{16} b_i - 2b_{18} - 2b_{19} - \sum_{i=20}^{29} b_i)\gamma_{10} \\ &\quad + (b_{16} - b_{17})\gamma_{11} + \cdots + (b_{10} - b_{11})\gamma_{17} + (b_2 - b_{10})\gamma_{18} \\ &\quad + (a - b_1 - b_2 - b_3)\gamma_{19} + (b_3 - b_4)\gamma_{20} + \cdots + (b_7 - b_8)\gamma_{24} + (b_8 - b_{29})\gamma_{25} \\ &\quad + (2a - b_1 - \sum_{i=3}^8 b_i - \sum_{i=31}^{33} b_i)\gamma_{26} + (3a - 2b_1 - b_2 - \sum_{i=10}^{17} b_i - \sum_{i=31}^{33} b_i)\gamma_{27}. \end{aligned}$$

Let Q denote the matrix for the intersection form of P with respect to the basis $\{u_1, \dots, u_{27}\}$. Then

$$\begin{aligned} K|_P \cdot \omega|_P &= Q^{-1}(K|_P, \omega|_P) \\ &= \frac{1}{585} \left(-5544a + 3309b_1 + 1082(b_2 + b_{10} + \cdots + b_{17}) \right. \\ &\quad \left. + 1153(b_3 + \cdots + b_8 + b_{29}) + 1168b_{18} + 601(b_{19} + \cdots + b_{22}) \right. \\ &\quad \left. + 670(b_{23} + \cdots + b_{28}) + 516b_{30} + 1067(b_{31} + \cdots + b_{33}) \right). \end{aligned}$$

Finally, we have that

$$\begin{aligned} K_X \cdot \omega_X &= (K_X)|_Z \cdot (\omega_X)|_Z = K|_Z \cdot \omega|_Z = K \cdot \omega - K|_P \cdot \omega|_P \\ &= \frac{1}{585} \left(3789a - 2724b_1 - 497(b_2 + b_{10} + \cdots + b_{17}) \right. \\ &\quad \left. - 568(b_3 + \cdots + b_8 + b_{29}) - 583b_{18} \right. \\ &\quad \left. - 16(b_{19} + \cdots + b_{22}) - 85(b_{23} + \cdots + b_{28}) \right. \\ &\quad \left. + 69b_{30} - 482(b_{31} + \cdots + b_{33}) \right) \\ &> \frac{1}{585} \left(a - \sum_{\substack{i=1 \\ i \neq 9}}^{33} b_i \right) > 0. \end{aligned}$$

□

Proposition 16. X is minimal.

Proof. Following the strategy of Ozsváth-Szabó in [31], we will show that X has a unique basic class (up to sign), which implies that X is minimal. The following calculations were completed in Matlab. Because the efficacy of this type of computation is well-documented (c.f. [31], [45],[23]), we will only highlight the main steps.

We first find the following basis for $H_2(Z; \mathbb{Z})$:

$$\begin{aligned} A_1 &= h - e_1 - e_{18} - e_{31}, & A_2 &= h - e_1 - e_{18} - e_{32}, & A_3 &= h - e_1 - e_{18} - e_{33}, \\ A_4 &= 3h - 3e_1 + 2e_{18} - \sum_{i=19}^{22} 2e_i - 2e_{30} - e_{31} - e_{32} - e_{33}, \\ A_5 &= 3e_{18} - \sum_{i=23}^{28} e_i + e_{30}, \\ A_6 &= 36h - 21e_1 - 7e_2 - \sum_{i=3}^8 8e_i - \sum_{i=10}^{17} 7e_i + 2e_{18} \\ &\quad + \sum_{i=23}^{28} e_i - 8e_{29} - e_{30} - e_{31} - e_{32} - e_{33}. \end{aligned}$$

Notice that A_1, A_2, A_3 and A_5 can be represented by spheres, A_4 can be represented by a torus, and A_6 can be represented by a surface of genus 387 ([24]). Moreover, $A_1^2, A_2^2, A_3^2 = -2$, $A_4^2 = -27$, $A_5 = -16$, and $A_6 = -48$.

We would like to find the number of basic classes L on X satisfying the additional criterion: $|L(A_i)| \leq -A_i^2$ and $L(A_i) \equiv A_i^2 \pmod{2}$. Such classes are called *adjunctive* classes. There are 9,317,700 possible adjunctive classes. If L a basic class, then by the Seiberg-Witten dimension formula, $d = \frac{L^2 - 3\sigma(X) - 2\chi(X)}{4} = \frac{L^2 - 4}{4} \geq 0$ and $d \equiv 0 \pmod{2}$. Thus $L^2 \geq 4$ and $L \equiv 4 \pmod{8}$. This restriction leaves us with 13,960 possible adjunctive basic classes.

Now, if L is a basic class on X , then by the gluing formula ([34]), there is a basic class $\tilde{L}$ on $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2$ inducing L such that $(\tilde{L}|_P)^2 = -27$. Moreover, the set of basic classes on $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2$ inducing L contains an element $\tilde{L}$ such that $u_i^2 + 2 \leq \tilde{L}_P(u_i) \leq -u_i^2$ for all $1 \leq i \leq 27$ (where u_i are the homology classes of the spheres in P shown in Figure 9b). There are 585 classes on P satisfying these conditions. Thus we have a total of 8,166,600 classes on $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2$ that could give rise to adjunctive basic classes on X . These are given by $(L|_Z, \tilde{L}|_P)$.

Let $H = (2, 0, 0, 0, 1, 1, 0, \dots, 0) \in H_2(\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2; \mathbb{Q})$, written in the basis $\{A_1, \dots, A_6, u_1, \dots, u_{27}\}$. Then $H(u_i) = 0$ for all i , $H^2 > 0$, and $H \cdot PD(h) > 0$. By the wall-crossing formula, if $\tilde{L}$ is a basic class on $\mathbb{CP}^2 \# 32\overline{\mathbb{CP}}^2$ inducing a basic class on X , then $\text{sign}(\tilde{L} \cdot H) \neq \text{sign}(\tilde{L} \cdot h)$. There are 1788 such classes. Finally, we find that only 2 of these 1788 classes are integral. They are $\pm K$ (the canonical class). Consequently, X has a two adjunctive basic classes.

Finally following the argument in [31], it is easy to see that every basic class on X must be adjunctive. Thus X has a unique basic class (up to sign). $\square$

Remark 17. It is worth highlighting that we derived our rational blowdown configuration only using the first two clusters in the positive factorization $(t_1^{14} t_a t_b t_4 t_6) \cdot (t_1 t_3 t_5 t_7) t_2^{-1} t_5 t_6^{-1} \cdot D_6 = t_{\delta_1} t_{\delta_2}$ in $\text{Mod}(\Sigma_3^2)$, where the key point is the topological configuration of the disjoint Dehn twist curves in each cluster. Curiously, if one can get a similar positive factorization but with another cluster, such as a factor *conjugate* to $(t_a t_3 t_5 t_7)$, rationally blowing down a disjoint (-4) -sphere there would yield an exotic $\mathbb{CP}^2 \# 4\overline{\mathbb{CP}}^2$. In our factorization, we can find two of the desired Dehn twist factors in D_6 , but not all four.

ACKNOWLEDGMENTS

The second author thanks UMass Amherst for their generous support and wonderful research environment during this project.

REFERENCES

- [1] Anar Akhmedov, Scott Baldridge, R. İnanç Baykur, Paul Kirk, and B. Doug Park, *Simply connected minimal symplectic 4-manifolds with signature less than -1* , J. Eur. Math. Soc. (JEMS) **12** (2010), no. 1, 133–161, DOI 10.4171/JEMS/192. MR2578606
- [2] Anar Akhmedov and B. Doug Park, *Exotic smooth structures on small 4-manifolds*, Invent. Math. **173** (2008), no. 1, 209–223, DOI 10.1007/s00222-008-0118-x. MR2403396
- [3] Anar Akhmedov and Sümeýra Sakallı, *Deformation of singular fibers of genus two fibrations and small exotic symplectic 4-manifolds*, Internat. J. Math. **30** (2019), no. 3, 1950017, 37, DOI 10.1142/S0129167X19500174. MR3941983
- [4] R. İnanç Baykur, *Minimality and fiber sum decompositions of Lefschetz fibrations*, Proc. Amer. Math. Soc. **144** (2016), no. 5, 2275–2284, DOI 10.1090/proc/12835. MR3460185
- [5] R. İnanç Baykur and Noriyuki Hamada, *Lefschetz fibrations with arbitrary signature*, J. Eur. Math. Soc. (JEMS), to appear.
- [6] R. İnanç Baykur and Mustafa Korkmaz, *Small Lefschetz fibrations and exotic 4-manifolds*, Math. Ann. **367** (2017), no. 3-4, 1333–1361, DOI 10.1007/s00208-016-1466-2. MR3623227
- [7] Adalet Cengel and Mustafa Korkmaz, *Low-slope Lefschetz fibrations*, J. Topol. Anal., to appear.
- [8] Zhi Jie Chen, *On the geography of surfaces. Simply connected minimal surfaces with positive index*, Math. Ann. **277** (1987), no. 1, 141–164, DOI 10.1007/BF01457284. MR884652
- [9] Zhi Jie Chen, *The existence of algebraic surfaces with preassigned Chern numbers*, Math. Z. **206** (1991), no. 2, 241–254, DOI 10.1007/BF02571340. MR1091940
- [10] Josef G. Dorfmeister, *Minimality of symplectic fiber sums along spheres*, Asian J. Math. **17** (2013), no. 3, 423–442, DOI 10.4310/AJM.2013.v17.n3.a2. MR3119794
- [11] Hisaaki Endo and Yusuf Z. Gurtas, *Lantern relations and rational blowdowns*, Proc. Amer. Math. Soc. **138** (2010), no. 3, 1131–1142, DOI 10.1090/S0002-9939-09-10128-4. MR2566578
- [12] Paul M. N. Feehan and Thomas G. Leness, *Witten’s conjecture for many four-manifolds of simple type*, J. Eur. Math. Soc. (JEMS) **17** (2015), no. 4, 899–923, DOI 10.4171/JEMS/521. MR3349302
- [13] Paul M. N. Feehan and Thomas G. Leness, *The $SO(3)$ monopole cobordism and superconformal simple type*, Adv. Math. **356** (2019), 106817, 38, DOI 10.1016/j.aim.2019.106817. MR4011024
- [14] Ronald Fintushel, Jongil Park, and Ronald J. Stern, *Rational surfaces and symplectic 4-manifolds with one basic class*, Algebr. Geom. Topol. **2** (2002), 391–402, DOI 10.2140/agt.2002.2.391. MR1917059
- [15] Ronald Fintushel and Ronald J. Stern, *Rational blowdowns of smooth 4-manifolds*, J. Differential Geom. **46** (1997), no. 2, 181–235. MR1484044
- [16] Ronald Fintushel and Ronald J. Stern, *Constructions of smooth 4-manifolds*, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998), Doc. Math. **Extra Vol. II** (1998), 443–452. MR1648094
- [17] Ronald Fintushel and Ronald J. Stern, *Knots, links, and 4-manifolds*, Invent. Math. **134** (1998), no. 2, 363–400, DOI 10.1007/s002220050268. MR1650308
- [18] Ronald Fintushel and Ronald J. Stern, *Families of simply connected 4-manifolds with the same Seiberg-Witten invariants*, Topology **43** (2004), no. 6, 1449–1467, DOI 10.1016/j.top.2004.03.002. MR2081432
- [19] Ronald Fintushel and Ronald J. Stern, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* , J. Amer. Math. Soc. **19** (2006), no. 1, 171–180, DOI 10.1090/S0894-0347-05-00500-X. MR2169045
- [20] Ronald Fintushel and Ronald J. Stern, *Pinwheels and nullhomologous surgery on 4-manifolds with $b^+ = 1$* , Algebr. Geom. Topol. **11** (2011), no. 3, 1649–1699, DOI 10.2140/agt.2011.11.1649. MR2821436
- [21] Michael Hartley Freedman, *The topology of four-dimensional manifolds*, J. Differential Geometry **17** (1982), no. 3, 357–453. MR679066

- [22] Robert E. Gompf, *A new construction of symplectic manifolds*, Ann. of Math. (2) **142** (1995), no. 3, 527–595, DOI 10.2307/2118554. MR1356781
- [23] Çağrı Karakurt and Laura Starkston, *Surgery along star-shaped plumbings and exotic smooth structures on 4-manifolds*, Algebr. Geom. Topol. **16** (2016), no. 3, 1585–1635, DOI 10.2140/agt.2016.16.1585. MR3523050
- [24] Bang-He Li, *Representing nonnegative homology classes of $\mathbb{CP}^2 \# n \overline{\mathbb{CP}}^2$ by minimal genus smooth embeddings*, Trans. Amer. Math. Soc. **352** (2000), no. 9, 4155–4169, DOI 10.1090/S0002-9947-99-02422-8. MR1637082
- [25] T. J. Li and A. Liu, *Symplectic structure on ruled surfaces and a generalized adjunction formula*, Math. Res. Lett. **2** (1995), no. 4, 453–471, DOI 10.4310/MRL.1995.v2.n4.a6. MR1355707
- [26] Paolo Lisca, *Lens spaces, rational balls and the ribbon conjecture*, Geom. Topol. **11** (2007), 429–472, DOI 10.2140/gt.2007.11.429. MR2302495
- [27] Marcos Mariño, Gregory Moore, and Grigor Peradze, *Superconformal invariance and the geography of four-manifolds*, Comm. Math. Phys. **205** (1999), no. 3, 691–735, DOI 10.1007/s002200050694. MR1711332
- [28] John D. McCarthy and Jon G. Wolfson, *Symplectic normal connect sum*, Topology **33** (1994), no. 4, 729–764, DOI 10.1016/0040-9383(94)90006-X. MR1293308
- [29] Naoyuki Monden, *Lefschetz fibrations with small slope*, Pacific J. Math. **267** (2014), no. 1, 243–256, DOI 10.2140/pjm.2014.267.243. MR3163482
- [30] Naoyuki Monden, *On the geography of Lefschetz fibrations*, Handbook of group actions. Vol. II, Adv. Lect. Math. (ALM), vol. 32, Int. Press, Somerville, MA, 2015, pp. 297–330. MR3382032
- [31] Peter Ozsváth and Zoltán Szabó, *On Park’s exotic smooth four-manifolds*, Geometry and topology of manifolds, Fields Inst. Commun., vol. 47, Amer. Math. Soc., Providence, RI, 2005, pp. 253–260. MR2189937
- [32] Heesang Park, Jongil Park, and Dongsoo Shin, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 3$* , Geom. Topol. **13** (2009), no. 2, 743–767, DOI 10.2140/gt.2009.13.743. MR2469529
- [33] Heesang Park, Jongil Park, and Dongsoo Shin, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 4$* , Geom. Topol. **13** (2009), no. 3, 1483–1494, DOI 10.2140/gt.2009.13.1483. MR2496050
- [34] Jongil Park, *Seiberg-Witten invariants of generalised rational blow-downs*, Bull. Austral. Math. Soc. **56** (1997), no. 3, 363–384, DOI 10.1017/S0004972700031154. MR1490654
- [35] Jongil Park, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , Invent. Math. **159** (2005), no. 3, 657–667, DOI 10.1007/s00222-004-0404-1. MR2125736
- [36] Jongil Park, *Exotic smooth structures on $3\mathbb{CP}^2 \# 8\overline{\mathbb{CP}}^2$* , Bull. Lond. Math. Soc. **39** (2007), no. 1, 95–102, DOI 10.1112/blms/bdl017. MR2303524
- [37] Jongil Park, András I. Stipsicz, and Zoltán Szabó, *Exotic smooth structures on $\mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$* , Math. Res. Lett. **12** (2005), no. 5-6, 701–712, DOI 10.4310/MRL.2005.v12.n5.a7. MR2189231
- [38] Ulf Persson, *Chern invariants of surfaces of general type*, Compositio Math. **43** (1981), no. 1, 3–58. MR631426
- [39] Ulf Persson, *An introduction to the geography of surfaces of general type*, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), Proc. Sympos. Pure Math., vol. 46, Amer. Math. Soc., Providence, RI, 1987, pp. 195–218, DOI 10.1090/pspum/046.1/927957. MR927957
- [40] Xavier Roulleau and Giancarlo Urzúa, *Chern slopes of simply connected complex surfaces of general type are dense in $[2, 3]$* , Ann. of Math. (2) **182** (2015), no. 1, 287–306, DOI 10.4007/annals.2015.182.1.6. MR3374961
- [41] Sümeýra Sakalli, *Symplectic 4-manifolds on the Noether line and between the Noether and half Noether lines*, Geom. Dedicata **215** (2021), 369–399, DOI 10.1007/s10711-021-00655-6. MR4330348
- [42] Jonathan Simone, *Symplectically replacing plumbings with Euler characteristic 2 4-manifolds*, J. Symplectic Geom. **18** (2020), no. 5, 1285–1318, DOI 10.4310/JSG.2020.v18.n5.a4. MR4174302
- [43] Andrew John Sommese, *On the density of ratios of Chern numbers of algebraic surfaces*, Math. Ann. **268** (1984), no. 2, 207–221, DOI 10.1007/BF01456086. MR744608
- [44] A. I. Stipsicz, *The geography problem of 4-manifolds with various structures*, Acta Math. Hungar. **87** (2000), no. 4, 267–278, DOI 10.1023/A:1006719917276. MR1771206

- [45] András I. Stipsicz and Zoltán Szabó, *An exotic smooth structure on $\mathbb{CP}^2 \# 6\overline{\mathbb{CP}^2}$* , Geom. Topol. **9** (2005), 813–832, DOI 10.2140/gt.2005.9.813. MR2140993
- [46] András I. Stipsicz, *Spin structures on Lefschetz fibrations*, Bull. London Math. Soc. **33** (2001), no. 4, 466–472, DOI 10.1017/S0024609301008232. MR1832559
- [47] Margaret Symington, *Symplectic rational blowdowns*, J. Differential Geom. **50** (1998), no. 3, 505–518. MR1690738
- [48] Margaret Symington, *Generalized symplectic rational blowdowns*, Algebr. Geom. Topol. **1** (2001), 503–518, DOI 10.2140/agt.2001.1.503. MR1852770
- [49] Shunsuke Tanaka, *On sections of hyperelliptic Lefschetz fibrations*, Algebr. Geom. Topol. **12** (2012), no. 4, 2259–2286, DOI 10.2140/agt.2012.12.2259. MR3020206
- [50] Ulf Persson, Chris Peters, and Gang Xiao, *Geography of spin surfaces*, Topology **35** (1996), no. 4, 845–862, DOI 10.1016/0040-9383(95)00046-1. MR1404912
- [51] Michael Usher, *Minimality and symplectic sums*, Int. Math. Res. Not., posted on 2006, Art. ID 49857, 17, DOI 10.1155/IMRN/2006/49857. MR2250015
- [52] Kouichi Yasui, *Small exotic rational surfaces without 1- and 3-handles*, Trans. Amer. Math. Soc. **362** (2010), no. 11, 5893–5907, DOI 10.1090/S0002-9947-2010-05205-5. MR2661501

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, MASSACHUSETTS 01003-9305

Email address: `inanc.baykur@umass.edu`

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, 06800 ANKARA, TURKEY

Email address: `korkmaz@metu.edu.tr`

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PENNSYLVANIA 15213

Email address: `jsimone@andrew.cmu.edu`