

Revision of the Indo-West Pacific crab genus *Soliella* (Brachyura: Xanthidae: Etisinae): “pseudocryptic species” and basinal speciation

Robert M. Lasley Jr^{1,2}, Jose Christopher E. Mendoza³, Gustav Paulay²

¹Marine Laboratory, University of Guam, Guam 96913

E-mail: rlasleyjr@gmail.com

²Division of Invertebrates, Florida Museum of Natural History, Gainesville, Florida 32601

³Lee Kong Chian Natural History Museum, Faculty of Science, National University of Singapore,
2 Conservatory Drive, 117377, Singapore

ABSTRACT

Etisine crabs are some of the most abundant cryptobionts in Indo-West Pacific coral reef systems. Despite their ecological importance and abundance in museum collections, several recent systematic studies have indicated family- to subspecies-level taxonomic problems. One such case involves the former chlorodielline genus *Soliella* Lasley, Klaus & Ng, 2015 (treated here as part of Etisinae), which currently comprises two valid species and three available names that have been in flux in recent literature. The validity of these taxa has only been cursorily discussed. To resolve species limits and distributions, a thorough morphological examination of hundreds of specimens was conducted, including scanning electron microscopy of male gonopods, along with analysis of sequence data of the mitochondrial marker cytochrome c oxidase subunit I (COI) from 84 exemplars across the distribution of the genus. The status of two species that have Indian Ocean versus Pacific Ocean distributions with overlap in the Indo-Australian Archipelago and adjacent regions is confirmed. While external morphology is not reliable for identification, a few discrete, although slight, differences in gonopod morphology were found, and these results are consistent with a “pseudocryptic species” designation. Speciation conforms to a previously published etisine model of allopatric differentiation followed by subsequent divergence of gonopod morphology upon secondary sympatry. This pattern, the biogeography of the two species, and the term “pseudocryptic species” are discussed.

Key words: Crustacea, Decapoda, Xanthoidea, Chlorodiellinae, *Pilodius*, taxonomy, systematics, coral reef

INTRODUCTION

The Indo-West Pacific (IWP) is the world's largest marine biogeographic region, characterized by many wide-ranging species that mostly disperse via long-lived, planktonic larvae (Briggs & Bowen, 2012; Kay, 1984; Myers, 1994; Forest & Guinot, 1961). Recent studies have indicated that many species previously thought to range across the IWP comprise mosaics of allo- or parapatric, cryptic lineages (e.g., Meyer *et al.*, 2005; Drew & Barber, 2009; Malay & Paulay, 2010; Titus *et al.*, 2018). One such study of the brachyuran crab clade "Chlorodiellinae" (now Etisinae) found that while some species have IWP-wide distributions with little genetic structuring, others are complexes of deeply-divergent allopatric lineages (Lasley *et al.* 2023). That study also uncovered a strong correlation between genetic distance (time), sympatry, and the divergence of genital morphology, highlighting the important roles of both allopatric genetic differentiation *and* genital divergence in the speciation process. Here the differentiation in one of these genera is examined in greater detail.

Members of the Etisinae Ortmann, 1893 are some of the most abundant crustacean cryptofauna in IWP coral reefs (Monteforte, 1987; Peyrot-Clausade, 1977, 1979, 1989). Despite their ecological importance, abundance and prevalence in museum collections, the taxonomy of this group has proved challenging and needs attention. The molecular phylogenetic study of the superfamily Xanthoidea Macleay, 1838, by Mendoza *et al.* (2022) greatly expanded the Etisinae, merging it with the subfamily Chlorodiellinae Ng & Holthuis, 2007 (sensu Ng *et al.*, 2008). Mendoza *et al.* (2022) further included three xanthine genera, *Leptodius* A. Milne-Edwards, 1863, *Macromedaeus* Ward, 1942, and *Neoxanthops* Guinot, 1968. Although a formal morphological diagnosis for this grouping has yet to be proposed, its members commonly share spoon-tipped chelae and, to a lesser extent, a dactylopropodal lock on the ambulatory legs. Nevertheless, most chlorodielline genera have been recovered in a subclade with high support, and this lineage has been the subject of recent systematic studies (Lai *et al.*, 2011; Lasley *et al.*, 2013, 2015, 2022, 2023; Mendoza *et al.*, 2022). Lasley *et al.* (2015) revised the genus-level taxonomy of 'Chlorodiellinae', and described two genera: *Luniella* Lasley, Klaus & Ng, 2015 and *Soliella* Lasley, Klaus & Ng, 2015.

Three nominal species are attributed to *Soliella*: *Pilodius flavus* Rathbun, 1894, *Chlorodopsis melanospinis* Rathbun, 1911, and *Chlorodopsis hawaiiensis* Edmondson, 1962. Clark & Galil (1993) considered all three to pertain to *S. flava*. Lasley *et al.* (2015) also recognized *S. melanospinis* as valid based on morphology of the male gonopod (G1) and sequence data from two specimens but did not evaluate *C. hawaiiensis*. The differences between the species' G1s remain unclear, as do external morphological differences, historical literature, and geographic distributions of these species.

To solidify the taxonomy of *Soliella* and investigate speciation in the genus, we conducted genus-level phylogenetic analyses using the DNA barcoding gene COI, and morphological examination, including scanning electron microscopy of male genital structures (first gonopod

or “G1”), of 100s of specimens. Historical records were also reviewed especially to assess the geographic distributions of the two species.

MATERIAL AND METHODS

Specimens for morphological and molecular analyses were obtained from the following institutions: Zoological Reference Collection of the Lee Kong Chian Natural History Museum, National University of Singapore, Singapore (ZRC); Florida Museum of Natural History, University of Florida, Gainesville, Florida, USA (UF); U.S. National Museum of Natural History, Smithsonian Institution, Washington D.C., USA (USNM); and American Museum of Natural History, New York, New York, USA (AMNH). Historical literature and material examined are covered in SM1.

Morphological examination was conducted using a dissecting microscope (Leica MZ16, Leica Biosystems, Wetzlar, Germany) and a scanning electron microscope (SEM) (Leica Stereoscan 440 at the USNM Imaging Laboratory). The right first and second male gonopods (G1, G2) were removed for examination unless they were damaged, in which case the left one was removed. G1s were prepared for SEM as described by Felgenhauer (1987) and Lasley et al. (2022). Geographic ranges were compiled from locality information from material examined and literature. These data were checked against locality information associated with COI sequences when possible. Occurrence maps were generated with the R package *ggplot2* (Wickham, 2016).

COI sequence data were obtained from Lasley et al. (2023), including sequences of *Pilodius maotieni* Serène, 1971, *Luniella spinipes* (Heller, 1860), and *Cyclodius granulatus* (Targioni Tozzetti, 1877) as outgroups (Lasley et al., 2015) (Table 1). Maximum likelihood trees were generated using RAxML-HPC BlackBox 8.2.12 (Stamatakis, 2014) in the computer cluster of CIPRES (CyberInfrastructure for phylogenetic RESearch project) (<http://www.phylo.org>; Miller et al., 2010). The GTRGamma + I model of nucleotide substitution was selected and the analysis was conducted with 1,000 bootstrap replicates. A Neighbor-Joining analysis using the Tamura-Nei genetic distance model was also performed in Geneious 8.1.9 with 1,000 bootstrap replicates. Between group mean P-distance between species was calculated using Mega version 11.0.13.

Nomenclature and terminology follow Dana (1851), Serène (1984), Ng et al. (2008) and Davie et al. (2015). Measurements provided (in millimeters) are of the maximum carapace width and length, respectively. The following abbreviations are used: G1, male first gonopod; G2, male second gonopod; stn., station; and coll., collected by. Works by Raoul Serène’s Vietnamese assistant, Nguyen Van Luom, have erroneously been referred to using one of his given names,

“Luom”, rather than his surname “Nguyen”, in previous studies. Here the name is used in full, “Nguyen Van Luom”, e.g., Serène & Nguyen Van Luom (1958), for clarity (Waterman, 1953).

RESULTS

Specimens grouped into two species based on G1 morphology and these corresponded to two reciprocally-monophyletic COI clades separated by 11.3 % P distance (Fig. 1). The *S. melanospinis* clade comprises individuals from the Western Indian Ocean to the Indo-Australia Archipelago and adjacent areas: the Scattered Islands, Reunion Island, Chagos Archipelago, Indonesia (Aceh and Bali), Ningaloo Reef (W. Australia), Taiwan, Okinawa, Palau, Heron Island (Great Barrier Reef), the Solomon Islands, and Vanuatu. The *S. flava* clade comprises individuals from the Indo-Australian Archipelago and adjacent areas to the Eastern Pacific Barrier: Christmas Island (Indian Ocean), the Philippines, Indonesia (Bali and Sulawesi), Guam, New Caledonia, Line Islands, Society Islands, Tuamotu Islands, and Hawaiian Islands (Fig. 2).

TAXONOMY

Superfamily Xanthoidea MacLeay, 1838

Family Xanthidae MacLeay, 1839

Subfamily Etisinae Ortmann, 1893

***Soliella* Lasley, Klaus & Ng, 2015**

Chlorodopsis, Rathbun, 1911: 226. – Balss, 1938: 58. – Serène & Nguyen Van Luom, 1958: 88; 1959: 336.

Pilodius, Balss, 1938: 56. – Forest & Guinot, 1961: 81. – Serène, 1984: 233. – Clark & Galil, 1993: 1121. – Ng et al., 2008: 197.

Soliella Lasley et al., 2015: 173

Diagnosis. Carapace transversely subhexagonal, dorsal surface granular, covered with short and long, light-coloured setae, regions well defined. Front sinuous, quadrilobate; submedian lobes broadly arched, separated by median, narrow, U-shaped notch, margins granular or spinose; lateral lobes distinct, narrow. Anterolateral margin with four lobes, each tipped with emergent, anteriorly directed spine surrounded by smaller accessory spines. Basal antennal article with distolateral extension reaching approximately halfway into orbital hiatus. Male thoracic sternum relatively broad; suture 3/4 distinct near lateral margins, interrupted medially; median line present on sternite 4 as short suture or shallow depression midway between anterior border of sternite and sterno-pleonal cavity, interrupted in exposed posterior part, reappearing

in sterno-pleonal cavity on posterior surface of sternite 4; tubercle of press-button locking mechanism located on anterior half of sternite 5. External, superior surfaces of chelipeds spinose, granular, with numerous long, simple, yellow setae. Ambulatory legs relatively stout; dactylopropodal lock present, well developed; tip of dactylus terminating in long, curved, chitinous claw and two subdistal, small, calcareous spines. Pleon relatively long, slender, tip of telson reaching beyond imaginary transverse line connecting sternal condyles of P1 coxae; pleonites 3–5 functionally fused, with distinct furrows delineating 3/4 and 4/5; telson subtriangular, basal width slightly greater than median length. G1 narrow, sinuous but not drastically curved; distal tip tubular or spatulate with numerous subdistal, proximally directed, spiniform setae. G2 ca. one-third length of G1, sigmoidally curved, terminal segment ca. one-fourth length of subterminal segment. Penis emerging at anterior portion of sternal condyle of P5 coxa.

Remarks. Lasley et al. (2015) provided a diagnosis of the genus and compared it with four genera that were, along with *Soliella*, previously classified in the subfamily Chlorodiellinae (= Etisinae in part): *Chlorodiella* Rathbun, 1897, *Cyclodius* Dana, 1851, *Pilodius* Dana, 1851, and *Luniella*. *Soliella* differs from these genera, most notably, in the morphology of its G1 (Fig. 3) (Serène 1984, figs. 144–158, 163–165, 167–172, 173–177). *Soliella* also differs from all species in these genera with the exceptions of *Luniella pubescens* (Dana, 1852), *Luniella scabricula* (Dana, 1852), and *Cyclodius paumotensis* (Rathbun, 1907), by the presence of long and short, light-colored setae on the carapace. Traditional characters, such as the shape and disposition of the basal antennal article and the form of the subterminal (bifid) spine of the ambulatory leg dactylus, that have been used to differentiate these genera are problematic and are not shared with the closest relatives of *Soliella*: *Cyclodius* and *Pilodius* (Ng & Yang, 1998; Clark & Ng, 1999; Lasley et al., 2015). The length of the distolateral extension of the basal antennal article varies in some genera (e.g., *Pilodius*) and with age (Lasley et al., 2015; Serène, 1984: 233, footnote by Crosnier). In *Soliella*, the basal antennal article has a distolateral extension that reaches approximately halfway into orbital hiatus (vs no extension in *Cyclodius* and usually reaching the orbital hiatus in *Pilodius*). *Soliella* has small, calcareous subterminal spines of the ambulatory leg dactylus, while the presence and length of subterminal spines vary in *Pilodius* and *Cyclodius* (Lasley et al., 2015). Relationships between *Soliella* and more distantly related genera that were previously assigned to Chlorodiellinae were reviewed in Lasley et al. (2015), e.g., *Tweedieia* Ward, 1935, *Vellodius* Ng & Yang, 1998, and *Sulcodius* Clark & Ng, 1999. All other etisine genera have been treated in Serène (1984). In view of the results from recent molecular phylogenetic studies on Xanthidae (Lai et al. 2011; Lasley et al. 2015; Mendoza et al., 2022), however, the diagnoses for the different genera in an expanded Etisinae will need to be re-evaluated and emended, with greater focus on thoracic sternal characters and other such non-traditional but informative characters.

Soliella flava (Rathbun, 1894)

Figs. 3A, C, 4, 5

Pilodius flavus Rathbun, 1894: 239; 1906: 860, fig. 21. – Edmondson, 1925: 43; 1933: 249; 1962: 275, fig. 22a, b. – Balss, 1938: 57. – Miyake, 1939: 215. – Forest & Guinot, 1961: 95. – Serène, 1968: 80; 1984: 235, 239 [key]. – Peyrot-Clausade, 1989: 111. – Clark & Galil, 1993: 1130 (in part), figs. 4A – G, 32B, 40D, 41A. – DeFelice et al., 1998: 16; 2002: 30, 72. Coles et al., 2002a: 271 (list); 2002b: 141, 194; 2008: 63 (list) – Ng et al., 2008: 197 (list). – Mendoza et al., 2014: 278.

Chlorodopsis flava, Serène & Nguyen Van Luom, 1959: 330, figs. 2C, 5F, pl. 1 fig. B, pl. 3 fig. B.

Chlorodopsis hawaiiensis Edmondson, 1962: 273, fig. 21a–e.

Soliella flava, Lasley et al., 2015: 174, suppl. figs. S1D, S3C, D, S5F.

Chlorodopsis melanodactylus, Miers, 1884: 531 (in part, from Etoile Island). Not *Pilodius melanodactylus* A. Milne-Edwards, 1873. [fide Clark & Galil, 1993].

Pilodius pubescens, De Man, 1902: 619. Not *Pilodius pubescens* Dana, 1852 [fide Balss, 1938].

?*Pilodius pubescens*, Nobili, 1907: 395. Not *Pilodius pubescens* Dana, 1852 [fide Balss, 1938].

Diagnosis. Carapace (Figs. 4, 5A–B) transversely subhexagonal, ca. 1.5 as broad as long; surface covered in short, stout and few long, light-colored setae; regions well defined, separated by distinct, smooth furrows; 1F indistinct; 2F distinct; 1M separated from 2F and inner branch of 3M by shallow furrow; 2M entire or feebly divided anteriorly, 3M entire; 4M indistinct; 1L indistinct; 1L and 2L partially confluent; 3L–6L distinct; 1P with defined anterior and posterior borders, lateral borders diffuse; 2P with transverse row of granules. Submedian lobes of front (Fig. 5C) broadly convex, margin lined with granules, separated by median V- or U-shaped notch; lateral lobes triangular, granulate, separated from submedian lobes by deep, triangular notch, separated from orbits/supraorbital margin by rounded, L-shaped notch. Supraorbital margin lined dorsolaterally with short spines or conical granules; infraorbital margin lined with conical granules. Anterolateral margin with four spinose lobes. Anterolateral angle of basal antennal article-slightly expanded, entering less than halfway into orbital hiatus. Pterygostomial region minutely granulate, with plumose setae diagonally from posterior to lateral surface. Male thoracic sternum (Fig. 5D) relatively broad, minutely granulate, with few long, scattered setae; tubercle of press-button locking mechanism located on anterior half of sternite 5; suture 3/4 distinct near lateral margins, interrupted medially; median line present on sternite 4 as short suture midway between anterior border of sternite and sterno-pleonal cavity, interrupted in exposed posterior part, reappearing in sterno-pleonal cavity on posterior surface of sternite 4, absent at level of sternites 5 and 6, present and complete at level of sternites 7 and 8. Chelipeds (Fig. 5E–F) subequal, covered with long, simple, light-colored setae, spinose; merus stout. Ambulatory legs (Fig. 5A) stout, setose; setae long, simple, light-colored; extensor margin of merus lined with long spines; dactylopropodal lock present, well developed; tip of dactylus terminating in long, curved, chitinous claw and two subdistal, small, calcareous spines. Male

pleon (Fig. 5D) moderately stout, with few long posterior setae; pleonites 3–5 functionally fused, with distinct furrows delineating 3/4 and 4/5; pleonite 6 subquadrate, ca. as broad as long; telson subtriangular, ca. as broad as long. G1 (Fig. 3A, C) slender, sinuous, distal 1/4 curved ventrally; apex pointing anteroventrally with ca. 20 subdistal, perpendicular to proximally-directed, stout, spiniform setae on the anterior surface; apical lobe almost tubular, opening facing anteriorly. G2 ca. one-third length of G1, sigmoidally curved, terminal segment ca. one-fourth length of subterminal segment.

Female morphology. Females are similar to males, except in having nearly equal chelipeds and in sexual characters. Sternopleonal cavity wide, with the median line obscured completely by the pleon; sutures 2/3, 6/7, and 7/8 complete; suture 3/4 indicated only near lateral margin; sutures 4/5 and 5/6 interrupted medially. Vulvae crescent-shaped, positioned on sternite 6 near suture 5/6. Pleon long and wide relative to male; tip of telson reaching imaginary line between midpoint of cheliped coxae; all pleonites freely articulated.

Type status. The female holotype (USNM17317) from the Hawaiian Islands was examined for this study (SM1).

Remarks. *Soliella flava* and *S. melanospinis* are difficult to differentiate based on external morphology. Rathbun (1894, 1911) described both species. In her description of *S. melanospinis*, Rathbun (1911) stated that *S. flava* has a less deeply areolated carapace, a dorsum devoid of spines, and an upper margin of the orbit (supraorbital margin) without spines (vs. less deeply areolated regions, a spinose dorsum, and upper margin of the orbit in *P. melanospinis*). Serène (1984) stated that the spination on the supraorbital margin was a good character for differentiation, but that the difference in the areolation of the carapace was difficult to assess. He also stated that the G1s are similar, although he had provided figures of the two in his previous publications with Nguyen Van Luom (Serène & Nguyen Van Luom, 1958: pl. 4 fig. f; 1959: figs. 2C, 2 bis M).

Edmondson (1962) described *Chlorodopsis hawaiiensis* without comparison with *S. flava* or *S. melanospinis*. He also provided illustrations of their G1s. Clark & Galil (1993) synonymized *S. melanospinis* and *S. hawaiiensis* with *S. flava*. However, Lasley *et al.* (2015, 2023) recovered two distinct, divergent species-level clades in *Soliella* in their molecular phylogenetic analyses, while there are three different G1 morphotypes illustrated in literature. Edmondson (1962: figs 21d, 22b) provided figures of the ladle-like G1 of *Chlorodopsis hawaiiensis* and the tubular G1 of *S. flava*, illustrating them with distinct morphologies albeit in a simplistic, even schematic, style (Fig. 3C). Rathbun's (1894) Hawaiian holotype of *S. flava* is female. Examination of many Hawaiian specimens (SM1), however, including those previously identified as *Chlorodopsis hawaiiensis* and *S. flava* makes it clear that the G1s show only slight variation that had been exaggerated in the figures of Edmondson (1962). These gonopod morphotypes fall within the *S. flava* COI clade in the present analysis. The third G1 morphotype was illustrated by Clark & Galil

(1993: fig. 4D–G) as *S. flava*, although their specimen is a paratype of *S. melanospinus* (Fig. 3D). This is the same morphotype as those illustrated by Serène & Nguyen Van Luom (1959: fig 2M) and Serène (1984: fig. 146), but in these studies, they are identified as *S. melanospinus*.

In summary, the external morphological characters of Rathbun (1911) are difficult to appreciate, but G1 morphology and phylogenetic analyses indicate that there are clearly two species. Although the depth of the furrows separating the carapace regions and spination of the supraorbital margin vary, *S. flava* specimens do generally have less defined carapace regions and a supraorbital margin with shorter spines or conical granules (vs less relatively deeply defined regions and supraorbital margin with larger spines in *S. melanospinus*). These characters, however, display too much variation, especially in small individuals, to be used without caution. The G1 morphology is the only reliable character for identification. *Soliella flava* has a G1 pointing anteroventrally with an apical lobe opening anteriorly and ca. 20 spiniform subdistal setae on the anterior surface (vs apex pointing ventrally with an apical lobe that is longitudinally hollowed with a sinuous anterior margin and ca. 12 subdistal setae; Fig. 3). The two G1 morphotypes correspond with the well-supported clades in the phylogenetic analyses.

Distribution. *Soliella flava* is reported from Christmas Island (Indian Ocean) and the Indo-Australian Archipelago to the Hawaiian Islands and French Polynesia (Fig. 2).

***Soliella melanospinus* (Rathbun, 1911)**

Figs. 3B, D, 6

Chlorodopsis melanospinus Rathbun, 1911: 226, pl. 18 fig. 11. – Balss, 1938: 62. – Serène &

Nguyen Van Luom, 1958: 108, pl. 1 fig. D, pl. 3 fig. b, pl. 4 fig. c; 1959: 302, fig. 2 bis M.

Pilodius melanospinus, Guinot, 1964: 67; 1967: 268. – Serène, 1968: 80; 1984: 242, figs. 143e, 146, pl. 33 fig. E.

Pilodius flavus, Clark & Galil, 1993: 1130 (in part). – Ng et al., 2008: 197 (list).

Chlorodopsis pilumnoides, Laurie, 1906: 406 (from Ceylon = Sri Lanka). Not *Pilodius pilumnoides* (White, 1848) [fide Clark & Galil, 1993].

Diagnosis. Carapace (Fig. 6A-B) transversely hexagonal, ca. 1.5 as broad as long; surface covered with short, stout light-colored setae and few long, light-colored setae; regions well defined, separated by wide, smooth, relatively deep furrows; 1F indistinct; 2F distinct; 1M separated from 2F and inner branch of 3M by shallow furrow; 2M entire or feebly divided anteriorly, 3M entire; 4M indistinct; 1L indistinct; 1L and 2L partially confluent; 3L-6L distinct; 1P with defined anterior and posterior borders, lateral borders diffuse; 2P with transverse row of granules. Submedian lobes of front (Fig. 6C) broadly convex, margin lined with granules,

separated by median V- or U-shaped notch; lateral lobes triangular, separated from submedian lobes by deep, triangular notch, separated from orbits/supraorbital margin by rounded, L-shaped notch. Supraorbital margin generally lined dorsolaterally with relatively long spines or conical granules; infraorbital margin lined with conical granules. Anterolateral margin with four spinose lobes. Anterolateral angle of basal antennal segment slightly expanded, entering less than halfway into orbital hiatus. Pterygostomial region minutely granulate, with plumose setae diagonally from posterior to lateral surface. Male thoracic sternum (Fig. 4D) relatively broad, minutely granulate, with few long, scattered setae; tubercle of press-button locking mechanism located on anterior half of sternite 5; suture 3/4 distinct near lateral margins, interrupted medially; median line present on sternite 4 as short suture midway between anterior border of sternite and sterno-pleonal cavity, interrupted in exposed posterior part, reappearing in sterno-pleonal cavity on posterior surface of sternite 4, absent at level of sternites 5 and 6, present and complete at level of sternites 7 and 8. Chelipeds (Fig. 6E-F) subequal, covered with long, simple, light-colored setae, spinose; merus stout. Ambulatory legs (Fig. 6A) stout, setose; setae long, simple, light-colored; extensor margin of merus lined with long spines; dactylopropodal lock present, well developed; tip of dactylus terminating in long, curved, chitinous claw and two subdistal, small, calcareous spines. Male pleon (Fig. 6D) moderately stout, few long posterior setae; pleonites 3–5 functionally fused, with distinct furrows delineating 3/4 and 4/5; pleonite 6 subquadrate, ca. broad as long; telson subtriangular ca. as broad as long. G1 (Fig. 3B, D) slender, sinuous, distal 1/4 curved ventrally; apex pointing ventrally with ca. 12 subdistal, perpendicular to proximally-directed, stout, spiniform setae on the anterior surface; apical lobe almost spatulate, longitudinally hollowed with sinuous anterior margin. G2 ca. one-third length of G1, sigmoidally curved, terminal segment ca. one-fourth length of subterminal segment.

Female morphology. Females are similar to males, except in having nearly equal chelipeds and in sexual characters. These characters are the same as those outlined for *Soliella flava* females (see above).

Remarks. See Remarks for *Soliella flava*.

Distribution. *Soliella melanospinis* occurs from the Western Indian Ocean to the Indo-Australian Archipelago and adjacent areas including Taiwan, Japan, Palau, the Solomon Islands, and Vanuatu (Fig. 2).

Type status. The male holotype (USNM 41268) from Saya del Malha Bank, Western Indian Ocean, was examined for this study (SM1).

Key to the Species of *Soliella*.

- G1 ultimately pointing anteroventrally with apical lobe opening anteriorly. Carapace regions relatively less defined. Supraorbital margin with relatively low spines or conical granules.....*S. flava*
- G1 apex pointing ventrally with an apical lobe that is longitudinally hollowed with a sinuous anterior margin (Pl. 45). Carapace regions relatively well defined. Supraorbital margin generally with longer spines.....*S. melanospinus*

DISCUSSION

The synonymy of *S. melanospinus* with *S. flava* by Clark & Galil (1993) reflects on the morphological similarity between the two species. The examination here further demonstrates this similarity: there are no external features that can reliably distinguish these two species. The two species, however, have discrete, although relatively slight, differences in G1 morphology (compare Fig. 3 with G1 figures of chlorodiellines in Serène 1984 and Lasley et al., 2023). Therefore, the term “pseudocryptic” is used because: 1) molecular data guided the discovery of the species distinctions, and 2) there are minor but reliable morphological differences in a previously lumped species. The agreement between reciprocal monophyly in COI (here) and other markers (Lasley et al., 2023) with discrete differences in genital structures substantiates the use of the term “species”, especially considering their sympatric ranges and that genital divergence in arthropods is commonly used to infer reproductive isolation (Eberhard, 1985).

Lasley et al. (2023) showed a correlation between secondary sympatry and divergence of G1s among species in the clade Chlorodiellinae, which includes *Soliella*. The implication is that these crabs differentiate first in allopatry (or technically parapatry if some degree of homogenizing gene flow was present) and secondary contact is accompanied, or allowed, by G1 divergence. For analysis in the study, lineages were categorized as, a) sharing a G1 morphology with its sibling lineage, or b) possessing a unique G1; and geographic distribution was categorized as allopatric, narrowly sympatric, or sympatric. “Narrowly sympatric” sibling lineages were defined as those having less than 10 percent overlap in total distribution. *Soliella melanospinus* and *S. flava* were coded as having unique G1s and sympatric distributions (Figs. 2, 3). This differs from many sibling lineages in Chlorodiellinae that show less genetic divergence, are allopatric, and share the same G1 morphology. Although the ranges of the two *Soliella* broadly overlap in the West Pacific, *S. melanospinus* is the sole species through most of the Indian Ocean while only *S. flava* is known from remote Oceania in the central Pacific. Allopatric divergence between the Indian and Pacific Ocean basins is the most prevalent geographic differentiation in IWP marine taxa (Barber et al., 2000; Malay & Paulay, 2010, Ahti et al., 2016). The distribution of *Soliella* is suggestive of a similar history of allopatric divergence, followed by secondary range overlap in the West Pacific, likely allowed or facilitated by genital divergence, but accompanied by little other morphological differentiation.

Our limited COI dataset indicates panmixia in both species—not-uncommon in marine organisms with long larval durations. Population genomic data could, however, indicate fine-scale divergence and/or directionality of gene flow. These data could shed more light on the geographic origins of, and processes that govern speciation in, these species.

ACKNOWLEDGEMENTS

We are grateful to numerous people that have aided in sequencing and other lab work, and the many more people that have helped in the field or provided specimens. We also thank Mandy Bemis and John Slapcinsky (UF) for their efforts in curation. We thank Peter Ng for his taxonomic expertise and input, which have been invaluable in unraveling chlorodielline systematics. This work was supported by the National Science Foundation [NSF DEB 1856245, GECCO 1457769], the Moore Foundation [Moorea BioCode], and the Sloan Foundation [MarBOL].

REFERENCES

Ahti, P. A., Coleman, R. R., DiBattista, J. D., Berumen, M. L., Rocha, L. A., & Bowen, B. (2016). Phylogeography of Indo-Pacific reef fishes: sister wrasses *Coris gaimard* and *C. cuvieri* in the Red Sea, Indian Ocean and Pacific Ocean, *Journal of Biogeography* 43(6), 1103-1115. <https://doi.org/10.1111/jbi.12712>

Barber, P. H., Palumbi, S. R., Erdmann, M. V., & Moosa, M. K. (2000). A marine Wallace's line? *Nature*, 406, 692-693. <https://doi.org/10.1038/35021135>

Briggs, J. C., & Bowen, B. W. (2012). A realignment of marine biogeographic provinces with particular reference to fish distributions. *Journal of Biogeography*, 39(1), 12-30. <https://doi.org/10.1111/j.1365-2699.2011.02613.x>

Clark, P. F., & Galil, B. S. (1993). A revision of the xanthid genus *Pilodius* Dana, 1851 (Crustacea: Brachyura: Xanthoidea). *Journal of Natural History*, 27(5), 1119-1206. <https://doi.org/10.1080/00222939300770691>

Dana, J. D. (1851). On the classification of the Cancroidea. *American Journal of Science and Arts, Series 2*, 12, 121-131.

Davie, P. J., Guinot, D., & Ng, P. K. L. (2015). Phylogeny of Brachyura pp. 921-979. In P. Castro, P. J. F. Davie, D. Guinot, F. R. Schram & J. C. Von Vaupel Klein (Eds.), *Decapoda: Brachyura*,

Treatise on Zoology – Anatomy, Taxonomy, Biology. The Crustacea, Complementary to the volumes translated from the French of the *Traité de Zoologie* [founded by P.-P. Grassé (†)] 9C-II: 639-1221. Brill, Leiden and Boston.

Drew, J., & Barber, P. H. (2009). Sequential cladogenesis of the reef fish *Pomacentrus moluccensis* (Pomacentridae) supports the peripheral origin of marine biodiversity in the Indo-Australian archipelago. *Molecular Phylogenetics and Evolution*, 53(1), 335.
<https://doi.org/10.1016/j.ympev.2009.04.014>

Edmondson, C. H. (1962). Xanthidae of Hawaii. *Occasional Papers of Bernice P. Bishop Museum*, 22(13), 215-309.

Felgenhauer, B. E. (1987). Techniques for preparing crustaceans for scanning electron microscopy. *Journal of Crustacean Biology*, 7, 71-76.
<https://doi.org/10.1163/193724087X00054>

Forest, J. & Guinot, D. (1961). Crustacés Décapodes Brachyoures de Tahiti et des Tuamotu. In: Expédition française sur les récifs coralliens de la Nouvelle-Calédonie. Volume préliminaire. Editions de la Fondation Singer Polignac, Paris, 9-11, 1-195, pls. 1-18.

Guinot, D. (1968). Recherches préliminaires sur les groupements naturels chez les Crustacés Décapodes Brachyoures. IV. Observations sur quelques genres de Xanthidae. *Bulletin du Muséum national d'Histoire naturelle, Paris*, 2e série 39, 695-727.

Heller C. (1860). Beiträge zur Crustaceen-Fauna des rothen Meeres. Erster Theil. – Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der kaiserlichen Akademie der Wissenschaften, 43(1), 297-374; pls. I-IV.

Kay, E. A. (1984). Patterns of speciation in the Indo-west Pacific. pp. 15-31 In P. H. Raven, F. J Radovsky, & S. H. Sohmer, (Eds.) Bio-geography of the tropical Pacific. Association of Systematic Collections and B. P. Bishop Museum. x + 221 pp.

Lasley, R. M., Lai, J. C., & Thoma, B. P. (2013). A new genus for *Chlorodiella longimana* (H. Milne Edwards) supported by morphology and molecular data, with a preliminary phylogeny of the Chlorodiellinae (Crustacea: Decapoda: Xanthidae). *Invertebrate Systematics*, 27(4), 379-390.
<http://dx.doi.org/10.1071/IS12075>

Lasley Jr, R. M., Klaus, S., & Ng, P. K. L. (2015). Phylogenetic relationships of the ubiquitous coral reef crab subfamily Chlorodiellinae (Decapoda, Brachyura, Xanthidae). *Zoologica Scripta*, 44(2), 165-178. <https://doi.org/10.1111/zsc.12094>

Lasley Jr, R. M., Smith, A., Paulay, G., & Ng, P. K. L. (2022). Revision of the coral reef crab genus *Tweedieia* Ward, 1935 (Decapoda: Brachyura: Xanthidae). *Journal of Crustacean Biology*, 42(1), ruab081. <https://doi.org/10.1093/jcbiol/ruab081>

Lasley Jr, R. M., Evans, N., Paulay, G., Michonneau, F., Windsor, A., & Ng, P. K. L. (2023). Allopatric mosaics in the Indo-West Pacific crab subfamily Chlorodiellinae reveal correlated patterns of sympatry, genetic divergence, and genitalic disparity. *Molecular Phylogenetics and Evolution*, 181, 107710. <https://doi.org/10.1016/j.ympev.2023.107710>

MacLeay, W. S. (1838). On the brachyurous decapod Crustacea brought from the Cape by Dr. Smith, In Smith A., Illustrations of the Zoology of South Africa; Consisting Chiefly of Figures and Descriptions of the Objects of Natural History Collected During an Expedition into the Interior of South Africa, in the Years 1834, 1835, and 1836; Fitted out by 'The Cape of Good Hope Association for Exploring Central Africa': Together with a Summary of African Zoology, and an Inquiry into the Geographical Ranges of Species in that Quarter of the Globe. Invertebratae. Smith, Elder & Co., London: 53-71, pls. 2, 3.

Malay, M. C. M. D., & Paulay, G. (2010). Peripatric speciation drives diversification and distributional pattern of reef hermit crabs (Decapoda: Diogenidae: Calcinus). *Evolution: International Journal of Organic Evolution*, 64(3), 634-662. <https://doi.org/10.1111/j.1558-5646.2009.00848.x>

Mendoza, J. C. E., Lasley R. M. Jr, & Ng P. K. L. (2014) New rock crab records from Christmas and Cocos (Keeling) Islands, Eastern Indian Ocean. *Raffles Bulletin of Zoology*, Suppl. No. 30, 274-300.

Mendoza, J. C., Chan, K. O., Lai, J. C., Thoma, B. P., Clark, P. F., Guinot, D., Felder, D. L., & Ng, P. K. L. (2022). A comprehensive molecular phylogeny of the brachyuran crab superfamily Xanthoidea provides novel insights into its systematics and evolutionary history. *Molecular Phylogenetics and Evolution*, 177, 107627. <https://doi.org/10.1016/j.ympev.2022.107627>

Meyer, C. P., Geller, J. B., & Paulay, G. (2005). Fine scale endemism on coral reefs: archipelagic differentiation in turbinid gastropods. *Evolution*, 59(1), 113-125. <https://doi.org/10.1111/j.0014-3820.2005.tb00899.x>

Miller, M. A., Pfeiffer, W., & Schwartz, T. (2011). The CIPRES science gateway: a community resource for phylogenetic analyses. In *Proceedings of the 2011 TeraGrid Conference: extreme digital discovery*, pp. 1-8.

Milne-Edwards, A. (1863). Monographie des Crustacés Fossiles de la Famille des Canceriens. *Annales des Sciences Naturelles*, 4e série 20, 273-324, pls. 5-12.

Monteforte, M. (1987). The decapod Reptantia and stomatopod crustaceans of a typical high island coral reef complex in French Polynesia (Tiahura, Moorea Island): zonation, community composition and trophic structure. *Atoll Research Bulletin*, 309, 1-37.
<https://doi.org/10.5479/si.00775630.309.1>

Myers, A. A. (1994). Biogeographic patterns in shallow-water marine systems and the controlling process at different scales. In P.S. Giller, A. G. Hildrew, & D. G. Raffaelli, (Eds.). *Aquatic Ecology, Scale, Pattern and Process* British Ecological Society Symposium, Blackwell Scientific, Oxford, pp. 547-574.

Ng, P. K. L., & Holthuis, L. B. (2007). Case 3394. *Etisus* H. Milne Edwards, 1834 and *Chlorodiella* Rathbun, 1897 (Crustacea, Decapoda, Brachyura): proposed conservation of the generic names by suppression of the generic name *Clorodius* A. G. Desmarest, 1823. *Bulletin of Zoological Nomenclature* 64, 19-24.

Ng, P. K. L., Guinot, D., and Davie, P. J. F. (2008). Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. *The Raffles Bulletin of Zoology* (Supplement 17), 1-286.

Ortmann, A. (1893). Die Decapoden-Krebse des Strassburger Museums, mit besonderer Berücksichtigung der von Herrn Dr. Döderlein bei Japan und bei den Liu-Kiu-Inseln gesammelten und zur Zeit im Strassburger Museum aufbewahrten Formen. VII. Theil. Abtheilung: Brachyura (Brachyura genuina Boas) II. Unterabtheilung: Cancroidea, 2. Section: Crancinea, 1. Gruppe: Cyclometopa. *Zoologische Jahrbücher. Abteilung für Systematik, Ökologie und Geographie der Tiere* 7(3), 411-495, pl. 17.

Peyrot-Clausade, M. (1977). Decapodes brachyures et anomoures (à l'exception des Paguridae) de la cryptofaune de Tiahura Ile de Moorea. Extrait des Cahiers du Pacifique, n°, 20, 211-222.

Peyrot-Clausade, M. (1979). Cryptofaune annelidienne et carcinologique des récifs de l'île de la Réunion et de l'île Maurice. *The Mauritius Institute Bulletin*, 8, 1-41.

Peyrot-Clausade, M. (1989). Crab crypto fauna (Brachyura and Anomura) of Tikehau, Tuamotu Archipelago, French Polynesia. *Coral Reefs*, 8, 109-117.

Rathbun, M. J. (1894). Scientific results of explorations by the U.S. Fish Commission steamer Albatross. No. XXIV.—Descriptions of new genera and species of crabs from the west coast of North America and the Sandwich Islands. *Proceedings of the United States National Museum* 16(933), 223-260.

Rathbun, M. J. (1911). No XI.—Marine Brachyura. In: The Percy Sladen Trust Expedition to the Indian Ocean in 1905 under the leadership of Mr. J. Stanley Gardiner, Volume III. *Transactions of the Linnaean Society of London*, (2)14(2), 191-261, pls. 15-20.

Serène, R. (1971). Observations préliminaires sur des Brachyures nouveaux ou mal connus du Sud-Est Asiatique (Crustacea Decapoda). *Bulletin du Muséum national d'Histoire naturelle*, (2)42(5), 903-918.

Serène, R. (1984). Crustacés Décapodes Brachyures de l'océan Indien occidental et de la mer Rouge, Xanthoidea: Xanthidae et Trapeziidae. Avec un addendum par Crosnier A.: Carpiliidae et Menippidae. *Faune tropicale*, 24, 1-243.

Serène, R. & Nguyen V. L. (1958). *Chlorodopsis* (Brachyure) du Viêtnam. *Annales de la Faculté des Sciences, Saigon*, 36, 87-147, pls. 1-4.

Serene, R. & Nguyen V. L. (1959). Note additionnelle sur les espèces de *Chlorodopsis* (Brachyures). *Annales de la Faculté des Sciences, Saigon*, 42, 301-340.

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30(9), 1312-1313.
<https://doi.org/10.1093/bioinformatics/btu033>

Targioni Tozzetti, A. (1877). Zoología del viaggio intorno al globo della R. Pirocorvetta Magenta durante gli anni 1865-68. Crostacei Brachiuri e Anomuri. Pubblicazioni del Reale Istituto di Studi Superiori Pratici e di Perfezionamento in Firenze, Sezione di Scienze Fisiche e Naturali. Vol. 1(24). Firenze: Reale Istituti di Studi Superiori, Sezione di Scienze Fisiche e Naturali. 257 pp, pls. 1-13.

Titus, B. M., Daly, M., Hamilton, N., Berumen, M. L., & Baeza, J. A. (2018). Global species delimitation and phylogeography of the circumtropical 'sexy shrimp' *Thor amboinensis* reveals a cryptic species complex and secondary contact in the Indo-West Pacific. *Journal of Biogeography*, 45(6), 1275-1287. <https://doi.org/10.1111/jbi.13231>

Ward, M. (1935). Notes on a collection of crabs from Christmas Island, Indian Ocean. *Bulletin of the Raffles Museum* 9, 5-28, pls. I-III.

Ward, M. (1942). Notes on the Crustacea of the Desjardins Museum, Mauritius Institute, with descriptions of new genera and species. *The Mauritius Institute Bulletin* 2, 49-108, pls. 5-6.

Waterman, T.H. (1953) Xiphosura from Xuong-Ha. *American Scientist*, 41(2), 292-302.

Wickham, H., Chang, W., & Wickham, M. H. (2016). Package ‘ggplot2’. Create elegant data visualizations using the grammar of graphics. *Version*, 2(1), 1-189.

CAPTIONS

Figure 1. Maximum likelihood phylogenetic tree of *Soliella* with bootstrap values from the RAxML analysis followed by bootstrap values from the Neighbor Joining analysis.

Figure 2. Geographic distributions of *Soliella flava* and *S. melanospinis*. Turquoise dots represent material examined. Pink dots represent additional localities recorded in literature.

Figure 3. First male gonopods (G1) of *Soliella* species; A, *Soliella flava* (Rathbun, 1894), G1, internal detail (scale = 200um), external detail (scale = 200um), and external full (scale = 1mm) (UF 12254); B, *Soliella melanospinis* (Rathbun, 1911), G1, internal detail (scale = 200um), external detail (scale = 200um), and external full (scale = 1mm) (ZRC 2013.1647); C, *S. flava* G1 (a) after Edmondson (1962: 21d) as *Chlorodopsis hawaiiensis*, (b) after Edmondson (1962: fig. 22b) as *Pilodius flavus*.

Figure 4. *Soliella flava* (Rathbun, 1894), holotype female, 9 × 6 (USNM 17317), Hawaiian Islands, dorsal view.

Figure 5. *Soliella flava* (Rathbun, 1894), male, 10.2 × 6.9 (USNM 1181377), Marshall Islands; A, dorsal view; B, carapace, dorsal view; C, frontal view; D, thoracic sternum; E, minor chela, external view; F, major chela, external view.

Figure 6. *Soliella melanospinis* (Rathbun, 1911), holotype male, 17.0 × 11.4 (USNM 41268), Saya del Malha Bank; A, dorsal view; B, carapace, dorsal view; C, frontal view; D, thoracic sternum; E, major chela, external view; F, minor chela, external view.