

Allosteric activation or inhibition of PI3K γ mediated through conformational changes in the p110 γ helical domain

Noah J Harris^{1†}, Meredith L Jenkins^{1†}, Sung-Eun Nam², Manoj K Rathinaswamy¹, Matthew AH Parson¹, Harish Ranga-Prasad¹, Udit Dalwadi², Brandon E Moeller¹, Eleanor Sheeky¹, Scott D Hansen³, Calvin K Yip^{2*}, John E Burke^{1,2*}

¹Department of Biochemistry and Microbiology, University of Victoria, Victoria, Canada; ²Department of Biochemistry and Molecular Biology, The University of British Columbia, Vancouver, Canada; ³Department of Chemistry and Biochemistry, Institute of Molecular Biology, University of Oregon, Eugene, United States

Abstract PI3K γ is a critical immune signaling enzyme activated downstream of diverse cell surface molecules, including Ras, PKC β activated by the IgE receptor, and G $\beta\gamma$ subunits released from activated GPCRs. PI3K γ can form two distinct complexes, with the p110 γ catalytic subunit binding to either a p101 or p84 regulatory subunit, with these complexes being differentially activated by upstream stimuli. Here, using a combination of cryo electron microscopy, HDX-MS, and biochemical assays, we have identified novel roles of the helical domain of p110 γ in regulating lipid kinase activity of distinct PI3K γ complexes. We defined the molecular basis for how an allosteric inhibitory nanobody potentially inhibits kinase activity through rigidifying the helical domain and regulatory motif of the kinase domain. The nanobody did not block either p110 γ membrane recruitment or Ras/G $\beta\gamma$ binding, but instead decreased ATP turnover. We also identified that p110 γ can be activated by dual PKC β helical domain phosphorylation leading to partial unfolding of an N-terminal region of the helical domain. PKC β phosphorylation is selective for p110 γ -p84 compared to p110 γ -p101, driven by differential dynamics of the helical domain of these different complexes. Nanobody binding prevented PKC β -mediated phosphorylation. Overall, this work shows an unexpected allosteric regulatory role of the helical domain of p110 γ that is distinct between p110 γ -p84 and p110 γ -p101 and reveals how this can be modulated by either phosphorylation or allosteric inhibitory binding partners. This opens possibilities of future allosteric inhibitor development for therapeutic intervention.

eLife assessment

This study presents **fundamental** new insight into the regulatory apparatus of PI3K γ , a kinase in signaling pathways that control the immune response and cancer. A suite of biophysical and biochemical approaches provide **convincing** evidence for new sites of allosteric control over enzyme activity. The rigorous findings provide structure and dynamic information that may be exploited in efforts to control PI3K γ activity in a therapeutic setting.

Introduction

The class I phosphoinositide 3 kinases (PI3Ks) are master regulators of myriad functions through their generation of the lipid signaling molecule phosphatidylinositol 3,4,5-trisphosphate (PIP₃) downstream of cell surface receptors (Burke and Williams, 2015; Rathinaswamy and Burke, 2020;

*For correspondence:

calvin.yip@ubc.ca (CKY);
jeburke@uvic.ca (JEB)

[†]These authors contributed
equally to this work

Competing interest: See page
18

Funding: See page 18

Preprint posted
12 April 2023

Sent for Review
12 April 2023

Reviewed preprint posted
23 May 2023

Reviewed preprint revised
12 June 2023

Version of Record published
07 July 2023

Reviewing Editor: Amy H
Andreotti, Iowa State University,
United States

© Copyright Harris, Jenkins
et al. This article is distributed
under the terms of the [Creative
Commons Attribution License](#),
which permits unrestricted use
and redistribution provided that
the original author and source
are credited.

Vanhaesebroeck et al., 2021; Vasan and Cantley, 2022). The class I PI3Ks can be further subdivided into the class IA and class IB subfamilies, with class IB PI3Ks being critical in immune signaling, and are composed of a single p110 γ catalytic subunit that can bind to either a p101 or p84 regulatory subunit (*Hawkins and Stephens, 2015; Lanahan et al., 2022; Okkenhaug, 2013*). The two PI3K γ complexes (either p110 γ -p84 or p110 γ -p101) play essential and independent roles in both the adaptive and innate immune systems. Class IB PI3K γ has shown promise as a therapeutic target, primarily as an immunomodulator of the tumor microenvironment leading to enhanced anti-tumor immune responses (*De Henau et al., 2016; Kaneda et al., 2016b*). Multiple isoform selective small molecule ATP competitive inhibitors of p110 γ are in clinical trials for multiple forms of human cancers (*Li et al., 2021*). However, all inhibitors currently developed toward p110 γ act as ATP competitive inhibitors, showing equal potency against both p110 γ -p84 and p110 γ -p101 complexes.

Detailed experiments on the role of p110 γ in mice show that knockout of both p101 and p84 leads to PIP₃ levels that are equivalent to knockout of p110 γ , showing that all cellular PI3K γ activities require the presence of either a p84 or p101 regulatory subunit (*Rynkiewicz et al., 2020*). The two complexes are differentially activated by membrane localized receptors, including G-protein coupled receptors (*Li et al., 2000; Stephens et al., 1997*), Ras (*Jin et al., 2020; Kurig et al., 2009*), toll-like receptors (*Luo et al., 2018*), and the IgE antigen receptor (*Laffargue et al., 2002; Walser et al., 2013*). This leads to the different complexes driving unique immune responses, with p110 γ -p101 involved in chemotaxis in neutrophils (*Bohnacker et al., 2009; Deladeriere et al., 2015*), and p110 γ -p84 involved in reactive oxide production. Differential activation of unique PI3K γ complexes downstream of GPCRs and Ras is caused by the ability of p101 to directly bind to G $\beta\gamma$ subunits downstream of activated GPCRs, with this being lost in p84, making p110 γ -p84 activation by G $\beta\gamma$ dependent on Ras-mediated membrane recruitment (*Rathinaswamy et al., 2023; Kurig et al., 2009; Rynkiewicz et al., 2020*). Activation of PI3K γ downstream of the IgE antigen receptor is driven by calcium-mediated activation of protein kinase C, leading to the selective phosphorylation and activation of p110 γ at S582 (*Walser et al., 2013*), with this putatively only occurring in p110 γ -p84 and not in p110 γ -p101. The full molecular mechanisms underlying how phosphorylation of p110 γ is selective for different p84 or p101 complexes, and how it activates lipid kinase activity are poorly understood.

Extensive biophysical approaches including cryo electron microscopy (cryo-EM), X-ray crystallography, and hydrogen-deuterium exchange mass spectrometry (HDX-MS) have provided extensive insight into the molecular underpinnings of how p110 γ associates with both p101 and p84, how they are differentially activated by Ras and GPCR signals, and how they can be activated on lipid membranes (*Pacold et al., 2000; Walker et al., 1999; Rathinaswamy et al., 2021a; Rathinaswamy et al., 2021c; Gangadhara et al., 2019; Vadas et al., 2013; Rathinaswamy et al., 2023*). The p110 γ catalytic subunit is composed of an adaptor binding domain (ABD), a Ras binding domain (RBD), a C2 domain, a helical domain, and a bi-lobal kinase domain (*Rathinaswamy et al., 2021a; Walker et al., 1999*). A set of helices positioned C-terminal to the activation loop in the kinase domain play a critical role in regulating activity, with this region referred to as the regulatory motif (*Rathinaswamy et al., 2021c*). The p110 γ isoform is unique in that it is inhibited in the absence of a regulatory subunit, with this driven by an autoinhibitory conformation of the regulatory motif, that is proposed to require membrane association to disrupt (*Gangadhara et al., 2019*). The regulatory motif is a common site of activating mutations in the other class I PI3K isoforms (*Jenkins et al., 2023*), with p110 γ having rare activating oncogenic mutants in this region (*Rathinaswamy et al., 2021c*). The p110 γ subunit interacts with either p84 or p101 at an interface composed of the C2 domain, and the linkers between the RBD-C2 and C2-helical domains (*Rathinaswamy et al., 2023*). The p110 γ -p84 complex forms a more dynamic complex compared to p110 γ -p101 (*Rathinaswamy et al., 2023; Shymanets et al., 2013*), however, no clear unique regulatory role of this difference in dynamics has been identified.

The fundamental roles of p110 γ in inflammatory processes have made it a therapeutic target in many pathological conditions, including asthma (*Campa et al., 2018*), arthritis (*Camps et al., 2005*), obesity (*Becattini et al., 2011; Breasson et al., 2017*), and cancer (*De Henau et al., 2016; Kaneda et al., 2016b; Kaneda et al., 2016a*). There are significant side effects from non-isoform selective PI3K γ inhibitors (*Bohnacker et al., 2017; Vanhaesebroeck et al., 2021*), which has driven the development of highly p110 γ selective small molecule inhibitors (*Bell et al., 2012; Evans et al., 2016; Gangadhara et al., 2019*). However, all p110 γ inhibitors will target both p110 γ -p101 and p110 γ -p84, so there is a potential for the development of allosteric inhibitors outside of the ATP binding site.

Initial promise has been reported for the development of class IA p110 α oncogene-specific allosteric inhibitors. However, further investigation of the molecular mechanisms underlying p110 γ regulation will be required for the discovery of regions that can be targeted for allosteric inhibitor development.

Here, we report critical roles of the helical domain of p110 γ in both activation and inhibition of lipid kinase activity. We characterized an allosteric inhibitory nanobody (NB7) that potently inhibits p110 γ activity. Cryo-EM was used to define the inhibitory interface, which is composed of the helical domain, the ABD-RBD linker, and the regulatory motif of the kinase domain of p110 γ . The region that the nanobody binds to is in close spatial proximity to a previously identified PKC β phosphorylation site (S582) in the helical domain, and oncogenic activating mutants in the regulatory motif. We fully characterized the activity and dynamics of stoichiometrically PKC β phosphorylated p110 γ , leading to the discovery of a novel additional phosphorylation site (either S594 or S595). PKC β phosphorylation was highly selective for p110 γ and p110 γ -p84, with limited phosphorylation of p110 γ -p101. HDX-MS analysis showed that phosphorylation of p110 γ leads to unfolding of the N-terminal region of the helical domain, and increased kinase activity. The presence of the inhibitory nanobody significantly blocks PKC β phosphorylation, while phosphorylation of p110 γ prevented binding to NB7. Overall, this work provides unique insight into the critical role of the helical domain in controlling p110 γ activity, and how phosphorylation and binding partners can modify this regulation. It also reveals a unique binding site located at the interface of the helical and kinase domain that can be targeted for future allosteric inhibitor design.

Results

Molecular mechanism of nanobody inhibition of p110 γ

We previously identified multiple nanobodies that inhibited the activity of p110 γ . One from this group (denoted NB7 throughout the manuscript) potently inhibited the membrane-mediated activation of p110 γ -p84 by both Ras and G $\beta\gamma$, with HDX-MS experiments mapping the NB7 binding interface to the RBD, helical and kinase domains (*Rathinaswamy et al., 2021b*). We originally hypothesized that NB7 worked by sterically inhibiting Ras binding to the RBD of p110 γ . To further explore the molecular mechanism of inhibition we purified all complexes of p110 γ (p110 γ apo, p110 γ -p84, p110 γ -p101) along with the NB7 nanobody. The SDS-PAGE of all proteins utilized in this study are shown in **Source data 1**.

To define the mechanism concerning how NB7 inhibits PI3K γ activity we analyzed how this nanobody inhibited all class IB PI3K γ complexes (p110 γ , p110 γ -p84, p110 γ -p101) upon activation by lipidated G $\beta\gamma$ subunits. Intriguingly, we found that all three forms of p110 γ were potently inhibited by NB7 (**Figure 1A**). While the IC₅₀ measured for the three complexes was different, this is likely mainly due to the dramatic difference in protein required to measure lipid kinase activity in vitro (~300 nM for p110 γ apo/p110 γ -p84, and ~10 nM for p110 γ -p101). This suggested that the mechanism of inhibition was not driven by a steric block of Ras association through the RBD, as previously proposed (*Rathinaswamy et al., 2021b*). We examined the binding of this nanobody to all complexes using biolayer interferometry (BLI). The nanobody bound equivalently tightly to all complexes, with ~2 nM potency for p110 γ , p110 γ -p84, and p110 γ -p101 (**Figure 1B**). We also tested binding of the nanobody to all class IA PI3Ks, and there was no detectable binding to p110 α , p110 β , and p110 δ (**Figure 1C**).

To further understand the mechanism by which this nanobody blocked lipid kinase activity we measured the bulk membrane recruitment dynamics of fluorescently labeled Dy647-p84-p110 γ on supported lipid bilayers (SLBs) using total internal reflection fluorescence (TIRF) microscopy. We found that the nanobody had no effect on membrane recruitment of p110 γ -p84 on bilayers containing membrane-tethered Ras (GTP) and G $\beta\gamma$ (**Figure 1D–F**). Membrane binding was not affected when the nanobody was spiked into samples containing membrane-associated Dy647-p84-p110 γ (**Figure 1D**). Similarly, pre-incubation of Dy647-p84-p110 γ with 500 nM NB7 did not perturb membrane association of the kinase when flowed over a supported membrane (**Figure 1E**).

We wanted to define the molecular mechanism of how nanobody NB7 was a potent allosteric inhibitor of lipid kinase activity. We purified the complex of nanobody NB7 bound to p110 γ -p84 to homogeneity by gel filtration. Using this sample, we obtained a cryo-EM reconstruction of the complex of nanobody (NB7)-bound p110 γ at 3.0 Å overall resolution from 149,603 particles (**Figure 2A–D**, **Figure 2—figure supplements 1 and 2** and **Supplementary file 1**). The density map was of sufficient




Figure 1. The inhibitory nanobody NB7 binds tightly to all p110γ complexes and inhibits kinase activity, but does not prevent membrane binding. **(A)** Cartoon schematic depicting nanobody inhibition of activation by lipidated Gβγ (1.5 μM final concentration) on 5% PIP₂ membrane (5% phosphatidylinositol 4,5-bisphosphate [PIP₂], 95% phosphatidylserine [PS]) activation. Lipid kinase assays showed a potent inhibition of lipid kinase activity with increasing concentrations of NB7 (3–3000 nM) for the different complexes. Experiments are carried out in triplicate (n=3) with each replicate shown. The y-axis shows lipid kinase activity normalized for each complex activated by Gβγ in the absence of nanobody. Concentrations of each protein were selected to give a lipid kinase value in the detectable range of the ATPase Transcreeper assay. The protein concentration of p110γ (300 nM), p110γ-p84 (330 nM), and p110γ-p101 (12 nM) was different due to intrinsic differences of each complex to be activated by lipidated Gβγ and is likely mainly dependent for the difference seen in NB7 response. **(B)** Association and dissociation curves for the dose response of His-NB7 binding to p110γ, p110γ-p84, and p110γ-p101 (50–1.9 nM) are shown. A cartoon schematic of biolayer interferometry (BLI) analysis of the binding of immobilized His-NB7 to p110γ is shown on the left. Dissociation constants (K_D) were calculated based on a global fit to a 1:1 model for the top three concentrations and averaged with error shown. Error was calculated from the association and dissociation value (n=3) with standard deviation shown. Full details are present in **Source data 1**. **(C)** Association and dissociation curves for His-NB7 binding to p110γ, p110α-p85, p110β-p85, and p110δ-p85. Experiments were performed in duplicate with a final concentration of 50 nM of each class I phosphoinositide 3 kinase (PI3K) complex. **(D)** Effect of NB7 on PI3Kγ recruitment to supported lipid bilayers containing H-Ras (GTP) and farnesyl-G as measured by total internal reflection fluorescence microscopy (TIRF-M). DY647-p84/p110γ displays rapid equilibration kinetics and is insensitive to the addition of 500 nM nanobody (black arrow, 250 s) on supported lipid bilayers containing H-Ras (GTP) and farnesyl-G. **(E)** Kinetics of 50 nM DY647-p84/p110γ membrane recruitment appears indistinguishable in the absence and presence of nanobody. Prior to sample injection, DY647-p84/p110γ was incubated for 10 min with 500 nM nanobody. **(F)** Representative TIRF-M images showing the localization of 50 nM DY647-p84/p110γ visualized in the absence or presence of 500 nM nanobody (+NB7). Membrane composition for panels C–E: 93% DOPC, 5% DOPS, 2% MCC-PE, Ras (GTP) covalently attached to MCC-PE, and 200 nM farnesyl-G.




Figure 2. Structure of p110 γ bound to inhibitory nanobody NB7. (A) Domain schematics of p110 γ with helical domain (blue), activation loop (orange), and regulatory motif (green) of p110 γ annotated. (B) Cryo electron microscopy (cryo-EM) density of the p110 γ -NB7 complex colored according to the schematic in (A). (C) Cartoon model of the structure of p110 γ bound to NB7 colored according to (A). (D) Schematic depicting the key features of p110 γ and the nanobody binding site, colored according to panel (A). (E) Domain schematic of NB7 complementarity determining regions (CDRs) and their sequences. (F) Zoom in on the binding interface of NB7, with the CDRs colored as in panel E, and the electron density of the CDRs contoured at 3 σ (blue mesh).

The online version of this article includes the following figure supplement(s) for figure 2:

Figure supplement 1. p110 γ -NB7 complex cryo electron microscopy (cryo-EM) analysis workflow.

Figure supplement 2. Density fit of p110 γ -NB7 complex.

Figure supplement 3. Comparison of full-length p110 γ bound to NB7 compared to p110 γ -p101.

quality to perform automated and manual construction of the p110 γ -NB7 complex, with unambiguous building of the interfacial contacts between NB7 and p110 γ (Figure 2—figure supplement 2). Nanobody binding did not induce any large-scale conformational changes of p110 γ , as the structure of p110 γ bound to NB7 was similar to the apo p110 γ crystal structure or p110 γ -p101 cryo-EM structure (Figure 2—figure supplement 3). The lowest local resolution was in the ABD, with increased B-factors of the ABD in the p110 γ -NB7 structure compared to p110 γ -p101 (Figure 2—figure supplement 3). This is consistent with the concept that ABD flexibility plays an important role in class I PI3K regulation (Liu et al., 2022).

The interface between NB7 and p110 γ was extensive, with ~1200 Å² of buried surface area, with interactions of the ABD-RBD linker, N-terminus of the helical domain, and the regulatory motif at the turn between k8 and k9 (1022-1026aa). This location in the regulatory motif is where both activating oncogenic (R1021C) and inhibitory loss of function mutants have been identified (R1021P) (*Takeda et al., 2019*), as well as a putative inhibitory phosphorylation site (T1024) (*Perino et al., 2011*). The resolution was sufficient to unambiguously build the three complementarity determining region (CDR) loops of NB7 that mediate target selectivity (*Figure 2E–F*). The interface is primarily hydrophobic, with only 8 hydrogen bonds, and 1 electrostatic interaction among the 33 interfacial residues of NB7. A pocket formed between the helical domain and the ABD-RBD linker forms the majority of the interface, with extensive interactions with the long CDR3. The CDR1 loop packed up against the N-terminal section of the helical domain, with the CDR2 loop forming the interface with the regulatory motif. Previous study of oncogenic mutants in the regulatory motif of p110 γ showed that increased dynamics mediated by these mutants increased kinase activity, putatively by breaking the autoinhibitory tryptophan lock in k12 of the regulatory motif (*Rathinaswamy et al., 2021c*). Therefore, rigidifying the regulatory motif likely explains the molecular basis for how it prevents kinase activity. The nanobody interface is distinct from the predicted G $\beta\gamma$ interface (*Rathinaswamy et al., 2023*) and the experimentally resolved Ras interface (*Pacold et al., 2000*), explaining why it can still be membrane recruited by these stimuli.

The interface of NB7 with p110 γ is distant from both the putative membrane binding surface and the catalytic machinery of the kinase domain. To further understand how this nanobody could potentially inhibit PI3K γ activity we examined any other potential modulators of PI3K γ activity localized in this region. There are two regulatory phosphorylation sites in the helical (*Walser et al., 2013*) and kinase domain (*Perino et al., 2011*) localized at the NB7 interface. This is intriguing as helical domain phosphorylation is activating, and kinase domain phosphorylation is inhibitory. This suggested that a critical role in the regulation of p110 γ is the dynamics of this kinase-helical interface. To fully define the role of NB7 in altering the dynamics of the helical domain we needed to study other modulators of helical domain dynamics.

p110 γ activation by helical domain phosphorylation

To further understand the potential role of helical domain dynamics in controlling p110 γ activity we examined the role of S582 helical domain phosphorylation by the protein kinase PKC β II (encoded by the gene *PRKCB2*, referred to as PKC β for simplicity throughout the manuscript) (*Walser et al., 2013*). S582 is located on the interior of the helical domain and would not be expected to be exposed when the N-terminal region of the helical domain is folded (*Figure 3A*). To understand this better at a molecular level, we purified a catalytic fragment of PKC β and performed protein phosphorylation reactions on p110 γ apo, p110 γ -p84, and p110 γ -p101. We identified a phosphorylated peptide containing S582, and surprisingly, we found an additional p110 γ phosphorylation site at either S594/S595 (*Figure 3B, Figure 3—figure supplement 1*). The modification at this site results in a single phosphorylation event, but due to CID MS/MS fragmentation we cannot determine which site is modified, and we will refer it as S594/S595 throughout the manuscript. The S594/S595 site is also located in the N-terminal region of the helical domain, and is even more buried than S582, and would not be expected to be exposed when this region is folded (*Figure 3A*). Dose-response curves of PKC β treatment were carried out for p110 γ (*Figure 3C*), p110 γ -p84 (*Figure 3D*), and p110 γ -p101 (*Figure 3E*). Both p110 γ and p110 γ -p84 showed similar dose-response curves for PKC β treatment, with similar curves for S582 and S594/S595. The p110 γ -p101 complex was only very weakly phosphorylated, with <100-fold lower levels compared to p110 γ and p110 γ -p84 (*Figure 3E*). This is consistent with the helical domain in p110 γ being more rigid when bound to p101, compared to either bound to p84 or p110 γ alone.

To provide additional insight into the molecular mechanisms underlying p110 γ phosphorylation we carried out HDX-MS experiments on p110 γ and phosphorylated p110 γ (90.8% phosphorylated S594/S595, 92% phosphorylated S582) (*Figure 4A*). The full data underlying the experiment is available in *Source data 1*, and data processing information is in *Supplementary file 2*. We have previously observed that the N-terminal region of the helical domain of apo p110 γ (residues spanning 557-630aa) shows isotope profiles that are consistent with EX1 hydrogen-deuterium (H/D) exchange kinetics (*Rathinaswamy et al., 2021b; Rathinaswamy et al., 2021c; Vadas et al., 2013; Walser et al., 2013*).




Figure 3. PKCβ leads to dual phosphorylation of internal sites in the helical domain, with selectivity for apo p110γ and p110γ-p84 over p110γ-p101. (A) Putative phosphorylation sites mapped on the structure of p110γ (PDB: 7MEZ) and cartoon schematic. The regions are colored based on domain schematics featured in Figure 2A. (B) Raw MS spectra of the unphosphorylated and phosphorylated peptide for a region spanning 579–592 (RYESLKHPKAYPKL) and 593–607 (FSSVKWGQEQEIVAKT). The putative phosphorylation sites in the sequence are shown in red, with the m/z theoretical (m/z t) and m/z experimental (m/z e) shown below each sequence. (C–E) Extracted traces and ratios of the intensity of extracted ion traces of different phosphorylation site peptides (top to bottom: S594/595 and S582) from (C) p110γ, (D) p110γ/p84, or (E) p110γ/p101 samples treated with increasing concentration of PKCβ according to the legend. The black traces in the ratio graphs are the intensity of the non-phosphorylated peptide, and the red traces in the ratio graphs are the intensity of the phosphorylated peptide.

The online version of this article includes the following figure supplement(s) for figure 3:

Figure supplement 1. MS/MS spectra of peptides spanning S582 and S594/S595 for both phosphorylated and unphosphorylated states.

This is indicative of cooperative unfolding of extended protein regions, with H/D exchange occurring faster than the refolding event. This region is where the PKCβ phosphorylation sites are located and may explain how the buried residues S582 and S594/S595 can be exposed to PKCβ. This is compatible with the observation that p110γ-p101 is protected from phosphorylation, as it does not show EX1 kinetics in this region, whereas both p110γ and p110γ-p84 do (Rathinaswamy et al., 2021a).




Figure 4. Activating phosphorylation at the helical domain leads to opening of the regulatory motif. **(A)** Hydrogen-deuterium exchange mass spectrometry (HDX-MS) comparing apo and phosphorylated p110γ. Significant differences in deuterium exchange are mapped on to the structure and cartoon of p110γ according to the legend (PDB: 7MEZ). **(B)** The graph of the #D difference in deuterium incorporation for p110γ, with each point representing a single peptide. Peptides colored in red are those that had a significant change in the mutants (greater than 0.4 Da and 5% difference at any time point, with a two tailed t-test $p < 0.01$). Error bars are SD ($n = 3$). **(C)** Representative bimodal distribution (EX1 kinetics) observed in the helical domain peptides of p110γ. **(D)** Representative p110γ peptides displaying increases in exchange in the phosphorylated state are shown. For all panels, error bars show SD ($n = 3$). **(E)** Measurement of ATP to ADP conversion of phosphorylated and non-phosphorylated p110γ (1000 nM final concentration) ATPase activity in the absence (left) and presence of PIP₂ membranes (5% phosphatidylinositol 4,5-bisphosphate [PIP₂], 95% phosphatidylserine [PS]) activation (right). Significance is indicated by **(<0.001%) and ***(<0.0001%).

When we compared phosphorylated p110γ (>90.8% as measured by mass spectrometry at both sites) to unphosphorylated p110γ we observed extensive increases in dynamics in the C2, helical domain, and kinase domain (**Figure 4A–D**). The largest increases in exchange upon phosphorylation were located in the N-terminal region of the helical domain, with the peptides directly adjacent to the phosphorylation site showing almost complete deuterium incorporation at the earliest time points of exchange. This is indicative of significant disruption of the alpha helical secondary structure in this region. When we examined the exchange profiles in this region, they still underwent EX1 kinetics (**Figure 4C**), however, phosphorylated p110γ was enriched in the more fully deuterated species. In addition to the regions in the helical domain, a portion of the regulatory motif of the kinase domain




Figure 5. Nanobody NB7 blocks PKC β phosphorylation, and phosphorylation prevents nanobody binding. **(A)** Extracted ion chromatograms for p110 γ , p110 γ -p101, and p110 γ bound to NB7 are shown for the S594 or S595 phosphorylation site in p110 γ . A bar graph showing the intensities of phosphorylated and non-phosphorylated p110 γ peptide (593-607) for p110 γ (black), p110 γ with NB7 (red), and p110 γ -p101 (purple) are shown to the right of the extracted ion chromatograms ($n=3$, right). In all experiments in panels A+B, PKC β was present at 500 nM. Significance is indicated by ***(<0.0001%). **(B)** Extracted ion chromatograms for p110 γ , p110 γ -p101, and p110 γ bound to NB7 are shown for the S582 phosphorylation site in p110 γ . A bar graph showing the intensities of phosphorylated and non-phosphorylated p110 γ peptide (579-592) p110 γ (black), p110 γ with NB7 (red), and p110 γ -p101 (purple) are shown to the right of the extracted ion chromatograms ($n=3$, right). Significance is indicated by *(<0.01%) and ***(<0.0001%). The putative phosphorylation site is shown in red in the sequence above the bar graphs for both panels A+B. **(C)** Cartoon schematic of biolayer interferometry (BLI) analysis of the binding of immobilized His-NB7 to phosphorylated and non-phosphorylated p110 γ . **(D)** Association curves for phosphorylated and non-phosphorylated p110 γ (25 nM) binding to His-NB7 are shown ($n=3$). **(E)** ATPase kinase activity assays comparing the activation/inhibition of phosphorylated and non-phosphorylated p110 γ (1000 nM) with or without nanobody (3000 nM final) in the absence of PIP $_2$ membranes. Significance is indicated by *(<0.05%) and NS (>0.05%).

also showed increased deuterium exposure. This included the k9-k12 helices that surround the activation loop of p110 γ . These increases in exchange were similar to those we had observed in a R1021C oncogenic activating mutant of *PIK3CG* (Rathinaswamy et al., 2021c).

To further explore the potential role of phosphorylation in mediating p110 γ activity, we examined the kinase activity of p110 γ under two conditions: basal ATP turnover, and with PIP $_2$ containing lipid membranes. The experiments in the absence of PIP $_2$ measure turnover of ATP into ADP and phosphate and is a readout of basal catalytic competency. Experiments with PIP $_2$ measured ATP consumed in the generation of PIP $_3$, as well as in non-productive ATP turnover. The p110 γ enzyme in the absence of stimulators is very weakly active toward PIP $_2$ substrate with only ~2-fold increased ATP turnover compared to in the absence of membranes. This is consistent with very weak membrane recruitment of p110 γ complexes in the absence of lipid activators (Rathinaswamy et al., 2023). PKC β -mediated phosphorylation enhanced the ATPase activity of p110 γ ~2-fold in both the absence and presence of membranes (Figure 4E). This suggests that the effect of phosphorylation is to change the intrinsic catalytic efficiency of phosphorylated p110 γ , with limited effect on membrane binding.

Nanobody decreases p110 γ phosphorylation

As NB7 bound at the interface of the helical and kinase domains that is exposed upon PKC β phosphorylation of p110 γ , we hypothesized that the nanobody would likely alter phosphorylation. We carried out PKC β phosphorylation of p110 γ , p110 γ -p101, and p110 γ bound to NB7. The presence of NB7 showed a significant decrease in p110 γ phosphorylation at both sites (Figure 5A–B). We also wanted to determine whether p110 γ phosphorylation reciprocally perturbed NB7 binding. BLI experiments

showed that there was no detectable binding of NB7 to phosphorylated p110 γ (**Figure 5C–D**), consistent with phosphorylation disrupting the N-terminal region of the p110 γ helical domain. In addition, lipid kinase assays using phosphorylated p110 γ showed no detectable difference in activity when measured in the absence and presence of nanobody (**Figure 5E**), consistent with NB7 being unable to bind to phosphorylated p110 γ .

Discussion

Here, we find that the helical domain is a central regulator of the p110 γ catalytic subunit of class IB PI3K γ , with modulation of helical dynamics through binding partners or PTMs able to either increase or decrease lipid kinase activity. These results expand on previous work defining the helical domain as a central regulator of class IA PI3Ks, where the nSH2 domain of the p85 regulatory subunits makes inhibitory interactions that significantly inhibit lipid kinase activity of all class IA catalytic subunits (p110 α , p110 β , and p110 δ) (**Mandelker et al., 2009**; **Miled et al., 2007**; **Burke and Williams, 2013**; **Burke, 2018**). This inhibitory interaction in class IA PI3Ks is disrupted in human cancers (helical hotspot




Figure 6. Comparison of nanobody binding site compared to p85 inhibition of class IA phosphoinositide 3 kinases (PI3Ks) and class IB activation sites. **(A)** Comparison of the nanobody NB7 binding site in p110 γ compared to the nSH2 inhibitory site in p110 α (PDB: 3HHM) (**Mandelker et al., 2009**). **(B)** Comparison of the nanobody NB7 binding site in p110 γ compared to the X-ray structure of the Ras binding site (PDB: 1HE8) (**Pacold et al., 2000**) and the AlphaFold model of G $\beta\gamma$ bound to p110 γ (**Rathinaswamy et al., 2023**). **(C)** Oncogenic mutations and post-translational modifications in spatial proximity to the nanobody binding site.

mutations in *PIK3CA* (Samuels et al., 2004) and immune disorders (helical mutations in *PIK3CD* in APDS1) (Angulo et al., 2013; Lucas et al., 2014). Class IB PI3Ks are uniquely compared to class IA PI3Ks, as they are not inhibited by p101 and p84 regulatory subunits, but instead potentiate GPCR activation. This lack of inhibition is due to the distinct binding interface of class IB PI3K regulatory subunits compared to class IA regulatory subunits, with only class IA regulatory subunits making direct inhibitory interactions with the kinase and helical domains of p110 catalytic subunits (Rathinaswamy et al., 2021a). Here, we show that a unique surface at the interface of the helical and kinase domains of p110 γ is a potential site for the development of novel allosteric inhibitors that modulate p110 γ activity.

The previously identified inhibitory nanobody (NB7) (Rathinaswamy et al., 2021b) bound with high affinity and inhibited all complexes of p110 γ . The nanobody interface is distinct from how the nSH2 inhibits class IA PI3K activity, as its binding site is on the opposite face of the helical domain (Figure 6A). The mechanism of inhibition is also distinct, as the nSH2-helical interaction plays a critical role in preventing membrane recruitment of inhibited class I PI3Ks, with removal of this interface either through pYXXM motif binding or oncogenic mutations leading to increased membrane recruitment (Burke et al., 2012; Burke et al., 2011; Zhang et al., 2011). Analysis of the nanobody binding site compared to the structure of HRas-p110 γ or the HDX-MS supported AlphaFold-multimer prediction of G β γ -p110 γ (Rathinaswamy et al., 2023) shows that nanobody binding does not sterically block complex formation (Figure 6B). This is consistent with it not blocking membrane recruitment by Ras/G β γ . The nanobody inhibited ATP turnover both in solution and on membranes, suggesting that it prevents formation of a catalytically competent conformation of p110 γ , but still allows for membrane recruitment. Further development of small molecule allosteric binding partners in this allosteric pocket between the kinase and helical domain may reveal the specific molecular interactions in this pocket that mediate inhibition.

Oncogenic mutations are frequent in the class IA PI3K encoded by *PIK3CA*, with this being the second most frequently mutated gene in human cancer (Lawrence et al., 2014). Mutations in p110 γ encoded by *PIK3CG* in cancer are less frequent, however, they can still provide insight into regulatory mechanisms that control activity. Oncogenic mutations in the kinase domain (R1021C) and helical domain (E581K) are in close proximity to the nanobody binding site, and both would be expected to disrupt the stability of the helical domain or regulatory motif of the kinase domain (Figure 6C). In addition to these mutations there are also multiple post-translational modifications that occur in this region, including inhibitory phosphorylation at T1024 (Perino et al., 2011) and activating phosphorylation at S582 (Walser et al., 2013). PKC β is activated downstream of the IgE receptor in mast cells (Walser et al., 2013), but the full details of how this activates PI3K γ have been unclear. We identified an additional PKC β phosphorylation site located in the helical domain (S594/S595) (Figure 6C). Both the S582 and S594/S595 sites are not surface accessible and would require a transient opening of the helical domain for kinase accessibility. HDX-MS analysis of the helical domain of p110 γ has shown that it is more dynamic than other class I PI3K isoforms (Burke and Williams, 2013; Walser et al., 2013), with the presence of the p101 regulatory subunit dramatically decreasing helical domain dynamics (Vadas et al., 2013). This putative mechanism of helical domain dynamics driving PKC β phosphorylation is consistent with our observation that p101 subunits decreased p110 γ phosphorylation >100-fold. PKC β phosphorylation of p110 γ leads to increased dynamics in both the helical and kinase domains with increased kinase activity, although only weakly compared to full activation by either membrane localized Ras or G β γ . This increase was observed with both membrane and soluble substrate, so likely is not driven by altered membrane recruitment.

Overall, our biophysical and biochemical analysis of modulators of helical domain dynamics reveal the critical role of this domain in regulating class IB PI3K γ activity. This raises possibilities for development of small molecule modulators that may either increase or decrease helical domain dynamics, leading to either activation or inhibition. The high-resolution structure of an allosteric inhibitor nanobody provides initial insight into which pockets can specifically be targeted. Multiple ATP competitive p110 γ selective inhibitors are in clinical trials for human cancers (Li et al., 2021), with many having significant side effects. The identification of novel inhibitory strategies provides new opportunities for targeting p110 γ dysregulation in human disease.

Methods

Plasmid generation

Plasmids encoding *Homo sapiens* p110 γ (human), *Mus musculus* p84 (mouse), *Sus scrofa* p101 (porcine), and G $\beta\gamma$ were used as previously described (Rathinaswamy et al., 2023). The plasmids encoding the class IA PI3Ks were also used as previously described (Dornan et al., 2017; Siempelkamp et al., 2017). The pDONR223-PRKCB2 (PKC β II, uniprot identifier: P05771-2) was a gift from William Hahn & David Root (Addgene plasmid #23746; <http://n2t.net/addgene:23746>; RRID:Addgene_23746) (Johannessen et al., 2010). The PKC β II construct contains an internal TEV site that cleaves the catalytic domains from the C1/C2 regulatory domains (TEV site inserted between residues 320 and 321 of PKC β). This construct was subcloned into a pACEBAC Sf9 expression vector for Sf9 protein production. All constructs were cloned to include a 10 $\times$ histidine tag, a 2 $\times$ strep tag, and a tobacco etch virus protease cleavage site on the N-terminus. For p110 γ and PKC β II this tag was included at the N-terminus, with this tag included at the N-terminus of either p84 or p101 for purification of p110 γ -p101 and p110 γ -p84. Full details of the plasmids are included in Appendix 1—key resources table.

Virus generation and amplification

The plasmids encoding genes for insect cell expression were transformed into DH10MultiBac cells (MultiBac, Geneva Biotech) to generate bacmid containing the genes of interest. Successful generation was identified by blue-white colony screening and the bacmid was purified using a standard isopropanol-ethanol extraction method. Bacteria were grown overnight (16 hr) in 3–5 mL 2 $\times$ YT (BioBasic #SD7019). Cells were spun down and the pellet was resuspended in 300 μ L of 50 mM Tris-HCl, pH 8.0, 10 mM EDTA, 100 mg/mL RNase A. The pellet was lysed by the addition of 300 μ L of 1% sodium dodecyl sulfate (SDS) (wt/vol), 200 mM NaOH, and the reaction was neutralized by addition of 400 μ L of 3.0 M potassium acetate, pH 5.5. Following centrifugation at 21130 RCF and 4°C (Rotor #5424R), the supernatant was mixed with 800 μ L isopropanol to precipitate bacmid DNA. Following centrifugation, the pelleted bacmid DNA was washed with 500 μ L 70% ethanol three times. The pellet was then air dried for 1 min and re-suspended in 50 μ L Buffer EB (10 mM Tris-Cl, pH 8.5; all buffers from QIAprep Spin Miniprep Kit, QIAGEN #27104). Purified bacmid was then transfected into Sf9 cells. Two mL of Sf9 cells at 0.6×10^6 cells/mL were aliquoted into a six-well plate and allowed to attach to form a confluent layer. Transfection reactions were prepared mixing 8–12 μ g of bacmid DNA in 100 μ L 1 $\times$ PBS and 12 μ g polyethyleneimine (Polyethyleneimine 'Max' MW 40,000, Polysciences #24765, USA) in 100 μ L 1 $\times$ PBS and the reaction was allowed to proceed for 20–30 min before addition to an Sf9 monolayer containing well. Transfections were allowed to proceed for 5–6 days before harvesting virus containing supernatant as a P1 viral stock.

Viral stocks were further amplified by adding P1 to Sf9 cells at $\sim 2 \times 10^6$ cells/mL (2/100 volume ratio). This amplification was allowed to proceed for 4–5 days and resulted in a P2 stage viral stock that was used in final protein expression. Harvesting of P2 viral stocks was carried out by centrifuging cell suspensions in 50 mL Falcon tubes at 2281 RCF (Beckman GS-15). To the supernatant containing virus, 5–10% inactivated fetal bovine serum (FBS; VWR Canada #97068-085) was added and the stock was stored at 4°C.

Expression and purification of PI3K γ , PI3K $\alpha/\beta/\delta$, and PKC β constructs

PI3K γ and PKC β constructs were expressed in Sf9 insect cells using the baculovirus expression system. Following 55 hr after infection with P2 viral stocks, cells were harvested by centrifuging at 1680 RCF (Eppendorf Centrifuge 5810R) and the pellets were snap-frozen in liquid nitrogen. The complex was purified through a combination of nickel affinity, streptavidin affinity, and size-exclusion chromatography.

Frozen insect cell pellets were resuspended in lysis buffer (20 mM Tris pH 8.0, 100 mM NaCl, 10 mM imidazole pH 8.0, 5% glycerol [vol/vol], 2 mM ME), protease inhibitor (Protease Inhibitor Cocktail Set III, Sigma) and sonicated for 2 min (15 s on, 15 s off, level 4.0, Misonix sonicator 3000). Triton X was added to the lysate to a final concentration of 0.1% and clarified by spinning at 15,000 RCF at 4°C for 45 min (Beckman Coulter JA-20 rotor). The supernatant was loaded onto a 5 mL HisTrap FF crude column (GE Healthcare) equilibrated in NiNTA A buffer (20 mM Tris pH 8.0, 100 mM NaCl, 20 mM imidazole pH 8.0, 5% [vol/vol] glycerol, 2 mM ME). The column was washed with high salt NiNTA A

buffer (20 mM Tris pH 8.0, 1 M NaCl, 20 mM imidazole pH 8.0, 5% [vol/vol] glycerol, 2 mM ME), NiNTA A buffer, 6% NiNTA B buffer (20 mM Tris pH 8.0, 100 mM NaCl, 250 mM imidazole pH 8.0, 5% [vol/vol] glycerol, 2 mM ME) and the protein was eluted with 100% NiNTA B. The eluent was loaded onto a 5 mL StrepTrap HP column (GE Healthcare) equilibrated in gel filtration buffer (20 mM Tris pH 8.5, 100 mM NaCl, 50 mM ammonium sulfate and 0.5 mM tris(2-carboxyethyl) phosphine [TCEP]). To purify *PI3K $\alpha/\beta/\delta$* , the purification protocol was performed as described above but instead the protein was eluted in *PI3K α* gel filtration buffer (20 mM HEPES 7.5, 150 mM NaCl, 0.5 mM TCEP). The column was washed with the corresponding gel filtration buffer and loaded with tobacco etch virus protease. After cleavage on the column overnight, the PI3K protein constructs were eluted in gel filtration buffer. The protein was concentrated in a 50,000 MWCO Amicon Concentrator (Millipore) to <1 mL and injected onto a Superdex 200 10/300 GL Increase size-exclusion column (GE Healthcare) equilibrated in gel filtration buffer. After size exclusion, the protein was concentrated, aliquoted, frozen, and stored at -80°C . For PKC β , the protein was eluted from the strep column in gel filtration buffer, and the eluate was then loaded on a 1 mL HisTrap FF column to remove his tagged LipTev. The flowthrough was collected, and the column was washed with 2 mL of gel filtration buffer. These fractions were pooled and concentrated and stored at -80°C .

To purify phosphorylated p110 γ , the purification protocol as described above was performed but PKC β was added to the strep column at a molar ratio of 1:3 (PKC β :p110 γ) along with LipTEV, 20 mM MgCl_2 , and 1 mM ATP and allowed to incubate on ice for 4 hr. The protein was eluted by adding 7 mL of gel filtration buffer and treated with a second dose of PKC β (same ratio as above) and allowed to incubate on ice for another 3 hr. For non-phosphorylated p110 γ , same protocol was followed with the exception in the addition of PKC β . Both the proteins were concentrated in a 50,000 MWCO Amicon Concentrator (Millipore) to <1 mL and injected onto a Superdex 200 10/300 GL Increase size-exclusion column (GE Healthcare) equilibrated in gel filtration buffer. The final phosphorylation level of the two sites was characterized by mass spectrometry, with these values being 92% and 90.8%, for S582 and S594/S595, respectively. After size exclusion, the protein was concentrated, aliquoted, frozen, and stored at -80°C .

Expression and purification of lipidated G for kinase activity assays

Full-length, lipidated human *G $\beta\gamma$* (*G β 1 γ 2*) was expressed in Sf9 insect cells and purified as described previously. After 65 hr of expression, cells were harvested, and the pellets were frozen as described above. Pellets were resuspended in lysis buffer (20 mM HEPES pH 7.7, 100 mM NaCl, 10 mM β ME, protease inhibitor [Protease Inhibitor Cocktail Set III, Sigma]) and sonicated for 2 min (15 s on, 15 s off, level 4.0, Misonix sonicator 3000). The lysate was spun at 500 RCF (Eppendorf Centrifuge 5810R) to remove intact cells and the supernatant was centrifuged again at 25,000 RCF for 1 hr (Beckman Coulter JA-20 rotor). The pellet was resuspended in lysis buffer and sodium cholate was added to a final concentration of 1% and stirred at 4°C for 1 hr. The membrane extract was clarified by spinning at 10,000 RCF for 30 min (Beckman Coulter JA-20 rotor). The supernatant was diluted three times with NiNTA A buffer (20 mM HEPES pH 7.7, 100 mM NaCl, 10 mM imidazole, 0.1% C12E10, 10 mM ME) and loaded onto a 5 mL HisTrap FF crude column (GE Healthcare) equilibrated in the same buffer. The column was washed with NiNTA A, 6% NiNTA B buffer (20 mM HEPES pH 7.7, 25 mM NaCl, 250 mM imidazole pH 8.0, 0.1% C12E10, 10 mM ME) and the protein was eluted with 100% NiNTA B. The eluent was loaded onto HiTrap Q HP anion exchange column equilibrated in Hep A buffer (20 mM Tris pH 8.0, 8 mM CHAPS, 2 mM dithiothreitol [DTT]). A gradient was started with Hep B buffer (20 mM Tris pH 8.0, 500 mM NaCl, 8 mM CHAPS, 2 mM DTT) and the protein was eluted in ~50% Hep B buffer. The eluent was concentrated in a 30,000 MWCO Amicon Concentrator (Millipore) to <1 mL and injected onto a Superdex 75 10/300 GL size-exclusion column (GE Healthcare) equilibrated in Gel Filtration buffer (20 mM HEPES pH 7.7, 100 mM NaCl, 10 mM CHAPS, 2 mM TCEP). Fractions containing protein were pooled, concentrated, aliquoted, frozen, and stored at -80°C .

Expression and purification of nanobody

Nanobody NB7-PIK3CG with a C-terminal 6 $\times$ His tag was expressed from a pMESy4 vector in the periplasm of WK6 *E. coli*. A 1 L culture was grown to OD600 of 0.7 in Terrific Broth containing 0.1% glucose and 2 mM MgCl_2 in the presence of 100 $\mu\text{g/mL}$ ampicillin and was induced with 0.5 mM isopropyl- β -D-thiogalactoside (IPTG). Cells were harvested the following day by centrifuging at 2500

RCF (Eppendorf Centrifuge 5810R) and the pellet was snap-frozen in liquid nitrogen. The frozen pellet was resuspended in 15 mL of TES buffer containing 200 mM Tris pH 8.0, 0.5 mM ethylenediaminetetraacetic acid (EDTA), and 500 mM sucrose and was mixed for 45 min at 4°C. To this mixture, 30 mL of TES buffer diluted four times in water was added and mixed for 45 min at 4°C to induce osmotic shock. The lysate was clarified by centrifuging at 14,000 rpm for 15 min (Beckman Coulter JA-20 rotor). Imidazole was added to the supernatant to final concentration of 10 mM loaded onto a 5 mL HisTrap FF crude column (GE Healthcare) equilibrated in NiNTA A buffer (20 mM Tris pH 8.0, 100 mM NaCl, 20 mM imidazole pH 8.0, 5% [vol/vol] glycerol, 2 mM β -mercaptoethanol [ME]). The column was washed with high salt NiNTA A buffer (20 mM Tris pH 8.0, 1 M NaCl, 20 mM imidazole pH 8.0, 5% [vol/vol] glycerol, 2 mM ME), followed by 100% NiNTA A buffer, then a 6% NiNTA B wash buffer (20 mM Tris pH 8.0, 100 mM NaCl, 250 mM imidazole pH 8.0, 5% [vol/vol] glycerol, 2 mM ME) and the protein was eluted with 100% NiNTA B. The eluent was concentrated in a 10,000 MWCO Amicon Concentrator (Millipore) to <1 mL and injected onto a Superdex 75 10/300 GL Increase size-exclusion column (GE Healthcare) equilibrated in gel filtration buffer (20 mM Tris pH 8.5, 100 mM NaCl, 50 mM ammonium sulfate, and 0.5 mM TCEP). Following size exclusion, the protein was concentrated, frozen, and stored at -80°C .

Lipid vesicle preparation for kinase activity assays

Lipid vesicles containing 5% brain phosphatidylinositol 4,5-bisphosphate (PIP₂) and 95% brain phosphatidylserine (PS) were prepared by mixing the lipids solutions in organic solvent. The solvent was evaporated in a stream of argon following which the lipid film was desiccated in a vacuum for 45 min. The lipids were resuspended in lipid buffer (20 mM HEPES pH 7.0, 100 mM NaCl, and 10% glycerol) and the solution was vortexed for 5 min followed by sonication for 15 min. The vesicles were then subjected to 10 freeze thaw cycles and extruded 11 times through a 100 nm filter (T&T Scientific: TT-002-0010). The extruded vesicles were sub-aliquoted and stored at -80°C . Final vesicle concentration was 2 mg/mL.

Kinase assays

All kinase assays were done using Transcreener ADP2 Fluorescence Intensity (FI) assays (Bellbrook labs) which measures ADP production. All assays contained ATP at a final concentration of 100 μM , and those with membranes used vesicles containing 5% PIP₂ and 95% PS at a final concentration of 0.5 mg/mL.

For assays measuring the inhibition by nanobody, 4 $\times$ kinase (final concentration: 330 nM for p110 γ , 300 nM for p110 γ /p84, and 12 nM for p110 γ /p101) was mixed with varying 4 $\times$ concentrations of nanobody (final concentration: 2 μM to 2.7 nM) or kinase buffer (20 mM HEPES pH 7.5, 100 mM NaCl, 3 mM MgCl₂, 0.03% CHAPS, 2 mM TCEP, and 1 mM EGTA) and allowed to sit on ice for 15 min. Two μL of protein mix was mixed with 2 μL of lipid solution containing G $\beta\gamma$ (1 μM final concentration), ATP (100 μM), PIP₂ lipid vesicles (0.5 mg/ml final concentration), and lipid buffer (25 mM HEPES pH 7, 5% glycerol, and 100 mM NaCl) and incubated at 20°C for 60 min.

For assays comparing the difference in activation between phosphorylated and non-phosphorylated p110 γ , 2 $\times$ kinase (final concentrations: 1 μM) was mixed with 2 $\times$ lipid solutions containing ATP (100 μM), and lipid buffer and either nanobody (3 μM final concentration), PIP₂ lipid vesicles (0.5 mg/mL final concentration), or both nanobody and lipid. The reaction was incubated at 20°C for 60 min.

After the 60 min incubation, all reactions were stopped with 4 μL of 2 $\times$ stop and detect solution containing Stop and Detect buffer (20 mM HEPES, 0.02% Brij-35, 40 mM EDTA pH 7.5), 8 nM ADP Alexa Fluor 594 Tracer and 93.7 $\mu\text{g/mL}$ ADP2 Antibody IRDye QC-1, covered and incubated at 20°C for 1 hr before reading the fluorescence. The fluorescence intensity was measured using a SpectraMax M5 plate reader at excitation 590 nm and emission 620 nm. All data was normalized against the appropriate measurements obtained for 100 μM ATP and 100 μM ADP with no kinase. The percent ATP turnover was interpolated using a standard curve (0.1–100 μM ADP). Interpolated values were then used to calculate the specific activity of the enzyme.

Biolayer interferometry

All BLI experiments were performed using the Octet K2 instrument (Fortebio Inc). For all experiments His-tagged nanobody (500 nM) was immobilized on an Anti-Penta-His biosensor for 600 s, and

the sensor was dipped into varying concentrations of the protein complex being measured. A dose response was carried out for p110 γ , p110 γ -p84, and p110 γ -p101 (50 nM – 1.9 nM), with association occurring for 600 s, followed by a 1200s dissociation in Octet Buffer (20 mM Tris pH 8.5, 100 mM NaCl, 50 mM ammonium sulfate, 0.1% bovine serum albumin, and 0.02% Tween 20). Experiments comparing class IA PI3K versus class IB PI3K γ used 50 nM of each class IA PI3K.

When comparing nanobody binding to phosphorylated and unphosphorylated p110 γ , we used a final concentration of 25 nM for both phosphorylated and non-phosphorylated p110 γ with association occurring for 600 s, followed by a 600 s dissociation. The K_D (dissociation constant) for the different p110 γ complexes was calculated from the binding curves based on their global fit to a 1:1 binding model using ForteBio data analysis 12.0 (Fortebio Inc).

SLB TIRF microscopy experiments

The membrane binding dynamics of Dy647-p84-p110 γ were measured in the absence and presence of nanobody 7 (NB7) using TIRF microscopy. As previously described (*Rathinaswamy et al., 2023*), SLBs were formed using 50 nm extruded small unilamellar vesicles containing the following lipids: 1,2-dioleoyl-*sn*-glycero-3-phosphocholine (18:1 DOPC, Avanti #850375C), 1,2-dioleoyl-*sn*-glycero-3-phospho-L-serine (18:1 DOPS, Avanti #840035C), 1,2-dioleoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[4-(p-maleimidomethyl)cyclohexane-carboxamide] (18:1 MCC-PE, Avanti #780201C). Lipid compositions reported in figure legends represent the molar percentage of each lipid species.

To create SLBs, a total concentration of 0.25 mM lipids was solvated in 1 $\times$ PBS (pH 7.4) and deposited on Piranha etched glass coverslips (25 $\times$ 75 mm) adhered to an IBIDI chamber. After a 30 min incubation, membranes were washed with 4 mL of 1 $\times$ PBS (pH 7.4) and then blocked for 10 min with 1 mg/mL beta casein (Thermo Fisher Science, Cat#37528) in 1 $\times$ PBS (pH 7.4) (Corning, Cat#46-013CM). To conjugate H-Ras to maleimide lipids (MCC-PE), blocked membranes were incubated with 30 μ M H-Ras (GDP) in buffer containing 1 $\times$ PBS (pH 7.4), 1 mM MgCl₂, 50 μ M GDP, and 0.1 mM TCEP for 2 hr. The membrane conjugation reaction was terminated after 2 hr with 1 $\times$ PBS (pH 7.4) containing 5 mM β -ME. Membranes were then washed and stored in 1 $\times$ PBS (pH 7.4) until performing the TIRF microscopy (TIRF-M) membrane binding experiments. H-Ras was purified as previously described (*Rathinaswamy et al., 2023*).

To perform the TIRF-M membrane binding assays, 200 nM farnesyl-*G* β γ was equilibrated into the supported membranes for 30 min. In parallel, nucleotide exchange of H-Ras (GDP) was performed by adding 50 nM guanine nucleotide exchange factor (SosCat) in 1 $\times$ PBS (pH 7.4), 1 mM MgCl₂, 50 μ M GDP. To measure membrane binding, Dy647-p84-p110 γ was diluted into the following buffer: 20 mM HEPES (pH 7.0), 150 mM NaCl, 50 μ M GTP, 1 mM ATP, 5 mM MgCl₂, 0.5 mM EGTA, 20 mM glucose, 200 μ g/mL beta casein (Thermo Scientific, Cat#37528), 20 mM BME, 320 μ g/mL glucose oxidase (Serva, #22780.01 *Aspergillus niger*), 50 μ g/mL catalase (Sigma, #C40-100MG Bovine Liver), and 2 mM Trolox. Trolox was prepared as previously described (*Hansen et al., 2019*). Perishable reagents (i.e., glucose oxidase, catalase, and Trolox) were added 10 min before image acquisition.

TIRF-M experiments were performed using an inverted Nikon Ti2 microscope with a 100 $\times$ Nikon (1.49 NA) oil immersion objective. The x-axis and y-axis positions were controlled using a Nikon motorized stage. Dy647-p84-p110 γ was excited with a 637 nm diode laser (OBIS laser diode, Coherent Inc Santa Clara, CA, USA) controlled with an acousto-optic tunable filter and laser launch built by Vortran (Sacramento, CA, USA). The power output measured through the objective for single particle imaging was 1–3 mW. Excitation light passing through quad multi-pass dichroic filter cube (Semrock). Fluorescence emission passed through Nikon emission filter wheel containing the following 25 mm ET700/75 M emission filters (Semrock) before being detected on iXion Life 897 EMCCD camera (Andor Technology Ltd., UK). All TIRF-M experiments were performed at room temperature (23°C). Microscope hardware was controlled using Nikon NIS elements. Data analysis was performed using ImageJ/Fiji and Prism graphing program.

Cryo-EM sample preparation and data collection

Three μ L of purified nanobody-bound p110 γ at 0.45 mg/mL was adsorbed onto C-Flat 2/2-T grids that were glow discharged for 25 s at 15 mA. Grids were then plunged into liquid ethane using a Vitrobot Mark IV (Thermo Fisher Scientific) with the following settings: –5 blot force, 1.5 s blot time, 100% humidity, and 4°C. Vitrified specimens were screened for ice and particle quality at the UBC

high-resolution macromolecular electron microscopy (HRMEM) facility using a 200 kV Glacios transmission electron microscope equipped with a Falcon 3EC direct electron detector (DED). Clipped grids were sent to the Pacific Northwest Cryo-EM Center (PNCC) where 7322 movies were collected using a Titan Krios equipped with a Gatan K3 DED and a BioQuantum K3 energy filter with a slit width of 20 eV. The movies were collected at a physical pixel size of 0.830 Å/pix and a total dose of 50 e⁻/Å² over 50 frames.

Cryo-EM image analysis

The data were processed using cryoSPARC v.3.3.2 (Punjani *et al.*, 2017). The movies were pre-processed by patch motion correction using default settings except Fourier cropping by a factor of 2, followed by patch CTF estimation using default settings. A 3D map of PI3K p110γ-p101 complex (EMD-23808) was used to create 2D projections for use as templates to auto-pick 1,463,553 particles. Particles were extracted with a box size of 380 pixels, Fourier cropped to a box size of 96 pixels and subjected to 2D classification. After discarding classes with obvious noise and no features, 795,162 particles were used for multiple rounds of ab initio reconstruction and heterogeneous refinement using four or five classes; 365,178 particles, which generated the two best 3D reconstruction, were used to carry out per-particle local-motion correction with 760 pixels box size later downsized to 380 pixels followed by several rounds of ab initio reconstruction and heterogeneous refinement using three or five classes; and 149,603 from best class were further refined by homogeneous refinement and a final non-uniform-refinement which generated a reconstruction with an overall resolution of 3.02 Å based on the Fourier shell correlation 0.143 criterion.

Building the structural model of p110γ-NB7

The previous structural model of full-length p110γ from the complex of p110γ-p101 (PDB: 7MEZ) (Rathinaswamy *et al.*, 2021a) was fit into the map using Chimera (Pettersen *et al.*, 2004). A model of the nanobody was generated using Alphafold2 using the Colabfold v1.5.2 server (Mirdita *et al.*, 2022). The CDR loops were removed from this initial model, and the remaining nanobody was fit into the map using Chimera. The final structure was built by iterative rounds of automated model building in Phenix, manual model building in COOT (Emsley *et al.*, 2010), and refinement in Phenix. real_space_refine using realspace, rigid body, and adp refinement with tight secondary structure restraints (Afonine *et al.*, 2012). This allowed for unambiguous building of the CDRs of the nanobody, and their interface with p110γ. The full refinement and validation statistics are shown in *Supplementary file 1*.

Phosphorylation analysis

For the dose-response phosphorylation of p110γ, p110γ/p84, and p110γ/p101, each protein or complex (750 nM) was mixed with ATP (200 μM), GFB (20 mM Tris pH 8.5, 100 mM NaCl, 50 mM ammonium sulfate and 0.5 mM TCEP), MgCl₂ (20 mM), and various amounts of PKCβ (4 μg, 800 ng, 160 ng, 32 ng, 6.4 ng, and 0 ng). Reactions were incubated for 3 hr on ice and quenched with 50 μL of ice-cold acidic quench buffer (0.7 M guanidine-HCl, 1% formic acid), followed by immediate freezing using liquid nitrogen and storage at −80°C.

For the experiment studying the effect of nanobody on phosphorylation, p110γ or p110γ/p101 (500 nM) was mixed with ATP (1 mM), GFB (20 mM Tris pH 8.5, 100 mM NaCl, 50 mM ammonium sulfate, and 0.5 mM TCEP), MgCl₂ (20 mM), with nanobody and PKCβ present at 1200 nM and 500 nM, respectively. Reactions were incubated for 1 hr at room temperature and quenched with 54 μL of ice-cold acidic quench buffer (0.7 M guanidine-HCl, 1% formic acid) followed by immediate freezing using liquid nitrogen and storage at −80°C.

Phosphorylation of all proteins was confirmed using mass spectrometry and PEAKS7 analysis. The LC-MS analysis of these samples was carried out using the same pipeline as used in the HDX-MS section. The phosphorylated and non-phosphorylated peptide ratios were determined by generating extracted ion chromatograms for each phosphorylated or non-phosphorylated peptide using their molecular formula and charge state in the Bruker Compass Data Analysis software. The area under each extracted curve was then extracted. The full MS quantification of each of the phosphorylated and non-phosphorylated peptide is provided in *Source data 1*.

Hydrogen-deuterium exchange mass spectrometry

Exchange reactions to assess differences in p110 γ upon phosphorylation were carried out at 20°C in 10 μ L volumes with final concentrations of 1.6 μ M for both apo and phosphorylated p110 γ . A total of two conditions were assessed: p110 γ apo and PKC β phosphorylated p110 γ . The H/D exchange reaction was initiated by the addition of 8 μ L D₂O buffer (94.3% D₂O, 100 mM NaCl, 20 mM HEPES pH 7.5) to the 2 μ L protein for a final D₂O concentration of 75.4%. Exchange was carried out over five time points (3 s on ice, and 3 s, 30 s, 300 s, and 3000 s at 20°C) and the reaction was quenched with addition of 60 μ L of ice-cold acidic quench buffer (0.7 M guanidine-HCl, 1% formic acid). After quenching, samples were immediately frozen in liquid nitrogen and stored at –80°C. All reactions were carried out in triplicate.

Protein digestion and MS/MS data collection

Protein samples for both HDX-MS and phosphorylation analysis were analyzed using the same LC-MS setup. Samples were rapidly thawed and injected onto an integrated fluidics system containing an HDx-3 PAL liquid handling robot and climate-controlled chromatography system (LEAP Technologies), a Dionex Ultimate 3000 UHPLC system, as well as an Impact HD QTOF Mass spectrometer (Bruker). The protein was run over two immobilized pepsin columns (Applied Biosystems; Poroszyme Immobilized Pepsin Cartridge, 2.1 mm $\times$ 30 mm; Thermo Fisher 2-3131-00; at 10°C and 2°C respectively) at 200 μ L/min for 3 min. The resulting peptides were collected and desalted on a C18 trap column (Acquity UPLC BEH C18 1.7 mm column [2.1 $\times$ 5 mm]; Waters 186003975). The trap was subsequently eluted in line with an ACQUITY 1.7 μ m particle, 100 $\times$ 1 mm² C18 UPLC column (Waters 186002352), using a gradient of 3–35% B (buffer A, 0.1% formic acid; buffer B, 100% acetonitrile) over 11 min immediately followed by a gradient of 35–80% B over 5 min. MS experiments acquired over a mass range from 150 to 2200 mass/charge ratio (m/z) using an electrospray ionization source operated at a temperature of 200°C and a spray voltage of 4.5 kV.

Peptide identification

Peptides were identified using data-dependent acquisition following tandem MS/MS experiments (0.5 s precursor scan from 150 to 2000 m/z ; twelve 0.25 s fragment scans from 150 to 2000 m/z). MS/MS datasets were analyzed using PEAKS7 (PEAKS), and a false discovery rate was set at 0.1% using a database of purified proteins and known contaminants. The same approach was used to identify phosphorylated and non-phosphorylated peptides for our in vitro phosphorylation experiments, with variable phosphorylation of STY residues was added to the search. The search parameters were set with a precursor tolerance of 20 parts per million, fragment mass error 0.02 Da, and charge states from 1 to 8, with a selection criterion of peptides that had a $-10\log p$ score of >24.03 for phosphorylated and >23.05 for non-phosphorylated. The MS/MS spectra of the PKC β phosphorylated peptides are included in **Figure 3—figure supplement 1**.

Mass analysis of peptide centroids and measurement of deuterium incorporation

HD-Examiner Software (Sierra Analytics) was used to automatically calculate the level of deuterium incorporation into each peptide. All peptides were manually inspected for correct charge state, correct retention time, and appropriate selection of isotopic distribution. Deuteration levels were calculated using the centroid of the experimental isotope clusters. HDX-MS results are presented with no correction for back exchange shown in **Source data 1**, with the only correction being applied correcting for the deuterium oxide percentage of the buffer used in the exchange (75.4%). Changes in any peptide at any time point greater than specified cut-offs (5% and 0.45 Da) and with an unpaired, two-tailed t-test value of $p<0.01$ was considered significant. A number of peptides in the helical domain showed isotope distributions consistent with EX1 H/D exchange. Attempts to define the relative percentages of each population using HDExaminer were extremely noisy, so representative EX1 profiles are shown in **Figure 4C**. The raw peptide deuterium incorporation graphs for a selection of peptides with significant differences are shown in **Figure 4D**, with the raw data for all analyzed peptides in **Source data 1**. To allow for visualization of differences across all peptides, we utilized number of deuterium difference (#D) plots (**Figure 4B**). These plots show the total difference in deuterium incorporation

over the entire H/D exchange time course, with each point indicating a single peptide. The data analysis statistics for all HDX-MS experiments are in Appendix 1—key resources table according to the guidelines of *Masson et al., 2019*. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (*Perez-Riverol et al., 2022*) with the dataset identifier PXD040765.

Acknowledgements

JEB is supported by the Canadian Institute of Health Research (CIHR, 168998), and the Michael Smith Foundation for Health Research (MSFHR, scholar 17686). CKY is supported by CIHR (FDN-143228, PJT-168907) and the Natural Sciences and Engineering Research Council of Canada (RGPIN-2018-03951). SDH is supported by an NSF CAREER Award (MCB-2048060). This research project was supported in part by the UBC High Resolution Macromolecular Cryo-Electron Microscopy Facility (HRMEM). A portion of this research was supported by NIH grant U24GM129547 and performed at the PNCC at OHSU and accessed through EMSL (grid.436923.9), a DOE Office of Science User Facility sponsored by the Office of Biological and Environmental Research. We appreciate help from Theo Humphreys and Rose Marie Haynes with data collection at PNCC. Competing Interests: The authors declare that they have no competing interests.

Additional information

Competing interests

John E Burke: JEB reports personal fees from Scorpion Therapeutics, Reactive therapeutics and Olema Oncology; and research grants from Novartis. The other authors declare that no competing interests exist.

Funding

Funder	Grant reference number	Author
Canadian Institutes of Health Research	FDN-143228	Calvin K Yip
Canadian Institutes of Health Research	PJT-168907	Calvin K Yip
Natural Sciences and Engineering Research Council of Canada	RGPIN-2018-03951	Calvin K Yip
National Science Foundation	MCB-2048060	Scott D Hansen
Canadian Institutes of Health Research	168998	John E Burke
Michael Smith Health Research BC	17686	John E Burke

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

Noah J Harris, Conceptualization, Validation, Investigation, Methodology, Writing – original draft, Writing – review and editing; Meredith L Jenkins, Conceptualization, Data curation, Formal analysis, Validation, Investigation, Methodology, Writing – original draft, Writing – review and editing; Sung-Eun Nam, Formal analysis; Manoj K Rathinaswamy, Matthew AH Parson, Formal analysis, Investigation, Methodology; Harish Ranga-Prasad, Udit Dalwadi, Brandon E Moeller, Eleanor Sheeky, Investigation, Methodology; Scott D Hansen, Formal analysis, Validation, Investigation, Methodology, Writing – original draft, Project administration, Writing – review and editing; Calvin K Yip, John E Burke, Conceptualization, Formal analysis, Supervision, Funding acquisition, Validation,

Investigation, Visualization, Methodology, Writing – original draft, Project administration, Writing – review and editing

Author ORCIDs

Meredith L Jenkins  <http://orcid.org/0000-0002-0685-5048>

Matthew AH Parson  <https://orcid.org/0000-0001-6270-559X>

Eleanor Sheeky  <https://orcid.org/0000-0002-1501-550X>

Scott D Hansen  <https://orcid.org/0000-0001-7005-6200>

Calvin K Yip  <http://orcid.org/0000-0003-1779-9501>

John E Burke  <http://orcid.org/0000-0001-7904-9859>

Peer review material

Reviewer #1 (Public Review): <https://doi.org/10.7554/eLife.88058.3.sa1>

Reviewer #2 (Public Review): <https://doi.org/10.7554/eLife.88058.3.sa2>

Author Response: <https://doi.org/10.7554/eLife.88058.3.sa3>

Additional files

Supplementary files

- Supplementary file 1. Cryo electron microscopy (cryo-EM) data collection, refinement, and validation statistics.
- Supplementary file 2. HDX-MS data collection and validation statistics.
- MDAR checklist
- Source data 1. Raw source data for figures.

Data availability

The model of the nanobody bound p110 γ has been deposited in the protein data bank (PDB) with the identifier 8DP0, with the EM data available at the EMDb with the identifier EMD-27627. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (**Perez-Riverol et al., 2022**) with the dataset identifier PXD040765. All source data underlying all data shown in the figures and figure supplements is available in **Source data 1**. All genetic material generated for this study is available from the lead contact by request.

The following datasets were generated:

Author(s)	Year	Dataset title	Dataset URL	Database and Identifier
Burke JE	2023	Allosteric activation or inhibition of PI3K γ mediated through conformational changes in the p110 γ helical domain	https://www.ebi.ac.uk/pride/archive/projects/PXD040765	PRIDE, PXD040765
Burke JE, Sung SE, Rathinaswamy MK, Yip CK	2023	Structure of p110 gamma bound to the Ras inhibitory nanobody NB7	https://www.rcsb.org/structure/8DP0	RCSB Protein Data Bank, 8DP0
Burke JE, Sung SE, Rathinaswamy MK, Yip CK	2023	Structure of p110 gamma bound to the Ras inhibitory nanobody NB7	https://www.ebi.ac.uk/emdb/EMD-27627	Electron Microscopy Data Bank, EMD-27627

References

- Afonine PV**, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, Terwilliger TC, Urzhumtsev A, Zwart PH, Adams PD. 2012. Towards automated crystallographic structure refinement with *p*henix.refine. *Acta Crystallographica. Section D, Biological Crystallography* **68**:352–367. DOI: <https://doi.org/10.1107/S0907444912001308>, PMID: 22505256
- Angulo I**, Vadas O, Garçon F, Banham-Hall E, Plagnol V, Leahy TR, Baxendale H, Coulter T, Curtis J, Wu C, Blake-Palmer K, Perisic O, Smyth D, Maes M, Fiddler C, Juss J, Cilliers D, Markelj G, Chandra A, Farmer G, et al. 2013. Phosphoinositide 3-kinase δ gene mutation predisposes to respiratory infection and airway damage. *Science* **342**:866–871. DOI: <https://doi.org/10.1126/science.1243292>, PMID: 24136356

- Becattini B**, Marone R, Zani F, Arsenijevic D, Seydoux J, Montani J-P, Dulloo AG, Thorens B, Preitner F, Wymann MP, Solinas G. 2011. PI3K γ within a nonhematopoietic cell type negatively regulates diet-induced thermogenesis and promotes obesity and insulin resistance. *PNAS* **108**:E854–E863. DOI: <https://doi.org/10.1073/pnas.1106698108>, PMID: 21949398
- Bell K**, Sunose M, Ellard K, Cansfield A, Taylor J, Miller W, Ramsden N, Bergamini G, Neubauer G. 2012. SAR studies around a series of triazolopyridines as potent and selective PI3K γ inhibitors. *Bioorganic & Medicinal Chemistry Letters* **22**:5257–5263. DOI: <https://doi.org/10.1016/j.bmcl.2012.06.049>, PMID: 22819766
- Bohnacker T**, Marone R, Collmann E, Calvez R, Hirsch E, Wymann MP. 2009. PI3K γ adaptor subunits define coupling to degranulation and cell motility by distinct PtdIns(3,4,5)P₃ pools in mast cells. *Science Signaling* **2**:ra27. DOI: <https://doi.org/10.1126/scisignal.2000259>, PMID: 19509406
- Bohnacker T**, Prota AE, Beaufils F, Burke JE, Melone A, Inglis AJ, Rageot D, Sele AM, Cmiljanovic V, Cmiljanovic N, Bargsten K, Aher A, Akhmanova A, Díaz JF, Fabbro D, Zvelebil M, Williams RL, Steinmetz MO, Wymann MP. 2017. Deconvolution of Buparlisib's mechanism of action defines specific PI3K and tubulin inhibitors for therapeutic intervention. *Nature Communications* **8**:14683. DOI: <https://doi.org/10.1038/ncomms14683>, PMID: 28276440
- Brasson L**, Becattini B, Sardi C, Molinaro A, Zani F, Marone R, Botindari F, Bousquenaud M, Ruegg C, Wymann MP, Solinas G. 2017. PI3K γ activity in leukocytes promotes adipose tissue inflammation and early-onset insulin resistance during obesity. *Science Signaling* **10**:eaaf2969. DOI: <https://doi.org/10.1126/scisignal.aaf2969>, PMID: 28720716
- Burke JE**, Vadas O, Berndt A, Finegan T, Perisic O, Williams RL. 2011. Dynamics of the phosphoinositide 3-kinase p110 δ Interaction with p85 α and membranes reveals aspects of regulation distinct from p110 α . *Structure* **19**:1127–1137. DOI: <https://doi.org/10.1016/j.str.2011.06.003>, PMID: 21827948
- Burke JE**, Perisic O, Masson GR, Vadas O, Williams RL. 2012. Oncogenic mutations mimic and enhance dynamic events in the natural activation of phosphoinositide 3-kinase P110 A (Pik3Ca). *PNAS* **109**:15259–15264. DOI: <https://doi.org/10.1073/pnas.1205508109>, PMID: 22949682
- Burke JE**, Williams RL. 2013. Dynamic steps in receptor tyrosine kinase mediated activation of class IA phosphoinositide 3-kinases (PI3K) captured by H/D exchange (HDX-MS). *Advances in Biological Regulation* **53**:97–110. DOI: <https://doi.org/10.1016/j.jbior.2012.09.005>, PMID: 23194976
- Burke JE**, Williams RL. 2015. Synergy in activating class I PI3Ks. *Trends in Biochemical Sciences* **40**:88–100. DOI: <https://doi.org/10.1016/j.tibs.2014.12.003>, PMID: 25573003
- Burke JE**. 2018. Structural basis for regulation of phosphoinositide kinases and their involvement in human disease. *Molecular Cell* **71**:653–673. DOI: <https://doi.org/10.1016/j.molcel.2018.08.005>, PMID: 30193094
- Campa CC**, Silva RL, Margaria JP, Pirali T, Mattos MS, Kraemer LR, Reis DC, Grosa G, Copperi F, Dalmarco EM, Lima-Júnior RCP, Aprile S, Sala V, Dal Bello F, Prado DS, Alves-Filho JC, Medana C, Cassali GD, Tron GC, Teixeira MM, et al. 2018. Inhalation of the prodrug PI3K inhibitor CL27c improves lung function in asthma and fibrosis. *Nature Communications* **9**:5232. DOI: <https://doi.org/10.1038/s41467-018-07698-6>, PMID: 30542075
- Camps M**, Rückle T, Ji H, Ardisson V, Rintelen F, Shaw J, Ferrandi C, Chabert C, Gillieron C, Françon B, Martin T, Gretener D, Perrin D, Leroy D, Vitte PA, Hirsch E, Wymann MP, Cirillo R, Schwarz MK, Rommel C. 2005. Blockade of PI3K γ suppresses joint inflammation and damage in mouse models of rheumatoid arthritis. *Nature Medicine* **11**:936–943. DOI: <https://doi.org/10.1038/nm1284>, PMID: 16127437
- De Henau O**, Rausch M, Winkler D, Campesato LF, Liu C, Cymerman DH, Budhu S, Ghosh A, Pink M, Tchaicha J, Douglas M, Tibbitts T, Sharma S, Proctor J, Kosmider N, White K, Stern H, Soglia J, Adams J, Palombella VJ, et al. 2016. Overcoming resistance to checkpoint blockade therapy by targeting PI3K γ in myeloid cells. *Nature* **539**:443–447. DOI: <https://doi.org/10.1038/nature20554>, PMID: 27828943
- Deladeriere A**, Gambardella L, Pan D, Anderson KE, Hawkins PT, Stephens LR. 2015. The regulatory subunits of PI3K γ control distinct neutrophil responses. *Science Signaling* **8**:ra8. DOI: <https://doi.org/10.1126/scisignal.2005564>, PMID: 25605974
- Dornan GL**, Siempelkamp BD, Jenkins ML, Vadas O, Lucas CL, Burke JE. 2017. Conformational disruption of PI3K δ regulation by immunodeficiency mutations in Pik3Cd and Pik3R1. *PNAS* **114**:1982–1987. DOI: <https://doi.org/10.1073/pnas.1617244114>
- Emsley P**, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of Coot. *Acta Crystallographica. Section D, Biological Crystallography* **66**:486–501. DOI: <https://doi.org/10.1107/S0907444910007493>, PMID: 20383002
- Evans CA**, Liu T, Lescarbeau A, Nair SJ, Grenier L, Pradeilles JA, Glenadel Q, Tibbitts T, Rowley AM, DiNitto JP, Brophy EE, O'Hearn EL, Ali JA, Winkler DG, Goldstein SI, O'Hearn P, Martin CM, Hoyt JG, Soglia JR, Cheung C, et al. 2016. Discovery of a selective Phosphoinositide-3-kinase (PI3K)- γ inhibitor (IPI-549) as an immuno-oncology clinical candidate. *ACS Medicinal Chemistry Letters* **7**:862–867. DOI: <https://doi.org/10.1021/acsmedchemlett.6b00238>, PMID: 27660692
- Gangadhara G**, Dahl G, Bohnacker T, Rae R, Gunnarsson J, Blaho S, Öster L, Lindmark H, Karabelas K, Pemberton N, Tyrchan C, Mogemark M, Wymann MP, Williams RL, Perry MWD, Papavoine T, Petersen J. 2019. A class of highly selective inhibitors bind to an active state of PI3K γ . *Nature Chemical Biology* **15**:348–357. DOI: <https://doi.org/10.1038/s41589-018-0215-0>, PMID: 30718815
- Hansen SD**, Huang WYC, Lee YK, Bieling P, Christensen SM, Groves JT. 2019. Stochastic geometry sensing and polarization in a lipid kinase-phosphatase competitive reaction. *PNAS* **116**:15013–15022. DOI: <https://doi.org/10.1073/pnas.1901744116>, PMID: 31278151
- Hawkins PT**, Stephens LR. 2015. PI3K signalling in inflammation. *Biochimica et Biophysica Acta* **1851**:882–897. DOI: <https://doi.org/10.1016/j.bbali.2014.12.006>, PMID: 25514767

- Jenkins ML, Ranga-Prasad H, Parson MAH, Harris NJ, Rathinaswamy MK, Burke JE. 2023. Oncogenic mutations of Plk3Ca lead to increased membrane recruitment driven by reorientation of the ABD, P85 and C-terminus. *Nature Communications* **14**:181. DOI: <https://doi.org/10.1038/s41467-023-35789-6>, PMID: 36635288
- Jin JR, Gogvadze E, Xavier AR, Bohnacker T, Voelzmann J, Wymann MP. 2020. PI3Ky regulatory protein p84 determines mast cell sensitivity to Ras inhibition—Moving towards cell specific PI3K targeting?. *Frontiers in Immunology* **11**:585070. DOI: <https://doi.org/10.3389/fimmu.2020.585070>, PMID: 33193405
- Johannessen CM, Boehm JS, Kim SY, Thomas SR, Wardwell L, Johnson LA, Emery CM, Stransky N, Cogdill AP, Barretina J, Caponigro G, Hieronymus H, Murray RR, Salehi-Ashtiani K, Hill DE, Vidal M, Zhao JJ, Yang X, Alkan O, Kim S, et al. 2010. COT drives resistance to RAF inhibition through MAP kinase pathway reactivation. *Nature* **468**:968–972. DOI: <https://doi.org/10.1038/nature09627>, PMID: 21107320
- Kaneda MM, Messer KS, Ralainirina N, Li H, Leem CJ, Gorjestani S, Woo G, Nguyen AV, Figueiredo CC, Foubert P, Schmid MC, Pink M, Winkler DG, Rausch M, Palombella VJ, Kutok J, McGovern K, Frazer KA, Wu X, Karin M, et al. 2016a. Pi3Ky is a molecular switch that controls immune suppression. *Nature* **539**:437–442. DOI: <https://doi.org/10.1038/nature19834>, PMID: 27642729
- Kaneda MM, Cappello P, Nguyen AV, Ralainirina N, Hardamon CR, Foubert P, Schmid MC, Sun P, Mose E, Bouvet M, Lowy AM, Valasek MA, Sasik R, Novelli F, Hirsch E, Varner JA. 2016b. Macrophage PI3Ky drives pancreatic ductal adenocarcinoma progression. *Cancer Discovery* **6**:870–885. DOI: <https://doi.org/10.1158/2159-8290.CD-15-1346>, PMID: 27179037
- Kurig B, Shymanets A, Bohnacker T, Prajwal BC, Ahmadian MR, Schaefer M, Gohla A, Harteneck C, Wymann MP, Jeanclos E, Nürnberg B. 2009. Ras is an indispensable coregulator of the class IB phosphoinositide 3-kinase p87/p110 γ . *PNAS* **106**:20312–20317. DOI: <https://doi.org/10.1073/pnas.0905506106>, PMID: 19906996
- Laffargue M, Calvez R, Finan P, Trifilieff A, Barbier M, Altruda F, Hirsch E, Wymann MP. 2002. Phosphoinositide 3-kinase gamma is an essential amplifier of mast cell function. *Immunity* **16**:441–451. DOI: [https://doi.org/10.1016/s1074-7613\(02\)00282-0](https://doi.org/10.1016/s1074-7613(02)00282-0), PMID: 11911828
- Lanahan SM, Wymann MP, Lucas CL. 2022. The role of Pi3Ky in the immune system: new insights and translational implications. *Nature Reviews. Immunology* **22**:687–700. DOI: <https://doi.org/10.1038/s41577-022-00701-8>, PMID: 35322259
- Lawrence MS, Stojanov P, Mermel CH, Robinson JT, Garraway LA, Golub TR, Meyerson M, Gabriel SB, Lander ES, Getz G. 2014. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature* **505**:495–501. DOI: <https://doi.org/10.1038/nature12912>, PMID: 24390350
- Li Z, Jiang H, Xie W, Zhang Z, Smrcka AV, Wu D. 2000. Roles of PLC- β 2 and - β 3 and PI3Ky in chemoattractant-mediated signal transduction. *Science* **287**:1046–1049. DOI: <https://doi.org/10.1126/science.287.5455.1046>, PMID: 10669417
- Li H, Prever L, Hirsch E, Gulluni F. 2021. Targeting PI3K/AKT/mTOR signaling pathway in breast cancer. *Cancers* **13**:3517. DOI: <https://doi.org/10.3390/cancers13143517>, PMID: 34298731
- Liu X, Zhou Q, Hart JR, Xu Y, Yang S, Yang D, Vogt PK, Wang MW. 2022. Cryo-EM structures of cancer-specific Helical and kinase domain mutations of PI3K α . *PNAS* **119**:e2215621119. DOI: <https://doi.org/10.1073/pnas.2215621119>, PMID: 36343266
- Lucas CL, Kuehn HS, Zhao F, Niemela JE, Deenick EK, Palendira U, Avery DT, Moens L, Cannons JL, Biancalana M, Stoddard J, Ouyang W, Frucht DM, Rao VK, Atkinson TP, Agharabimi A, Hussey AA, Folio LR, Olivier KN, Fleisher TA, et al. 2014. Dominant-activating germline mutations in the gene encoding the PI(3)K catalytic subunit p110 δ result in T cell senescence and human immunodeficiency. *Nature Immunology* **15**:88–97. DOI: <https://doi.org/10.1038/ni.2771>, PMID: 24165795
- Luo L, Wall AA, Tong SJ, Hung Y, Xiao Z, Tarique AA, Sly PD, Fantino E, Marzolo MP, Stow JL. 2018. TLR Crosstalk activates Lrp1 to recruit Rab8A and PI3Ky for suppression of inflammatory responses. *Cell Reports* **24**:3033–3044. DOI: <https://doi.org/10.1016/j.celrep.2018.08.028>, PMID: 30208326
- Mandelker D, Gabelli SB, Schmidt-Kittler O, Zhu J, Cheong I, Huang CH, Kinzler KW, Vogelstein B, Amzel LM. 2009. A frequent kinase domain mutation that changes the interaction between PI3K α and the membrane. *PNAS* **106**:16996–17001. DOI: <https://doi.org/10.1073/pnas.0908444106>, PMID: 19805105
- Masson GR, Burke JE, Ahn NG, Anand GS, Borchers C, Brier S, Bou-Assaf GM, Engen JR, Englander SW, Faber J, Garlish R, Griffin PR, Gross ML, Guttman M, Hamuro Y, Heck AJR, Houde D, Iacob RE, Jørgensen TJD, Kaltashov IA, et al. 2019. Recommendations for performing, interpreting and reporting hydrogen deuterium exchange mass spectrometry (HDX-MS) experiments. *Nature Methods* **16**:595–602. DOI: <https://doi.org/10.1038/s41592-019-0459-y>, PMID: 31249422
- Miled N, Yan Y, Hon WC, Perisic O, Zvelebil M, Inbar Y, Schneidman-Duhovny D, Wolfson HJ, Backer JM, Williams RL. 2007. Mechanism of two classes of cancer mutations in the phosphoinositide 3-kinase catalytic subunit. *Science* **317**:239–242. DOI: <https://doi.org/10.1126/science.1135394>, PMID: 17626883
- Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022. Colabfold: making protein folding accessible to all. *Nature Methods* **19**:679–682. DOI: <https://doi.org/10.1038/s41592-022-01488-1>, PMID: 35637307
- Okkenhaug K. 2013. Signaling by the Phosphoinositide 3-kinase family in immune cells. *Annual Review of Immunology* **31**:675–704. DOI: <https://doi.org/10.1146/annurev-immunol-032712-095946>, PMID: 23330955
- Pacold ME, Suire S, Perisic O, Lara-Gonzalez S, Davis CT, Walker EH, Hawkins PT, Stephens L, Eccleston JF, Williams RL. 2000. Crystal structure and functional analysis of Ras binding to its effector phosphoinositide 3-kinase gamma. *Cell* **103**:931–943. DOI: [https://doi.org/10.1016/s0092-8674\(00\)00196-3](https://doi.org/10.1016/s0092-8674(00)00196-3), PMID: 11136978
- Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A, Vizcaino JA. 2022. The PRIDE

- database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids Research* **50**:D543–D552. DOI: <https://doi.org/10.1093/nar/gkab1038>, PMID: 34723319
- Perino A**, Ghigo A, Ferrero E, Morello F, Santulli G, Baillie GS, Damilano F, Dunlop AJ, Pawson C, Walser R, Levi R, Altruda F, Silengo L, Langeberg LK, Neubauer G, Heymans S, Lembo G, Wymann MP, Wetzker R, Houslay MD, et al. 2011. Integrating cardiac Pip3 and cAMP signaling through a PKA anchoring function of p110 Γ . *Molecular Cell* **42**:84–95. DOI: <https://doi.org/10.1016/j.molcel.2011.01.030>, PMID: 21474070
- Pettersen EF**, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera—a visualization system for exploratory research and analysis. *Journal of Computational Chemistry* **25**:1605–1612. DOI: <https://doi.org/10.1002/jcc.20084>, PMID: 15264254
- Punjani A**, Rubinstein JL, Fleet DJ, Brubaker MA. 2017. cryoSPARC: Algorithms for rapid unsupervised Cryo-EM structure determination. *Nature Methods* **14**:290–296. DOI: <https://doi.org/10.1038/nmeth.4169>, PMID: 28165473
- Rathinaswamy MK**, Burke JE. 2020. Class I Phosphoinositide 3-kinase (PI3K) regulatory subunits and their roles in signaling and disease. *Advances in Biological Regulation* **75**:100657. DOI: <https://doi.org/10.1016/j.jbior.2019.100657>, PMID: 31611073
- Rathinaswamy MK**, Dalwadi U, Fleming KD, Adams C, Stariha JTB, Pardon E, Baek M, Vadas O, DiMaio F, Steyaert J, Hansen SD, Yip CK, Burke JE. 2021a. Structure of the phosphoinositide 3-kinase (PI3K) P110 Γ -P101 complex reveals molecular mechanism of GPCR activation. *Science Advances* **7**:eabj4282. DOI: <https://doi.org/10.1126/sciadv.abj4282>, PMID: 34452907
- Rathinaswamy MK**, Fleming KD, Dalwadi U, Pardon E, Harris NJ, Yip CK, Steyaert J, Burke JE. 2021b. HDX-MS-Optimized approach to characterize Nanobodies as tools for biochemical and structural studies of class IB phosphoinositide 3-kinases. *Structure* **29**:1371–1381. DOI: <https://doi.org/10.1016/j.str.2021.07.002>, PMID: 34348129
- Rathinaswamy MK**, Gaieb Z, Fleming KD, Borsari C, Harris NJ, Moeller BE, Wymann MP, Amaro RE, Burke JE. 2021c. Disease-related mutations in PI3K γ disrupt regulatory C-terminal dynamics and reveal a path to selective inhibitors. *eLife* **10**:e64691. DOI: <https://doi.org/10.7554/eLife.64691>, PMID: 33661099
- Rathinaswamy MK**, Jenkins ML, Duewell BR, Zhang X, Harris NJ, Evans JT, Stariha JTB, Dalwadi U, Fleming KD, Ranga-Prasad H, Yip CK, Williams RL, Hansen SD, Burke JE. 2023. Molecular basis for differential activation of p101 and p84 complexes of PI3K γ by Ras and GPCRs. *Cell Reports* **42**:112172. DOI: <https://doi.org/10.1016/j.celrep.2023.112172>, PMID: 36842083
- Robert X**, Gouet P. 2014. Deciphering key features in protein structures with the new ENDscript server. *Nucleic Acids Research* **42**:W320–W324. DOI: <https://doi.org/10.1093/nar/gku316>, PMID: 24753421
- Rynkiewicz NK**, Anderson KE, Suire S, Collins DM, Karanasios E, Vadas O, Williams R, Oxley D, Clark J, Stephens LR, Hawkins PT. 2020. G $\beta\gamma$ is a direct regulator of endogenous P101/P110 Γ and P84/P110 Γ PI3K γ complexes in mouse neutrophils. *Science Signaling* **13**:eaaz4003. DOI: <https://doi.org/10.1126/scisignal.aaz4003>, PMID: 33144519
- Samuels Y**, Wang Z, Bardelli A, Silliman N, Ptak J, Szabo S, Yan H, Gazdar A, Powell SM, Riggins GJ, Willson JKV, Markowitz S, Kinzler KW, Vogelstein B, Velculescu VE. 2004. High frequency of mutations of the *PIK3CA* gene in human cancers. *Science* **304**:554. DOI: <https://doi.org/10.1126/science.1096502>, PMID: 15016963
- Shymanets A**, Bucher K, Beer-Hammer S, Harteneck C, Nürnberg B. 2013. P87 and P101 subunits are distinct regulators determining class IB phosphoinositide 3-kinase (PI3K) specificity. *The Journal of Biological Chemistry* **288**:31059–31068. DOI: <https://doi.org/10.1074/jbc.M113.508234>, PMID: 24014027
- Siempelkamp BD**, Rathinaswamy MK, Jenkins ML, Burke JE. 2017. Molecular mechanism of activation of class IA phosphoinositide 3-kinases (PI3Ks) by membrane-localized HRas. *The Journal of Biological Chemistry* **292**:12256–12266. DOI: <https://doi.org/10.1074/jbc.M117.789263>, PMID: 28515318
- Stephens LR**, Eguinoa A, Erdjument-Bromage H, Lui M, Cooke F, Coadwell J, Smrcka AS, Thelen M, Cadwallader K, Tempst P, Hawkins PT. 1997. The G $\beta\gamma$ sensitivity of a PI3K is dependent upon a tightly associated adaptor, p101. *Cell* **89**:105–114. DOI: [https://doi.org/10.1016/S0092-8674\(00\)80187-7](https://doi.org/10.1016/S0092-8674(00)80187-7), PMID: 9094719
- Takeda AJ**, Maher TJ, Zhang Y, Lanahan SM, Bucklin ML, Compton SR, Tyler PM, Comrie WA, Matsuda M, Olivier KN, Pittaluga S, McElwee JJ, Long Priel DA, Kuhns DB, Williams RL, Mustillo PJ, Wymann MP, Koneti Rao V, Lucas CL. 2019. Human PI3K γ deficiency and its microbiota-dependent mouse model reveal immunodeficiency and tissue Immunopathology. *Nature Communications* **10**:4364. DOI: <https://doi.org/10.1038/s41467-019-12311-5>, PMID: 31554793
- Vadas O**, Dbouk HA, Shymanets A, Perisic O, Burke JE, Abi Saab WF, Khalil BD, Harteneck C, Bresnick AR, Nürnberg B, Backer JM, Williams RL. 2013. Molecular determinants of PI3K γ -mediated activation downstream of G-protein-coupled receptors (GPCRs). *PNAS* **110**:18862–18867. DOI: <https://doi.org/10.1073/pnas.1304801110>, PMID: 24190998
- Vanhaesebroeck B**, Perry MWD, Brown JR, André F, Okkenhaug K. 2021. PI3K inhibitors are finally coming of age. *Nature Reviews. Drug Discovery* **20**:741–769. DOI: <https://doi.org/10.1038/s41573-021-00209-1>, PMID: 34127844
- Vasan N**, Cantley LC. 2022. At a crossroads: how to translate the roles of PI3K in oncogenic and metabolic signalling into improvements in cancer therapy. *Nature Reviews. Clinical Oncology* **19**:471–485. DOI: <https://doi.org/10.1038/s41571-022-00633-1>, PMID: 35484287
- Walker EH**, Perisic O, Ried C, Stephens L, Williams RL. 1999. Structural insights into phosphoinositide 3-kinase catalysis and signalling. *Nature* **402**:313–320. DOI: <https://doi.org/10.1038/46319>, PMID: 10580505

- Walser R**, Burke JE, Gogvadze E, Bohnacker T, Zhang X, Hess D, Küenzi P, Leitges M, Hirsch E, Williams RL, Laffargue M, Wymann MP. 2013. PKC β phosphorylates Pi3Ky to activate it and release it from GPCR control. *PLOS Biology* **11**:e1001587. DOI: <https://doi.org/10.1371/journal.pbio.1001587>, PMID: 23824069
- Zhang X**, Vadas O, Perisic O, Anderson KE, Clark J, Hawkins PT, Stephens LR, Williams RL. 2011. Structure of lipid kinase P110 β /P85 β elucidates an unusual Sh2-domain-mediated inhibitory mechanism. *Molecular Cell* **41**:567–578. DOI: <https://doi.org/10.1016/j.molcel.2011.01.026>, PMID: 21362552

Appendix 1

Appendix 1—key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (<i>Escherichia coli</i>)	<i>E. coli</i> XL10-GOLD KanR Ultracompetent Cells	Agilent	200317	
Strain, strain background (<i>Escherichia coli</i>)	BL21 <i>E. coli</i> C41 (DE3) RIPL	PMID: 24876499	C41	
Strain, strain background (<i>Spodoptera frugiperda</i>)	Sf9 Insect Cells	Expression systems	94-001S	
Strain, strain background (<i>Escherichia coli</i>)	<i>E. coli</i> DH10EMBacY Competent Cells	Geneva Biotech	DH10EMBacY	
Recombinant DNA reagent	pMultiBac-Gβ1/Gγ2	PMID:34452907	pOP737	
Recombinant DNA reagent	pACEBac1-hsp110γ	PMID:34452907	MR30	
Recombinant DNA reagent	pMultiBac-hsp110γ-ssp101	PMID:34452907	MR22	
Recombinant DNA reagent	pMultiBac-hsp110γ-mmp84	PMID:34452907	MR24	
Recombinant DNA reagent	pFastBac HRas G12V	PMID:34452907	BS9	
Recombinant DNA reagent	biGBac hsp110γ/ybbr-hsp84	PMID:36842083	HP28	
Recombinant DNA reagent	biGBac hsp110γ/ybbr-hsp101	PMID:36842083	HP29	
Recombinant DNA reagent	his6-GST-PrescissionProtease-SNAP-RBD(K65E)	PMID:34452907	pSH936	
Recombinant DNA reagent	his6TEV-HRas(1-184aa) C118S, C181S	PMID:34452907	pSH414	
Recombinant DNA reagent	his6-G2, SNAP-G1 (DUAL FastBac)	PMID:34452907	pSH651	
Recombinant DNA reagent	pACEBAC-PKCβII (internal tev cleavage site)	This paper	pMR56	
Recombinant DNA reagent	pFASTBac p110α	PMID: 28515318	pOV1181	
Recombinant DNA reagent	pFASTBac p110β	PMID: 28515318	pOV1182	
Recombinant DNA reagent	pFASTBac p110δ	PMID: 28515318	pOV1183	
Recombinant DNA reagent	pFASTBac p85β	This paper	EX21	
Sequence-based reagent	Fwd primer for amplifying KD of PKCII	Sigma	MR51F	GTATTTTCAGGGCgccggtaccACGA CCAACACTGTCTCCAAATTG
Sequence-based reagent	Rvs primer for amplifying KD of PKCII	Sigma	MR51R	gactcgagcggccgcTTATAGCTCTT GACTTCGGGTTTAAAAATTG
Sequence-based reagent	Fwd primer for amplifying N term of PKCII	Sigma	MR52F	CCATCACggatctggcgtagt ATGGCTGACCCGCTGCG
Sequence-based reagent	Rvs primer for amplifying N term of PKCII	Sigma	MR52R	GCCCTGAAAATACAGGTTTTCC TTTTCTCCGGGACCTTGTTCCC

Appendix 1 Continued on next page

Appendix 1 Continued

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Sequence-based reagent	Fwd primer for adding stop codon to PKCII	Sigma	MR56F	AGTCAAGAGCTAAgcggccgctcgagtctagagcctgc
Sequence-based reagent	Rvs primer for adding stop codon to PKCII	Sigma	MR56R	gactcgagcggccgcTTAGCTCTTGACTTCGGGTTTAAAAATTCA
Commercial assay or kit	Transcreener ADP2 FI Assay (1000 Assay, 384 Well)	BellBrook Labs	3013-1K	
Chemical compound, drug	Deuterium oxide 99.9%	Sigma	151882	
Chemical compound, drug	Guanosine 5'-diphosphate (GDP) sodium salt hydrate	Sigma	G7127-100MG	
Chemical compound, drug	Guanosine 5'-triphosphate (GTP) sodium salt hydrate	Sigma	G8877-250MG	
Chemical compound, drug	Sodium deoxycholate	Sigma	D6750	
Chemical compound, drug	Polyoxyethylene (10) lauryl ether	Sigma	P9769	
Chemical compound, drug	CHAPS, Molecular Biology Grade	EMD Millipore	220201	
Chemical compound, drug	Phosphatidylserine (Porcine Brain)	Avanti	840032C	
Chemical compound, drug	Phosphatidylethanolamine (Egg yolk)	Sigma	P6386	
Chemical compound, drug	Cholesterol	Sigma	47,127U	
Chemical compound, drug	Phosphatidylcholine (Egg yolk)	Avanti	840051C	
Chemical compound, drug	Phosphatidylinositol-4,5-bisphosphate (Porcine Brain)	Avanti	840046	
Chemical compound, drug	Sphingomyelin (Egg yolk)	Sigma	S0756	
Chemical compound, drug	1,2-Dioleoyl- <i>sn</i> -glycero-3-phosphocholine (DOPC)	Avanti	850375C	
Chemical compound, drug	1,2-Dioleoyl- <i>sn</i> -glycero-3-phospho-L-serine (18:1, DOPS)	Avanti	840035C	
Chemical compound, drug	1,2-Dioleoyl- <i>sn</i> -glycero-3-phosphoethanolamine- <i>N</i> -[4-(<i>p</i> -maleimidomethyl)cyclohexane-carboxamide] (18:1 MCC-PE)	Avanti	780201C	
Chemical compound, drug	10 mg/mL beta casein solution	Thermo Fisher	37528	
Chemical compound, drug	10× PBS (pH 7.4)	Corning	46-013CM	
Chemical compound, drug	glucose oxidase from <i>Aspergillus niger</i> (225 U/mg)	Biophoretics	B01357.02	
Chemical compound, drug	Catalase	Sigma	C40-100MG Bovine Liver	
Chemical compound, drug	Trolox	Cayman Chemicals	10011659	
Chemical compound, drug	Dyomics 647 maleimide dye	Dyomics	647P1-03	
Chemical compound, drug	Coenzyme A	Sigma	C3019	

Appendix 1 Continued on next page

Appendix 1 Continued

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Chemical compound, drug	Sulfuric acid	Sigma	58105-2.5L-PC	
Software, algorithm	COOT-0.9.4.1	CCP4	https://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot/	
Software, algorithm	Phenix-1.19.1	Open source	https://www.phenix-online.org/	
Software, algorithm	PDBePISA (Proteins, Interfaces, Structures and Assemblies)	EMBL-EBI	https://www.ebi.ac.uk/pdbe/pisa/pistart.html	
Software, algorithm	ESPrIpt 3.0	Robert and Gouet, 2014	https://espript.ibcp.fr	
Software, algorithm	HDExaminer	Sierra Analytics	http://massspec.com/hdexaminer	
Software, algorithm	GraphPad Prism 7	GraphPad	https://www.graphpad.com	
Software, algorithm	PyMOL	Schroedinger	http://pymol.org	
Software, algorithm	Compass Data Analysis	Bruker	https://www.bruker.com	
Software, algorithm	ChimeraX	UCSF	https://www.rbvi.ucsf.edu/chimerax/	
Software, algorithm	ImageJ/Fiji	ImageJ	https://imagej.net/software/fiji/	
Software, algorithm	Nikon NIS elements	Nikon	https://www.microscope.healthcare.nikon.com/products/software/nis-elements	
Software, algorithm	cryoSPARC v.3.3.2	Structura Biotechnology	https://cryosparc.com/	
Other	Sf9 insect cells for expression	Expression Systems	94-001S	Sf9 cell line used for protein expression (Methods)
Other	Insect cell media	Expression Systems	96-001-01	Sf9 cell media (Methods)
Other	Hellmanex III cleaning solution	Fisher	14-385-864	Cleaning solution for TIRF experiments (PMID:36842083)
Other	Six-well sticky-side chamber	IBIDI	80608	Side chamber used for TIRF experiments (Methods)
Other	C-Flat 2/2T grids	Electron Microscopy Sciences	CFT-223C	Grids used for EM studies (Methods)
Other	PDB coordinate file for p110 γ -NB7 structure	PDB	8DP0	PDB to use for p110-NB7 structure
Other	EM density file for p110 γ -NB7 complex	EMD	EMD-27627	EM density file to use for p110-NB7
Other	HDX-MS and phosphorylation proteomics data	PRIDE	PXD040765	Database and number where HDX data was uploaded