

Interfacial electric fields catalyze Ullmann coupling reactions on gold surfaces

Ilana B. Stone^{1†}, Rachel L. Starr^{1†}, Norah Hoffmann^{1‡}, Xiao Wang^{2‡}, Austin M. Evans,¹ Colin Nuckolls¹, Tristan H. Lambert³, Michael L. Steigerwald¹, Timothy C. Berkelbach^{1,2*}, Xavier Roy^{1*}, Latha Venkataraman^{1,4*}

¹ Department of Chemistry, Columbia University, New York, New York, 10027, United States

² Center for Computational Quantum Physics, Flatiron Institute, New York, New York, 10010, United States

³ Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853, United States

⁴ Department of Applied Physics, Columbia University, New York, New York, 10027, United States

† Equal contribution

‡ Equal contribution

Corresponding Authors:

Timothy Berkelbach – tcb2112@columbia.edu

Xavier Roy – xr2114@columbia.edu

Latha Venkataraman – lv2117@columbia.edu

*The electric fields created at solid-liquid interfaces are important in heterogeneous catalysis. Here we describe the Ullmann coupling of aryl iodides on rough gold surfaces, which we monitor *in situ* using the scanning tunneling microscope-based break junction (STM-BJ) and *ex situ* using mass spectrometry and fluorescence spectroscopy. We find that this Ullmann coupling reaction occurs only on rough gold surfaces in polar solvents, the latter of which implicates interfacial electric fields. These experimental observations are supported by density functional theory calculations that elucidate the roles of surface roughness and local electric fields on the reaction. More broadly, this touchstone study offers a facile method to access and probe in real time an increasingly prominent yet incompletely understood mode of catalysis.*

Electric fields are ubiquitous at the interface of a metal and a liquid—for example, in the double layer of electrolyte solutions or in the solvent’s reaction field in response to a molecular adsorbate.¹⁻⁵ These interfacial electric fields can be measured by vibrational Stark shift spectroscopy of probe molecules and are highly variable depending on the ionic concentrations,⁶ the surface morphology, the nature of adsorbates, and the dielectric properties of the solvent. Such interfacial fields offer new opportunities to promote chemical reactivity without the use of externally applied voltages or other stimuli.^{1, 7-9} In this work, we demonstrate that rough Au surfaces, when submerged in a highly polar solvent, can act as mesoscopic active sites for on-surface aryl iodide dehalogenation, and subsequent aryl–aryl bond formation via an Ullmann-type coupling reaction. The traditional solution-phase Ullmann coupling is a long-standing strategy to form biaryls from aryl halides using stoichiometric elemental Cu in polar solvents.¹⁰⁻¹¹ While still heavily used in traditional solution-phase organic synthesis, the Ullmann coupling has more recently emerged as a stalwart of on-surface chemistry.¹²⁻¹⁵ In these reactions, aryl bromides or iodides are deposited on single-crystal Cu/Ag/Au surfaces in ultra-high vacuum and heated to cleave the aryl–halide bond and form the biaryl product.¹⁶⁻¹⁷ These reactions offer an attractive strategy for the synthesis of covalent two-dimensional polymers, but leave products bound to surfaces, and thus preclude the diversification of reactions and limit the range of accessible chemical substrates that can be probed.¹⁸

In this work, we report the Ullmann coupling of aryl iodides on a Au surface submerged in solution under ambient conditions. The surface also serves as a scanning tunneling microscope-based break junction (STM-BJ) substrate, thereby enabling *in situ* determination of product formation by monitoring single-molecule junction conductance. STM-BJ measurements provide molecule-specific signatures (i.e. the most probable molecular conductance and molecular plateau

length) that can be used to identify carbon-carbon bond formation *in situ*.¹⁹ Our observations are consistent with *ex situ* fluorescence spectroscopy and mass spectrometry measurements that confirm biaryl formation across a range of aryl iodide precursors, including those with strongly coordinating groups such as thiomethyls, pyridines, and amines. Notably, in addition to a Au surface, we find that the Ullmann coupling requires an Au surface that is rough at least on the 10-100 nm scale, as measured by AFM (Supporting Information Figure S1), and a highly polar solvent. Furthermore, the reaction works primarily with aryl iodides. Finally, we probe the mechanism of the Ullmann coupling and propose that interfacial electric fields contribute to the reaction catalysis, which is distinct from the mechanisms commonly invoked for solution-based Ullmann couplings.^{12, 16, 20-24}




Figure 1. (a) Molecular structures of aryl iodide **I1** and tetraphenyl **D1**. (b) Logarithmically binned 1D histograms showing a molecular conductance peak for **D1** that increases over time in a solution of **I1** measured at 100 mV in PC. (c) 2D conductance-displacement histogram created by overlaying all **D1** traces taken at 100 mV from the 15-18 hr measurement.

We first characterize, *in situ*, the formation of the homo-coupled dimer (**D1**, 4,4'-diamino-4,4'-quaterphenylene) when a solution of 4-iodo-4'-amino-biphenyl (**I1**) is subject to STM-BJ measurements from a solution deposited on a Au surface (Figure 1a). A 0.1 mM solution of **I1** in propylene carbonate (PC), a highly polar solvent with a dielectric constant of ~64 at 25 °C,²⁵ is

placed on a rough Au STM-BJ substrate prepared by evaporating 70-100 nm of Au on a steel puck (see SI Figure S1 for SEM and AFM images). The reaction is monitored by performing STM-BJ measurements over a period of 18 hours using an insulated Au STM tip (see SI for details),²⁶ while applying a bias voltage of 100 mV. Figure 1b presents conductance histograms created over the measurement period showing a molecular conductance peak centered at $3.7 \times 10^{-5} G_0$ ($G_0 = 2e^2/h$ is the conductance quantum). This conductance value is comparable to previous measurements of tetraphenyl diamine **D1** (SI Figure S2a).²⁷ The 2D conductance-displacement histogram shown in Figure 1c reveals that this conductance peak also features a molecular plateau extending to ~0.7 nm, which is considerably longer than the length expected for a biphenyl (see Figure S2 for additional 2D histogram).²⁷ The longer plateau length observed *in situ* is consistent with that of **D1**,²⁷ indicating that **D1** is formed under these conditions (Figure 1c). The lower conducting **D1** forms via an Ullmann coupling of two **I1** molecules, which results in a peak whose height increases over time as is seen in Figure 1b.

We next demonstrate that the coupling reaction happens on the Au surface without performing STM-BJ measurements.²⁸ An aliquot of a solution of **I1** and TBAPF₆ prepared in PC was deposited on an Au substrate. Samples of the solution were subsequently collected after 2, 5 and 10 hours and measured by fluorescence spectroscopy. If the formation of **D1** required the STM measurements, we would not detect any product with this experiment through fluorescence measurements. Figures 2a and 2b show fluorescence spectra measured using different substrates and conditions with those of an **I1** solution that was not exposed to the Au substrate and a solution of the product **D1** synthesized *ex situ*. These spectra are measured using 284 nm and 340 nm photoexcitation, respectively, with wavelengths chosen based on the electronic absorption spectra of **I1** and **D1** measured *ex situ*, (SI Figure S3a). The spectrum reveals that after 2 hours of exposure

to the Au substrate in an ionic solution, in addition to the **I1** peak at 380 nm, a smaller peak appears at 455 nm corresponding to **D1** (Figure 2a, light blue). Over the course of 10-12 hours, the emission peak consistent with that of **D1** increases in intensity (Figure 2a, dark blue), indicating that dimerization is occurring when **I1** is exposed to an interfacial field on a rough Au substrate. Starting with a 100 μ M solution of **I1**, a solution was obtained that has 6.25 μ M **D1** after 12 hours (Figure 2b, blue), showing that the conversion is significant.




Figure 2. (a). Emission spectra ($\lambda_{\text{ex}} = 284$ nm) of a solution of **I1** placed on a rough Au-coated steel substrate over time (blue traces) overlaid with reference peaks for **I1** (yellow trace) and an ex situ synthesized solution of **D1** (red trace). These traces are normalized to have the same peak height at 380 nm. (b). Emission spectra ($\lambda_{\text{ex}} = 340$ nm) showing a peak for **D1** that appears in solution left on a rough Au-coated steel substrate (blue trace) overlaid with reference peaks for **I1** (yellow trace) and **D1** (red trace). Also shown are data from a solution is left on a smooth Au-coated mica substrate (green trace) and with an added suspension of Au dust (purple trace). (c) Selected ion chromatogram (LC-MS) of the **I1** standard (red), the **D1** standard (blue), and the solution subject to STM-BJ measurements for 12 hours (green). Masses obtained from these spectra are noted in the legend and shown in SI Figure S4. (d) Molecular structures of aryl iodides studied here **I1-I4**.

To elucidate the role of the Au substrate in the homocoupling of the aryl iodides, we show, using *ex situ* fluorescence spectroscopy, that the coupling reaction does not proceed directly on a steel substrate without Au coating (SI Figure S3c,d), indicating that the Au catalyzes the reaction. Similarly, when the rough Au-coated substrate is replaced with an atomically flat Au-coated mica substrate, no dimer formation is observed over 12 hours by fluorescence spectroscopy (Figure 2b, green trace) or after 12 hours of measurement in the STM-BJ setup (SI Figure S3b). We note here that although the tip and Au-coated mica substrate are locally roughened during the STM-BJ measurement, the number of undercoordinated sites created is likely insufficient to generate a detectable amount of the product. Finally, we also observe dimer formation on an Au-coated glass substrate, which produces a rough gold layer also catalyzes the reaction (SI Figure S3b). Our experiments therefore indicate that a rough Au surface is required and while also eliminating the possibility that steel on its own is catalyzing the reaction. The rough surface thus provides the active catalytic sites, similar to surface-catalyzed reactions under UHV that happen primarily at facets or defects.²⁹⁻³¹ These results are further supported by analysis of solutions subjected to STM-BJ measurements via Liquid Chromatography Mass Spectrometry (LC-MS) spectra shown in Figure 2c. We find that the chromatogram shows a peak at 3.6 minutes when the product **D1** is present in the solution while we see a peak at 3.1 minutes when the reactant **I1** is present. The reactant and product are identified by their respective masses extracted from these chromatograms and shown in SI Figure S4.

We next evaluate the role of the solvent and ions in solution. Although the reaction proceeds in the polar solvent PC and in PC with an added electrolyte (Figure 2b), when a solution of **I1** in the nonpolar solvent trichlorobenzene is deposited on the Au substrate, no peak corresponding to **D1** is observed in either STM-BJ measurements or fluorescence spectroscopy

(SI Figure S5), demonstrating that the solvent polarity is critical. As shown by Dawlaty and co-workers,¹ molecules at the interface of a metal and solvent experience a reaction field whose magnitude is a function of the solvent dielectric constant. Specifically, they show that for a solvent with a dielectric constant over 50 (as is the case with PC), the interfacial fields can be as large as 2V/nm. Our experimental observations suggest that such interfacial electric fields are contributors to the catalysis of the Ullman coupling. We note that identical results are obtained when solutions are created with an added electrolyte (such as tetrabutyl ammonium hexafluorophosphate (TBAPF₆) at a 0.1 M concentration) indicating that added ions do not alter the interfacial fields significantly, consistent with previous measurements.³²

Finally, when the Au surface is replaced with Au dust (diameter: 500-850 nm) suspended in a reaction mixture which contains **I1** dissolved in PC with 0.1 M TBAPF₆, the fluorescence spectra (Figure 2b) indicate no dimer formation. Under these conditions, **I1** is exposed to a rough surface comparable in total surface area to the Au substrate, yet no coupling reaction is observed.

These results could indicate weaker reaction fields at the interface of a small metallic sphere; however, we cannot rule out other possibilities, including different surface morphologies, fewer undercoordinated sites or inhibited surface diffusion, adsorbed moieties that prevent the reactants from interacting with Au.

The observed homocoupling reaction is generalizable with respect to both the linker and molecular backbone. Four additional aryl iodides (Figure 2d) were studied under the same conditions as **I1**, and all appear to homocouple. The coupling of both the thiomethyl- and pyridyl-terminated biphenyl iodides (**I2** and **I3**, respectively) can be probed *in situ* through STM-BJ measurements because they are equipped with anchoring groups necessary for metallic contact. Similar to the dimerization of **I1**, when each aryl iodide is measured, a lower molecular

conductance peak with a longer molecular plateau appears and continues to grow over time (SI Figure S6). We verify the identity of these products by comparison of the 1D and 2D conductance histograms for **D2** and **D3** synthesized *ex situ* and measured independently (Figure S6). LC-MS spectra of the extracted solutions also confirms biaryl coupling (SI Figure S7). We note that while these specific iodides were selected for their aurophilic substituents, they demonstrate that both strongly electron withdrawing and electron donating groups are tolerated in this system. Furthermore, even the larger molecule **I4** homocouples on the rough Au surface despite its steric encumbrance, as verified by electronic absorption spectroscopy of the extracted solution (SI Figure S7). We note that this particular coupling cannot be probed using the STM-BJ because the large twist angle between adjacent anthracene units in the product decreases its conductance below the measurable limit of our instrument.³³⁻³⁴

We propose a general scheme for the coupling of aryl iodides in an interfacial field (Figure 3a). We probe this reaction scheme with mechanistic and computational studies described further below. Two aryl iodides undergo an oxidative addition forming two independent C–Au–I constructs at active sites on the rough Au substrate. If two oxidative addition events occur near one another, a small amount of surface migration may occur before a reversible reductive elimination can release the aryl iodides back into solution.³⁵⁻³⁷ Once the active Au–C bonds are close enough, a C–C bond can form between the activated aryl species, presumably through a mechanism that resembles that of thermally driven reactions on metal surfaces.^{16, 21, 38} The resulting tetraphenyl product is then released into solution, likely leaving iodine chemisorbed to the Au surface. We note that this double oxidative addition mechanism differs from the proposed mechanisms for solution-based Ullmann couplings in which the two halides have distinct roles in

the coupling process: one aryl iodide free in solution is activated by the metal catalyst, thus forming an organometallic complex with which the second free aryl iodide can interact.³⁹⁻⁴⁰




Figure 3. (a) Schematic of the key steps for the proposed Ullmann coupling mechanism on a rough Au surface: (1) binding on the surface, (2) oxidative addition, (3) surface migration and (4) dimerization. The yellow sphere denotes a gold adatom and the red sphere denotes the different substituents used in this study. (b) The DFT reaction energy over the course of the dehalogenation process, from reactant (reaction coordinate = 0) to transition state (TS) to product (reaction coordinate = 1) of iodobenzene. The calculations are performed on an Au(111) surface (green) and with an additional Au adatom (pink). The blue bars at the TS denote the change of the activation energy due to an applied electric field in the range of ± 3 V/nm (close-up shown in insets at right). Insets at left show the TS geometries and the black arrows indicate the direction of a positive electric field.

Within this reaction scheme, oxidative addition is proposed to be the key step catalyzed by the rough Au surface and the interfacial field. We can probe this step by comparing the reactivity of aryl iodides with that of a preformed Au complex. We synthesized the Au complex **Au2** (SI Figure S8a), which has a preformed Au–C bond and a labile PPh₃ ligand that readily dissociates *in situ*.⁴¹ This Au complex can be understood as an analog of the product of an oxidative addition of biphenyl iodide **I2** to Au, as the dissociation of PPh₃ replaces the oxidative addition step such

that starting with **Au2** instead of an aryl iodide bypasses the oxidative addition step. Indeed, when **Au2** is measured in the junction in a non-polar solvent, a molecular conductance peak for **D2** appears immediately (SI Figure S8b). Furthermore, when **Au2** is left on a smooth mica substrate in polar solvent, **D2** is observed in the solution via fluorescence spectroscopy (SI Figure S8c). Together, these results prove that when the oxidative addition step is removed from the reaction scheme altogether, neither a rough surface nor a polar solvent is necessary to the coupling. As such, we conclude that the oxidative addition is the rate limiting step in this transformation and is the key process that is assisted by the rough surface and its interfacial electric field.

For further insights into the oxidative addition process, we turn to electronic structure calculations using density functional theory (DFT); full computational details are provided in the SI. For simplicity, we study the dehalogenation of iodobenzene, whose behavior is representative of other aryl halides. We emphasize that because of the idealized nature of our calculations (especially the surface morphology and absence of explicit solvent), we do not expect quantitative agreement with experimental findings, but rather qualitative insights. Using the nudged elastic band method with 6-12 images, we calculated the minimum energy reaction pathway on an Au(111) surface and on an Au(111) with a single Au adatom (Figure 3b), as a model for surface roughness. On Au(111), we find an activation energy of 0.76 eV (17.5 kcal/mol) and a reaction energy of -0.4 eV (-9 kcal/mol), in good agreement with previous calculations.⁴² In contrast, in the presence of an Au adatom, which represents an extreme under-coordinated structure, we find a significantly reduced activation energy of 0.18 eV (3.5 kcal/mol) and more negative reaction energy of -0.6 eV (-13 kcal/mol).⁴³ The reduced activation energy supports the experimental observation that a rough Au surface, where a fraction of Au atoms on step-edges are undercoordinated (though not as undercoordinated as the adatom shown in Figure 3b) is essential

for catalyzing the Ullmann coupling. The difference in activation energies can be understood by comparing the transition state geometries (Figure 3b, inset). Compared to Au(111), the Au adatom is more readily inserted into the C–I bond.

Next, we analyze the potential impact of interfacial electric fields on the dehalogenation. In the direction perpendicular to the surface, we apply an electric field of -3 V/nm and $+3$ V/nm, which is comparable to the interfacial field strengths previously measured^{1, 44} and calculated⁴⁵ at metal-solvent interfaces. Over this range, we find that the electric field modifies the activation energy by about 0.05 – 0.1 eV (~ 0.5 – 2 kcal/mol). We see that a positive electric field (i.e. pointing up from the surface) decreases the activation energy on a pristine surface but increases the activation energy in the presence of an adatom modestly. This contrasting behavior is explained by a difference in the relative dipole moments of the reactant and transition state structures (all dipole moments are directed away from the surface and are thus stabilized by a field pointing in the same direction): on the pristine surface, the dipole moment of the reactant is smaller than that of the transition state, and in the presence of an adatom, the reverse is true. More discussion about this behavior and its implications for a linear free energy relationship can be found in Hoffman et al.⁴⁶ These results indicate that both surface roughness and local electric fields are important in determining the kinetics of surface-mediated oxidative addition. We expect an interfacial reaction field to point in the direction of the dipole moment, which is away from the surface.¹ Our calculation results show that a field in this direction could either raise or lower the activation energy, depending on the precise local morphology as indicated in Figure 3b.




Figure 4. (a). Schematic showing that cross-coupled product **CC1** forms from a mixture of **I1** and **B1** but not from a mixture of **I1** and **T1**. (b). Logarithmically binned 1D histograms showing the *in situ* cross-coupling of **I1** and **B1** (dark blue = 1000 traces, light blue = 100 traces), as confirmed by the alignment with the histogram of the product **CC1** (green) synthesized *ex situ*. Also shown in pink is the dimerization product **D1** also measured in the same experiment.

To demonstrate that the coupling reaction may involve two independent oxidative additions, we compare the reactivity of a bromide- and triflate-terminated biphenyl (**B1** and **T1**, respectively) (Figure 4a), which are less prone to oxidative addition than aryl iodides. Indeed, DFT calculations of the dehalogenation of bromobenzene in the presence of an Au adatom predict an activation energy of 0.5 eV (12 kcal/mol), which is much larger than that for iodobenzene (SI Figure S9). Consistent with our hypothesis of multiple oxidative additions to the Au surface, these less reactive reactants do not homocouple in the junction over the time scale studied. Although these reactants are unlikely to form two C–M–X constructs near each other, when surrounded by reactive iodides, only one such construct is necessary. Based on the hypothesis that two oxidative additions are necessary for aryl–aryl coupling, we suspect an Au-bound intermediate is more likely to be formed by a bromide than a triflate, the latter being less prone to oxidative addition, and thus **B1** is more likely to cross-couple with **I1** than **T1** (Figure 4a). We note that the methyl group *meta*

to the linker on **B1** and **T1** is used to distinguish between the cross-coupling product **CC1** and the homocoupling product **D1**.

When **B1** and **I1** are exposed to the reaction conditions described above, dimerization is observed. A molecular conductance peak for **D1** appears at 3.7×10^{-5} G₀ in the 1D histogram after 12 hours (Figure 4b, pink peak). After 18 hours, this peak becomes broader and asymmetric, with a shoulder on the left side (Figure 4b, dark blue trace). The first 100 traces after the appearance of the shoulder were isolated, and two distinct peaks were resolved at 3.7×10^{-5} G₀ and 1.9×10^{-5} G₀ (Figure 4b, light blue trace). A lower conductance peak is expected for the cross-coupled product, **CC1** (Figure 4a), due to the presence of the methyl group and consequently larger twist angle between the adjacent phenyl rings.³⁴ We therefore assign this new conductance peak to the putative product of the cross-coupling reaction, **CC1**. To further validate these results, we synthesized **CC1** ex situ and compared its conductance value (Figure 4b, green peak) with that of the molecule formed in situ. The alignment of the two peaks supports our hypothesis that the lower conducting peak at 1.9×10^{-5} G₀ can be assigned to the cross-coupled product (Figure 4b, green trace). Finally, when we repeat these experiments with **T1**, no such peak corresponding to **CC1** appears (SI Figure S10). The order of reactivity of aryl bromides and sulfonates is highly dependent on the mechanism of the reaction, and aryl bromides are more prone to oxidative addition than aryl sulfonates.⁴⁷ Although the results with **I1** alone do not exclude the possibility of the aryl iodide interacting with C-M-X to form **D1**, these control measurements indicate strongly that the coupling follows two oxidative additions. This is because **T1** should be as reactive (if not more reactive) towards that reaction **T1** + C-M-X = **CC1** than the aryl iodide. Yet, we do not observe any **CC1** product in our mixed measurements with **T1** and **I1**. By contrast, **B1** is less reactive towards the same reaction (**B1** + C-M-X = **CC1**), yet we see the **CC1** product in the mixed measurements of **I1** and **B1**.

However, the reactivity of two oxidative additions forming the product (i.e. two C-M-X forming **D1** or **CC1**) would be highest for **I1** and lowest for **T1** with **B1** having an intermediate reactivity. Our experimental results suggest that in this system, the bromide is more reactive than the triflate, although it is possible that the cross-coupling product with **T1** was simply never observed in the six measurements taken. The order of reactivity observed in the cross-coupling experiments is therefore consistent with our hypothesis that two independent oxidative addition steps are involved in the formation of aryl-aryl coupling products.

In this report, we have shown that the Ullmann-type coupling of aryl iodides can be catalyzed by a rough Au surface submerged in a polar solvent under ambient conditions. Through a combination of *in situ* STM-BJ measurements, *ex situ* mass spectrometry, fluorescence spectroscopy, and first-principles calculations, we have gained mechanistic insight into these transformations that occur by a double oxidative-addition mechanism. Taken together, this proof-of-principle study also suggests that STM-BJ may provide a valuable method to interrogate surface-based reactivity in real-time. More broadly, this investigation connects interfacial electric fields to chemical reactivity and reiterates the promise of electric-field catalysis that could be leveraged for a wide number of molecular transformations.

Acknowledgments: This work is supported primarily by NSF CHE-2023568 CCI Phase I: Center for Chemistry with Electric Fields. A.M.E. is supported by the Schmidt Science Fellows, in partnership with the Rhodes Trust. I.B.S. was supported in part by NIH R35 GM127135. We thank Lisa Wei for Fluorescence measurements taken on an instrument provided by NSF Award 1828491. We thank Ren Wiscons and Boyuan Zhang for SEM and AFM images shown in the SI and Claudia Prindle for help with the Mass-Spectra. The Flatiron Institute is a division of the Simons Foundation.

References:

- (1) Sorenson, S. A.; Patrow, J. G.; Dawlaty, J. M., Solvation Reaction Field at the Interface Measured by Vibrational Sum Frequency Generation Spectroscopy, *J. Am. Chem. Soc.* **2017**, *139*, 2369-2378.
- (2) Oklejas, V.; Sjostrom, C.; Harris, J. M., SERS Detection of the Vibrational Stark Effect from Nitrile-Terminated SAMs to Probe Electric Fields in the Diffuse Double-Layer, *J. Am. Chem. Soc.* **2002**, *124*, 2408-2409.
- (3) Delley, M. F.; Nichols, E. M.; Mayer, J. M., Interfacial Acid–Base Equilibria and Electric Fields Concurrently Probed by In Situ Surface-Enhanced Infrared Spectroscopy, *J. Am. Chem. Soc.* **2021**, *143*, 10778-10792.
- (4) Steinmann, S. N.; Seh, Z. W., Understanding electrified interfaces, *Nat. Rev. Mat.* **2021**, *6*, 289-291.
- (5) Dutta Dubey, K.; Stuyver, T.; Kalita, S.; Shaik, S., Solvent Organization and Rate Regulation of a Menshutkin Reaction by Oriented External Electric Fields are Revealed by Combined MD and QM/MM Calculations, *J. Am. Chem. Soc.* **2020**, *142*, 9955-9965.
- (6) Smith, A. M.; Lee, A. A.; Perkin, S., The Electrostatic Screening Length in Concentrated Electrolytes Increases with Concentration, *The Journal of Physical Chemistry Letters* **2016**, *7*, 2157-2163.
- (7) Wesley, T. S.; Román-Leshkov, Y.; Surendranath, Y., Spontaneous Electric Fields Play a Key Role in Thermochemical Catalysis at Metal–Liquid Interfaces, *ACS Central Science* **2021**, *7*, 1045-1055.
- (8) Ryu, J.; Surendranath, Y., Tracking Electrical Fields at the Pt/H₂O Interface during Hydrogen Catalysis, *J. Am. Chem. Soc.* **2019**, *141*, 15524-15531.

(9) Voegtle, M. J.; Pal, T.; Pennathur, A. K.; Menachekanian, S.; Patrow, J. G.; Sarkar, S.; Cui, Q.; Dawlaty, J. M., Interfacial Polarization and Ionic Structure at the Ionic Liquid–Metal Interface Studied by Vibrational Spectroscopy and Molecular Dynamics Simulations, *The Journal of Physical Chemistry B* **2021**, *125*, 2741-2753.

(10) Ullmann, F.; Bielecki, J., Ueber Synthesen in der Biphenylreihe, *Berichte der deutschen chemischen Gesellschaft* **1901**, *34*, 2174-2185.

(11) Lin, H.; Sun, D., Recent Synthetic Developments and Applications of the Ullmann Reaction. A Review, *Organic Preparations and Procedures International* **2013**, *45*, 341-394.

(12) Grill, L.; Dyer, M.; Lafferentz, L.; Persson, M.; Peters, M. V.; Hecht, S., Nano-architectures by covalent assembly of molecular building blocks, *Nat. Nanotechnol.* **2007**, *2*, 687-691.

(13) Cai, J.; Ruffieux, P.; Jaafar, R.; Bieri, M.; Braun, T.; Blankenburg, S.; Muoth, M.; Seitsonen, A. P.; Saleh, M.; Feng, X.; Mullen, K.; Fasel, R., Atomically precise bottom-up fabrication of graphene nanoribbons, *Nature* **2010**, *466*, 470-3.

(14) Lafferentz, L.; Eberhardt, V.; Dri, C.; Africh, C.; Comelli, G.; Esch, F.; Hecht, S.; Grill, L., Controlling on-surface polymerization by hierarchical and substrate-directed growth, *Nat. Chem.* **2012**, *4*, 215-220.

(15) Clair, S.; de Oteyza, D. G., Controlling a Chemical Coupling Reaction on a Surface: Tools and Strategies for On-Surface Synthesis, *Chem. Rev.* **2019**, *119*, 4717-4776.

(16) Xi, M.; Bent, B. E., Mechanisms of the Ullmann Coupling Reaction in Adsorbed Monolayers, *J. Am. Chem. Soc.* **1993**, *115*, 7426-7433.

(17) Batra, A.; Cvetko, D.; Kladnik, G.; Adak, O.; Cardoso, C.; Ferretti, A.; Prezzi, D.; Molinari, E.; Morgante, A.; Venkataraman, L., Probing the mechanism for graphene nanoribbon formation on gold surfaces through X-ray spectroscopy, *Chem. Sci.* **2014**, *5*, 4419.

(18) Grill, L.; Hecht, S., Covalent on-surface polymerization, *Nat. Chem.* **2020**, *12*, 115-130.

(19) Kamenetska, M.; Koentopp, M.; Whalley, A. C.; Park, Y. S.; Steigerwald, M. L.; Nuckolls, C.; Hybertsen, M. S.; Venkataraman, L., Formation and Evolution of Single-Molecule Junctions, *Phys. Rev. Lett.* **2009**, *102*, 126803.

(20) Lewis, E. A.; Murphy, C. J.; Pronschinske, A.; Liriano, M. L.; Sykes, E. C. H., Nanoscale insight into C–C coupling on cobalt nanoparticles, *Chemical Communications* **2014**, *50*, 10035-10037.

(21) Dong, L.; Liu, P. N.; Lin, N., Surface-Activated Coupling Reactions Confined on a Surface, *Acc. Chem. Res.* **2015**, *48*, 2765-2774.

(22) Wang, W.; Shi, X.; Wang, S.; Van Hove, M. A.; Lin, N., Single-Molecule Resolution of an Organometallic Intermediate in a Surface-Supported Ullmann Coupling Reaction, *J. Am. Chem. Soc.* **2011**, *133*, 13264-13267.

(23) Monopoli, A.; Cotugno, P.; Palazzo, G.; Ditaranto, N.; Mariano, B.; Cioffi, N.; Ciminale, F.; Nacci, A., Ullmann Homocoupling Catalysed by Gold Nanoparticles in Water and Ionic Liquid, *Advanced Synthesis & Catalysis* **2012**, *354*, 2777-2788.

(24) Li, G.; Liu, C.; Lei, Y.; Jin, R., Au25 nanocluster-catalyzed Ullmann-type homocoupling reaction of aryl iodides, *Chemical Communications* **2012**, *48*, 12005-12007.

(25) Payne, R.; Theodorou, I. E., Dielectric properties and relaxation in ethylene carbonate and propylene carbonate, *The Journal of Physical Chemistry* **1972**, *76*, 2892-2900.

(26) Nagahara, L. A.; Thundat, T.; Lindsay, S. M., Preparation and Characterization of STM Tips for Electrochemical Studies, *Rev. Sci. Instrum.* **1989**, *60*, 3128-3130.

(27) Zang, Y.; Pinkard, A.; Liu, Z.-F.; Neaton, J. B.; Steigerwald, M. L.; Roy, X.; Venkataraman, L., Electronically transparent Au-N bonds for molecular junctions, *J. Am. Chem. Soc.* **2017**, *139*, 14845-14848.

(28) Zang, Y.; Zou, Q.; Fu, T.; Ng, F.; Fowler, B.; Yang, J.; Li, H.; Steigerwald, M. L.; Nuckolls, C.; Venkataraman, L., Directing isomerization reactions of cumulenes with electric fields, *Nat. Commun.* **2019**, *10*, 4482.

(29) Hla, S.-W.; Bartels, L.; Meyer, G.; Rieder, K.-H., Inducing All Steps of a Chemical Reaction with the Scanning Tunneling Microscope Tip: Towards Single Molecule Engineering, *Phys. Rev. Lett.* **2000**, *85*, 2777-2780.

(30) McCarty, G. S.; Weiss, P. S., Footprints of a Surface Chemical Reaction: Dissociative Chemisorption of p-Diiodobenzene on Cu{111}†, *The Journal of Physical Chemistry B* **2002**, *106*, 8005-8008.

(31) Weiss, P. S.; Kamna, M. M.; Graham, T. M.; Stranick, S. J., Imaging Benzene Molecules and Phenyl Radicals on Cu{111}, *Langmuir* **1998**, *14*, 1284-1289.

(32) Capozzi, B.; Chen, Q.; Darancet, P.; Kotiuga, M.; Buzzeo, M.; Neaton, J. B.; Nuckolls, C.; Venkataraman, L., Tunable Charge Transport in Single-Molecule Junctions via Electrolytic Gating, *Nano Lett.* **2014**, *14*, 1400-1404.

(33) Becker, H. D.; Langer, V.; Sieler, J.; Becker, H. C., Molecular conformations of 9,9'-bianthryl, di-9-anthrylmethane, and some related twisted anthracene derivatives, *The Journal of Organic Chemistry* **1992**, *57*, 1883-1887.

(34) Venkataraman, L.; Klare, J. E.; Nuckolls, C.; Hybertsen, M. S.; Steigerwald, M. L., Dependence of single-molecule junction conductance on molecular conformation, *Nature* **2006**, *442*, 904-907.

(35) Joost, M.; Amgoune, A.; Bourissou, D., Reactivity of Gold Complexes towards Elementary Organometallic Reactions, *Angew. Chem. Int. Ed.* **2015**, *54*, 15022-15045.

(36) Huang, B.; Hu, M.; Toste, F. D., Homogeneous Gold Redox Chemistry: Organometallics, Catalysis, and Beyond, *Trends in Chemistry* **2020**, *2*, 707-720.

(37) Rocchigiani, L.; Bochmann, M., Recent Advances in Gold(III) Chemistry: Structure, Bonding, Reactivity, and Role in Homogeneous Catalysis, *Chem. Rev.* **2021**, *121*, 8364-8451.

(38) Lackinger, M., Surface-assisted Ullmann coupling, *Chemical Communications* **2017**, *53*, 7872-7885.

(39) Ullmann, F., Ueber eine neue Bildungsweise von Diphenylaminderivaten, *Berichte der deutschen chemischen Gesellschaft* **1903**, *36*, 2382-2384.

(40) Ullmann, F., Ueber symmetrische Biphenylderivate, *Justus Liebigs Annalen der Chemie* **1904**, *332*, 38-81.

(41) Cheng, Z. L.; Skouta, R.; Vazquez, H.; Widawsky, J. R.; Schneebeli, S.; Chen, W.; Hybertsen, M. S.; Breslow, R.; Venkataraman, L., In situ formation of highly conducting covalent Au-C contacts for single-molecule junctions, *Nat. Nanotechnol.* **2011**, *6*, 353-357.

(42) Bjork, J.; Hanke, F.; Stafstrom, S., Mechanisms of Halogen-Based Covalent Self-Assembly on Metal Surfaces, *J. Am. Chem. Soc.* **2013**, *135*, 5768-5775.

(43) Corma, A.; Juárez, R.; Boronat, M.; Sánchez, F.; Iglesias, M.; García, H., Gold catalyzes the Sonogashira coupling reaction without the requirement of palladium impurities, *Chemical Communications* **2011**, *47*, 1446-1448.

(44) Shi, H.; Cai, Z.; Patrow, J.; Zhao, B.; Wang, Y.; Wang, Y.; Benderskii, A.; Dawlaty, J.; Cronin, S. B., Monitoring Local Electric Fields at Electrode Surfaces Using Surface Enhanced

Raman Scattering-Based Stark-Shift Spectroscopy during Hydrogen Evolution Reactions, *ACS Applied Materials & Interfaces* **2018**, *10*, 33678-33683.

(45) Goldsmith, Z. K.; Secor, M.; Hammes-Schiffer, S., Inhomogeneity of Interfacial Electric Fields at Vibrational Probes on Electrode Surfaces, *ACS Central Science* **2020**, *6*, 304-311.

(46) Hoffmann, N. M.; Wang, X.; Berkelbach, T. C., Linear Free Energy Relationships in Electrostatic Catalysis, *ACS Catalysis* **2022**, 8237-8241.

(47) McMillen, D. F.; Golden, D. M., Hydrocarbon bond dissociation energies, *Annu. Rev. Phys. Chem.* **1982**, *33*, 493-532.