


Neuron

Mapping of neuronal and glial primary cilia contactome and connectome in the human cerebral cortex

Graphical abstract



Authors

Jun Yao Wu, Su-Ji Cho,
Katherine Descant, ..., Ryohei Yasuda,
Jeff W. Lichtman, E.S. Anton

Correspondence

ryohei.yasuda@mpfi.org (R.Y.),
jeff@mcb.harvard.edu (J.W.L.),
anton@med.unc.edu (E.S.A.)

In brief

The unique pattern of primary cilia contacts with the surrounding cortical circuitry may enable cilia signaling to serve as a mechanism through which local environmental signals can shape and refine neuronal circuits. Disruptions in the primary cilia connectome may thus contribute to circuit dysfunction in ciliopathies and other human brain disorders.

Highlights

- Cortical neuronal and astroglial cilia are structurally and connectomically diverse
- Unlike neuronal cilia, astrocyte cilia are mostly in pockets or embedded inside soma
- The contactome of each cilium enables unique access to surrounding neural circuitry
- Human neuronal primary cilia are components of tetrapartite synapses



NeuroResource

Mapping of neuronal and glial primary cilia contactome and connectome in the human cerebral cortex

Jun Yao Wu,^{1,7} Su-Ji Cho,^{1,7} Katherine Descant,^{1,7} Peter H. Li,² Alexander Shapson-Coe,⁶ Michal Januszewski,² Daniel R. Berger,⁶ Cailyn Meyer,¹ Cristine Casingal,¹ Ariba Huda,¹ Jiaqi Liu,¹ Tina Ghashghaei,¹ Mikayla Brenman,¹ Michelle Jiang,¹ Joseph Scarborough,¹ Art Pope,² Viren Jain,² Jason L. Stein,³ Jiami Guo,⁴ Ryohei Yasuda,^{5,*} Jeff W. Lichtman,^{6,*} and E.S. Anton^{1,8,*}

¹UNC Neuroscience Center and the Department of Cell Biology and Physiology, University of North Carolina School of Medicine, Chapel Hill, NC 27599, USA

²Google Research, Mountain View, CA 94043, USA

³UNC Neuroscience Center and the Department of Genetics, University of North Carolina School of Medicine, Chapel Hill, NC 27599, USA

⁴Department of Cell Biology and Anatomy, Hotchkiss Brain Institute, University of Calgary, Calgary, AB, Canada

⁵Max Planck Florida Institute for Neuroscience, Jupiter, FL 33458, USA

⁶Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA 02138, USA

⁷These authors contributed equally

⁸Lead contact

*Correspondence: ryohei.yasuda@mpfi.org (R.Y.), jeff@mcb.harvard.edu (J.W.L.), anton@med.unc.edu (E.S.A.)

<https://doi.org/10.1016/j.neuron.2023.09.032>

SUMMARY

Primary cilia act as antenna receivers of environmental signals and enable effective neuronal or glial responses. Disruption of their function is associated with circuit disorders. To understand the signals these cilia receive, we comprehensively mapped cilia's contacts within the human cortical connectome using serial-section EM reconstruction of a 1 mm³ cortical volume, spanning the entire cortical thickness. We mapped the "contactome" of cilia emerging from neurons and astrocytes in every cortical layer. Depending on the layer and cell type, cilia make distinct patterns of contact. Primary cilia display cell-type- and layer-specific variations in size, shape, and microtubule axoneme core, which may affect their signaling competencies. Neuronal cilia are intrinsic components of a subset of cortical synapses and thus a part of the connectome. This diversity in the structure, contactome, and connectome of primary cilia endows each neuron or glial cell with a unique barcode of access to the surrounding neural circuitry.

INTRODUCTION

The primary cilium, a microtubule (MT)-based antennae-like organelle, is present in cerebral cortical neurons and, when defective, leads to clinically significant disorders known generally as ciliopathies. A distinguishing feature of ciliopathies is disruptions in neural circuit formation and function. These disruptions are associated with a range of disorders including autism, intellectual disabilities, mood disorders, obesity, and epilepsy.^{1–30} Correspondingly, the expression of genes encoding cilia-associated proteins is significantly dysregulated in sets of patients with major psychiatric disorders including schizophrenia (SCZ), autism spectrum disorder, bipolar disorder, and major depressive disorder.³¹ Further, copy-number variation of the complement component 4A gene confers significant SCZ risk in humans, and disruption of primary cilia-related processes in excitatory (E) neurons is a key contributor to this risk.³² Likewise, disruption of ciliary function in interneurons (INs) and E

neurons can lead to altered synaptic connectivity and E/inhibitory (I) balance.^{5,33,34} Recent studies also reveal that novel axo-ciliary synapses can regulate the epigenetic state of post-synaptic neurons.³⁵ Collectively, these studies imply a role for primary cilia in neuronal function and circuit dynamics.

However, how primary cilia of diverse neurons and glia in the human cortex are organized, how they interact with adjacent cells, and how they are incorporated into the brain connectome are unknown. Primary cilia signaling may serve as a non-traditional synaptic signaling mechanism through which local environmental signals can shape and refine neuronal circuits in health and disease. Nonetheless, their signaling partners are poorly defined. We, therefore, sought to precisely map primary cilia interactions within the human cortical connectome.

We examined the distinct ultrastructural features of neuronal (both E and I) and astroglial primary cilia in every layer of the human cerebral cortex and their entire interactions with their surrounding cellular milieu using the human brain serial-section




Figure 1. Examination of primary cilia in projection neurons, interneurons, and astrocytes of every cerebral cortical layer

Primary cilium of interneurons (A), projection neurons (B), and astrocytes (C) from all different layers of the cerebral cortex were analyzed. Neurons and glia examined are shown. I–VI, different cortical layers; WM, white matter.

electron microscope (EM) connectome.³⁶ Our analysis reveals neuronal- and glial-type- and layer-specific diversity in primary cilia characteristics and their contactome. These characteristics may subserve their functions in cortical circuitry. This comprehensive, ultrastructural map of the nature and interactions of neuronal and astroglial primary cilia within the human cerebral cortex supports the functional significance of primary cilia in the ongoing maintenance and function of cortical neuronal circuitry in humans. Neuronal primary cilium, in addition to functioning as an antenna, sensing neuromodulators in the local environment, also influences neuronal functions via specialized contacts it makes with synapses and adjacent cells. We find that neuronal primary cilia are an intrinsic component of synaptic structure in a subset of cortical synapses. Primary cilia thus help create small, local modulatory signaling ensembles within larger neuronal circuits.

RESULTS

Organization of neuronal and glial primary cilia in the cerebral cortex

To evaluate the primary cilia contactome and connectome of the human cortical neurons, we first mapped the primary cilium of INs and projection neurons (PNs) in all 6 layers of the cerebral cortex (Figures 1, 2, and 3; Table S1; Videos S1 and S2A). Primary cilia occupy 0.03% of cortical volume (Figure S1A). The total cellular density or volume of layers does not correlate with ciliary volume (Figures S1B–S1D). We then analyzed the length, orientation, morphology, and intra-ciliary structural features of neuronal primary cilia. In both the PNs and INs, upper-layer primary cilia were significantly longer than the lower-layer ones (layer 1 IN: $6.3 \pm 0.3 \mu\text{m}$, layer 6 IN: $4.9 \pm 0.4 \mu\text{m}$; layer 2 PN: $8.4 \pm 0.2 \mu\text{m}$, layer 6 PN: $5.5 \pm 0.2 \mu\text{m}$; Figures 4A–4D). The average length of IN and PN cilia are 5.8 ± 0.2 and $7.4 \pm 0.6 \mu\text{m}$, respectively. The majority of primary cilia are oriented

into the plane of the cortical wall (Figures 4E and 4F) and are located in closer proximity to the neuron's dendrites than its axon (Figures 4G and 4H). Distinct neuron-specific differences are evident in cilia shape. Compared with the smoother, tubular primary cilia of PNs, IN cilia tend to be more complex, with a convoluted or beaded ciliary membrane sheath (Figures 4I–4K). These outpocketings of the ciliary membrane (arrow, Figure 4I) are reminiscent of budding ciliary ectosomes, the signaling vesicles released from cilia.³⁷

We next analyzed whether the cilia of INs and PNs in different layers have distinguishing membrane specializations and cytoskeletal organization. Ciliary pockets are indicative of active vesicle transport and docking at the base of the cilium.^{38,39} In general, depending on the cortical layer and neuronal subtype, 24%–56% of neuronal primary cilia have a ciliary pocket membrane specialization at their base (arrow, Figures 4L–4N). Furthermore, basal body organization is thought to be a key modulator of ciliary membrane extension and MT axoneme orientation and stability.^{40,41} A subset of cortical neurons in all layers display an irregular basal body structure (arrow, Figures 4O–4Q), instead of the well-organized mother centriole-based, MT-organizing basal body at the base (arrowhead, Figure 4O). Layer 4 INs and layer 2 PNs have the highest percentage of irregular basal bodies (26% and 38%, respectively).

The MT core of neuronal primary cilia dynamically varies across neuronal types. In cortical PNs, MT filaments extend up to 70%–86% of the ciliary length, with the layer 4 PN cilia displaying the shortest MT core extension ($70\% \pm 3.58\%$ of the ciliary length; Figures 4R–4T). Similarly, MT filaments extend up to 61%–88% of IN cilia length, with layer 1 IN cilia exhibiting the shortest MT core extension ($61\% \pm 3.12\%$ of the ciliary length; Figures 4R–4T). The neuronal primary cilia MT architecture is highly diverse in contrast to the commonly observed $9 + 0$ MT arrangement in primary cilia elsewhere.²⁸ The MT axoneme of




Figure 2. Cortical interneuronal primary cilia

Sample images of the primary cilium (arrow) of human MT core units of human cortical interneurons from all six cortical layers (A–F). Cells are oriented so that axons are pointing downward. Scale bars, 1.6 μ m. I–VI, cortical layers.

neuronal cilia gradually changes from base to tip with devolving numbers of MT core units. We measured MT core units at the base, middle, and tip of neuronal primary cilia. A rich variety in MT axoneme architecture across different IN and PN populations is evident (Figures 4U–4W). Importantly, in the majority of cortical neurons of all subtypes, the 9 + 0 MT core orga-

nization is not present at the midpoint of the ciliary length; instead, 7 or fewer MT core filaments are present at the midpoint (Figures 4U–5W; Videos S3 and S4).

Occasionally, we noticed oddities in neuronal ciliary organization, such as a cilium emanating from a major dendrite at a significant distance away from the soma (Figures S2A and




Figure 3. Cortical projection neuronal primary cilia

Sample images of the primary cilium (arrow) of human cortical projection neurons from cortical layers II–VI (A–E). There are no projection neurons in layer I. Cells are oriented so that apical dendrites are pointing upward, and axons are pointing downward. Scale bars, 1.85 μm . II–VI, cortical layers.

S2B) or an IN with two basal body structures and two cilia (Figures S2C and S2D). We did not detect a primary cilium only in 0.48% (9/1,890) of neurons, all of which were INs.

We then similarly mapped primary cilia of astrocytes in all 6 layers of the cerebral cortex (Table S2). Astrocyte cilia have features that are rather distinct from neuronal cilia. In contrast to neurons, astrocyte cilia are often embedded in deep ciliary pockets (greater than 50% of the ciliary axoneme length is inside a membrane pocket in $78\% \pm 3\%$ of cortical astrocytes). The average length of the astrocyte cilium is $3.97 \pm 0.09 \mu\text{m}$, shorter than neuronal cilia (IN_{cilia} : $5.8 \pm 0.2 \mu\text{m}$, PN_{cilia} : $7.4 \pm 0.6 \mu\text{m}$; Figure S3A). The majority of them are oriented into the

cortical wall (Figures S3B and S3C) and are of smooth tubular shape (Figure S3D). $\sim 40\%$ or more of astrocytes in every cortical layer display an irregular basal body structure (Figure S3E). Unlike neurons, in $10.58\% \pm 2\%$ of cortical astrocytes, the ciliary axoneme was found to be still within the ciliary sheath embedded inside the astrocyte cell soma (Figures S3F–S3H; Video S5). These astrocytes do not have externally visible cilia. We did not detect such cilia in any neurons. A $9 + 0$ MT structure is mostly absent at the midpoint of astrocyte cilium (Figures S3I–S3K). The majority of them contain fewer than 4 MT core units at the midpoint (Figures S3I–S3K). No ciliary structures were detected in mature myelinating oligodendrocytes.




Figure 4. Diversity in the structure and organization of neuronal primary cilia

(A–D) Diversity in the length of neuronal primary cilia. Quantification of the average length of interneuronal (A) and projection neuronal (B) primary cilia. (C) Sample images of interneuronal primary cilia from layers I and VI. (D) Sample images of projection neuronal primary cilia from layers II and VI. Data shown are mean $\pm$ SEM. One-way ANOVA: IN, $F_{5, 286}[\text{cilia length}] = 3.33$, $p = 0.0061$; PN, $F_{4, 303}[\text{cilia length}] = 29.85$, $p < 1E-16$. Two-way ANOVA: $F_{1, 540}[\text{cilia length, IN vs. PN}] = 86.24$, $p < 1E-16$. Scale bars: 2 μm in (C) and (D). IN, interneuron; PN, projection neuron; I–VI, cortical layers.

(legend continued on next page)

Neuronal and glial primary cilia contactome in the cerebral cortex

To explore how neuronal primary cilia interact with the surrounding cells in the human cerebral cortex, we mapped every cell that came in membrane contact with the primary cilium of INs and PNs in all 6 layers of the cerebral cortex (Figures 5 and 6; Table S3; Videos S2B, S6, S7, S8, and S9). We first analyzed the cell type, cell domain, and the layer position of these ciliary contacts. On average, an individual IN primary cilium is contacted by 41 ± 0.4 other cells, whereas 47 ± 0.5 cells contact a PN cilium ($p < 0.0001$, t test). Only rarely does a neuronal cilium contact the same cell twice or more. IN cilia, across layers, make a comparably similar number of contacts, whereas upper-layer PN cilia make a significantly higher number of contacts than lower-layer PN cilia (layer 2 PN cilia: 57 ± 1.6 , layer 6 PN cilia: 39 ± 0.7 , $p < 0.0001$ [t test]; Figures 5, 6, S4A, and S4B). The length of a neuronal cilium positively correlates with the number of contacts it makes with other neural cells in the cortex (r , 0.5974, $p = 1E-14$).

In general, PN cilia make significantly more contact with other PNs, whereas IN cilia make contact with both INs and PNs (PN cilia contacts [#PNs/#INs]: 6.86 ± 0.37 , IN cilia contacts [#PNs/#INs]: 1.97 ± 0.08 , $p < 0.0001$ [t test]; Figures S4A–S4D; Table S4). This pattern is seen in all cortical layers even though the ratio of PN to IN numbers varies layer by layer (0.0068 [L1], 1.83 [L2], 1.73 [L3], 3.80 [L4], 3.46 [L5], and 6.29 [L6]).³⁶

Cortical neuronal cilia make substantially more membrane contacts with axons than dendrites (~ 10 -fold higher). Compared with IN cilia, PN cilia predominantly contact other E axons than I axons (PN cilia contacts [E/I axons]: 6.81 ± 0.37 vs. IN cilia contacts [E/I axons]: 1.84 ± 0.09 , $p < 0.0001$ [t test]; Figures S4G–S4J; Table S5). In contrast, both IN and PN cilia make significantly more contact with E, spiny dendrites than with I dendrites (IN cilia: 0.27 ± 0.03 [I] vs. 2.04 ± 0.10 [spiny]; PN cilia: 0.26 ± 0.03 [I] vs. 1.93 ± 0.10 [spiny]; $p < 0.0001$ [t test]; Figures S4K and S4L).

PNs within the six-layered human cortex are hierarchically organized based on their projection patterns.⁴² The intratelence-

phalic, upper-layer neurons (2–3) project to other domains of the cerebral hemisphere and striatum, whereas the extratelencephalic, deeper-layer neurons (5–6) project to thalamus and subcortical targets. The ratio of I neurons contacting the subcortically projecting, deeper-layer (L5–L6) PNs' cilia is significantly higher than the intracortically projecting, upper-layer (L2–L3) PNs' cilia (E/I: $4.04 \pm 0.46_{L5-L6}$ vs. $7.53 \pm 1.71_{L2-L3}$, $p < 0.0001$ [t test]). Similar to PNs, cortical IN diversity can be further distinguished based on molecular, morphological, connectivity, and activity patterns.⁴³ Morphologically, cortical INs can be broadly divided into basket cells, bipolar cells, neurogliaform cells, Martinotti cells, and chandelier cells. Primary cilia of chandelier cells, which make unique axo-axonic synapses on the axon initial segments (AISs) of cortical PNs,⁴⁴ contact 1.6-fold more cells and almost twice as many PNs as compared with primary cilia of other cortical INs ($p < 0.01$, Tukey's multiple comparison test; Figures S5A–S5D).

Compared with neuronal cilia-neuron contacts, neuronal cilia-glial contacts are ~ 15 -fold less frequent (Figures S4A and S4B). Both IN and PN cilia are contacted primarily by astrocytes than other glial cell types (i.e., microglia and oligodendroglia). However, PN cilia make significantly more glial contacts ($+22.3\% \pm 2.7\%$) than IN cilia ($p < 0.0001$, t test; Figures S4A, S4B, S4E, and S4F).

To clarify the distinctiveness of neuronal ciliary contacts with the surrounding cortical cellular milieu, we mapped the cellular contacts of adjacent axons of the same phenotype (i.e., E or I) and length as the respective neuronal cilia (Figures S6A, S6B, S6D, and S6E; Table S6). The patterns of cellular contacts of these axonal domains are significantly different and did not form the same pattern, based on layers or neuronal types, noticed for neuronal cilia (Figures S6A, S6B, S6D, and S6E; Tables S4 and S5), thus indicating the specificity of the neuronal ciliary contactome. The distinctiveness of the cellular contacts made by a cilia-sized domain of the respective neuronal AIS further supports this specificity (Figures S6C and S6F; Tables S7, S8A, and S8B).


(E and F) Quantification of neuronal primary cilia orientation. Orientation of neuronal primary cilia toward the pial surface, white matter, or into the cortical wall was quantified (E and F). Sample images of cilia orientation are shown in inset (top left, F). (E') and (F') show the layer location of interneurons (E') and projection neurons (F') with different cilia orientations. Cell soma of these neurons are highlighted in colors corresponding to different cilia orientation. One-way ANOVA (orientation): IN, $F_{2, 15} = 7.26$, $p = 0.0062$; PN, $F_{2, 12} = 5.72$, $p = 0.018$.

(G and H) The position of neuronal primary cilia relative to the position of the dendrites, axon, or cell soma was examined in interneurons and projection neurons. For projection neurons, the apical dendrite was used as the landmark, whereas any dendrite was used for interneurons. Sample images of cilia (arrow) position are shown in inset (top left, H). I–VI, cortical layers. One-way ANOVA (position): IN, $F_{3, 20} = 14.67$, $p = 0.0000278$; PN, $F_{3, 16} = 12.66$, $p = 0.00017$.

(I–K) Diversity of cortical neuronal primary cilia shape. (I) Sample cilia images (EM and 3D reconstruction) of each category of shape (smooth, beaded, mixed). Arrows point to ciliary membrane outpocketings. (J and K) Quantification of neuronal cilia shape. Shape of interneuronal cilia and projection neuronal cilia from different layers were quantified. IN, interneuron; PN, projection neuron; I–VI, cortical layers. Two-way ANOVA: $F_{2, 27[\text{cilia type}]} = 20.22$, $p = 4.3E-6$. Post hoc $p_{[\text{beaded cilia, IN vs. PN}]} = 0.0064$. Scale bars, 1.5 μm .

(L–Q) Organization of neuronal primary cilia. Quantification of cilia extending from a ciliary pocket at the base (L–N) and basal body organization (normal or irregular) of cilia (O–Q). Sample cilia images of each category are shown in (L) (normal or ciliary pocket [arrow]) and (O) (normal [arrowhead] or irregular-shaped basal body [arrow]). Interneuronal cilia and projection neuronal cilia from different layers were quantified (M, N, P, and Q). IN, interneuron; PN, projection neuron; I–VI, cortical layers. One-way ANOVA (cilia pocket organization): IN, $F_{1, 5} = 16.68$, $p = 0.002$; PN, $F_{1, 4} = 25.52$, $p = 0.00098$. One-way ANOVA (basal body organization): IN, $F_{1, 5} = 19.28$, $p = 0.0014$; PN, $F_{1, 4} = 29.04$, $p = 0.00065$. Scale bars: 1 μm in (L) and (O).

(R–W) Changing dynamics of axoneme MT core of neuronal cilia. Sample images of a cilium with MT filaments extending through the majority of its length (arrow, R [left]) and a cilium with MT filaments extending only midway through its length (arrow, R [right]). Quantification of the extension of MT filaments within interneuronal (S) or projection neuronal (T) cilia from different layers. (U) Sample images of cross sections from the base, middle, and tip of a neuronal cilium. Cross sections containing 9–8, 7–4, and 3 or fewer MT doublet filaments are classified as type 1, 2, and 3, respectively. Quantification of the MT type at the midpoint of interneuronal (V) or projection neuronal (W) cilia from different layers. IN, interneuron; PN, projection neuron; I–VI, cortical layers. Data shown are mean $\pm$ SEM. Two-way ANOVA: $F_{2, 27[\text{MT organization-midpoint}]} = 254.9$, $p < 1E-16$, post hoc $p_{[\text{IN vs. PN}]} = 0.03$. Scale bars: 1.4 μm in (R) and 0.7 μm in (U).



(legend on next page)

To further explore the distinct nature of the neuronal cilia contactome, we mapped the primary cilia contactome of the astrocytes in every cortical layer (Figures 1C and 7; Videos S10 and S11). Astrocytes intimately associate with neurons and modulate their function.⁴⁵ In contrast to neurons, the majority of astrocyte cilia are embedded in deep ciliary pockets. Astrocyte cilia make a significantly fewer number of contacts compared with neurons (7.6 ± 0.12 [vs. IN_{cilia} : 41 ± 0.4 or PN_{cilia} : 47 ± 0.5 , $p < 0.0001$, Tukey's multiple comparison test]); the majority of them are with PNs and their axons (Figure 7; Tables S9A and S9B). The distinctly different pattern of the astrocyte cilia contactome as compared with the neuronal cilia contactome indicates the uniqueness and diversity of neuronal and glial ciliary interactions with the surrounding cortical circuitry.

The diversity of neuronal and glial ciliary interactions with the cortical connectome

We then examined the defining features of neuronal ciliary membrane contacts with other cells of the cortical connectome. In particular, we examined if there are membrane specializations (i.e., synapse, gap junction, tight junction, adherens junction, etc.) at the sites of contact between a neuronal cilium and nearby cells, and if any preferential distribution of organelles (vesicles, lysosomes, mitochondria, endoplasmic reticulum, Golgi, etc.) are evident at these sites of contact.

Three types of neuronal cilia-axon contacts are evident. First, some of the contacts between axons and cilia are gap junction-like. These contacts are distinguished by a narrow membrane separation of 2–4 nm with electron-dense intercellular bridges in the middle (arrow, Figures S7A–S7D), characteristics similar to known gap junctions.^{46–49} Consistent with the presence of gap junctions in cilia, expression of a neuronal gap junction hemichannel, connexin-36,⁵⁰ is detected in subsets of ($9.6\% \pm 0.7\%$) human cortical neuronal primary cilia (Figure S7E). The second type of axon-ciliary contact is characterized by a dense accumulation of vesicles (arrow, Figures S7F and S7H) on the axonal side of cilia-axon contacts (Figures S7F–S7L). Reminiscent of presynaptic axon compartments, mitochondria are also present in axons at these sites (asterisk, Figure S7F). Whether these axon-cilia contacts are the cortical analogs of the recently described axo-ciliary synapses between serotonergic brainstem neurons and hippocampal CA1 pyramidal neurons³⁵ remains to be established. Intriguing are the third kind of neuronal cilia-axon contacts. These ciliary contacts are immediately adjacent to the synapses (Figure 8). Primary cilia of both INs and PNs of every cortical layer can be found intimately associated with synapses (Figure 8; Video S12), often forming a triangular partnership in which the primary ciliary compartment is intercalated next to the synaptic cleft. Primary ciliary tips are also found in tripartite (axon-dendrite-astrocyte) synaptic clefts (Figure 8). 51% of neuronal primary cilia engage in synaptic contacts. The majority

of these synapses are E (67%) as is the case for synapses generally in this sample. These observations indicate that primary cilia are intrinsic components of synaptic structure in subsets of cortical synapses. In contrast, control axonal segments adjacent to the respective neuronal cilia rarely make such contacts (neuronal cilia: 0.6 ± 0.09 vs. axon control: 0.05 ± 0.03 , $p < 0.0001$ [t test]). Compared with simple, tubular neuronal cilia, complex, beaded neuronal cilia make a significantly higher number of axo-ciliary associations, synaptic partnerships, and gap junction-like contacts (smooth cilia: 1.53 ± 0.19 , beaded cilia: 4.16 ± 0.42 , $p < 0.0001$ [t test]). Furthermore, 10% of neuronal cilia made only synaptic contacts, 14% made only axo-ciliary type contacts, and 7% formed only gap junction-like contacts. The majority of cilia made two or all three types of contacts.

Neuronal primary cilia also make numerous contacts with dendritic spines (Figures S8A–S8D). Large mitochondria, indicative of active Ca^{2+} dynamics, are seen in 32.5% of cilia-contacting dendritic spines (asterisk, Figures S8A–S8C).⁵¹ In contrast to neurons, both astrocyte (Figures S8E–S8G; Video S13) and microglial (Figures S8H and S8I) processes encircle the neuronal cilium at the sites of contact, often fully embedding them into their network of processes.

Akin to neuronal cilia, a subset of astrocyte cilia associates with synaptic clefts (Figures S9A, S9B, and S9E), and can form axo-ciliary contacts (Figures S9C–S9E). Astrocyte cilia makes significantly fewer such contacts than neuronal cilia (neuronal cilia: 1.78 ± 0.16 vs. astrocyte cilia: 0.0297 ± 0.016 , $p < 0.001$ [t test]). Further, astrocyte primary cilia also form contacts with dendritic spines (Figure S9F). As like neuronal cilia, astrocyte cilia are often engulfed by astrocyte processes (Figure S9G).

The majority of neuronal ciliary contacts emerge from other neurons or glia in the same or adjacent layer (99.9% of IN or PN cilia contacts with other neurons; 99.8% of IN or PN cilia contacts with glia) (Figures S10A and S10B). However, occasional long-distance contacts spanning multiple layers, between a neuronal cilium and dendritic projections from neurons a few layers away, were also noticed (Figures S2E and S2F). Overall, this predominantly local nature of the neuronal ciliary connectome is in contrast to the extensive, multi-layer, and region-spanning nature of the respective neuron's synaptic connectome (Figure S11). As with neurons, the majority of astrocyte ciliary contacts (99.9% of contacts with neurons; 100% of contacts with other glia) originate from other neurons and glia in the same or an adjacent layer (Figure S10C).

Relationship between cilia and neuronal structural and functional features

Since neuronal primary cilia are thought to modulate the growth, connectivity, and function of neurons, we analyzed whether neuronal ciliary characteristics and the corresponding functional (e.g., synaptic input or output) or structural features of neurons (e.g., spininess, volume, etc.) are correlated. During image

Figure 5. Cortical interneuronal primary cilia contactome

Primary cilia of cortical interneurons from all six cortical layers (arrow, columns 1 and 2; A–F) and all the axons and dendrites of other neurons (column 3) as well as the non-neural cells contacting them (column 4). Column 1 shows electron micrographs of the relevant cilium (arrow) and its contacting cells. I–VI, cortical layers. Scale bars: 2 μ m (column 1) and 5 μ m (columns 2–4).




Figure 6. Cortical projection neuronal primary cilia contactome

Primary cilia of cortical projection neurons from layers II–VI (arrow, columns 1 and 2; A–E) and all the axons and dendrites of other neurons (column 3) as well as the non-neural cells contacting them (column 4). Column 1 shows electron micrographs of the relevant cilium (arrow) and its contacting cells. II–VI, cortical layers. Scale bars: 2 μ m (column 1) and 5 μ m (columns 2–4).

segmentation, automated predictions of the neuronal subcompartment composition (axon, dendrite, dendritic spine, AIS, etc.) were generated at points along each segment.³⁶ We used

these predictions to quantify neuronal characteristics, for example, the spininess of a neuron is defined by the ratio of spine predictions to dendrite predictions along the segment. The




Figure 7. Cortical astrocyte primary cilia contactome

(A–F) Primary cilia (arrow) of astrocytes from layers I–VI (arrow, columns 1 and 2; A–F) and all the axons and dendrites of other neurons as well as the non-neural cell processes contacting them (column 3). Column 1 shows electron micrographs of the relevant cilium (arrow) and its contacting cells. Ciliary pockets are outlined in column 1 (white box). In columns 2 and 3 astrocyte cell body is shown in a translucent background so as not to obstruct the cilium. Insets in column 2 show the whole astrocyte.

(G and H) Quantification of astrocyte primary cilia contactome. Cell types (G) and different cell domains (H) contacting the primary cilium of different cortical layer astrocytes were quantified. Data shown are mean $\pm$ SEM. Two-way ANOVA (cell types): $F_{6, 30} = 85.93$, $p = 1.1E-15$. Two-way ANOVA (cell domains): $F_{10, 50} = 102$, $p = 1E-15$. I–VI, cortical layers. Scale bars: $0.45 \mu\text{m}$ (column 1) and $0.8 \mu\text{m}$ (columns 2–3).

number and type of synaptic connections and volume (voxels) of each neuron were also quantified during image segmentation and annotation.³⁶ We found a moderate but significant positive correlation between the cilia length and the spininess of a neuron ($r, 0.41$, $p < 1E-16$; Figure S12A). The spininess of a neuron positively correlates with the total number of neuronal ciliary contacts ($r, 0.352$, $p < 1E-16$; Figure S12B), in particular with the number of E axonal contacts ($r, 0.614$, $p < 1E-16$; Figure S12C). In addition, the number of incoming synapses to neurons positively correlates with the number of E axonal contacts of cilia ($r, 0.43$, $p < 1E-16$; Figure S12D). In contrast, both the spininess and the number of incoming synapses to neurons negatively correlate with the number of neuronal ciliary I axonal contacts ($r, -0.39$ spininess [$p < 1E-16$], -0.34 number of incoming synapses [$p = 1E-16$]; Figures S12E and S12F). These observations provide a rationale to further examine the intentional or incidental nature of the correlation between the patterns within the ciliary connectome and the functional characteristics of cortical neurons.

DISCUSSION

Disrupted primary cilia signaling and the resultant changes in neuronal circuit formation and function lead to brain malformations and neurodevelopmental disorders, including developmental delay, autism, intellectual disabilities, mood disorders, SCZ, obesity, ataxia, apraxia, and epilepsy. While these observations implicate primary cilia signaling as a potential non-synaptic mechanism through which environmental signals may shape and refine neuronal circuits, how neuronal primary cilia are organized within the cortical connectome to impact neural circuit function in the human cerebral cortex remain unresolved. Here, using serial EM reconstructed human cortex, we precisely and comprehensively delineated primary cilia organization and the connectome that animates ciliary signaling mechanisms in a diverse array of neurons and astrocytes in all layers of the human temporal cortex.

The length of a neuronal cilium positively correlates with the number of contacts it makes. Cortical neuronal cilia can make several types of functional associations with axons, dendrites, and synapses to effect changes in neuronal function. First, neuronal cilia can form intimate contacts with the synaptic cleft, thus exposing neuronal ciliary membrane receptors to synaptically released neurotransmitters (NTs), neuromodulators, second messengers, and other neuroactive components. A perisynaptic cilium




Figure 8. Neuronal primary cilia as an intrinsic component of synaptic structure

(A and B) Neuronal primary cilia tips are located right next to synapses. Primary cilia highlighted in yellow, and purple indicate primary cilia from inhibitory (A) and excitatory neurons (B), respectively. Astrocytes, axons, and dendrites are colored in green, blue, and red, respectively. Cortical layers (I–VI) and synapse type (excitatory [E] and inhibitory [I]) are indicated in each panel. IN, interneuron; PN, projection neuron.

(C) Electron micrograph of two synapses with a contacting projection neuronal primary cilium in the middle.

(D) Electron micrograph of two synapses (E and I) contacted by a projection neuronal primary cilium.

(E) Quantification of synaptic association of neuronal cilia. For each cilium, the number of ciliary contacts with *bona fide* chemical synapses was quantified. Synapse type (E and I) was also noted. Similar quantifications were made for adjacent control axonal segments of the same length and phenotype as the respective cilium. IN, interneuron; PN, projection neuron. Data shown are mean $\pm$ SEM. IN_{cilia} vs. $IN_{axon\ control}$, $p < 0.05$ (t test). PN_{cilia} vs. $PN_{axon\ control}$, $p < 0.05$ (t test).

(F) Using the cilia localized signaling machinery, synaptic primary cilium may sense and respond to synaptic activity and environment. Unlike the previously known tripartite synapse (axon, dendrite, astrocyte), the integration of neuronal primary cilium leads to the formation of a tetrapartite (pre, post, astrocyte, cilium) synapse. Please see [supplemental information](#) for Neuroglancer links to images in (A)–(D).

may bind synaptically released NTs. NTs such as norepinephrine, serotonin, dopamine, and somatostatin that can activate different ciliary G-protein-coupled receptors (GPCRs) are widely expressed in the mature cerebral cortex. Furthermore, the ion channels in cilium (e.g., PKD1L1, PKD2L1, PKD2, TRPC1 and TRPC4, TRPM2 and TRPM3, and TRPV4)^{52–58} may help maintain the appropriate ionic milieu of the synaptic cleft necessary to facilitate proper synaptic function. Single-cell RNA-seq-based expression analysis of these cilia-associated receptors, ion channels, and second messengers in diverse human cortical neurons indicate distinct patterns of neuronal expression (Figure S13),^{21,23,59–62} thus suggesting a neuron-type-specific diversity in the signaling competence of cortical neuronal primary cilia. Compellingly, 75% of the known cilia-associated GPCRs are differentially ex-

pressed in the frontal cortices of a major neural circuit disorder, SCZ.³¹

Neuronal cilia also make numerous contacts with axonal segments that are full of synaptic vesicles. These contacts are distinct from classical chemical synapses. A few of these cilia-axonal contacts have mitochondria (Figure S7) in addition to vesicles, suggestive of a presynaptic style architecture. However, post-synaptic density-type membrane contrast enhancement is not consistently seen on the ciliary side of these contacts, though this should be further explored with different EM fixation methods designed to enhance contrast. Neuronal cilia make comparatively more axo-ciliary-type associations than contacts with chemical synapses (1.18 ± 0.12 vs. 0.6 ± 0.09 , $p < 0.001$, t test). This intriguing pattern of ciliary interactions with different

synaptic compartments may provide an avenue of functional flexibility to ciliary signaling.

Aside from the interactions with chemical synapses and axo-ciliary contacts, gap-junction-like, electrical synaptic contacts between cilia and axons are evident as well. These gap junctional contacts may enable direct diffusion of second messengers and small metabolites, such as Ca^{2+} , cAMP, ATP, glucose, and inositol triphosphate, between a neuronal primary cilium and the axonal terminal.^{63–65} Due to their ability to electrically couple cells, these gap junctions may also directly facilitate the spreading of presynaptic electrical currents to the post-synaptic neuron via its primary cilium. However, it is important to note that due to the current resolution ($4 \times 4 \text{ nm}^2$) of the EM dataset, our observations might underestimate the electrical synapses associated with cilia because of their small size (1–20 nm). Enhanced electron-dense labeling methods during the EM procedure coupled with molecular tagging of gap junction proteins will help resolve this deficit in the future. Further, human cortical neurons from both sexes were used in our cilia-gap junction co-localization studies, but the influence of sex on the pattern of co-localization is unknown.

Of these associations of neuronal primary cilia with the activity-related compartments of other neurons, the primary cilium as an intrinsic component of subsets of cortical synapses is compelling (Figure 8). A single neuronal primary cilium can contact up to ~5 synapses, both E and I. Synaptically released ligands can act on the ciliary membrane receptors, generate Ca^{2+} transients,^{5,66} and thus provide neurons an additional ability to monitor and modulate their functional homeostasis in response to local activity. However, several questions remain: how do cilia-derived signals or ciliary invagination of synaptic structures affect the function, maturation, maintenance, and sculpting of synaptic circuits? Contact by primary cilia may regulate critical aspects of synapse formation, function, maintenance, or elimination, as the other cellular partner of synapses, i.e., astrocytes, do.⁴⁵ Nonetheless, primary cilia do not intercalate with all synapses; thus, it will be important to establish if the primary cilia-synapse association is a dynamic process that depends on neuronal or astroglial activity. Other related questions, including whether the primary cilia's association with synapses is part of the key epochs of cortical circuit development and plasticity, such as the critical period, long-term potentiation, long-term depression, and homeostatic scaling, remain completely open.

Astrocyte-derived signals may not only directly regulate the synapses^{45,67,68} but may also fine-tune neuronal activity via the adjacent primary cilia. For example, astrocyte-released signals may regulate NT or neuromodulator receptor localization on the ciliary membrane at synaptic sites. Furthermore, synaptic activity can trigger astrocytic Ca^{2+} rise and the secretion of neuroactive gliotransmitter molecules, such as glutamate, D-serine, taurine, TNF- α , and ATP.^{45,69–71} Similarly, microglia can also release modulators such as ATP in response to circuit activity.^{72–74} The ciliary membrane is known to be decorated with receptors for these gliotransmitter ligands (e.g., P2YR2 for ATP) and thus may enable glial modulation of neuronal or glial homeostasis via primary cilia.

Primary cilia can form contacts with axons, dendrites, astrocytes, and microglia but rarely associate with neuronal AIS or

blood vessels. Neuronal primary cilia contact 10-fold or more axons than dendrites, even though the numerical ratio of axonal to dendritic processes in the cortical volume is ~1.8:1.³⁶ Similarly, despite the presence of twice as many glia than neurons, only ~6% of neuronal ciliary contacts are with glia. Further, there are nearly 3.7 times as many oligodendrocytes than astrocytes in the cortical volume,³⁶ but neuronal cilia associate with ~14-fold or more astrocytes than with oligodendrocytes (Table S5). However, other than cell numbers, the volume occupied by the different cell types and cellular domains in each layer may also be an important contributing factor to the pattern of ciliary contactome we observed. Further evaluation of the functional relevance of these unique features of ciliary association with the surrounding cells in the cortical connectome and examination of the intentional vs. incidental nature of these associations will help determine how neuronal and glial cells form and organize their contacts with neuronal primary cilia to subserve a circuit modulating function.

In addition to rapidly modulating neural circuits through its rich and diverse interactions with the cortical circuit, neuronal cilia may also generate a direct link to the nucleus to modulate neuronal activity via transcriptional changes. Ciliary receptor activation can trigger changes in the accessibility of gene regulatory elements and chromatin landscape in neurons.^{35,75,76} The positive correlation between the cilia length and the spininess or the number of incoming synapses in a neuron raises the intriguing possibility that neuronal ciliary signaling may impact the synaptic competence of a neuron via such cilia-mediated gene regulation.

Structurally, it is remarkable that almost all cortical neuronal cilia do not extend a 9 + 0 MT axoneme all the way to the tip. The majority of cortical neuronal cilia do not display a 9 + 0 MT doublet-based axoneme core at the ciliary midpoint. This may help create neuronal ciliary domains unencumbered by a rigid MT axoneme core to facilitate efficient ciliary signaling vesicle (i.e., ectosome)³⁷ release. Importantly, the changing MT core along the neuronal ciliary length may also modulate the anterograde and retrograde intraflagellar transport of signaling molecules within a cilium and thus its signaling competence.^{77–80} The layer- and neuronal-type-specific variations in the MT axoneme in cortical neurons may thus underlie the functional diversity in neuronal primary cilia. A similar variation in 9 + 0 MT axoneme core is also evident in astrocyte cilia. However, unlike neurons, the majority of astrocyte cilia (~75%) are in deep membrane pockets, without access to surrounding circuitry. Further, the presence of a ciliary axoneme embedded within the ciliary sheath inside an astrocyte soma in subsets (~10%) of adult human cortical astrocytes suggests an active intracellular pathway-mediated ciliogenesis in these cells.^{81,82} Such diversity between neuronal and glial cilia may help modulate the distinct roles primary cilia play in neurons vs. glia in the human cortex.

Collectively, these observations suggest that there exists a neuronal cell-type-specific ciliary signaling avenue that can exert a dynamic influence on the functional state of neurons. By coincidence or design, each cortical neuronal or astroglial primary cilium interacts with a unique subset of other cortical neurons and glia. This contactome gives rise to the ciliary connectome. This remarkable diversity in the individual neuronal and glial ciliary connectome and their fine structure may underlie how primary cilia activity is transformed into changes in neural circuit function. The primary

cilia connectome endows a neuron or astrocyte with a unique barcode of association with the surrounding neural circuitry. It raises the intriguing possibility that this diversity enables primary cilia of different cortical neurons and glia to sample and fine-tune local neuronal circuit activity and neuronal homeostasis in a unique manner. When perturbed in ciliopathies or other human brain disorders, alterations in this unique neuromodulation may trigger cortical neural circuit dysfunction.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **RESOURCE AVAILABILITY**
 - Lead contact
 - Materials availability
 - Data and code availability
- **EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS**
 - Serial EM reconstruction of human cortical volume
 - Human cortical neuronal cells
- **METHOD DETAILS**
 - Serial EM reconstruction of primary cilia in human cortical volume
 - Analysis of primary cilia in neurons and glia of all cerebral cortical layers
 - Immunohistochemistry
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
 - General statistical analysis

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.neuron.2023.09.032>.

ACKNOWLEDGMENTS

This research was supported by NIH grants MH132710 (to R.Y., J.W.L., and E.S.A.) and NS116859 (to E.S.A.); National Institute of Mental Health Conte Center Award P50 MH094271 (to J.W.L.); NIH Brain Initiative grants U19 NS104653, U24 NS109102, and U01 EB026996 (to J.W.L.); the Stanley Center at the Broad Institute (J.W.L.); and New York Stem Cell Foundation Robertson Investigator Award (to J. G.). We thank Yukako Yokota, Tanae Lewis, and Troy Ghashghaei for insightful discussions.

AUTHOR CONTRIBUTIONS

J.W.L. and E.S.A. designed the experiments and supervised the project. J.Y.W., S.-J.C., K.D., C.M., C.C., A.S.-C., D.B., C.M., A.H., J.L., T.G., M.B., J.S., M. Januszewski, P.H.L., V.J., A.P., M. Jiang, J.S., J.G., and E.S.A. conducted the experiments and analyzed the data. J.W.L., R.Y., and E.S.A. wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: December 11, 2022

Revised: July 25, 2023

Accepted: September 22, 2023

Published: October 27, 2023

REFERENCES

1. Alvarez Retuerto, A.I., Cantor, R.M., Gleeson, J.G., Ustaszewska, A., Schackwitz, W.S., Pennacchio, L.A., and Geschwind, D.H. (2008). Association of common variants in the Joubert syndrome gene (AHI1) with autism. *Hum. Mol. Genet.* 17, 3887–3896. <https://doi.org/10.1093/hmg/ddn291>.
2. Arellano, J.I., Guadiana, S.M., Breunig, J.J., Rakic, P., and Sarkisian, M.R. (2012). Development and distribution of neuronal cilia in mouse neocortex. *J. Comp. Neurol.* 520, 848–873. <https://doi.org/10.1002/cne.22793>.
3. Guemez-Gamboa, A., Coufal, N.G., and Gleeson, J.G. (2014). Primary cilia in the developing and mature brain. *Neuron* 82, 511–521.
4. Guo, J., Higginbotham, H., Li, J., Nichols, J., Hirt, J., Ghukasyan, V., and Anton, E.S. (2015). Developmental disruptions underlying brain abnormalities in ciliopathies. *Nat. Commun.* 6, 7857. <https://doi.org/10.1038/ncomms8857>.
5. Guo, J., Otis, J.M., Higginbotham, H., Monckton, C., Cheng, J., Asokan, A., Mykytyn, K., Caspary, T., Stuber, G.D., and Anton, E.S. (2017). Primary cilia signaling shapes the development of interneuronal connectivity. *Dev. Cell* 42, 286–300.e4. <https://doi.org/10.1016/j.devcel.2017.07.010>.
6. Guo, J., Otis, J.M., Suciu, S.K., Catalano, C., Xing, L., Constable, S., Wachten, D., Gupton, S., Lee, J., Lee, A., et al. (2019). Primary cilia signaling promotes axonal tract development and is disrupted in Joubert syndrome-related disorders models. *Dev. Cell* 51, 759–774.e5. <https://doi.org/10.1016/j.devcel.2019.11.005>.
7. Green, J.A., and Mykytyn, K. (2014). Neuronal primary cilia: an underappreciated signaling and sensory organelle in the brain. *Neuropsychopharmacology* 39, 244–245.
8. Higginbotham, H., Eom, T.Y., Mariani, L.E., Bachleda, A., Hirt, J., Ghukasyan, V., Cusack, C.L., Lai, C., Caspary, T., and Anton, E.S. (2012). Arl13b in primary cilia regulates the migration and placement of interneurons in the developing cerebral cortex. *Dev. Cell* 23, 925–938. <https://doi.org/10.1016/j.devcel.2012.09.019>.
9. Hildebrandt, F., Benzing, T., and Katsanis, N. (2011). Ciliopathies. *N. Engl. J. Med.* 364, 1533–1543. <https://doi.org/10.1056/NEJMra1010172>.
10. Hilgendorf, K.I., Johnson, C.T., and Jackson, P.K. (2016). The primary cilium as a cellular receiver: organizing ciliary GPCR signaling. *Curr. Opin. Cell Biol.* 39, 84–92. <https://doi.org/10.1016/j.ceb.2016.02.008>.
11. Jacoby, M., Cox, J.J., Gayral, S., Hampshire, D.J., Ayub, M., Blockmans, M., Pernot, E., Kisseleva, M.V., Compère, P., Schiffrmann, S.N., et al. (2009). INPP5E mutations cause primary cilium signaling defects, ciliary instability and ciliopathies in human and mouse. *Nat. Genet.* 41, 1027–1031.
12. Jovasevic, V., Zhang, H., Sananbenesi, F., Guedea, A.L., Soman, K.V., Wiktorowicz, J.E., Fischer, A., and Radulovic, J. (2021). Primary cilia are required for the persistence of memory and stabilization of perineuronal nets. *iScience* 24, 102617.
13. Kirschen, G.W., Liu, H., Lang, T., Liang, X., Ge, S., and Xiong, Q. (2017). The radial organization of neuronal primary cilia is acutely disrupted by 5 seizure and ischemic brain injury. *Front. Biol. (Beijing)* 12, 124–138.
14. Lancaster, M.A., and Gleeson, J.G. (2009). The primary cilium as a cellular signaling center: lessons from disease. *Curr. Opin. Genet. Dev.* 19, 220–229. <https://doi.org/10.1016/j.gde.2009.04.008>.
15. Lee, C.H., Kang, G.M., and Kim, M.S. (2022). Mechanisms of weight control by primary cilia. *Mol. Cells* 45, 169–176. <https://doi.org/10.14348/molcells.2022.2046>.
16. Marley, A., and von Zastrow, M. (2012). A simple cell-based assay reveals that diverse neuropsychiatric risk genes converge on primary cilia. *PLoS One* 7, e46647. <https://doi.org/10.1371/journal.pone.0046647>.
17. Meffre, J., Chaumont-Dubel, S., la Cour, C., Loiseau, F., Watson, D.J., Dekeyne, A., Veno, S., M., Rivet, J., Gaven, F., Déléris, P., et al. (2012).

HT6 receptor recruitment of mTOR as a mechanism for perturbed cognition in schizophrenia. *EMBO Mol. Med.* 4, 1043–1056.

18. Mitchison, H.M., and Valente, E.M. (2017). Motile and non-motile cilia in human pathology: from function to phenotypes. *J. Pathol.* 241, 294–309. <https://doi.org/10.1002/path.4843>.
19. Mullins, N., Forstner, A.J., O'Connell, K.S., Coombes, B., Coleman, J.R.I., Qiao, Z., Als, T.D., Bigdeli, T.B., Bérte, S., Bryois, J., et al. (2021). Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. *Nat. Genet.* 53, 817–829.
20. Novarino, G., Akizu, N., and Gleeson, J.G. (2011). Modeling human disease in humans: The ciliopathies. *Cell* 147, 70–79. <https://doi.org/10.1016/J.CELL.2011.09.014>.
21. Liu, S., Trupiano, M.X., Simon, J., Guo, J., and Anton, E.S. (2021). The essential role of primary cilia in cerebral cortical development and disorders. *Curr. Top. Dev. Biol.* 142, 99–146. <https://doi.org/10.1016/bs.ctdb.2020.11.003>.
22. Park, S.M., Jang, H.J., and Lee, J.H. (2019). Roles of primary cilia in the developing brain. *Front. Cell. Neurosci.* 13, 218. <https://doi.org/10.3389/fncel.2019.00218>.
23. Reiter, J.F., and Leroux, M.R. (2017). Genes and molecular pathways underpinning ciliopathies. *Nat. Rev. Mol. Cell Biol.* 18, 533–547. <https://doi.org/10.1038/nrm.2017.60>.
24. Sarkisian, M.R., and Guadiana, S.M. (2015). Influences of primary cilia on cortical morphogenesis and neuronal subtype maturation. *Neuroscientist* 21, 136–151. <https://doi.org/10.1177/1073858414531074>.
25. Sattar, S., and Gleeson, J.G. (2011). The ciliopathies in neuronal development: a clinical approach to investigation of Joubert syndrome and Joubert syndrome-related disorders. *Dev. Med. Child Neurol.* 53, 793–798. <https://doi.org/10.1111/j.1469-8749.2011.04021.x>.
26. Stubbs, T., Koemeter-Cox, A., Bingman, J.I., Zhao, F., Kalyanasundaram, A., Rowland, L.A., Periasamy, M., Carter, C.S., Sheffield, V.C., Askwith, C.C., and Mykytyn, K. (2022). Disruption of dopamine receptor 1 localization to primary cilia impairs signaling in striatal neurons. *J. Neurosci.* 42, 6692–6705. <https://doi.org/10.1523/JNEUROSCI.0497-22.2022>.
27. Thomas, S., Boutaud, L., Reilly, M.L., and Benmerah, A. (2019). Cilia in hereditary cerebral anomalies. *Biol. Cell* 111, 217–231. <https://doi.org/10.1111/boc.201900012>.
28. SenGupta, P. (2017). Cilia and sensory signaling: The journey from “animalcules” to human disease. *PLoS Biol.* 15, e2002240. <https://doi.org/10.1371/journal.pbio.2002240>.
29. Vaisse, C., Reiter, J.F., and Berbari, N.F. (2017). Cilia and obesity. *Cold Spring Harbor Perspect. Biol.* 9, a028217. <https://doi.org/10.1101/cshperspect.a028217>.
30. Valente, E.M., Rosti, R.O., Gibbs, E., and Gleeson, J.G. (2014). Primary cilia in neurodevelopmental disorders. *Nat. Rev. Neurol.* 10, 27–36. <https://doi.org/10.1038/nrneurol.2013.247>.
31. Alhassen, W., Chen, S., Vawter, M., Robbins, B.K., Nguyen, H., Myint, T.N., Saito, Y., Schulmann, A., Nauli, S.M., Civelli, O., et al. (2021). Patterns of cilia gene dysregulations in major psychiatric disorders. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 109, 110255. <https://doi.org/10.1016/j.pnpbp.2021.110255>.
32. Kim, M., Haney, J.R., Zhang, P., Hernandez, L.M., Wang, L.K., Perez-Cano, L., Loohuis, L.M.O., de la Torre-Ubieta, L., and Gandal, M.J. (2021). Brain gene co-expression networks link complement signaling with convergent synaptic pathology in schizophrenia. *Nat. Neurosci.* 24, 799–809. <https://doi.org/10.1038/s41593-021-00847-z>.
33. Kumamoto, N., Gu, Y., Wang, J., Janoschka, S., Takemaru, K., Levine, J., and Ge, S. (2012). A role for primary cilia in glutamatergic synaptic integration of adult-born neurons. *Nat. Neurosci.* 15, 399. <https://doi.org/10.1038/nn.3042>.
34. Tereshko, L., Gao, Y., Cary, B.A., Turrigiano, G.G., and SenGupta, P. (2021). Ciliary neuropeptidergic signaling dynamically regulates excitatory synapses in postnatal neocortical pyramidal neurons. *eLife* 10, e65427. <https://doi.org/10.7554/eLife.65427>.
35. Sheu, S.H., Upadhyayula, S., Dupuy, V., Pang, S., Deng, F., Wan, J., Walpita, D., Pasolli, H.A., Houser, J., Sanchez-Martinez, S., et al. (2022). A serotonergic axon-cilium synapse drives nuclear signaling to alter chromatin accessibility. *Cell* 185, 3390–3407.e18. <https://doi.org/10.1016/j.cell.2022.07.026>.
36. Shapson-Coe, A., Januszewski, M., Berger, D.R., Pope, A., Wu, Y., Blakely, T., Schalek, R.L., Li, P., Wang, S., Maitin-Shepard, J., et al. (2021). A connectomic study of a petascale fragment of human cerebral cortex. <https://doi.org/10.1101/2021.05.29.446289>.
37. Nager, A.R., Goldstein, J.S., Herranz-Pérez, V., Portran, D., Ye, F., Garcia-Verdugo, J.M., and Nachury, M.V. (2017). An actin network dispatches ciliary GPCRs into extracellular vesicles to modulate signaling. *Cell* 168, 252–263.e14. <https://doi.org/10.1016/j.cell.2016.11.036>.
38. Molla-Herman, A., Ghossoub, R., Blisnick, T., Meunier, A., Serres, C., Silbermann, F., Emmerson, C., Romeo, K., Bourdoncle, P., Schmitt, A., et al. (2010). The ciliary pocket: an endocytic membrane domain at the base of primary and motile cilia. *J. Cell Sci.* 123, 1785–1795. <https://doi.org/10.1242/jcs.059519>.
39. Benmerah, A. (2013). The ciliary pocket. *Curr. Opin. Cell Biol.* 25, 78–84. <https://doi.org/10.1016/j.ceb.2012.10.011>.
40. Ishikawa, T., Ueno, H., Omori, T., and Kikuchi, K. (2021). Cilia and centrosomes: ultrastructural and mechanical perspectives. *Semin. Cell Dev. Biol.* 110, 61–69. <https://doi.org/10.1016/j.semcdb.2020.03.007>.
41. Vertii, A., Hung, H.F., Hehnl, H., and Doherty, S. (2016). Human basal body basics. *Cilia* 5, 13. <https://doi.org/10.1186/s13630-016-0030-8>.
42. Harris, K.D., and Shepherd, G.M. (2015). The neocortical circuit: themes and variations. *Nat. Neurosci.* 18, 170–181. <https://doi.org/10.1038/nn.3917>.
43. Lim, L., Mi, D., Llorca, A., and Marín, O. (2018). Development and functional diversification of cortical interneurons. *Neuron* 100, 294–313. <https://doi.org/10.1016/j.neuron.2018.10.009>.
44. Gallo, N.B., Paul, A., and Van Aelst, L. (2020). Shedding light on chandelier cell development, connectivity, and contribution to neural disorders. *Trends Neurosci.* 43, 565–580. <https://doi.org/10.1016/j.tins.2020.05.003>.
45. Allen, N.J., and Eroglu, C. (2017). Cell biology of astrocyte-synapse interactions. *Neuron* 96, 697–708. <https://doi.org/10.1016/j.neuron.2017.09.056>.
46. Rash, J.E., Yasumura, T., and Dudek, F.E. (1998). Ultrastructure, histological distribution, and freeze-fracture immunocytochemistry of gap junctions in rat brain and spinal cord. *Cell Biol. Int.* 22, 731–749. <https://doi.org/10.1006/cbir.1998.0392>.
47. Shivers, R.R., and McVicar, L.K. (1995). Gap junctions revealed by freeze-fracture electron microscopy. *Microsc. Res. Tech.* 31, 437–445. <https://doi.org/10.1002/jemt.1070310512>.
48. Wu, L., Dong, A., Dong, L., Wang, S.Q., and Li, Y. (2019). PARIS, an optogenetic method for functionally mapping gap junctions. *eLife* 8, e43366. <https://doi.org/10.7554/eLife.43366>.
49. Zihni, C., Mills, C., Matter, K., and Balda, M.S. (2016). Tight junctions: from simple barriers to multifunctional molecular gates. *Nat. Rev. Mol. Cell Biol.* 17, 564–580. <https://doi.org/10.1038/nrm.2016.80>.
50. Söhl, G., Maxeiner, S., and Willecke, K. (2005). Expression and functions of neuronal gap junctions. *Nat. Rev. Neurosci.* 6, 191–200.
51. Hirabayashi, Y., Kwon, S.K., Paek, H., Pernice, W.M., Paul, M.A., Lee, J., Erfani, P., Raczkowski, A., Petrey, D.S., Pon, L.A., and Polleux, F. (2017). ER-mitochondria tethering by PDZD8 regulates Ca²⁺ dynamics in mammalian neurons. *Science* 358, 623–630. <https://doi.org/10.1126/science.aan6009>.
52. Delling, M., DeCaen, P.G., Doerner, J.F., Febvay, S., and Clapham, D.E. (2013). Primary cilia are specialized calcium signalling organelles. *Nature* 504, 311–314.

53. DeCaen, P.G., Delling, M., Vien, T.N., and Clapham, D.E. (2013). Direct recording and molecular identification of the calcium channel of primary cilia. *Nature* 504, 315–318.
54. Doerner, J.F., Delling, M., and Clapham, D.E. (2015). Ion channels and calcium signaling in motile cilia. *eLife* 4, e11066. <https://doi.org/10.7554/eLife.11066>.
55. Lishko, P., and Kirichok, Y. (2015). Signaling the differences between cilia. *eLife* 4, e12760. <https://doi.org/10.7554/eLife.12760>.
56. Mick, D.U., Rodrigues, R.B., Leib, R.D., Adams, C.M., Chien, A.S., Gygi, S.P., and Nachury, M.V. (2015). Proteomics of primary cilia by proximity labeling. *Dev. Cell* 35, 497–512. <https://doi.org/10.1016/j.devcel.2015.10.015>.
57. Pablo, J.L., DeCaen, P.G., and Clapham, D.E. (2017). Progress in ciliary ion channel physiology. *J. Gen. Physiol.* 149, 37–47. <https://doi.org/10.1085/jgp.201611696>.
58. Tajhya, R., and Delling, M. (2020). New insights into ion channel-dependent signalling during left-right patterning. *J. Physiol.* 598, 1741–1752. <https://doi.org/10.1113/JP277835>.
59. van Dam, T.J., Wheway, G., Slaats, G.G., SYSCILIA Study Group, Huynen, M.A., and Giles, R.H. (2013). The SYSCILIA gold standard (SCGSv1) of known ciliary components and its applications within a systems biology consortium. *Cilia* 2, 7. <https://doi.org/10.1186/2046-2530-2-7>.
60. van Dam, T.J.P., Kennedy, J., van der Lee, R., de Vrieze, E., Wunderlich, K.A., Rix, S., Dougherty, G.W., Lambacher, N.J., Li, C., Jensen, V.L., et al. (2019). CiliaCarta: An integrated and validated compendium of ciliary genes. *PLoS One* 14, e0216705. <https://doi.org/10.1371/journal.pone.0216705>.
61. Ishikawa, H., Thompson, J., Yates, J.R., 3rd, and Marshall, W.F. (2012). Proteomic analysis of mammalian primary cilia. *Curr. Biol.* 22, 414–419. <https://doi.org/10.1016/j.cub.2012.01.031>.
62. Hodge, R.D., Bakken, T.E., Miller, J.A., Smith, K.A., Barkan, E.R., Graybuck, L.T., Close, J.L., Long, B., Johansen, N., Penn, O., et al. (2019). Conserved cell types with divergent features in human versus mouse cortex. *Nature* 573, 61–68.
63. Dong, A., Liu, S., and Li, Y. (2018). Gap junctions in the nervous system: Probing functional connections using new imaging approaches. *Front. Cell. Neurosci.* 12, 320. <https://doi.org/10.3389/fncel.2018.00320>.
64. Evans, W.H., and Martin, P.E. (2002). Gap junctions: structure and function (Review). *Mol. Membr. Biol.* 19, 121–136. <https://doi.org/10.1080/09687680210139839>.
65. Miller, A.C., and Pereda, A.E. (2017). The electrical synapse: molecular complexities at the gap and beyond. *Dev. Neurobiol.* 77, 562–574. <https://doi.org/10.1002/dneu.22484>.
66. Su, S., Phua, S.C., DeRose, R., Chiba, S., Narita, K., Kalugin, P.N., Katada, T., Kontani, K., Takeda, S., and Inoue, T. (2013). Genetically encoded calcium indicator illuminates calcium dynamics in primary cilia. *Nat. Methods* 10, 1105–1107. <https://doi.org/10.1038/nmeth.2647>.
67. Bushong, E.A., Martone, M.E., Jones, Y.Z., and Ellisman, M.H. (2002). Protoplasmic astrocytes in CA1 stratum radiatum occupy separate anatomical domains. *J. Neurosci.* 22, 183–192. <https://doi.org/10.1523/JNEUROSCI.22-01-00183.2002>.
68. Oberheim, N.A., Takano, T., Han, X., He, W., Lin, J.H., Wang, F., Xu, Q., Wyatt, J.D., Pilcher, W., Ojemann, J.G., et al. (2009). Uniquely hominid features of adult human astrocytes. *J. Neurosci.* 29, 3276–3287. <https://doi.org/10.1523/JNEUROSCI.4707-08.2009>.
69. Harada, K., Kamiya, T., and Tsuboi, T. (2015). Gliotransmitter release from astrocytes: Functional, developmental, and pathological implications in the brain. *Front. Neurosci.* 9, 499. <https://doi.org/10.3389/fnins.2015.00499>.
70. Koizumi, S. (2021). Glial purinergic signals and psychiatric disorders. *Front. Cell. Neurosci.* 15, 822614. <https://doi.org/10.3389/fncel.2021.822614>.
71. Savtchouk, I., and Volterra, A. (2018). Gliotransmission: Beyond black-and-white. *J. Neurosci.* 38, 14–25. <https://doi.org/10.1523/JNEUROSCI.0017-17.2017>.
72. Ginhoux, F., and Garel, S. (2018). The mysterious origins of microglia. *Nat. Neurosci.* 21, 897–899.
73. Squarzone, P., Oller, G., Hoeftel, G., Pont-Lezica, L., Rostaing, P., Low, D., Bessis, A., Ginhoux, F., and Garel, S. (2014). Microglia modulate wiring of the embryonic forebrain. *Cell Rep.* 8, 1271–1279.
74. Thion, M.S., and Garel, S. (2017). On place and time: microglia in embryonic and perinatal brain development. *Curr. Opin. Neurobiol.* 47, 121–130.
75. Hilgendorf, K.I., Johnson, C.T., Mezger, A., Rice, S.L., Norris, A.M., Demeter, J., Greenleaf, W.J., Reiter, J.F., Kopinke, D., and Jackson, P.K. (2019). Omega-3 fatty acids activate ciliary FFAR4 to control adipogenesis. *Cell* 179, 1289–1305.e21.
76. Kong, J.H., Siebold, C., and Rohatgi, R. (2019). Biochemical mechanisms of vertebrate hedgehog signaling. *Development* 146, dev166892. <https://doi.org/10.1242/dev.166892>.
77. Klena, N., and Pigino, G. (2022). Structural biology of cilia and intraflagellar transport. *Annu. Rev. Cell Dev. Biol.* 38, 103–123. <https://doi.org/10.1146/annurev-cellbio-120219-034238>.
78. Mallet, A., and Bastin, P. (2022). Restriction of intraflagellar transport to some microtubule doublets: An opportunity for cilia diversification? *Bioessays* 44, e2200031. <https://doi.org/10.1002/bies.202200031>.
79. Jordan, M.A., and Pigino, G. (2021). The structural basis of intraflagellar transport at a glance. *J. Cell Sci.* 134, jcs247163. <https://doi.org/10.1242/jcs.247163>.
80. Sun, S., Fisher, R.L., Bowser, S.S., Pentecost, B.T., and Sui, H. (2019). Three-dimensional architecture of epithelial primary cilia. *Proc. Natl. Acad. Sci. USA* 116, 9370–9379. <https://doi.org/10.1073/pnas.1821064116>.
81. Sorokin, S.P. (1968). Reconstructions of centriole formation and ciliogenesis in mammalian lungs. *J. Cell Sci.* 3, 207–230.
82. Ganga, A.K., Kennedy, M.C., Oguchi, M.E., Gray, S., Oliver, K.E., Knight, T.A., De La Cruz, E.M., Homma, Y., Fukuda, M., and Breslow, D.K. (2021). Rab34 GTPase mediates ciliary membrane formation in the intracellular ciliogenesis pathway. *Curr. Biol.* 31, 2895–2905.e7. <https://doi.org/10.1016/j.cub.2021.04.075>.
83. Stein, J.L., de la Torre-Ubieta, L., Tian, Y., Parikshak, N.N., Hernández, I.A., Marchetto, M.C., Baker, D.K., Lu, D., Hinman, C.R., Lowe, J.K., et al. (2014). A quantitative framework to evaluate modeling of cortical development by neural stem cells. *Neuron* 83, 69–86. <https://doi.org/10.1016/j.neuron.2014.05.035>.
84. Konopka, G., Wexler, E., Rosen, E., Mukamel, Z., Osborn, G.E., Chen, L., Lu, D., Gao, F., Gao, K., Lowe, J.K., and Geschwind, D.H. (2012). Modeling the functional genomics of autism using human neurons. *Mol. Psychiatry* 17, 202–214. <https://doi.org/10.1038/mp.2011.60>.
85. Barral, D.C., Staiano, L., Guimas Almeida, C., Cutler, D.F., Eden, E.R., Futter, C.E., Gallone, A., Marques, A.R.A., Medina, D.L., Napolitano, G., et al. (2022). Current methods to analyze lysosome morphology, positioning, motility and function. *Traffic* 23, 238–269. <https://doi.org/10.1111/tra.12839>.
86. Nakagawa, N., Plestant, C., Yabuno-Nakagawa, K., Li, J., Lee, J., Huang, C.W., Lee, A., Krupa, O., Adhikari, A., Thompson, S., et al. (2019). Memo1-mediated tiling of radial glial cells facilitates cerebral cortical development. *Neuron* 103, 836–852.e5. <https://doi.org/10.1016/j.neuron.2019.05.049>.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit polyclonal anti-connexin 36	ThermoFisher	Cat#710663; RRID: AB_2576616
Mouse anti-ar113b	Biolegend	Cat# 857602; RRID: AB_2801216
Chicken anti- β tubulin	Millipore	Cat# 9354; RRID: AB_570918
AlexaFluor goat anti-mouse 647	Invitrogen	Cat# A21236; RRID: AB_2535805
AlexaFluor goat anti-rabbit Cy3	Invitrogen	Cat# A10520; RRID: AB_2534029
AlexaFluor goat anti-chicken 488	Invitrogen	Cat# A11039; RRID: AB_2534096
Experimental models: Organisms/strains		
Adults human cerebral cortex	Shapson-Coe et al. ³⁶	N/A
Human neural progenitor cells	Stein et al. ⁸³	N/A
Chemicals, peptides, and recombinant proteins		
DAPI	Invitrogen	Cat# D21490
Osmium tetroxide-thiocarbohydrazide	EMS, Sigma	Cat# 19150, Cat# T2137
Uranyl acetate	EMS	Cat# 22400
Software and algorithms		
Excel	Microsoft	https://products.office.com/en-us/excel
SAS	SAS Institute	https://oda.sas.com
GraphPad	GraphPad Software Inc.	https://graphpad.com
Neuroglancer	Shapson-Coe et al. ³⁶	https://github.com/google/neuroglancer

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to, and will be fulfilled by, the lead contact, E.S. Anton (anton@med.unc.edu).

Materials availability

This study did not generate unique reagents.

Data and code availability

Serial EM dataset used in this study is available at Serial EM Data: <https://h01-release.storage.googleapis.com/landing.html>. Microscopy data reported in this paper will be shared by the [lead contact](#) upon request. Any additional information required to reanalyze the data reported in this paper is available from the [lead contact](#) upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Serial EM reconstruction of human cortical volume

A cortical thickness slab (2.5cm X 0.8cm) of histologically normal, unaffected left anterior temporal lobe from a 45-year-old female with a history of simple and complex partial seizures, resistant to drug treatment, was removed during surgical ablation of an epileptic focus in her left medial temporal lobe. This normal tissue was used for the generation of serial EM dataset.³⁶ The human brain tissue sample was collected and processed according to the Massachusetts General Hospital, Harvard University, and NIH guidelines on the use of human tissue.

Human cortical neuronal cells

The fetal tissue (GW15-18) harvesting and neurosphere collection were done as described previously^{83,84} and cells were derived from frozen pHNPC stocks. Neuronal cells from both males and females were used. Cells were maintained at 37°C/5%

CO₂ in Neurobasal A (Invitrogen) supplemented with 10% BIT (Stem Cell Technologies), Primocin (Invivogen), GlutaMAX (Gibco), and heparin (1 µg/mL; Sigma)] with freshly added EGF, FGF2, PDGF (each at 20 ng/mL; Invitrogen), and LIF (2 ng/mL; Invitrogen).

METHOD DETAILS

Serial EM reconstruction of primary cilia in human cortical volume

Serial EM reconstruction of human cortical volume is described in Shapson-Coe et al.³⁶ Briefly, a full cortical thickness slab (2.5cm X 0.8cm) of histologically normal human temporal lobe from a 45-year-old female was harvested during hippocampal resection surgery to treat epilepsy. The sample was stained with osmium, trimmed down to a rectangular area of 4584 X 1975 micrometers spanning all cortical layers from layer 1 to superficial white matter, sectioned at 30nm, and imaged at 4X4 nm resolution in multibeam (61 electron beams) scanning electron microscope mSEM (Zeiss). Images from more than 5292 slices were aligned, stitched, rendered, and segmented.³⁶ The total data set has a volume of ~ 1 mm³ volume and contains ~1.4 PB of image data, containing the three-dimensional structure of 56,000 cells, hundreds of millions of neurites, and >100 million synaptic connections. Training data based on human tracing of >700 cells were used to obtain a machine-learning algorithm to locate each cilium in the tissue. Primary cilia in different cortical neurons and astrocytes in all 6 layers of the cerebral cortex were reconstructed from this data set. These cilia annotations and the rest of this human connectomic data are available at Connectomic Data: <https://h01-release.storage.googleapis.com/landing.html>.

Analysis of primary cilia in neurons and glia of all cerebral cortical layers

We randomly identified 50 or more non-spiny local interneurons (INs) and pyramidal-shaped spiny projection neurons (PNs) with primary cilia in each layer (Table S1). The primary cilium of each neuron was confirmed by its characteristic microtubule-based axoneme and basal body anchor. We then analyzed (1) if there are neuron type-specific differences in the structure, shape, content, and cell domain origin of the primary cilium, (2) a complete census of all the cellular contacts ['contactome'] made by individual neuronal cilium, (3) if these neuronal cilia form specific, distinguishable, functional contacts ['connectome'] with other neurons (e.g., with axons, dendritic spines, synapses, and AIS [axon initial segments]), glia (i.e., astrocytes, oligodendroglia, and microglia), or blood vessels in the environment, (4) the distinguishing features and diversity of primary cilium-neuron or glial contact sites, and (5) the neuronal layer and subtype-specific characteristics of primary cilia. Further, we examined the contactome and connectome of astroglial primary cilia in every cortical layer. Due to the complex morphology of astrocytes, astrocyte cilia were identified by first locating the basal bodies in these cells and then tracing the microtubule (MT) based cilia emanating from them. The following number of neurons and glia were analyzed: Interneurons (Layer I-50, Layer II-63, Layer III-59, Layer IV-57, Layer V-51, Layer VI-12); Projection neurons (Layer II-55, Layer III-55, Layer IV-50, Layer V-50, Layer VI-98); Astrocytes (Layer I-38, Layer II-32, Layer III-32, Layer IV-33, Layer V-31, Layer VI-36).

If >75 % of the length of ciliary surface was smooth or beaded, it was binned as smooth or beaded cilium, respectively. If a cilium had both membrane characteristics and neither (smooth or beaded) covered at least 75% of its length, it was binned as a mixed type. A cilium was considered to be near an axon or a dendrite if the ciliary base is less than 5 µm from the base of the axon/dendrite. For cilia orientation, a cilium is classified as "toward pia" or "toward white matter" if it is angled at 45 degrees or greater away from the cell soma in the direction of pia or white matter, respectively. Otherwise, its direction is noted as 'into the cortical wall'. Cell fragments that were less than 10µm in size could not be classified (for example, as axon, dendrite, soma, or glia, etc.) and were binned as undefined cells. Organelles such as lysosomes were defined as described in Barral et al.⁸⁵ Lysosomes range in size from 200 to 600 nm are often electron dense and have membrane whorls.

As a control for the specificity of neuronal primary ciliary interactions with the surrounding milieu, we identified a segment of the axon of the same phenotype (excitatory or inhibitory) that is immediately adjacent to the neuron on the other side of the cilium and of the same length as the respective neuronal cilium. We then mapped all the cells and cellular subcompartments that are in contact with that segment of the axon the same way we did for the neuronal cilium. Patterns of control axonal contacts were then compared with the patterns of neuronal ciliary contacts. Similar quantification was also done for the axon initial segments of a subset of interneurons and projection neurons in layers I and VI, respectively.

Immunohistochemistry

Human cortical neuronal cells (4 weeks *in vitro*) were generated from human neural precursors (GW15-18).^{83,86} The following primary antibodies were used to immunohistochemically label these cells: Arl13B (mouse, 857602, Biolegend), Connexin 36 (rabbit, 710663, ThermoFisher), and neuron-specific class III β-tubulin [Tuj1] (chicken, ab9354, Millipore). Appropriate Cy2, Cy3, or Alexa dye-conjugated secondary antibodies (Jackson ImmunoResearch, Molecular Probes) were used to detect primary antibody binding. DAPI (Sigma-Aldrich, D9542) was used as a nuclear counterstain. 2914 neurons from 8 independent experiments were analyzed. Images were obtained using a Zeiss LSM780 confocal microscope.

QUANTIFICATION AND STATISTICAL ANALYSIS

General statistical analysis

Excel, SAS, GraphPad, or R language were used for data analysis. Statistical significance was determined by two-tailed Student's *t* test for comparisons between two groups and by ANOVA with Tukey-Kramer's test for comparisons among multiple groups. Correlation between groups was tested by Pearson's or Point Biserial correlation.^{4–6,8,86} No *a priori* power analyses were performed. All data are expressed as mean $\pm$ standard error of the mean (SEM). Statistical details, including *p* values, are indicated in the text or figure legends. The number of cells analyzed (*n*) is indicated in the relevant methods sections ([Analysis of primary cilia in neurons and glia of all cerebral cortical layers](#) and [Immunohistochemistry](#)) or the figure legends.