

## Open Access

<https://doi.org/10.48130/grares-0024-0002>

Grass Research 2024, 4: e004

# Leaf senescence in forage and turf grass: progress and prospects

Kangning Zhang<sup>1</sup>, Hongli Xie<sup>1</sup>, Jiangqi Wen<sup>2</sup>, Jing Zhang<sup>3</sup>, Zeng-Yu Wang<sup>1</sup>, Bin Xu<sup>3\*</sup>  and Maofeng Chai<sup>1\*</sup> <sup>1</sup> Key Laboratory of National Forestry and Grassland Administration on Grassland Resources and Ecology in the Yellow River Delta, College of Grassland Science, Qingdao Agricultural University, Qingdao 266109, China<sup>2</sup> Institute for Agricultural Biosciences, Oklahoma State University, Ardmore, Oklahoma 74078-1010, USA<sup>3</sup> College of Agro-grassland Science, Nanjing Agricultural University, Nanjing 210095, China\* Corresponding authors, E-mail: [binxu@njau.edu.cn](mailto:binxu@njau.edu.cn); [chimu2100@126.com](mailto:chimu2100@126.com)

## Abstract

Leaf senescence is a complex biological process regulated by development, phytohormones, and various environmental factors. For forage and turf grasses, controlling leaf senescence can greatly improve forage quality, the amenity of lawn and turf, and the grasses' stress tolerances. Leaf senescence involves a multitude of gene regulation and metabolic changes, including the alteration of chlorophyll metabolism. Here, we summarized the recent progress of studies on leaf senescence in major forage and turf grass species, such as *Medicago truncatula*, *M. sativa*, *Lolium perenne*, *Panicum virgatum*, and *Agrostis stolonifera*, to provide an insight into the development of effective methods for delaying leaf senescence in grass species.

**Citation:** Zhang K, Xie H, Wen J, Zhang J, Wang ZY, et al. 2024. Leaf senescence in forage and turf grass: progress and prospects. *Grass Research* 4: e004 <https://doi.org/10.48130/grares-0024-0002>

## A diagram under leaf senescence

For forage and turf grass species, regulation of leaf senescence has a significant impact on both the biomass yield and quality of forages and the amenity of turfgrasses. Preventing or delaying precocious leaf senescence is important to improve forage yield, enhance forage quality, and maintain a long-term green lawn and turf. Leaves are the primary photosynthetic organs sustaining the energy supply in plants. Senescence is the final stage of leaf growth and development. It involves the degradation of chlorophyll (Chl), carbohydrates, fats, proteins, and nucleic acids and the remobilization of organic and inorganic nutrients to newly developed tissues and storage organs within the same plant<sup>[1]</sup>. In agricultural production, precocious leaf senescence causes a decrease in crop photosynthetic efficiency and yield<sup>[2,3]</sup>.

Multiple endogenous and external factors influence the progression of leaf senescence, such as plant and leaf age, growth and development, plant hormones, light, drought, salt stress, pathogen infection, and herbivore attacks. When the environment is unfavorable for plants, nutrients are transferred from old leaves to new leaves, seeds, and other sink organs (e.g., stolons/rhizomes/shoot crowns) to sustain their survival. Therefore, the progression of leaf senescence is rarely at a constant speed but rather a complex and ever-changing process. The occurrence of various interactions among the developmental processes of leaf senescence, growth of the whole plant, plant reproduction, and environmental adaptation underscores the importance of leaf senescence in the overall life cycle of plants<sup>[4]</sup>.

Color changes in leaves are the most remarkable sign of leaf senescence. With the gradual degradation of Chl and accumulation of carotenoids and anthocyanins, leaf color transitions from green to yellow or red. In the early stages of leaf senescence, the chloroplast membrane is damaged; however,

the chloroplast size remains unchanged, whereas in the later stages, chloroplasts rapidly decrease in size, and the thylakoid membranes disintegrate. Through intracellular autophagy, the damaged chloroplasts collapse and are then transported to vacuoles for further breakdown by hydrolases<sup>[5]</sup>. As the cells progress from senescence to death, they still require energy to sustain their activities. In the early stages of senescence, cellular energy depends on the mitochondrial supply because of Chl degradation and reduced photosynthesis. Therefore, mitochondria remain intact during the later stages of senescence, thereby maintaining their ability to provide energy to cells. In leaves, up to 70% of the total nitrogen content are in chloroplasts. During leaf senescence, proteins in chloroplasts degrade to amino acids that are redistributed to other plant organs to enhance nitrogen use efficiency<sup>[6]</sup>. The selective catabolism of amino and fatty acids supplements  $\alpha$ -ketoglutarate and glutamate in cells, providing a carbon chain for nitrogen remobilization, which facilitates nitrogen transport in aging leaves<sup>[7]</sup>.

From the perspective of molecular genetics, leaf senescence involves chromatin remodeling, DNA transcription, and post-transcriptional, translational, and post-translational fine-tuning mechanisms. For example, the regulation of chromatin encompasses nucleosome assembly, disassembly, and rearrangement, DNA methylation, and post-translational covalent modifications of histones (including acetylation, ubiquitination, methylation, phosphorylation, and overall acylation), which together influence the gene expression of specific or groups of senescence-associated genes (SAGs)<sup>[8–12]</sup>.

## External factors impacting grass leaf senescence

For grasses used for forage and turf purposes, both external abiotic (e.g., light, atmosphere, temperature, soil, water, and

mowing, etc.) and biotic factors (e.g., animal feeding, tramping, pathogen, and insect attacks) impact the progress of leaf senescence.

## Light

For plants, light serves as both an energy source and a crucial signal for growth and stress response<sup>[13]</sup>. Dense planting or shading by neighboring plants can alter the intensity and quality of light, leading to changes in plant growth patterns and hastening leaf senescence<sup>[14]</sup>. Prolonged exposure to low-light conditions has a negative impact on turfgrass, inducing leaf senescence, and ultimately affecting plant growth and overall turf quality<sup>[15]</sup>.

## Atmospheric pollution and elevated CO<sub>2</sub> level

Tropospheric ozone, also known as ground-level ozone (O<sub>3</sub>), is a significant air pollutant in the lower boundary layers. Prolonged exposure to high levels of O<sub>3</sub> imposes detrimental effects on photosynthesis and carbon allocation, causing premature leaf senescence and ultimately lower grass yield and quality<sup>[16,17]</sup>. Despite efforts to control anthropogenic emissions, additional sources and factors can also contribute to increasing O<sub>3</sub> levels<sup>[18]</sup>. For example, increased O<sub>3</sub> exposure triggers senescence in *Briza maxima* leaves, resulting in a higher ratio of senescent to green biomass, increased fiber content (particularly lignin concentration), and nutritive value for herbivores<sup>[19]</sup>. Alfalfa (*Medicago sativa*) varieties with a slower rate of leaf senescence were found to have greater tolerance to O<sub>3</sub><sup>[20]</sup>.

An increase in atmospheric CO<sub>2</sub> is the primary cause of global climate change. Over the past century, the rate of CO<sub>2</sub> accumulation has steadily increased because of land cover changes and fossil fuel combustion<sup>[21]</sup>. Elevated CO<sub>2</sub> (eCO<sub>2</sub>) could have positive effects on plants, including an increased C:N ratio, production of secondary metabolites, and higher phytohormone concentrations<sup>[22]</sup>. eCO<sub>2</sub> extends the lifespan of forages during autumn, resulting in an extended growing season and higher biomass yield<sup>[23,24]</sup>. However, eCO<sub>2</sub> also adversely affects the nutritional quality of crops and seeds<sup>[22]</sup>. Similar changes have been observed in grasses, whereby increased levels of CO<sub>2</sub> enhance grass productivity at the cost of decreased nutritional quality of the grasses with higher lignin but lower protein contents<sup>[25]</sup>.

## High temperature

Greenhouse gas emissions are increasing the average air temperature such that the global mean annual temperatures are predicted to rise 0.3–4.8 °C by the year 2,100 with more frequent and severe heat waves<sup>[26]</sup>. Elevated temperature benefits the early growth stages of annual crops; however, it also leads to early leaf senescence before the final harvest. For example, the increased temperatures enhance leaf photosynthesis and the above-ground biomass of reed canary grass (*Phalaris arundinacea*) in the early growing stage but eventually result in early senescence and lower biomass at the end of the growing season<sup>[27,28]</sup>. Furthermore, severe heat stress cause Chl degradation and severe membrane lipid peroxidation, accelerating leaf senescence<sup>[29,30]</sup>.

## Soil water content

Limited soil water availability is the main cause of drought stress on plants<sup>[31]</sup>. Grasses grown under drought conditions show a rapid decline in fresh biomass, leaf water potential, and

Chl levels, which is normally coupled with early initiation of senescence<sup>[32,33]</sup>. Grasses exhibiting prolonged greenness in arid and semi-arid habitats often display high resilience to drought<sup>[34]</sup>.

Extremely high soil water content often caused by flooding is associated with low soil oxygen content that imposes disruptive stress on most land plants<sup>[35]</sup>. For lowland-grown grasses, the recovery ability from submergence is directly associated with the maintenance of green photosynthetically active leaves. Water-logging-intolerant grasses, such as *Dactylis glomerata* L. and *Bromus catharticus* Vahl, exhibit premature leaf senescence and persistent partial stomatal closure during recovery, whereas flood-tolerant species *Phalaris aquatica* L. maintains leaf greenness and stomatal conductance during recovery<sup>[36]</sup>.

## Saline-alkali and toxic metal

The worldwide area of saline-alkali land is approximately 0.34 × 10<sup>9</sup> ha. Soil salinization reduces soil fertility, adversely affects plant productivity, growth, and development and damages agricultural productivity and ecosystem diversity<sup>[37]</sup>. Premature leaf yellowing is a significant symptom in *M. truncatula* under salt or alkali stress<sup>[38,39]</sup>.

High toxic heavy metal content in top soils due to mining, exhaustion emissions, sewage irrigation, and other activities is another plant abiotic stress factor<sup>[40]</sup>. When *Miscanthus × giganteus* is planted in soil contaminated with Pb, Cu, or Zn, its leaves show decreases in the photosynthetic rate and total photosynthetic capacity, noticeable precocious leaf senescence, and reduction in the plant's growth rate and biomass yield<sup>[41]</sup>.

## Fungi and bacteria

Plants, along with their associated microbiomes, operate as intricate, multispecies organisms. These associations give them mutual benefits in most cases; however, once pathogenic microbes are dominant, plants often die fast or senesce early<sup>[42]</sup>. For example, the necrotrophic pathogen *Phoma medicaginis* var. *medicaginis* causes alfalfa to develop black spring stems and leaf spots. The symptoms of this infection include black spots on the leaves that later turn yellow<sup>[43]</sup>. In some cases, certain symbiotic bacteria like *Clavicipitaceae* (specifically, *Claviceps purpurea*) form mutualistic relationships with drunken horse grass (*Achnatherum inebrians*). These microbes can prolong the life cycle of leaves, enhance the activity of enzymes, such as superoxide dismutase, peroxidase, and catalase, and augment proline contents. Additionally, these bacteria reduce malondialdehyde contents while increasing the Chl content and photosynthetic rate in leaves<sup>[44]</sup>. Additionally, treating alfalfa with exogenous PopW, a harpin protein derived from *Ralstonia solanacearum*, improves its drought tolerance with altered endogenous hormone levels and expression of genes related to drought stress and leaf senescence<sup>[45]</sup>.

## Turf and grassland management

Effective management techniques, such as fertilization, irrigation, mowing, and grazing management, can significantly postpone the leaf senescence of forage and turf grasses<sup>[46–50]</sup>. For example, removing switchgrass panicles can help the grass maintain higher Chl content, photosynthetic efficiency, and activity of key enzymes, such as phosphoenolpyruvate carboxylase and ribulose-1,5-diphosphate carboxylase. This process also increases the content of important plant hormones, like

## Review of leaf senescence in forage and turf grass

zeatin nucleosides, gibberellins, and indoleacetic acid in flag leaves, thereby delaying leaf senescence<sup>[51]</sup>.

## Leaf SAGs in forage grasses

In recent years, a number of leaf SAGs were cloned and functionally characterized in forage and turf grass. Here, we summarize SAGs cloned from two leguminous and three grass species in the Poaceae family.

### Medicago truncatula


Alfalfa is an allogamous, autotetraploid plant with a complex genetic background. *M. truncatula* is a diploid, self-fertile plant closely related to alfalfa and has been used as the leguminous model plant. According to the comparative transcriptomic data, a total of 545 SAGs were identified from naturally senescent leaves<sup>[52,53]</sup> and up to 1546 SAGs from salt stress-, alkali stress-, and dark-induced leaf senescence<sup>[38,39]</sup>. During leaf senescence, Chl degradation is coupled with the remobilization of nutrients in leaves. The deletion of the gene *STAYGREEN* (*SGR*, encoding a Chl  $\alpha$  catabolic enzyme) in *M. truncatula* enables the plant to maintain its green leaves and retain a significant amount of Chl during senescence. Furthermore, the induced expression of the *MtSGR* gene has been observed in leaf senescence caused by various stresses<sup>[38,39,53,54]</sup>.

In *M. truncatula*, mutation of the tyrosyl-DNA phosphodiesterase 1 (*MtTdp1 $\alpha$* ) gene leads to early leaf senescence. The *MtTdp1* protein has two isoforms, *MtTdp1 $\alpha$*  (*Medtr7g050860*) and *MtTdp1 $\beta$*  (*Medtr8g095490*)<sup>[55]</sup>. Transcriptome analysis reveals that the *MtTdp1 $\alpha$*  downstream genes encompass multiple SAGs, including the *MtSGR* and *SAG101* (an orthologue of *Arabidopsis SENESCENCE-RELATED GENE 1* [*SRG1*]). Silencing the *MtTdp1 $\alpha$*  gene leads to an upregulation of proteolytic genes and genes encoding different subtypes of glutathione S-transferase, as well as the concurrent downregulation of genes involved in photosynthesis. Furthermore, *MtTdp1 $\alpha$*  plays a role in maintaining genomic integrity<sup>[56]</sup>. In the transcriptome induced by darkness, the expression of *MtTdp1 $\alpha$*  is inhibited<sup>[53]</sup>. The decreased expression of *MtTdp1 $\alpha$*  caused by darkness weakens the repair function of the genome and promotes the expression of senescence-related genes and leaf senescence (Fig. 1).

### Alfalfa (*Medicago sativa*)

Alfalfa is an important leguminous forage crop cultivated worldwide. The biomass and quality of alfalfa can be affected by leaf senescence. Yuan et al. conducted a transcriptome analysis on mature and senescent leaves and identified 1,250 differentially expressed genes, of which 713 were upregulated, and 537 were downregulated<sup>[57]</sup>. Calderini et al. constructed a vector containing the *SAG12* promoter and a cytokinin biosynthetic gene (*IPT*) that encodes an isopentenyl transferase. This vector was then used to genetically modify alfalfa and inhibit leaf senescence<sup>[58]</sup>. The results revealed that senescence occurs at a slower rate in alfalfa *SAG12-IPT* plants, as indicated by higher Chl levels.

Alfalfa transgenic plants with *SGR* gene silenced by RNA interference (RNAi) retained over 50% of their Chl content during senescence and higher crude protein content<sup>[54]</sup>. Another study showed that *MtTMT* expression is upregulated in the dark. Overexpression of  $\gamma$ -tocopherol methyltransferase (*MtTMT*) to increase their  $\alpha$ -tocopherol contents also delayed



**Fig. 1** Regulation mechanism of leaf senescence in *Medicago truncatula*.


leaf senescence and increased crude protein content in transgenic alfalfa<sup>[59]</sup>.

*MicroRNA156* plays a crucial role in enhancing alfalfa tolerance to abiotic stress. At least seven *MsSPL* (SQUAMOSA promoter binding protein-like) genes are regulated by *miR156* through transcript cleavage. For example, the *miR156/SPL* module is involved in drought- and flooding-induced leaf senescence; when *miR156* is overexpressed, or its target gene *SPL13* is silenced, the onset of leaf senescence caused by flooding is delayed<sup>[60]</sup>. Similarly, overexpression of *miR156* leads to increased drought resistance in alfalfa, whereas silencing its target gene *SPL9* delays drought-induced leaf senescence<sup>[61]</sup>. Thus, the *miR156/SPL* module may play multiple regulatory roles in stress responses, including stress-induced leaf senescence.

Furthermore, *MsSAG113*, a negative regulator of the abscisic acid (ABA) signaling, is controlled by various hormones, such as ABA, salicylic acid, and methyl jasmonate (MeJA). Overexpression of *MsSAG113* can accelerate the senescence of dark-induced and aged alfalfa leaves, with its downstream genes mostly related to hormone-regulated genes. Therefore, *MsSAG113* may play a crucial role in leaf senescence and the hormone regulation network of alfalfa. *MsC3H-39* is a CCCH-like zinc-finger protein that directly recognizes and binds to the upstream region of *MsSAG113*, enhancing its expression. Additionally, the transient expression of *MsC3H-39* promotes alfalfa leaf senescence<sup>[62]</sup>, which implies that *MsC3H-39* directly regulates *MsSAG113* expression and contributes to leaf senescence (Fig. 2).

### Perennial ryegrass (*Lolium perenne*)

Hormones play crucial roles in regulating leaf senescence in perennial ryegrass. Application of exogenous 6-benzylaminopurine increases the activity of antioxidant enzymes such as superoxide dismutase, catalase, ascorbate peroxidase, monodehydroascorbate reductase, and glutathione reductase and inhibits the accumulation of  $\text{Na}^+$  in cells, delaying salt-induced leaf senescence<sup>[63]</sup>. Similarly, exogenous application of melatonin enhances the activity and transcription levels of superoxide dismutase and catalase, activating the superoxide dismutase enzymatic antioxidant pathway and reducing the



**Fig. 2** Regulation mechanism of leaf senescence in *Medicago sativa* L.

transcription of genes associated with Chl degradation (*LpSGR*, *LpNYC1*, *LpNOL*, *LpPPH*, *LpPAO*, and *LpRCCR1*) and senescence markers (*LpSAG12.1*, *Lph36*, and *Lpl69*). Up- or down-regulation of Chl catabolic genes, such as *LpNYC1*, *LpNOL*, *LpSGR*, and *LpPPH*, may accelerate or suppress leaf senescence<sup>[64–66]</sup>. Moreover, melatonin inhibits dark-induced leaf senescence by downregulating Chl degradation<sup>[67]</sup>. Additionally, exogenous application of melatonin upregulates the expression of genes involved in cytokinin biosynthesis (*LpIPT2* and *LpOG1*) and their signal response transcription factors (type-B ARR), downregulates the expression of genes involved in ABA biosynthesis (*LpZEP* and *LpNCED1*) and signal transduction (*LpABI3* and *LpABI5*), increases the content of endogenous cytokinins, and reduces the ABA content to alleviate heat stress-induced leaf senescence<sup>[68]</sup>. The effect of strigolactone on leaf senescence in grass has received little attention; however, exogenous application of a strigolactone analog (GR24) accelerates the downregulation of genes related to Chl biosynthesis (*LpHEMA*, *LpGSA*, *LpHEMB*, and *LpCHLH*) and promotes the upregulation of genes associated with senescence (*LpSAG12.1*, *Lph36*, and *Lpl69*) and Chl degradation (*LpNOL*, *LpPPH*, *LpPAO*, and *LpRCCR1*) to promote dark-induced leaf senescence<sup>[69]</sup> (Fig. 3).

In the Chl catabolic pathway, the *LpSGR* gene, which is responsible for magnesium dechelation of Chl *a*, is highly expressed in leaves undergoing senescence due to development or darkness. This gene plays a crucial role in Chl degradation, consequently affecting various metabolic processes. In perennial ryegrass, RNAi targeting *LpSGR* leads to the inhibition of Chl degradation, increasing Chl content and enhancing photochemical efficiency in senescent leaves<sup>[70]</sup>. However, when subjected to heat stress, the suppression of *LpSGR* results in the downregulation of genes encoding photosystem proteins and ROS-scavenging enzymes and the upregulation of genes encoding ROS-generating enzymes, thereby accelerating leaf senescence in *LpSGR*-RNAi lines<sup>[71]</sup>.


Similarly, the Chl *b* reductases, non-yellow coloring 1 (NYC1)-like (NOL) and NYC1, locate upstream of SGR, whereas pheophytinase (PPH), pheophorbide a oxygenase (PAO), and red Chl catabolite reductase (RCCR) are enzymes downstream of SGR in the Chl catabolic pathway. Previous studies have

demonstrated that *LpNYC1* is involved in the ABA and ethylene signaling pathways, which can be directly regulated by *LpABI5*, *LpABF3*, and *LpEIN3*<sup>[72]</sup>. Decreased *LpNOL* expression or the addition of exogenous Chl promotes Chl accumulation, exhibits an antioxidant effect during heat stress, and delays leaf senescence<sup>[73–75]</sup>. *LpPPH* overexpression accelerates Chl degradation. Additionally, *LpPPH*-mediated Chl breakdown may be positively regulated by ABA and ethylene but negatively regulated by cytokinin<sup>[76]</sup>. Overexpression of *LpNAL* leads to a phenotype characterized by delayed leaf senescence or retention of green color, whereas knocking down *LpNAL* using RNAi accelerates leaf senescence. Furthermore, transcriptome analysis has revealed that *LpNAL* plays a role in photosynthesis, antioxidant metabolism, ABA and ethylene synthesis and signaling, and Chl catabolism (Fig. 4).

### Switchgrass (*Panicum virgatum*)

Switchgrass is a warm-season perennial grass species utilized as both forage for livestock and biofuel feedstock. Improving nitrogen (N) remobilization from aboveground to underground organs during annual shoot senescence is an important goal for the sustainable production of switchgrass as a biofuel crop. Transcriptome and proteome analysis has identified genes that are differentially expressed in various organs during development, such as those involved in protein degradation, nitrogen remobilization, transcriptional processes, and photosynthesis-related proteins<sup>[77]</sup>. During senescence, the increase in total N content in underground organs is consistent with the decrease in N content in aboveground organs. The contents of 14-3-3-like proteins and glutathione S-transferase proteins in late-senescent leaves are significantly higher than those in early-senescent leaves, indicating that a higher abundance of these proteins might delay leaf senescence<sup>[78]</sup>.

An in-depth understanding of mechanisms controlling mineral uptake, distribution, and remobilization is important for sustainable mineral production. A previous study identified genes associated with mineral transporters in switchgrass and observed relative changes in transporter expression during different growing seasons. Genes from the same family exhibit specific expressions at different stages. Moreover, members of the MRS2/MGT family are responsible for magnesium transport;

*Lolium perenne* L.**Fig. 3** Mechanism of exogenous hormone application regulating leaf senescence of *Lolium perenne* L.


however, the *MRS2/MGT* genes expressed during the active shoot growth phase (Pavirv00000269 and Pavirv00025330) differ from those expressed during early senescent (Pavirv00023490 and Pavirv00047094) and fully senescent (Pavirv00069828 and Pavirv00014024) stages<sup>[79]</sup>.

Transcriptome analyses of switchgrass flag leaves revealed the enrichment of transporters during senescence, as well as dynamic changes in other genes and related metabolites during senescence, emphasizing the role of NAC transcription factors in remobilization<sup>[80]</sup>. In another study, Yang et al. found that the transcription level of *PvNAC1* was upregulated by darkness and age<sup>[81]</sup>. Additionally, *PvNAC1* overexpression significantly increased *AtSAG12* expression, protein degradation, and nitrogen cycle-related genes, significantly reduced Chl concentration, and promoted Arabidopsis leaf senescence. Conversely, *PvNAC2* overexpression in switchgrass resulted in increased aboveground biomass, accompanied by increased transcript levels of the nitrogen metabolism gene *PvGDH* in leaves and the nitrate and ammonium transporter genes *PvNRT1*, *PvNRT2*, and *PvAMT2* in roots. In addition to the NAC family, the WRKY family is also involved in switchgrass leaf senescence. According to an analysis of the WRKY family in switchgrass, the expression levels of 23 genes are increased when flag leaves begin to senesce<sup>[82]</sup>.

CCCH-type zinc-finger proteins have been implicated in multiple biological processes and environmental responses in plants. In this context, a switchgrass CCCH-type zinc-finger protein, known as PvSSG, was found to inhibit the binding efficiency of PvNAP1 and PvNAP2 to DNA through protein–protein interactions. Additionally, PvSSG reduces the activation of downstream genes related to Chl degradation by PvNAP1 and PvNAP2 and exhibits inhibitory effects on leaf senescence<sup>[83]</sup> (Fig. 5).


**Creeping bentgrass (*Agrostis stolonifera*)**

Creeping bentgrass is a cool-season grass species that actively grows during spring and fall when temperatures range from 15°C to 24°C and declines during hot summer. The decline in turf quality during the summer poses a significant problem for creeping bentgrass in areas with warm climates, particularly in transition zones<sup>[30]</sup>. Moreover, heat stress reduces Chl content, protease activity, amino acid and total protein content, and antioxidant activity in creeping bentgrass leaves

**Fig. 4** Regulation mechanism of leaf senescence in *Lolium perenne* L.

while increasing membrane lipid peroxidation<sup>[30,84,85]</sup>. Leaf senescence induced by heat stress is negatively correlated with ethylene and ABA accumulation but positively correlated with cytokinin production<sup>[86]</sup>. The application of exogenous cytokinins decreases the heat-induced membrane lipid peroxidation and protease activity and maintains higher leaf Chl content, Fv/Fm, soluble protein content, and antioxidant enzyme activity. The application of cytokinins also delays leaf senescence and improves turf quality under heat stress<sup>[84,87–89]</sup>. Several other exogenous reagents have a similar effect on delaying leaf senescence (Table 1).

Cytokinin can effectively inhibit leaf senescence. To obtain transgenic plants with stay green phenotype, researchers transferred the Isopentenyl transferase (*IPT*) gene, which is responsible for catalyzing a crucial step in *de novo* cytokinin biosynthesis, into creeping bentgrass. These transgenic plants can delay leaf senescence induced by shade, heat, and drought stress<sup>[90,91]</sup>. Under stressful conditions, the production of cytokinins in transgenic lines increases, which can help maintain high Chl contents. This positive effect also extends to osmotic regulation, antioxidant enzyme activity, protein abundance, metabolic accumulation, and the preservation of high photosynthesis levels<sup>[92–95]</sup>. Transcriptomic and proteomic



**Fig. 5** Regulation mechanism of leaf senescence in *Panicum virgatum*.

**Table 1.** Mechanism of exogenous application of various reagents to inhibit leaf senescence of creeping glumes.

Reagents	Treatment	Effects	Ref.
Aminoethoxyvinylglycine	Heat	Regulating chlorophyll metabolic activities	[88, 111]
Protease inhibitors in the cysteine, serine, or aspartic classes		Suppressing proteolysis	[112]
Glutamate		Suppressing chlorophyll degradation. Activating amino acid metabolism involved in energy production, antioxidant defense, and nitrogen balance	[113]
$\beta$ -sitosterol		Alleviating the membrane lipid peroxidation and the enzymatic antioxidant system	[114]
Chitosan		Up-regulating chlorophyll biosynthesis related genes. Down-regulating chlorophyll degradation and senescence related genes	[115]
Spermine		Increasing the activity of antioxidant enzymes Maintaining higher chlorophyll content, net photosynthetic rate, photochemical efficiency, and performance index on absorption basis. Promoting osmotic adjustment ability and antioxidant enzyme activities to enhance the scavenging capacity of reactive oxygen species. Upregulating transcriptions of heat shock protein genes helping to maintain normal synthesis and functions of proteins	[116]
Biostimulants (TurfVigor and CPR)		Promoting both shoot and root growth	[117]
Trinexapac-ethyl		Improving the chlorophyll content and photosynthetic rate	[117]
$\gamma$ -aminobutyric acid, proline, and ammonium nitrate		Enhancing chlorophyll content and photochemical efficiency. Suppressing chlorophyll-degrading enzyme activities under heat stress	[118]
Nitrogen		Enhancing protein abundance in photosynthesis and amino acid metabolism and stress defense systems (heat shock protection and antioxidants)	[89]
Zeatin riboside		Reducing heat-induced membrane lipid peroxidation Reducing protease activity. Maintaining higher leaf chlorophyll content, Fv/Fm, soluble protein content and antioxidant enzyme activity	[84, 87–89]
$\gamma$ -Aminobutyric acid	Drought	Promoting energy production and conversion, antioxidant defense, and DHN3 accumulation	[119]
Melatonin		Increasing photochemical efficiency, chlorophyll content and relative water content. Suppressing leaf electrolyte leakage, lipid peroxidation and hydrogen peroxide production. Up-regulating the cytokinin-signaling and synthesis genes. Down-regulating the chlorophyll-degradation genes and enzyme activities	[120]

analyses have been used to identify genes encoding proteins related to energy production, metabolism, stress defense, signalling, protein synthesis and transport, and membrane transport, which may play significant roles in the overexpression of *IPT* in creeping bentgrass<sup>[96]</sup>.

Nitrogen deficiency inhibits plant growth and induces leaf senescence by regulating multiple metabolic processes. Specifically, nitrogen deficiency increases the activity of glutamine

synthetase transferase (glutamate dehydrogenase) and decreases the content of amino acids and soluble proteins in leaves<sup>[97, 98]</sup>. However, transgenic lines overexpressing *IPT* can still inhibit leaf senescence induced by nitrogen or phosphorus deficiency<sup>[99]</sup>. Therefore, enhancing multiple stress resistance in grasses through exogenous hormones or regulating hormone-related genes may be one of the most efficient methods of delaying senescence.

Furthermore, the membrane proteins involved in ATP metabolism, light harvesting, and photosynthetic photochemical reactions, as well as proteins related to the efficient processing of photorespiratory products and reactive oxygen species, may play important roles in regulating leaf senescence in creeping bentgrass under heat stress<sup>[100]</sup>. The main reason for the heat-induced decrease in leaf Chl contents is the expression of Chl degradation-related genes and increased chlorophyll degradation enzyme activity<sup>[101]</sup>. For example, when *SGR* is silenced in creeping bentgrass, transgenic plants display a green phenotype<sup>[102]</sup>.

### Other species

Research on leaf senescence has also been conducted on other grass species. Studies have investigated the transcriptome analysis of mature and senescent leaves in red clover (*Trifolium pratense* L.) and the heat-induced leaf senescence transcriptome of tall fescue (*Festuca arundinacea*) using RNA-Seq<sup>[103,104]</sup>. Similarly, transcriptome analysis of dark-induced leaf senescence has been reported in Bermudagrass (*Cynodon dactylon* L.)<sup>[15]</sup>. *Zoysia japonica* has been studied to understand the aging process induced by factors such as age, darkness, and salt, using RNA sequencing<sup>[105]</sup>. Several genes, including *ZjPPH*<sup>[106]</sup>, *ZjNOL*<sup>[107,108]</sup>, *ZjSGR*<sup>[109]</sup> and *ZjNYC1*<sup>[110]</sup>, have been successfully cloned in *Zoysia japonica* and overexpressed in *Arabidopsis* to investigate their roles in senescence.

For convenience, Table 2 presents a comprehensive list of the genes discussed in this paper.

### Conclusions and prospects

Premature leaf senescence reduces the amenity of turf and significantly affects the biomass yield and feedstock quality of forage grasses. The stay-green genotypes have been successfully identified in species like ryegrass and alfalfa and have been integrated into breeding programs to minimize yield losses in plants grown under unfavorable environmental conditions. Compared to model plant species (e.g., *Arabidopsis*

**Table 2.** List of the gene involved in leaf senescence in grass.

Gene	Species	Effect	Ref.
<i>SGR</i>	<i>Medicago truncatula</i>	Promote	[54]
	<i>Medicago sativa</i>	Promote	[54]
<i>IPT</i>	<i>Lolium perenne</i>	Promote (except heat stress)	[71]
	<i>Agrostis stolonifera</i>	Promote	[102]
<i>Tdp1α</i>	<i>Zoysia japonica</i>	Promote	[109]
	<i>Medicago sativa</i>	Delay	[58]
<i>TMT</i>	<i>Agrostis stolonifera</i>	Delay	[90,91]
	<i>Medicago truncatula</i>	Delay	[56]
<i>miR156</i>	<i>Medicago sativa</i>	Delay	[59]
	<i>Medicago sativa</i>	Delay	[60]
<i>SPL9</i>	<i>Medicago sativa</i>	Promote	[61]
	<i>Medicago sativa</i>	Promote	[62]
<i>SAG113</i>	<i>Lolium perenne</i>	Promote	[72]
	<i>Zoysia japonica</i>	Promote	[110]
<i>NOL</i>	<i>Lolium perenne</i>	Promote	[73,74,75]
	<i>Zoysia japonica</i>	Promote	[107,108]
<i>PPH</i>	<i>Lolium perenne</i>	Promote	[76]
	<i>Zoysia japonica</i>	Promote	[106]
<i>NAL</i>	<i>Lolium perenne</i>	Delay	[73]
	<i>Panicum virgatum</i>	Promote	[81]
<i>NAC1</i>	<i>Panicum virgatum</i>	Promote	[81]
	<i>Panicum virgatum</i>	Delay	[83]
<i>SSG</i>	<i>Panicum virgatum</i>		

and rice), systematic approaches to unravel the molecular mechanisms underlying leaf senescence are still to be carried out in forage grasses, many questions related remain unanswered.

Outstanding future questions:

- (1) What are the specific mechanisms that regulate leaf senescence in forage and turf grasses?
- (2) What are the most effective methods for delaying leaf senescence in these grasses?
- (3) How can the knowledge gained from research on leaf senescence be applied to molecular breeding?

With the identification of more key genes regulating leaf senescence, combined with the CRISPR gene-editing technology, functional 'stay-green' traits will be achieved in forage or turf grass breeding programs for higher quality, higher yield, and improved stress tolerance.

### Author contributions

The authors confirm contribution to the paper as follows: study conception and design: Chai M, Zhang K; data analysis: Wen J, Zhang J, Wang ZY, Chai M; draft manuscript preparation: Zhang K, Xie H, Xu B, Chai M. All authors reviewed the results and approved the final version of the manuscript.

### Data availability

All data generated or analyzed during this study are included in this published article.

### Acknowledgments

This study was supported by the Natural Science Foundation of Shandong Province, China (ZR2023MC120), the National Natural Science Foundation of China (No. 32271755 and No. 31972958) and by National Key Research and Development Program of China (2023YFF1001403).

### Conflict of interest

The authors declare that they have no conflict of interest. Bin Xu is the Editorial Board member of *Grass Research* who was blinded from reviewing or making decisions on the manuscript. The article was subject to the journal's standard procedures, with peer-review handled independently of this Editorial Board member and the research groups.

### Dates

Received 17 December 2023; Revised 9 February 2024; Accepted 13 February 2024; Published online 28 February 2024

### References

1. Koyama T. 2014. The roles of ethylene and transcription factors in the regulation of onset of leaf senescence. *Frontiers in Plant Science* 5:650
2. Zhu K, Tao H, Xu S, Li K, Zafar S, et al. 2019. Overexpression of salt-induced protein (*salT*) delays leaf senescence in rice. *Genetics and Molecular Biology* 42:80–86
3. Jan S, Abbas N, Ashraf M, Ahmad P. 2019. Roles of potential plant hormones and transcription factors in controlling leaf senescence and drought tolerance. *Protoplasma* 256:313–29

4. Woo HR, Kim HJ, Lim PO, Nam HG. 2019. Leaf senescence: systems and dynamics aspects. *Annual Review of Plant Biology* 70:347–76
5. Tamary E, Nevo R, Naveh L, Levin-Zaidman S, Kiss V, et al. 2019. Chlorophyll catabolism precedes changes in chloroplast structure and proteome during leaf senescence. *Plant Direct* 3:e00127
6. Buet A, Costa ML, Martínez DE, Guiamet JJ. 2019. Chloroplast protein degradation in senescing leaves: proteases and lytic compartments. *Frontiers in Plant Science* 10:747
7. Chrobok D, Law SR, Brouwer B, Lindén P, Ziolkowska A, et al. 2016. Dissecting the metabolic role of mitochondria during developmental leaf senescence. *Plant Physiology* 172:2132–53
8. Ostrowska-Mazurek A, Kasprzak P, Kubala S, Zaborowska M, Sobieszcuk-Nowicka E. 2020. Epigenetic landmarks of leaf senescence and crop improvement. *International Journal of Molecular Sciences* 21:5125
9. Wang X, Gao J, Gao S, Song Y, Yang Z, et al. 2019. The H3K27me3 demethylase REF6 promotes leaf senescence through directly activating major senescence regulatory and functional genes in *Arabidopsis*. *PLoS Genetics* 15:e1008068
10. Chen L, Xiang S, Chen Y, Li D, Yu D. 2017. *Arabidopsis* WRKY45 interacts with the DELLA protein RGL1 to positively regulate age-triggered leaf senescence. *Molecular Plant* 10:1174–89
11. Sakuraba Y, Kim D, Han SH, Kim SH, Piao W, et al. 2020. Multilayered regulation of membrane-bound ONAC054 is essential for abscisic acid-induced leaf senescence in rice. *The Plant Cell* 32:630–49
12. An J, Zhang X, Bi S, You C, Wang X, et al. 2019. MdbHLH93, an apple activator regulating leaf senescence, is regulated by ABA and MdBt2 in antagonistic ways. *New Phytologist* 222:735–51
13. Roeber VM, Bajaj I, Rohde M, Schmülling T, Cortleven A. 2021. Light acts as a stressor and influences abiotic and biotic stress responses in plants. *Plant, Cell & Environment* 44:645–64
14. Li Z, Zhao T, Liu J, Li H, Liu B. 2023. Shade-induced leaf senescence in plants. *Plants* 12:1550
15. Fan J, Lou Y, Shi H, Chen L, Cao L. 2019. Transcriptomic analysis of dark-induced senescence in bermudagrass (*Cynodon dactylon*). *Plants* 8:614
16. Emberson LD, Pleijel H, Ainsworth EA, van den Berg M, Ren W, et al. 2018. Ozone effects on crops and consideration in crop models. *European Journal of Agronomy* 100:19–34
17. Dawney L, Mills G. 2009. Relative effects of elevated background ozone concentrations and peak episodes on senescence and above-ground growth in four populations of *Anthoxanthum odoratum* L. *Environmental Pollution* 157:503–10
18. Nguyen DH, Lin C, Vu CT, Cheruiyot NK, Nguyen MK, et al. 2022. Tropospheric ozone and NO<sub>x</sub>: a review of worldwide variation and meteorological influences. *Environmental Technology & Innovation* 28:102809
19. Sanz J, Bermejo V, Muntífering R, González-Fernández I, Gimeno BS, et al. 2011. Plant phenology, growth and nutritive quality of *Briza maxima*: responses induced by enhanced ozone atmospheric levels and nitrogen enrichment. *Environmental Pollution* 159:423–30
20. Renaud JP, Allard G, Mauffette Y. 1997. Effects of ozone on yield, growth, and root starch concentrations of two alfalfa (*Medicago sativa* L.) cultivars. *Environmental Pollution* 95:273–81
21. Sun Z, Wang X, Yamamoto H, Tani H, Nie T. 2020. The effects of spatiotemporal patterns of atmospheric CO<sub>2</sub> concentration on terrestrial gross primary productivity estimation. *Climatic Change* 163:913–30
22. Roy S, Mathur P. 2021. Delineating the mechanisms of elevated CO<sub>2</sub> mediated growth, stress tolerance and phytohormonal regulation in plants. *Plant Cell Reports* 40:1345–65
23. Blumenthal DM, Mueller KE, Kray JA, LeCain DR, Pendall E, et al. 2018. Warming and elevated CO<sub>2</sub> interact to alter seasonality and reduce variability of soil water in a semiarid grassland. *Ecosystems* 21:1533–44
24. Reyes-Fox M, Steltzer H, Trlica MJ, McMaster GS, Andales AA, et al. 2014. Elevated CO<sub>2</sub> further lengthens growing season under warming conditions. *Nature* 510:259–62
25. Abdalla Filho AL, Costa Junior GT, Lima PMT, Soltangheisi A, Abdalla AL, et al. 2019. Fiber fractions, multielemental and isotopic composition of a tropical C<sub>4</sub> grass grown under elevated atmospheric carbon dioxide. *PeerJ* 7:e5932
26. Jagadish SVK, Way DA, Sharkey TD. 2021. Plant heat stress: concepts directing future research. *Plant, Cell & Environment* 44:1992–2005
27. Ge ZM, Kellomäki S, Zhou X, Peltola H, Wang KY, et al. 2012. Seasonal physiological responses and biomass growth in a bioenergy crop (*Phalaris arundinacea* L.) under elevated temperature and CO<sub>2</sub>, subjected to different water regimes in boreal conditions. *BioEnergy Research* 5:637–48
28. Ge ZM, Zhou X, Kellomäki S, Biassi C, Wang KY, et al. 2012. Carbon assimilation and allocation (<sup>13</sup>C labeling) in a boreal perennial grass (*Phalaris arundinacea*) subjected to elevated temperature and CO<sub>2</sub> through a growing season. *Environmental and Experimental Botany* 75:150–58
29. Yang Z, Miao Y, Yu J, Liu J, Huang B. 2014. Differential growth and physiological responses to heat stress between two annual and two perennial cool-season turfgrasses. *Scientia Horticulturae* 170:75–81
30. Xu Q, Huang B. 2004. Antioxidant metabolism associated with summer leaf senescence and turf quality decline for creeping bentgrass. *Crop Science* 44:553–60
31. Faiz MA, Zhang Y, Zhang X, Ma N, Aryal SK, et al. 2022. A composite drought index developed for detecting large-scale drought characteristics. *Journal of Hydrology* 605:127308
32. Guenni O, Romero E, Guédez Y, Macías MP, Infante D. 2016. Survival strategies of *Centroserma molle* and *C. macrocarpum* in response to drought. *Tropical Grasslands-Forrajes Tropicales* 5:1
33. Ahkami AH, Wang W, Wietsma TW, Winkler T, Lange I, et al. 2019. Metabolic shifts associated with drought-induced senescence in *Brachypodium*. *Plant Science* 289:110278
34. Jardine EC, Thomas GH, Osborne CP. 2021. Traits explain sorting of C<sub>4</sub> grasses along a global precipitation gradient. *Ecology and Evolution* 11:2669–80
35. Bates PD. 2022. Flood inundation prediction. *Annual Review of Fluid Mechanics* 54:287–315
36. Striker GG, Ploschuk RA. 2018. Recovery from short-term complete submergence in temperate pasture grasses. *Crop and Pasture Science* 69:745–53
37. Zhang K, Chang L, Li G, Li Y. 2023. Advances and future research in ecological stoichiometry under saline-alkali stress. *Environmental Science and Pollution Research* 30:5475–86
38. Dong S, Pang W, Liu Z, Li H, Zhang K, et al. 2022. Transcriptome analysis of leaf senescence regulation under alkaline stress in *Medicago truncatula*. *Frontiers in Plant Science* 13:881456
39. Dong S, Sang L, Xie H, Chai M, Wang ZY. 2021. Comparative transcriptome analysis of salt stress-induced leaf senescence in *Medicago truncatula*. *Frontiers in Plant Science* 12:666660
40. He H, Li Y, He L. 2018. The central role of hydrogen sulfide in plant responses to toxic metal stress. *Ecotoxicology and Environmental Safety* 157:403–08
41. Andrejić G, Šinčar-Sekulić J, Prica M, Dželatović Ž, Rakić T. 2019. Phytoremediation potential and physiological response of *Miscanthus × giganteus* cultivated on fertilized and non-fertilized flotation tailings. *Environmental Science and Pollution Research* 26:34658–69
42. Harman G, Khadka R, Doni F, Uphoff N. 2020. Benefits to plant health and productivity from enhancing plant microbial symbionts. *Frontiers in Plant Science* 11:610065
43. Fan Q, Creamer R, Li Y. 2018. Time-course metabolic profiling in alfalfa leaves under *Phoma medicaginis* infection. *PLoS One* 13:e0206641

## Review of leaf senescence in forage and turf grass

44. Zhang H, Li X, White JF, Wei X, He Y, et al. 2022. *Epichloë* endophyte improves ergot disease resistance of host (*Achnatherum inebrians*) by regulating leaf senescence and photosynthetic capacity. *Journal of Plant Growth Regulation* 41:808–17

45. Demirkol G. 2021. PopW enhances drought stress tolerance of alfalfa via activating antioxidative enzymes, endogenous hormones, drought related genes and inhibiting senescence genes. *Plant Physiology and Biochemistry* 166:540–48

46. Lü X, Reed SC, Yu Q, Han X. 2016. Nutrient resorption helps drive intra-specific coupling of foliar nitrogen and phosphorus under nutrient-enriched conditions. *Plant and Soil* 398:111–20

47. Liu D, Song C. 2008. Responses of marsh wetland plant *Calamagrostis angustifolia* to exogenous nitrogen input. *Ying Yong Sheng Tai Xue Bao* 19:2599–604

48. Farias LD, Zanine AD, Ferreira DD, Ribeiro MD, de Souza AL, et al. 2019. Effects of nitrogen fertilization and seasons on the morphogenetic and structural characteristics of Piata (*Brachiaria brizantha*) grass. *Revista De La Facultad De Ciencias Agrarias* 51:42–54

49. Magalhães JA, Socorro de Souza Carneiro M, Andrade AC, de Andrade AP, Bakke OA, et al. 2013. Morphogenetic and structural characteristics of andropogon grass under irrigation and fertilization. *Semina-Ciencias Agrarias* 34:2427–35

50. Maranhao SR, Pompeu RCF, de Araujo RA, Lopes MN, Cândido MJD, et al. 2021. Morphophysiology of tropical grasses under different water supply in two growing seasons: II. BRS Massai and BRS Tamani grasses. *Semina: Ciências Agrárias* 42:301–18

51. Zhao C, Yue Y, Wu J, Scullion J, Guo Q, et al. 2023. Panicle removal delays plant senescence and enhances vegetative growth improving biomass production in switchgrass. *Biomass and Bioenergy* 174:106809

52. De Michele R, Formentin E, Todesco M, Toppo S, Carimi F, et al. 2009. Transcriptome analysis of *Medicago truncatula* leaf senescence: similarities and differences in metabolic and transcriptional regulations as compared with *Arabidopsis*, nodule senescence and nitric oxide signalling. *New Phytologist* 181:563–75

53. Mahmood K, Torres-Jerez I, Krom N, Liu W, Udvardi MK. 2022. Transcriptional programs and regulators underlying age-dependent and dark-induced senescence in *Medicago truncatula*. *Cells* 11:1570

54. Zhou C, Han L, Pislaru C, Nakashima J, Fu C, et al. 2011. From model to crop: functional analysis of a STAY-GREEN gene in the model legume *Medicago truncatula* and effective use of the gene for alfalfa improvement. *Plant Physiology* 157:1483–96

55. Macovei A, Balestrazzi A, Confalonieri M, Carbonera D. 2010. The tyrosyl-DNA phosphodiesterase gene family in *Medicago truncatula* Gaertn.: bioinformatic investigation and expression profiles in response to copper- and PEG-mediated stress. *Planta* 232:393–407

56. Donà M, Confalonieri M, Minio A, Biggiogera M, Buttafava A, et al. 2013. RNA-Seq analysis discloses early senescence and nucleolar dysfunction triggered by *Tdp1α* depletion in *Medicago truncatula*. *Journal of Experimental Botany* 64:1941–51

57. Yuan J, Sun X, Guo T, Chao Y, Han L. 2020. Global transcriptome analysis of alfalfa reveals six key biological processes of senescent leaves. *PeerJ* 8:e8426

58. Calderini O, Bovone T, Scotti C, Pupilli F, Piano E, et al. 2007. Delay of leaf senescence in *Medicago sativa* transformed with the *ipt* gene controlled by the senescence-specific promoter SAG12. *Plant Cell Reports* 26:611–15

59. Jiang J, Jia H, Feng G, Wang Z, Li J, et al. 2016. Overexpression of *Medicago sativa* TMT elevates the  $\alpha$ -tocopherol content in *Arabidopsis* seeds, alfalfa leaves, and delays dark-induced leaf senescence. *Plant Science* 249:93–104

60. Feyissa BA, Amyot L, Nasrollahi V, Papadopoulos Y, Kohalmi SE, et al. 2021. Involvement of the miR156/SPL module in flooding response in *Medicago sativa*. *Scientific Reports* 11:3243

61. Hanly A, Karagiannis J, Lu QSM, Tian L, Hannoufa A. 2020. Characterization of the role of *SPL9* in drought stress tolerance in *Medicago sativa*. *international Journal of Molecular Sciences* 21:6003

62. Li S, Xie H, Zhou L, Dong D, Liu Y, et al. 2022. Overexpression of MsSAG113 gene promotes leaf senescence in alfalfa via participating in the hormone regulatory network. *Frontiers in Plant Science* 13:1085497

63. Ma X, Zhang J, Huang B. 2016. Cytokinin-mitigation of salt-induced leaf senescence in perennial ryegrass involving the activation of antioxidant systems and ionic balance. *Environmental and Experimental Botany* 125:1–11

64. Zhang J, Xing J, Lu Q, Yu G, Xu B, et al. 2019. Transcriptional regulation of chlorophyll-catabolic genes associated with exogenous chemical effects and genotypic variations in heat-induced leaf senescence for perennial ryegrass. *Environmental and Experimental Botany* 167:103858

65. Zhang J, Li H, Jiang Y, Li H, Zhang Z, et al. 2020. Natural variation of physiological traits, molecular markers, and chlorophyll catabolic genes associated with heat tolerance in perennial ryegrass accessions. *BMC Plant Biology* 20:520

66. Gan L, Han L, Yin S, Jiang Y. 2020. Chlorophyll metabolism and gene expression in response to submergence stress and subsequent recovery in perennial ryegrass accessions differing in growth habits. *Journal of Plant Physiology* 251:153195

67. Zhang J, Li H, Xu B, Li J, Huang B. 2016. Exogenous melatonin suppresses dark-induced leaf senescence by activating the superoxide dismutase-catalase antioxidant pathway and down-regulating chlorophyll degradation in excised leaves of perennial ryegrass (*Lolium perenne* L.). *Frontiers in Plant Science* 7:1500

68. Zhang J, Shi Y, Zhang X, Du H, Xu B, et al. 2017. Melatonin suppression of heat-induced leaf senescence involves changes in abscisic acid and cytokinin biosynthesis and signaling pathways in perennial ryegrass (*Lolium perenne* L.). *Environmental and Experimental Botany* 138:36–45

69. Hu Q, Ding F, Li M, Zhang X, Zhang S, et al. 2021. Strigolactone and ethylene inhibitor suppressing dark-induced leaf senescence in perennial ryegrass involving transcriptional downregulation of chlorophyll degradation. *Journal of the American Society for Horticultural Science* 146:79–86

70. Xu B, Yu G, Li H, Xie Z, Wen W, et al. 2019. Knockdown of STAY-GREEN in perennial ryegrass (*Lolium perenne* L.) leads to transcriptomic alterations related to suppressed leaf senescence and improved forage quality. *Plant and Cell Physiology* 60:202–12

71. Zhang J, Li H, Huang X, Xing J, Yao J, et al. 2022. STAYGREEN-mediated chlorophyll a catabolism is critical for photosystem stability during heat-induced leaf senescence in perennial ryegrass. *Plant, Cell & Environment* 45:1412–27

72. Xu B, Li H, Li Y, Yu G, Zhang J, et al. 2018. Characterization and transcriptional regulation of chlorophyll b reductase gene NON-YELLOW COLORING 1 associated with leaf senescence in perennial ryegrass (*Lolium perenne* L.). *Environmental and Experimental Botany* 149:43–50

73. Lei S, Yu G, Rossi S, Yu J, Huang B. 2021. *LpNOL*-knockdown suppression of heat-induced leaf senescence in perennial ryegrass involving regulation of amino acid and organic acid metabolism. *Physiologia Plantarum* 173:1979–91

74. Yu G, Xie Z, Zhang J, Lei S, Lin W, et al. 2021. *NOL*-mediated functional stay-green traits in perennial ryegrass (*Lolium perenne* L.) involving multifaceted molecular factors and metabolic pathways regulating leaf senescence. *The Plant Journal* 106:1219–32

75. Yu G, Xie Z, Chen W, Xu B, Huang B. 2022. Knock down of NON-YELLOW COLOURING 1-like gene or chlorophyllin application enhanced chlorophyll accumulation with antioxidant roles in suppressing heat-induced leaf senescence in perennial ryegrass. *Journal of Experimental Botany* 73:429–44

76. Zhang J, Yu G, Wen W, Ma X, Xu B, et al. 2016. Functional characterization and hormonal regulation of the *PHEOPHYTINASE* gene *LpPPH* controlling leaf senescence in perennial ryegrass. *Journal of Experimental Botany* 67:935–45

77. Yang J, Worley E, Ma Q, Li J, Torres-Jerez I, et al. 2016. Nitrogen remobilization and conservation, and underlying senescence-associated gene expression in the perennial switchgrass *Panicum virgatum*. *New Phytologist* 211:75–89

78. Paudel B, Das A, Tran M, Boe A, Palmer NA, et al. 2016. Proteomic responses of switchgrass and prairie cordgrass to senescence. *Frontiers in Plant Science* 7:293

79. Palmer NA, Saathoff AJ, Waters BM, Donze T, Heng-Moss TM, et al. 2013. Global changes in mineral transporters in tetraploid switchgrasses (*Panicum virgatum* L.). *Frontiers in Plant Science* 4:549

80. Palmer NA, Donze-Reiner T, Horvath D, Heng-Moss T, Waters B, et al. 2015. Switchgrass (*Panicum virgatum* L) flag leaf transcriptomes reveal molecular signatures of leaf development, senescence, and mineral dynamics. *Functional & Integrative Genomics* 15:1–16

81. Yang J, Worley E, Torres-Jerez I, Miller R, Wang M, et al. 2015. *PvNAC1* and *PvNAC2* are associated with leaf senescence and nitrogen use efficiency in switchgrass. *BioEnergy Research* 8:868–80

82. Rinerson CI, Scully ED, Palmer NA, Donze-Reiner T, Rabara RC, et al. 2015. The WRKY transcription factor family and senescence in switchgrass. *BMC Genomics* 16:912

83. Xie Z, Yu G, Lei S, Wang H, Xu B. 2022. STRONG STAYGREEN inhibits DNA binding of PvNAP transcription factors during leaf senescence in switchgrass. *Plant Physiology* 190:2045–58

84. Liu X, Huang B. 2002. Cytokinin effects on creeping bentgrass response to heat stress: II. leaf senescence and antioxidant metabolism. *Crop Science* 42:466–72

85. He Y, Liu X, Huang B. 2005. Changes in protein content, protease activity, and amino acid content associated with heat injury in creeping bentgrass. *Journal of the American Society for Horticultural Science* 130:842–47

86. Xu Y, Huang B. 2007. Heat-induced leaf senescence and hormonal changes for thermal bentgrass and turf-type bentgrass species differing in heat tolerance. *Journal of the American Society for Horticultural Science* 132:185–92

87. Veerasamy M, He Y, Huang B. 2007. Leaf senescence and protein metabolism in creeping bentgrass exposed to heat stress and treated with cytokinins. *Journal of the American Society for Horticultural Science* 132:467–72

88. Xu Y, Huang B. 2009. Effects of foliar-applied ethylene inhibitor and synthetic cytokinin on creeping bentgrass to enhance heat tolerance. *Crop Science* 49:1876–84

89. Jespersen D, Huang B. 2015. Proteins associated with heat-induced leaf senescence in creeping bentgrass as affected by foliar application of nitrogen, cytokinins, and an ethylene inhibitor. *Proteomics* 15:798–812

90. Xing J, Xu Y, Tian J, Gianfagna T, Huang B. 2009. Suppression of shade- or heat-induced leaf senescence in creeping bentgrass through transformation with the *ipt* gene for cytokinin synthesis. *Journal of the American Society for Horticultural Science* 134:602–09

91. Merewitz E, Gianfagna T, Huang B. 2010. Effects of *SAG12-ipt* and *HSP18.2-ipt* expression on cytokinin production, root growth, and leaf senescence in creeping bentgrass exposed to drought stress. *Journal of the American Society for Horticultural Science* 135:230–39

92. Xu Y, Gianfagna T, Huang B. 2010. Proteomic changes associated with expression of a gene (*ipt*) controlling cytokinin synthesis for improving heat tolerance in a perennial grass species. *Journal of Experimental Botany* 61:3273–89

93. Merewitz EB, Du H, Yu W, Liu Y, Gianfagna T, et al. 2011. Elevated cytokinin content in *ipt* transgenic creeping bentgrass promotes drought tolerance through regulating metabolite accumulation. *Journal of Experimental Botany* 63:1315–28

94. Merewitz EB, Gianfagna T, Huang B. 2011. Protein accumulation in leaves and roots associated with improved drought tolerance in creeping bentgrass expressing an *ipt* gene for cytokinin synthesis. *Journal of Experimental Botany* 62:5311–33

95. Merewitz EB, Gianfagna T, Huang B. 2011. Photosynthesis, water use, and root viability under water stress as affected by expression of *SAG12-ipt* controlling cytokinin synthesis in *Agrostis stolonifera*. *Journal of Experimental Botany* 62:383–95

96. Merewitz E, Xu Y, Huang B. 2016. Differentially expressed genes associated with improved drought tolerance in creeping bentgrass overexpressing a gene for cytokinin biosynthesis. *PLoS One* 11:e0166676

97. Jiang Z, Xu C, Huang B. 2011. Enzymatic metabolism of nitrogen in leaves and roots of creeping bentgrass under nitrogen deficiency conditions. *Journal of the American Society for Horticultural Science* 136:320–28

98. Xu C, Jiang Z, Huang B. 2011. Nitrogen deficiency-induced protein changes in immature and mature leaves of creeping bentgrass. *Journal of the American Society for Horticultural Science* 136:399–407

99. Zhang Y, Liang C, Xu Y, Gianfagna T, Huang B. 2010. Effects of *ipt* gene expression on leaf senescence induced by nitrogen or phosphorus deficiency in creeping bentgrass. *Journal of the American Society for Horticultural Science* 135:108–15

100. Jespersen D, Xu C, Huang B. 2015. Membrane proteins associated with heat-induced leaf senescence in a cool-season grass species. *Crop Science* 55:837–50

101. Jespersen D, Zhang J, Huang B. 2016. Chlorophyll loss associated with heat-induced senescence in bentgrass. *Plant Science* 249:1–12

102. Hwang OJ, Han Y, Paek NC, Kim JI. 2014. Antisense expression of a staygreen gene (SGR) delays leaf senescence in creeping bentgrass. *Rapid Communication in Photoscience* 3:28–31

103. Chao Y, Xie L, Yuan J, Guo T, Li Y, et al. 2018. Transcriptome analysis of leaf senescence in red clover (*Trifolium pratense* L.). *Physiology and Molecular Biology of Plants* 24:753–65

104. Qian Y, Cao L, Zhang Q, Amee M, Chen K, et al. 2020. SMRT and Illumina RNA sequencing reveal novel insights into the heat stress response and crosstalk with leaf senescence in tall fescue. *BMC Plant Biology* 20:366

105. Wang L, Doan PPT, Chuong NN, Lee HY, Kim JH, et al. 2023. Comprehensive transcriptomic analysis of age-, dark-, and salt-induced senescence reveals underlying mechanisms and key regulators of leaf senescence in *Zoysia japonica*. *Frontiers in Plant Science* 14:1170808

106. Teng K, Yue Y, Zhang H, Li H, Xu L, et al. 2021. Functional characterization of the pheophytinase gene, *ZjPPH*, from *Zoysia japonica* in regulating chlorophyll degradation and photosynthesis. *Frontiers in Plant Science* 12:786570

107. Guan J, Teng K, Yue Y, Guo Y, Liu L, et al. 2022. *Zoysia japonica* chlorophyll b reductase gene *NOL* participates in chlorophyll degradation and photosynthesis. *Frontiers in Plant Science* 13:906018

108. Dong D, Yang Z, Ma Y, Li S, Wang M, et al. 2022. Expression of a chlorophyll b reductase gene from *Zoysia japonica* causes changes in leaf color and chlorophyll morphology in *Agrostis stolonifera*. *International Journal of Molecular Sciences* 23:6032

109. Teng K, Chang Z, Li X, Sun X, Liang X, et al. 2016. Functional and RNA-sequencing analysis revealed expression of a novel staygreen gene from *Zoysia japonica* (*ZjSGR*) caused chlorophyll degradation and accelerated senescence in *Arabidopsis*. *Frontiers in Plant Science* 7:1894

## Review of leaf senescence in forage and turf grass

## Review of leaf senescence in forage and turf grass

110. Teng K, Tan P, Guan J, Dong D, Liu L, et al. 2021. Functional characterization of the chlorophyll b reductase gene *NYC1* associated with chlorophyll degradation and photosynthesis in *Zoysia japonica*. *Environmental and Experimental Botany* 191:104607

111. Rossi S, Huang B. 2023. Heat-induced leaf senescence in creeping bentgrass suppressed by aminoethoxyvinylglycine involving regulation of chlorophyll metabolism. *Journal of the American Society for Horticultural Science* 148:126–33

112. Rossi S, Huang B. 2024. Protease inhibitors suppressed leaf senescence in creeping bentgrass exposed to heat stress in association with inhibition of protein degradation into free amino acids. *Plant Growth Regulation* 102:65–75

113. Rossi S, Chapman C, Yuan B, Huang B. 2021. Glutamate acts as a repressor for heat-induced leaf senescence involving chlorophyll degradation and amino acid metabolism in creeping bentgrass. *Grass Research* 1:4

114. Rossi S, Huang B. 2021. Sitosterol-mediated antioxidant regulation to enhance heat tolerance in creeping bentgrass. *Journal of the American Society for Horticultural Science* 147:18–24

115. Huang C, Tian Y, Zhang B, Hassan MJ, Li Z, et al. 2021. Chitosan (CTS) alleviates heat-induced leaf senescence in creeping bentgrass by regulating chlorophyll metabolism, antioxidant defense, and the heat shock pathway. *Molecules* 26:5337

116. Liang L, Cao Y, Wang D, Peng Y, Zhang Y, et al. 2021. Spermine alleviates heat-induced senescence in creeping bentgrass by regulating water and oxidative balance, photosynthesis, and heat shock proteins. *Biologia Plantarum* 65:184–92

117. Xu Y, Huang B. 2010. Responses of creeping bentgrass to trinexapac-ethyl and biostimulants under summer stress. *HortScience* 45:125–31

118. Rossi S, Chapman C, Huang B. 2020. Suppression of heat-induced leaf senescence by  $\gamma$ -aminobutyric acid, proline, and ammonium nitrate through regulation of chlorophyll degradation in creeping bentgrass. *Environmental and Experimental Botany* 177:104116

119. Li Z, Huang T, Tang M, Cheng B, Peng Y, et al. 2019. iTRAQ-based proteomics reveals key role of  $\gamma$ -aminobutyric acid (GABA) in regulating drought tolerance in perennial creeping bentgrass (*Agrostis stolonifera*). *Plant Physiology and Biochemistry* 145:216–26

120. Ma X, Zhang J, Burgess P, Rossi S, Huang B. 2018. Interactive effects of melatonin and cytokinin on alleviating drought-induced leaf senescence in creeping bentgrass (*Agrostis stolonifera*). *Environmental and Experimental Botany* 145:1–11



Copyright: © 2024 by the author(s). Published by Maximum Academic Press, Fayetteville, GA. This article is an open access article distributed under Creative Commons Attribution License (CC BY 4.0), visit <https://creativecommons.org/licenses/by/4.0/>.