

ARTICLE

Landscape change and alien invasions drive shifts in native lady beetle communities over a century

Kayla I. Perry^{1,2}  | Christie A. Bahlai²  | Timothy J. Assal³  |
Christopher B. Riley⁴  | Katherine J. Turo⁴  | Leo Taylor⁴  | James Radl⁴ |
Yvan A. Delgado de la flor⁴  | Frances S. Sivakoff⁵ | Mary M. Gardiner⁴ 

¹Department of Entomology, The Ohio State University, Wooster, Ohio, USA

²Department of Biological Sciences, Kent State University, Kent, Ohio, USA

³Department of Geography, Kent State University, Kent, Ohio, USA

⁴Department of Entomology, The Ohio State University, Columbus, Ohio, USA

⁵Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Marion, Ohio, USA

Correspondence

Kayla I. Perry

Email: perry.1864@osu.edu

Present addresses

Timothy J. Assal, National Operations Center, Bureau of Land Management, Denver Federal Center, Building 85 Denver, Colorado 80225-0047, USA; Christopher B. Riley, Bartlett Tree Research Laboratories, 2498 Briggs Road, Berryville, VA 22611, USA; Katherine J. Turo, Department of Ecology, Evolution, and Natural Resources, Rutgers University, 14 College Farm Road, New Brunswick, New Jersey 08901, USA; Leo Taylor, The Office of Diversity, Equity, and Inclusion, The Ohio State University, 2120 Fyffe Road, Columbus, Ohio 43210, USA; and Yvan A. Delgado de la flor, Air Program, Environmental Monitoring Branch, California Department of Pesticide Regulation, 1001 I Street, Sacramento, California 95814, USA.

Funding information

National Science Foundation, Grant/Award Numbers: 2045721, DBI number 2045721; USDA NIFA, Grant/Award Number: 2018-67012-28011

Handling Editor: Juan C. Corley

Abstract

Understanding causes of insect population declines is essential for the development of successful conservation plans, but data limitations restrict assessment across spatial and temporal scales. Museum records represent a source of historical data that can be leveraged to investigate temporal trends in insect communities. Native lady beetle decline has been attributed to competition with established alien species and landscape change, but the relative importance of these drivers is difficult to measure with short-term field-based studies. We assessed distribution patterns for native lady beetles over 12 decades using museum records, and evaluated the relative importance of alien species and landscape change as factors contributing to changes in communities. We compiled occurrence records for 28 lady beetle species collected in Ohio, USA, from 1900 to 2018. Taxonomic beta-diversity was used to evaluate changes in lady beetle community composition over time. To evaluate the relative influence of temporal, spatial, landscape, and community factors on the captures of native species, we constructed negative binomial generalized additive models. We report evidence of declines in captures for several native species. Importantly, the timing, severity, and drivers of these documented declines were species-specific. Land cover change was associated with declines in captures, particularly for *Coccinella novemnotata* which declined prior to the arrival of alien species. Following the establishment and spread of alien lady beetles, processes of species loss/gain and turnover shifted communities toward the dominance of a few alien species beginning in the 1980s.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Ecological Applications* published by Wiley Periodicals LLC on behalf of The Ecological Society of America.

Because factors associated with declines in captures were highly species-specific, this emphasizes that mechanisms driving population losses cannot be generalized even among closely related native species. These findings also indicate the importance of museum holdings and the analysis of species-level data when studying temporal trends in insect populations.

KEYWORDS

Coccinellidae, exotic, invasive, ladybird, land cover, land use, non-native, species decline

INTRODUCTION

Global biodiversity loss is a growing threat to the ecosystem services and functions on which humans depend (Cardoso et al., 2020; Ceballos et al., 2015; Dirzo & Raven, 2003; Mace et al., 2012). Given the importance of insects for ecosystem services such as pollination, pest suppression, and nutrient cycling (Losey & Vaughan, 2006; Yang & Gratton, 2014), understanding the causes of documented spatiotemporal changes in insect populations is a critical research focus. Numerous recent studies have pointed to declines in the abundance, richness, and biomass of insects (Hallmann et al., 2017; Seibold et al., 2019), including bees (Gixti et al., 2009; Potts et al., 2010), beetles (Brooks et al., 2012; Homburg et al., 2019), leafhoppers and planthoppers (Schuch et al., 2012), and butterflies and moths (Conrad et al., 2006; Habel et al., 2016; Maes & Van Dyck, 2001; Warren et al., 2021). Although population declines have been frequently reported, relatively stable or increasing populations of insects also have been documented (Fox et al., 2014; Schowalter et al., 2021; Shortall et al., 2009). For example, observations of moths in Great Britain identified highly species-specific temporal patterns over a 40-year period, as 260 species declined in frequency, 160 species increased, and 253 species remained unchanged (Fox et al., 2014). Complexity in the responses of insects has generated scientific debate about whether observed declines are generalizable across insect taxa and guilds and at larger spatial scales (Didham et al., 2020; Saunders et al., 2020; Simmons et al., 2019; Thomas et al., 2019). Although many challenges to studying insect population trends exist (Didham et al., 2020; Montgomery et al., 2020), understanding the magnitude and drivers of declines in insect species is critical for biodiversity conservation management and the maintenance of insect-based ecosystem services and functions.

One primary limitation to understanding insect populations over time stems from data deficiencies such as low taxonomic resolution, geographic restrictions, and short time series (Didham et al., 2020; Sánchez-Bayo & Wyckhuys, 2019; Thomas et al., 2019). However, specimen records from natural history museums and other institutions can be used to investigate trends in insect populations

over greater spatial and temporal scales (Kharouba et al., 2019; Lister, 2011). Although specimen record-based data have their own set of biases and challenges such as geographic, taxonomic, and temporal biases that result in nonrandom collections (Boakes et al., 2010; Meineke & Daru, 2021), biological collections represent a unique source of historical data that document the occurrence of species in time and space. For example, museum records revealed that 11 bumble bee species native to eastern North America and Canada have experienced substantial (>50%) declines in captures over the past century, while eight species have remained stable or increased in collections (Colla et al., 2012). Importantly, specimen records can be used as baseline measures for investigating the impacts of anthropogenic stressors such as the establishment of alien species, environmental degradation, and climate change on patterns of biodiversity (Johnson et al., 2011; Kharouba et al., 2019; Suarez & Tsutsui, 2004; Winker, 2004). For instance, moth assemblages collected on Mount Kinabalu in Borneo in 2007 were compared to historical records collected from the same sites in 1965, revealing upward shifts in the altitudinal distribution of species in response to changes in temperature (Chen et al., 2009). Although the use of collections data is limited in its ability to track the absolute abundance of a species due to sampling effort variation (Ries et al., 2019), historical specimen data facilitate understanding of species' responses to anthropogenic change by helping to distinguish signals of decline from natural population variability, especially when considering relative observations in groups of similar taxa. Therefore, specimen records are invaluable resources that can be used to address biodiversity conservation initiatives, including documenting changes in communities of beneficial insects.

Lady beetles (Coleoptera: Coccinellidae) are a family of charismatic insect species that are commonly collected and contribute broadly to pest suppression by consuming aphids, scales, psyllids, mites, fungi, and other pests (Evans, 2009; Hodek et al., 2012; Hodek & Honěk, 2009; Weber & Lundgren, 2009). Because lady beetles are capable of rapidly colonizing habitats to exploit ephemeral prey resources, these species have been used widely in biological control programs in agricultural systems (Caltagirone & Doult, 1989; Koch, 2003; Obrycki & Kring, 1998; Rondoni

et al., 2021). A number of authors have noted declines in native lady beetle populations across the United States (Alyokhin & Sewell, 2004; Harmon et al., 2007) and Europe (Brown & Roy, 2018; Roy et al., 2012), which may contribute to an overall loss of resilience of the biological control services offered by this community (Bahlai et al., 2021). For example, the historically widespread native aphidophagous species *Hippodamia convergens* Guerin has declined in the US states of Michigan (Gardiner et al., 2009), Ohio (Gardiner et al., 2012), Wisconsin (Steffens & Lumen, 2015), and Minnesota (Steffens & Lumen, 2015), as well as the Canadian Province of Manitoba (Turnock et al., 2003). Likewise, the nine-spotted lady beetle *Coccinella novemnotata* Herbst was once common in eastern North America, but had not been collected for over a decade until a community scientist “rediscovered” it (Losey et al., 2007). Anthropogenic activities such as the establishment of alien species and landscape change have been hypothesized as potential drivers of native lady beetle decline (Alyokhin & Sewell, 2004; Bahlai et al., 2015; Gardiner et al., 2012; Harmon et al., 2007; Honěk et al., 2014; Roy et al., 2016).

Several studies have observed that the decline of native aphidophagous species coincided with the establishment and spread of alien lady beetle species, particularly the Asian species *Harmonia axyridis* (Pallas) and the European species *Coccinella septempunctata* (Linnaeus) (Roy et al., 2012, 2016; Steffens & Lumen, 2015; Turnock et al., 2003). Following their establishment, alien lady beetles became the dominant species within many native communities (Alyokhin & Sewell, 2004; Bahlai et al., 2015; Harmon et al., 2007). Because of their numerical and behavioral dominance, direct and indirect competitive interactions with alien species are hypothesized as a cause of declines in native lady beetles (Li et al., 2021; Pell et al., 2008). For example, intraguild predation has been documented among native and alien lady beetles in the field (Brown et al., 2015; Gagnon et al., 2011; Ortiz-Martínez et al., 2020; Thomas et al., 2013), wherein native eggs and larvae were more likely to be the intraguild prey for alien species (Katsanis et al., 2013; Snyder et al., 2004; Turnipseed et al., 2015). In the field, native species experienced greater egg predation from a guild of shared predators than alien lady beetle species, which could translate to apparent competition (Smith & Gardiner, 2013). These asymmetric interactions may largely benefit alien species at the expense of native lady beetle populations, but the extent and context of these effects on native species are difficult to quantify. The short time scales of many studies and the lack of data from lady beetle communities before the establishment of alien species limits understanding of the impacts of these invaders.

Landscape change that results in the loss, fragmentation, and degradation of natural habitat also has been

hypothesized as a key driver contributing to population declines of insect species (Potts et al., 2010; Wagner, 2020; Wagner et al., 2021), including lady beetles (Honěk et al., 2017). Land cover change resulting from increased urbanization and agricultural intensification can influence the structure and composition of lady beetle communities (Gardiner et al., 2009, 2021; Grez et al., 2019; Parker et al., 2020; Woltz & Landis, 2014). For example, native and alien species were less abundant in isolated urban greenspaces that were embedded in landscapes dominated by impervious surfaces and built infrastructure (Parker et al., 2020). In agricultural landscapes, native and alien lady beetles were more abundant in fields surrounded by higher crop diversity and more semi-natural habitat types such as grasslands and forests (Woltz & Landis, 2014). Loss and fragmentation of natural habitat in landscapes such as those dominated by urban and agricultural land cover may differentially affect species depending on their life history traits such as phenology, dispersal ability, overwintering biology, and food requirements (Zaviezo et al., 2006). However, the impacts of landscape change on native lady beetle populations may occur gradually over time, making species responses difficult to detect over short time scales. For example, gradual directional change in the species composition of native lady beetle communities was documented over a 118-year period in Missouri, USA, using museum specimen records (Diepenbrock et al., 2016). Because there was no evidence that the establishment of alien lady beetle species affected the rate of change, Diepenbrock et al. (2016) hypothesized that these long-term community changes may be related to altered land use patterns. The substantial year-to-year variation in the abundance of lady beetle species (Elliott et al., 1996; Honěk et al., 2014) requires longer time series data such as those used by Diepenbrock et al. (2016) to detect changes in populations caused by landscape change and to distinguish these effects from other anthropogenic stressors (Bahlai et al., 2021; Bahlai & Zipkin, 2020).

While several hypotheses have been proposed, the causes of declines in native lady beetle populations remain under debate. Importantly, these hypotheses are not mutually exclusive, and it is likely that causes of declines in native species are multifaceted, wherein multiple mechanisms are responsible for the observed patterns in lady beetle populations. Although causes of declines in native species are often studied independently, these drivers may interact to influence native populations (Didham et al., 2007). For example, habitat modification that transitions natural habitat to more highly disturbed urban and agricultural habitats may differentially benefit some lady beetle species over others (Grez et al., 2013), with implications for direct and indirect competitive interactions among native and alien species.

Understanding the magnitude and drivers of declines in native lady beetle populations will require comprehensive

time-series data documenting community responses that can then be used to assess the contribution of the establishment of alien species and landscape change simultaneously. To understand the relative importance of the establishment of alien species and landscape change as contributors of native species decline, we compiled historical occurrence records of lady beetles collected in Ohio, USA, from museums and other institutions across the United States. Although we cannot directly assess changes in abundance with occurrence records, these data allow the investigation of declines in captures of lady beetles over time and the comparison of relative observations among species. Our goals were to: (1) characterize long-term changes in land cover and lady beetle communities which have occurred over time in Ohio by assessing temporal patterns of beta-diversity; and (2) evaluate the relative importance of the establishment and spread of alien lady beetle species and landscape change in Ohio as stressors contributing to patterns of captures in several native lady beetle species over time.

METHODS

Lady beetle specimens and data requests

To investigate long-term changes in native lady beetle communities within Ohio, USA, we used historic occurrence records for native and alien species within the tribe Coccinellini and four additional commonly collected species within the tribes Hyperaspini, Chilocorini, and Psyllorini. Targeted Coccinellini genera were *Adalia*, *Anatis*, *Anisosticta*, *Aphidecta*, *Calvia*, *Ceratomegilla*, *Coccinella*, *Coelophora*, *Coleomegilla*, *Cycloneda*, *Harmonia*, *Hippodamia*, *Macronaemia*, *Mulsantina*, *Myzia*, *Naemia*, *Neoharmonia*, *Olla*, and *Propylea*, and the four additional species were *Brachiacantha ursina* (Fabricius), *Chilocorus stigma* (Say), *Hyperaspis undulata* (Say), and *Psyllobora vigintimaculata* (Say). We contacted 59 institutions based within the United States, the majority of which are hosted by the Entomological Collections Network (ENC, 2020). Ohio lady beetle records were compiled from 25 institutions with assistance of their curators (see Appendix S1: Table S1 and Section S1).

Specimen loans were requested from museums and institutions that had unidentified lady beetle records from Ohio. Any unidentified lady beetle species that were unaccounted for within the collections of these institutions were not included within our dataset. Lady beetle species identifications were determined using Gordon (1985). All lady beetle species were characterized by their status (native or alien to North America) and their primary diet (aphidophagous, coccidophagous, or fungivorous) which

was based on most frequently reported prey (aphids, scales, or fungi) (Angalet et al., 1979; Dixon & Dixon, 2000; Gordon, 1985; Hodek et al., 2012; Majerus, 2016; Michaud, 2001; Staines, 2008).

Land use and land cover change analysis

To assess the influence of landscape change on lady beetle communities, we analyzed historical land use and land cover (LULC) data from four points in time (1938, 1970, 1992, 2016). Annual historical LULC data were obtained from the US Geological Survey (Sohl et al., 2018) for the years 1938, 1970, and 1992. These historical LULC backcasts were modeled using numerous historical datasets and created explicitly to extend the National Land Cover Database (NLCD) to earlier time periods, prior to the availability of remote-sensing data (Sohl et al., 2016). The earliest data available are from 1938, which corresponds with the start of historical reporting in the Agricultural Census (Sohl et al., 2016). The NLCD and historical backcast data were converted into four primary land use classes: agriculture, developed, forest, and other (i.e., non-target; all other land cover classes found in Ohio) (see Appendix S2: Tables S1 and S2). The contemporary landscape of Ohio was assessed with the 2016 NLCD (Dewitz, 2019; Wickham et al., 2021). To assess change over time, we calculated the total area of each of the four cover classes (i.e., agriculture, developed, forest, and other), along with the percentage of area occupied by each class, for each county at each of the four time periods. Although the LULC data have different spatial resolutions (NLCD = 30 m; historical backcasts = 250 m) they were not resampled, as the derived metrics (e.g., total area of cover class, percentage of county area) are relative, and therefore comparable across time. Analyses were completed in R (R Core Team, 2020) using the “raster” and “sf” R packages (Hijmans, 2020; Pebesma, 2018).

There was an increase in forest cover from 1938 to 1970, and it has held steady since the early 1990s (see Appendix S2: Figure S2). Most of the increase occurred in the eastern and southeastern regions of the state (Figure 1). There was a steady increase in developed land around existing population centers, with a larger increase in suburban areas since the early 1990s. The increase in forest and developed lands came at the expense of agricultural lands, yet this cover class remains the dominant land cover in many counties of the state (Figure 1; see Appendix S2: Figure S2). During each time period, there was high variability in the amount of agricultural and forested land across Ohio counties (see Appendix S2: Figure S2).

Land cover data were compiled at the county and decade resolution to align with the spatial and temporal




FIGURE 1 Spatial extent of agriculture (a), developed land (b), and forest (c) in Ohio for each time period (columns).

resolution available for the museum specimen data. For each county, we used an assumed linear progression between available timepoints to estimate the percentage land cover of a given class in each decade. The 1930 land cover data were extrapolated using the same procedure with available data from 1938 to 1970. Because lady beetle data were generally recorded at the county level, we computed the geographic centroids of each Ohio county based on present-day county boundaries as defined by the Ohio Department of Transportation (<http://ogrip-eohio.opendata.arcgis.com/datasets/odot-county-boundaries>).

Statistical analysis

Lady beetle museum records were included in the analyses if they were collected within Ohio, and we could obtain county-level location data and the year of collection. All analyses were conducted in R (R Core Team, 2020). The

analysis code and development history are available in Bahlai and Perry (2024).

To characterize changes in land cover and lady beetle communities over time, differences in land cover and lady beetle community composition were visualized using nonmetric multidimensional scaling (NMDS) analyses. Raw land cover composition data (agriculture, developed, forest, and other/non-target) were compiled at the county level for the four time periods collected (i.e., 1938, 1970, 1992, 2016). Lady beetle specimen data were compiled at the county level and pooled across decades to create four time periods that corresponded with the available landscape data (before 1938, 1939–1970, 1971–1992, 1993–2016). Two lady beetle analyses were conducted, one with all species and one with native species only. NMDS analyses were conducted using the “vegan” package (Oksanen et al., 2011).

To further investigate long-term changes in native lady beetle communities within the region, community

composition was assessed by partitioning incidence-based measures of taxonomic beta-diversity for total and aphidophagous lady beetle species. Aphidophagous lady beetles were investigated separately because the established alien species primarily feed on aphid prey, and it is hypothesized that native aphidophagous species would experience strong competition with alien species due to this overlap in resource use. Due to low county-level counts, county records of lady beetle species were pooled across 10-year intervals such that patterns of community composition were assessed at the spatial scale of the state of Ohio across 11 decades from the 1900s through the 2010s. Descriptive patterns of total and aphidophagous lady beetle taxonomic beta-diversity were evaluated by calculating total Sorenson dissimilarity (β_{SOR}) using the *beta.temp* function in the R package “betapart” (Baselga & Orme, 2012). Total Sorensen dissimilarity was decomposed into a turnover component (β_{SIM} ; reflects species replacement) and a nestedness component (β_{SNE} ; reflects species loss or gain) (Baselga, 2010).

To evaluate the relative importance of temporal, spatial, landscape, and community factors on the captures of five key native lady beetle species (*Adalia bipunctata*, *C. novemnotata*, *Coleomegilla maculata*, *H. convergens*, the four most abundant native aphidophagous species, and *C. stigma*, the most abundant coccidophagous species), we constructed negative binomial generalized additive models (GAMs) for each species using the “mgcv” package (Wood, 2017), and then used an adaptive model selection procedure in combination with available contextual data to determine the relative importance of the factors. For all GAMs, lady beetle records were pooled at the county spatial scale and the decade temporal scale. Because the absolute number of records was very sparse earlier in our study period, we restricted these analyses to specimens captured in or after 1930 for these analyses. For *A. bipunctata* and *C. novemnotata*, which were extremely rare or absent from collections in later decades, data were culled at 1990 to restrict these analyses to times when these species were present. We computed several community variables for each decade-county combination, in addition to absolute captures of each coccinellid species, including the captures of two dominant invaders, *C. septempunctata* and *H. axyridis*, the total lady beetles captured, total alien species captured, and the proportion of the community captures belonging to alien species. All GAMs constructed included an offset variable of the structure $\log(1 + \text{total lady beetles captured})$ to account for variability in sampling effort (with the exception of the model describing sampling effort over time).

First, to describe the relative abundance of each key species over time, corrected for varied sampling intensity,

a negative binomial GAM was constructed for each species with decade of capture as the independent variable, constrained to 3 knots and a smoothing parameter of 0.5. For the spatial analyses, data were aggregated into three-decade groups, and the captures of each species were modeled using a negative binomial GAM with a Gaussian process smooth and a combination of latitude and longitude, both in aggregate and then by decade group. Latitude and longitude were included in the model to control for differences in the spatial distribution of lady beetle collections within the state and to account for spatial autocorrelation. To examine the interaction of temporal, spatial, landscape, and community factors on captures of each species, we used a modified backward-stepping model selection applied to negative binomial GAMs. First, a global model was constructed that included decade, longitude, latitude, percentage land cover in agriculture, forest, and developed uses, the total captures of alien species, and the sampling offset. Because of a high degree of autocorrelation and non-independence between the aggregate alien species metrics, each of these variables was substituted into the model separately, and the variables with the best performance (determined by Akaike information criterion [AIC]) were selected for further analysis. Captures of *C. septempunctata* and *H. axyridis* were considered together in the same model to test if target native species were exhibiting differential responses to the two alien species. After the substitution-based model selection was completed, the remaining model selection was completed using backward selection by systematically dropping each variable with the lowest explanatory power, using AIC as the decision criterion. The selected model was then subjected to concurvity analysis and if the “worst case” concurvity estimate exceeded 0.8 for any parameter, the parameter was eliminated from the model and backward model selection was resumed (Ross, 2019).

RESULTS

We compiled 4194 lady beetle museum records representing 28 species collected in Ohio, USA, from 1900 to 2018 (Table 1; see Appendix S2: Table S3 and Figures S3–S6). Total collections of native species varied from year to year, with high numbers collected in the 1930s and 1980s (Figure 2a). The most common native species represented in these collections were *C. maculata* (Degeer) (16.4% of total records), *H. convergens* Guérin (9.5%), *Hippodamia parenthesis* (Say) (8.4%), *P. vigintimaculata* (Say) (7.6%), *Cycloneda munda* (Say) (7.2%), *B. ursina* (Fabricius) (6.1%), and *A. bipunctata* (Linnaeus) (5.2%). Records documented the presence of the aphidophagous alien species

TABLE 1 Ohio lady beetle species records from 1900 to 2018 compiled from 25 institutions across the United States.

Tribe	Lady beetle species	Records	First record	Last record	Status	Primary diet
Hyperaspidini	<i>Brachiacantha ursina</i> (Fabricius)	255	1902	2015	Native	Aphidoidea, Coccoidea
	<i>Hyperaspis undulata</i> (Say)	61	1905	2014	Native	Coccoidea
Chilocorini	<i>Chilocorus stigma</i> (Say)	111	1901	2016	Native	Coccoidea, Aphidoidea
Coccinellini	<i>Adalia bipunctata</i> (Linnaeus)	218	1901	1996	Native	Aphidoidea
	<i>Anatis labiculata</i> (Say)	176	1907	2012	Native	Coccoidea, Aphidoidea
	<i>Anatis mali</i> (Say)	23	1932	2013	Native	Coccoidea, Aphidoidea
	<i>Anisosticta bitriangularis</i> (Say)	16	1923	2012	Native	Aphidoidea, Pollen
	<i>Coccinella novemnotata</i> Herbst	169	1901	1985	Native	Aphidoidea
	<i>Coccinella septempunctata</i> (Linnaeus)	102	1978	2017	Alien	Aphidoidea
	<i>Coccinella transversoguttata</i> Mulsant	51	1921	1986	Native	Aphidoidea
	<i>Coccinella trifasciata</i> Linnaeus	19	1911	1971	Native	Aphidoidea
	<i>Coccinella undecimpunctata</i> Linnaeus	1	1953	1953	Alien	Aphidoidea
	<i>Coleomegilla maculata</i> (Degeer)	690	1901	2018	Native	Aphidoidea, Pollen
	<i>Cycloneda munda</i> (Say)	302	1901	2017	Native	Aphidoidea
	<i>Harmonia axyridis</i> (Pallas)	470	1993	2018	Alien	Aphidoidea
	<i>Hippodamia convergens</i> Guerin	400	1903	2015	Native	Aphidoidea
	<i>Hippodamia glacialis</i> (Fabricius)	39	1905	2016	Native	Aphidoidea
	<i>Hippodamia parenthesis</i> (Say)	353	1900	2017	Native	Aphidoidea
	<i>Hippodamia quindecimmaculata</i> Mulsant	3	1905	1935	Native	Aphidoidea
	<i>Hippodamia tredecimpunctata</i> (Linnaeus)	155	1903	2014	Native	Aphidoidea
	<i>Hippodamia variegata</i> (Goeze)	10	2000	2016	Alien	Aphidoidea
	<i>Mulsantina luteodorsa</i> J. Chapin	1	2008	2008	Native	Aphidoidea
	<i>Mulsantina picta</i> (Randall)	51	1924	2015	Native	Aphidoidea
	<i>Myzia pullata</i> (Say)	30	1934	2012	Native	Aphidoidea
	<i>Neoharmonia venusta</i> (Melsheimer)	119	1902	2012	Native	Coccoidea
	<i>Olla v-nigrum</i> (Mulsant)	21	1932	2004	Native	Aphidoidea, Psylloidea
	<i>Propylea quatuordecimpunctata</i> (Linnaeus)	27	2003	2016	Alien	Aphidoidea
Psylloborini	<i>Psyllobora vigintimaculata</i> (Say)	321	1902	2016	Native	Fungi (Erisiphaceae)

Note: Data collection focused on native and alien species within the tribe Coccinellini and four additional species in the tribes Hyperaspidini, Chilocorini, and Psylloborini. Lady beetle species were characterized based on their status (native or alien to Ohio, USA) and their primary diet (aphids, scales, or fungi).

C. septempunctata (Linnaeus) (2.4% of total records; first detected in 1978), *Coccinella undecimpunctata* Linnaeus (0.02%; single record in 1953), *H. axyridis* (Pallas) (11.2%; first detected in 1993), *Hippodamia variegata* (Goeze) (0.2%; first detected in 2000), and *Propylea quatuordecimpunctata* (Linnaeus) (0.6%; first detected in 2003). Collections of alien species began to increase in the 1980s, with numbers surpassing natives in the 2000s and 2010s (Figure 2a). The proportion of native lady beetles in collections, including *A. bipunctata*, *C. novemnotata*, and *H. convergens*, has decreased in Ohio since the 1970s as alien lady beetles such

as *C. septempunctata* and *H. axyridis* have increased (Figure 2b).

Temporal patterns of landscape and lady beetle community change

NMDS analyses identified parallel shifts in land cover and lady beetle composition that occurred from 1992 to 2016 (Figure 3). Land cover composition was similar among the 1938, 1970, and 1992 time periods, until substantial changes occurred in the 2016 time period (Figure 3a).




FIGURE 2 Number of collections (a) and proportion of collections (b) of native and alien lady beetle records from Ohio by decade from 1900 to 2018.

This shift corresponded to a decrease in agricultural land cover and an increase in forest and developed land cover across these time periods. Total and native lady beetle communities displayed a similar temporal response. Lady beetle species composition was similar among the first two time periods (before 1938 and 1938–1970), shifted slightly from 1971 to 1992, and then substantially changed in the final time period from 1993 to 2016 (Figure 3b,c).

Lady beetle species composition

Descriptive patterns of lady beetle taxonomic beta-diversity (β_{SOR}) across decades within the state of Ohio were the result of species turnover (β_{SIM}) and nestedness (β_{SNE}), but the strongest contributor to changes in lady beetle community composition shifted over time (Figure 4). Nestedness was the primary contributor to patterns of lady beetle beta-diversity before the establishment of alien species. Following the establishment of *C. septempunctata* and *H. axyridis* in the late 1970s and early 1990s, respectively, an increase in total beta-diversity was observed driven by increased species nestedness. From 1980 to 2018, species turnover increasingly became the stronger contributor, especially for aphidophagous species, and was the dominant driver of lady beetle beta-diversity by the 2010s (Figure 4).

Responses of native lady beetle species

Relative captures of all lady beetle species varied with time in models accounting only for sampling effort and using a normal error structure (see Appendix S3: Figure S1). However, these simple negative binomial models explained limited variation in the data. Native species were captured less per sampling effort later in the

study period, but the steepness and timing of declines in captures varied by species (see Appendix S3: Figure S1A–E). The two alien species exhibited more variable patterns. Captures of *C. septempunctata* initially increased in effort-corrected captures but then decreased during the four decades it has been present in Ohio (see Appendix S3: Figure S1F). Conversely, captures of *H. axyridis* increased over the three decades since its establishment (see Appendix S3: Figure S1G). For *A. bipunctata* and *C. novemnotata*, statistically significant temporal patterns were not observed within the modeled time period, as both species had relatively stable capture frequency prior to the 1980s, but became extremely rare and then absent in later years of data collection. To provide meaningful model fits for the time when these species were present, years with zero-biased data were not included in their respective models. Sampling effort varied dramatically by year and location. Patterns of captures over time were not spatially static, as several native species also exhibited spatiotemporal dependencies over the study (i.e., changing spatial distributions with time; see Appendix S3: Table S1 and Figures S2–S6).

Species-specific patterns emerged in the negative binomial models that accounted for landscape change, invasion, and spatiotemporal factors, and model fit was much improved in all cases compared with models that did not account for these potential factors (Figure 5; see Appendix S3: Table S1). Captures of *A. bipunctata* exhibited a steep negative response to the increasing proportion of alien species in the community but also were positively associated with agriculture and developed land covers. Captures of *C. novemnotata* had a strong negative trend over time as well as spatial dependencies but only appeared to respond negatively to agricultural cover. Since captures of this native species had already begun to decline prior to invasions by *C. septempunctata* and *H. axyridis*, there was limited co-occurrence of *C. novemnotata* and any




FIGURE 3 Changes in land cover composition by county (a), total lady beetle species composition by county (b), and native lady beetle species composition by county (c) across four time periods in Ohio. Shaded ellipses represent 99% confidence intervals around centroids. Centroids adjacent in time to each other are connected with a solid black line.

of the alien species (only five captures of *C. novemnotata* were recorded after 1980). Captures of *C. maculata* were relatively consistent in the earlier period of the study, with a slight increase peaking in the 1970s and 1980s, but a net decrease since. Our models suggest a relatively neutral effect of small numbers of alien lady beetles on *C. maculata*, but a negative impact as alien species become dominant (i.e., >50% of the lady beetles collected). Additionally, agricultural land cover had a slight positive association with captures of *C. maculata*. Captures of *H. convergens* had a negative trend over time and exhibited a differential response to the two dominant alien species: a

positive association with *C. septempunctata* and a negative association with *H. axyridis*, as well as a positive association with higher values of developed land cover. Captures of *C. stigma* exhibited spatial dependencies, a positive association with forested habitats, and a negative association with the proportion of alien lady beetles in the community.

DISCUSSION

Specimen records can be harnessed to understand long-term biodiversity trends across various spatial scales in




FIGURE 4 Descriptive patterns of taxonomic beta-diversity across decades for all lady beetle species (a) and aphidophagous lady beetle species (b) collected in Ohio. Total Sorensen dissimilarity (β_{SOR}) among decades was partitioned into two additive components: Turnover (β_{SIM} ; reflects species replacement) and nestedness (β_{SNE} ; reflects species loss or gain). Therefore, $\beta_{\text{SOR}} = \beta_{\text{SIM}} + \beta_{\text{SNE}}$.

response to anthropogenic threats (Lister, 2011; Meineke et al., 2019; Shaffer et al., 1998; Suarez & Tsutsui, 2004). Using historic land cover data and occurrence records of lady beetles collected in Ohio, USA, we documented parallel shifts in land cover and lady beetle composition that occurred from 1992 to 2016, aligning with decreased agricultural land cover and the establishment and spread of alien lady beetle species. Investigation of these patterns revealed evidence of declines in captures for several native species, but the timing and severity of declines as well as the relative importance of alien species and land cover change as long-term factors were species-specific. Observed changes in species composition that began in the 1980s indicated processes of species loss/gain and turnover that shifted communities toward dominance of a few alien lady beetle species.

Shifts in lady beetle species composition

As collections of some native species declined over time in Ohio, captures of alien lady beetle species increased following their establishment and spread. Beginning in the 2000s, alien species comprised over 60% of the total specimens collected, indicating a shift in lady beetle community structure and alien species dominance within the state. This shift in total and native lady beetle species composition first became evident in the 1971–1992 time period, which coincided with the establishment and spread of *C. septempunctata*, and this was followed by a more substantial shift in species composition during the 1993–2016 time period when *H. axyridis* became

established and dominant. These patterns were further supported by changes in lady beetle beta-diversity across the state that began in the 1980s. From the 1980s to the 2000s, total beta-diversity of lady beetle species increased across decades, indicating that lady beetle communities became increasingly dissimilar within the state during the period that coincided with alien lady beetle establishment and spread. This increase in beta-diversity was initially driven by processes of species loss or gain (i.e., species nestedness), which is consistent with the establishment and spread of alien species within native lady beetle communities. From the 2000s to 2010s, the contribution of species turnover to patterns of beta-diversity increased compared with previous decades, especially for aphidophagous species, which suggests replacement of natives by alien species. High species replacement indicates that a similar number of lady beetle species were collected during these decades, but a low number were shared as species were replaced over time. Changes in lady beetle species composition from the 1900s to the 1970s were primarily driven by species nestedness. These patterns were likely the result of the loss and gain of uncommon native species in collections throughout the state, as well as the loss of common native species such as *A. bipunctata* and *C. novemnotata* which began to decline prior to alien species arrival.

A combination of species nestedness and species turnover that resulted in high dissimilarity in lady beetle species composition during the 1980s, 1990s, and 2000s is indicative of spatiotemporal loss and replacement of native species. By the 2010s, lady beetle communities had become more similar to each other in terms of species




FIGURE 5 Matrix of partial generalized additive model (GAM) predictions of responses for five key native species to spatiotemporal, invasion, and landscape parameters. Negative binomial GAMs used Ohio museum collections, 1930–2018 data on lady beetle captures, and were adjusted for sampling effort. Predictions were constructed by generating data that held all other parameters constant and varying the parameter of interest, and then substituting these data into the best-fit model determined by model selection (see Appendix S3: Table S1). Solid lines, shading represent 95% confidence intervals.

composition as indicated by the decrease in total beta-diversity and were dominated by alien lady beetle species. Similar shifts in lady beetle communities have been observed in response to the establishment of alien species (Alyokhin & Sewell, 2004; Bahlai et al., 2013; Elliott et al., 1996). For example, a shift in species dominance with the establishment of *C. septempunctata* in the 1980s was observed in potato fields in Maine, USA (Alyokhin & Sewell, 2004). Within 4 years of its establishment in southwestern Michigan, USA, *H. axyridis* was the dominant lady beetle species collected and was found in diverse habitats, including agricultural and old fields, and poplar plantations (Colunga-Garcia & Gage, 1998). Although the composition of lady beetle communities has become dominated by alien species, Bahlai et al. (2013) documented that their potential to suppress pest populations in agricultural and natural habitats had remained relatively consistent over time. As significant shifts in lady beetle communities have occurred across the Midwestern US, reliance on alien species to maintain pest suppression may be required for management, but further declines of native species may affect long-term resilience of this ecosystem service (Bahlai et al., 2021).

Drivers of native lady beetle decline

Using museum specimens collected over 12 decades, we documented declines in captures of the native species *A. bipunctata*, *C. novemnotata*, *H. convergens*, *C. maculata*, and *C. stigma*. The native species *A. bipunctata*, *C. novemnotata*, and *H. convergens* were once widely distributed across much of the United States (Gordon, 1985), with early surveys in Ohio recording these species across the state (Bubna, 1902; Dury, 1879). Now, these species are considered rare or potentially extirpated in much of eastern North America (Elliott et al., 1996; Gardiner et al., 2012; Wheeler & Hoebeke, 1995). For example, no captures of *C. novemnotata* or *A. bipunctata* were found after 1985 and 1996, respectively, suggesting their populations were below the detection threshold or extirpated within the state of Ohio. Major drivers of declines in native biodiversity remain the establishment and spread of alien species and habitat loss from land use change (Lövei, 1997; Newbold et al., 2015; Seibold et al., 2019; Vitousek et al., 1997; Wilcove et al., 1998). Our findings revealed that captures of native lady beetles responded to both stressors.

As alien lady beetle species increased in dominance, declines in captures of *C. maculata*, *H. convergens*, and *C. stigma* were associated with these invaders, although the magnitude and timing of the effects were species-specific. For example, collections of *H. convergens*

declined with captures of *H. axyridis* but increased with captures of *C. septempunctata*. These divergent patterns highlight that alien lady beetle species are not ecologically equivalent, and instead, can have different impacts on native species. Moreover, *C. stigma* is coccidophagous, and declines in captures of this native species suggests that aphidophagous alien lady beetles may affect non-aphidophagous native species to some extent. The magnitude of decline in native lady beetles is predicted to be linked to competitive interactions and degree of niche overlap with alien lady beetle species (Bahlai et al., 2015). The competitive exploitation hypothesis posits that decline in populations of native species would occur following the establishment of functionally similar alien species that are superior competitors (Snyder, 2009). In this study, collections of alien species were predominantly *C. septempunctata* and *H. axyridis*, both of which feed on a wide variety of aphid prey and are efficient competitors (Angalet et al., 1979; Bahlai et al., 2013; Hodek et al., 2012; Koch, 2003). Decline of *C. stigma* immediately following the establishment of *H. axyridis* has been reported in southwestern Michigan, USA, further suggesting potential competitive interactions among these species (Colunga-Garcia & Gage, 1998). Although *H. axyridis* is primarily aphidophagous (Koch, 2003), this species also feeds on scale insects in arboreal environments (McClure, 1986), which suggests some degree of dietary and habitat overlap with *C. stigma*. *C. maculata* is known to be a strong aphidophagous competitor (Long & Finke, 2014), and this species remains one of the most frequently collected native lady beetles in croplands, grasslands, and home gardens (Gardiner et al., 2021; Smith & Gardiner, 2013). For example, *C. maculata* was the most commonly collected native lady beetle in residential gardens across the state of Ohio by community scientists (Gardiner et al., 2012, 2021). However, decline in *C. maculata* populations were reported during a long-term (two decadal-scale) study in southwestern Michigan (Bahlai et al., 2015). *C. maculata* feeds on pollen in addition to aphid prey (Dixon & Dixon, 2000; Hodek et al., 2012; Majerus, 2016), and this pollen feeding was hypothesized to reduce competition with alien species (Bahlai et al., 2015). However, there is evidence that *H. axyridis* and *C. septempunctata* also feed on pollen as a supplemental food resource when arthropod prey is scarce (Berkvens et al., 2008, 2010; Ricci et al., 2005), which may suggest the competitive impacts of alien species on *C. maculata* have been underestimated.

We found evidence that declines in captures of the aphidophagous species *A. bipunctata* and *C. novemnotata* began prior to the arrival and dominance of alien lady beetle species. Declines in captures of *A. bipunctata* began in the 1960s, with the last collection in Ohio in

1996. Since the arrival of *C. septempunctata* in 1978, there were only 13 individuals of *A. bipunctata* collected from the state. Although our findings indicated that alien species likely contributed to declines in captures of *A. bipunctata* to some degree, declines in captures prior to the arrival of alien species suggests other contributing factors such as landscape change. Captures of *C. novemnotata* began to decline in the 1950s, with the last Ohio collection occurring in 1985. Since the arrival of *C. septempunctata* in Ohio, only 12 individuals of *C. novemnotata* were recorded. This finding contrasts with many studies across the United States that have hypothesized declines of *C. novemnotata* were related to competitive displacement by *C. septempunctata* (Snyder & Evans, 2006; Staines et al., 1990; Ugine et al., 2018; Wheeler & Hoebeke, 1995).

Land cover change was an important factor for several native lady beetles, although responses were highly species-specific. For example, positive, neutral, and negative responses to the percentage of agricultural land cover were documented. Captures of *C. novemnotata* decreased with the percentage of agricultural land cover, while collections of *A. bipunctata* increased. Since the 1930s, the amount of agricultural land cover has decreased within the state, but the spatial extent of these changes was highly variable and not consistent across all Ohio counties. This loss of agriculture was primarily in the eastern region of the state, while western Ohio remained dominated by agricultural land cover throughout the study period. Along with these changes in the extent of agricultural land cover, landscapes have experienced a shift from more diversified cropping systems to highly managed crop monocultures since the mid-1900s (Crossley et al., 2021). *C. novemnotata* has a broad ecological niche (Losey et al., 2007), and populations have been found in a variety of cultivated crops (Wheeler & Hoebeke, 1995), including alfalfa (Goodarzy & Davis, 1958; Pimentel & Wheeler, 1973), corn (Smith, 1971), cotton (Bell & Whitcomb, 1964), soybeans (Dumas et al., 1964), and fruit trees (Oatman et al., 1964; Putman, 1964). Simplification of agricultural landscapes due to loss of natural habitat, reductions in crop diversity, and insecticide use may impact temporal prey availability and diversity as well as refuge and overwintering habitats that are required for predatory insect species such as *C. novemnotata* (Rusch et al., 2016).

Of the species investigated, *C. stigma* was the only native lady beetle associated with forest land cover, with captures increasing as the percentage of forest increased at the county level. This response is likely linked with the ecology of this species. Scale insects are common pests of trees in managed and natural forests. Further, *C. stigma* is reported to oviposit eggs in bark cracks and crevices as well as overwinter as adults in the leaf litter layer

(Mayer & Allen, 1983). Similar positive associations with forests have been reported in other native lady beetle surveys. For instance, the amount of forest habitat at a 2 km landscape scale was positively associated with the abundance and species diversity of native lady beetles within residential gardens (Gardiner et al., 2021). We hypothesize that the conservation value of forests for native lady beetles may be underestimated, but additional research is required to understand when and how native species are utilizing these habitats.

Conclusions

Using specimen records collected over 118 years in Ohio, USA, we documented shifts in lady beetle species composition beginning in the 1980s as communities became increasingly dominated by alien species. Additionally, we documented a parallel shift in land cover composition during the same time periods that resulted from the replacement of agriculture by forest and developed land. Because of uneven sampling inherent to museum collections, total records of native lady beetles varied from year to year. Therefore, we were unable to make inferences about changes in absolute abundance for a given species, and instead, make our interpretations from proportional capture rates at the community-level. Unknown sampling effort in early museum records is a limitation of any studying utilizing museum data (Bartomeus et al., 2013), and species records are often more numerous within recent decades (Colla et al., 2012), with some evidence that this is due to increased sampling effort (Jeppsson et al., 2010). Our method likely provides a conservative estimate of individual species decline, as it assumes the net community of lady beetles, inclusive of alien species, has remained roughly stable, and changing rates of captures across the community through time are associated with sampling effort. Although occupancy models have been used to attempt to account for wide variation in sampling intensity in similar museum records (Erickson & Smith, 2021), these methods are generally dependent on large amounts of metadata to properly parameterize the models to give authentic estimates of absolute abundance (Guzman et al., 2021). Likewise, comprehensive land cover data at the spatial and temporal resolution required for other modeling techniques are not readily available for studies focused on time periods before contemporary remote sensing technology. Given the limitations to the available specimen and land cover data in our case, our methodology and results were constrained and must be interpreted with these constraints in mind, including that they cannot provide evidence that native lady beetle populations are declining in abundance.

Despite these data limitations, we detected evidence of shifts in dominance of lady beetle communities from native to alien species and declines in captures of several native lady beetle species, including several aphidophagous species and a coccidophagous species, via decreasing representation in the collected community of lady beetles.

The use of long-term specimen records facilitated investigation of the relative importance of the establishment of alien species and landscape change as contributing factors to the decline in captures of native lady beetles. Although the establishment of alien species and landscape change have been identified as major drivers of spatiotemporal patterns in insect populations (Fox et al., 2014; Sánchez-Bayo & Wyckhuys, 2019; Seibold et al., 2019), the causes of declines are likely more complex and multifaceted (Homburg et al., 2019; Wagner, 2020; Wagner et al., 2021). Our study underscores this complexity by documenting how captures of closely related native lady beetle species displayed opposing, species-specific responses to alien species and land cover change, as well as variation in captures over time and space that were not well explained by either of these factors. For several native species investigated, the dominance of alien lady beetles was identified as a major contributor to declines in captures, but other native species responded strongly to land cover change and captures began to decline prior to alien species establishment. Importantly, we are unable to disentangle the effects of historical changes in land cover with insecticide use and more recent intensification and simplification of agricultural landscapes on these observed trends in captures. Some native lady beetle species also responded strongly to spatiotemporal factors which may be the manifestation of climate change signals on the community, although this driver cannot be explicitly tested due to the resolution at which historical specimen data are available.

Although alien species and landscape change were important predictors, declines in captures of native lady beetles were highly species-specific, emphasizing that mechanisms driving population losses cannot be generalized even among closely related species. Additionally, this finding highlights the importance of species-level data when investigating temporal trends in insect populations. Species-specific responses of native lady beetles to these factors have important implications for conservation management programs. Increased dominance of alien lady beetles in native communities suggests that reliance on these species may be required to maintain successful pest management in the future (Bahlai et al., 2021). The importance of landscape change as a driver structuring the distributions of lady beetle populations suggests biodiversity conservation management is required at the landscape scale. Landscape scale management will need to balance the

opposing needs of native species to be effective, as there is not one inclusive strategy that will aid in native lady beetle conservation. Due to the differential responses of native species to land cover type, prioritizing habitat diversity at the landscape scale, especially in agricultural systems, may be an important component in future conservation management. Native lady beetle species are key predators of aphids, scales, psyllids, mites, fungi, and other pests (Evans, 2009; Hodek & Honěk, 2009; Weber & Lundgren, 2009), contributing broadly to biological control in agricultural systems (Caltagirone & Douth, 1989; Obrycki & Kring, 1998; Rondoni et al., 2021). Further declines of native species in response to alien lady beetles, landscape change, or other drivers such as climate change may disrupt the long-term resilience of this ecosystem service. Understanding how these major anthropogenic drivers influence long-term trends in native lady beetle populations will inform the conservation of this ecologically and economically important family of insects.

AUTHOR CONTRIBUTIONS

Mary M. Gardiner, Kayla I. Perry, and Christie A. Bahlai conceived, designed, and implemented the study. Kayla I. Perry, Christopher B. Riley, Katherine J. Turo, Leo Taylor, James Radl, Yvan A. Delgado de la flor, and Frances S. Sivakoff compiled the specimen data. Kayla I. Perry, Christie A. Bahlai, and Timothy J. Assal analyzed and interpreted the data. Kayla I. Perry wrote the first draft of the manuscript. All authors reviewed and edited the manuscript.

ACKNOWLEDGMENTS

We acknowledge the institutions and their curators for their time and assistance with compiling lady beetle specimen records: California Academy of Sciences (Susan Gin and Christopher C. Grinter); Natural History Museum of Los Angeles County (Brian Brown and Weiping Xie); Peabody Museum, Yale University (Lawrence Gall); Biodiversity Research Collections, University of Connecticut (Jane O'Donnell); Smithsonian National Museum of Natural History (Natalie Vandenburg); Collection of Arthropods, University of Georgia (Joseph V. McHugh and E. Richard Hoebeke); Gantz Family Collections Center, The Field Museum (Rebekah Shuman Baquiran); Illinois Natural History Survey, University of Illinois (Thomas McElrath); Louisiana State Arthropod Museum, Louisiana State University (Victoria Moseley Bayless); A. J. Cook Arthropod Research Collection, Michigan State University (Gary L. Parsons); Museum of Zoology, University of Michigan (Erika Tucker); University of Minnesota (Robin Thomson); Wilbur R. Enns Entomology Museum, University of Missouri

(Kristin B. Simpson); Mississippi Entomological Museum, Mississippi State University (Terence L. Schiefer); C. A. Triplehorn Insect Collection, The Ohio State University (Luciana Musetti); Cleveland Museum of Natural History (Nicole Gunter); Carnegie Museum of Natural History (Robert Androw and Robert Davidson); The Frost Entomological Museum, Pennsylvania State University; Clemson University Arthropod Collection, Clemson University (Michael Ferro); Severin-McDaniel Insect Research Collection, South Dakota State University (Louis Hesler); Brigham Young University (Shawn M. Clark); Crane Hollow Nature Preserve (Gary and Holly Coovert); Agricultural Technical Institute Teaching Collection, The Ohio State University (Jon Van Gray). Ohio lady beetle records from Boonshoft Museum of Discovery were provided by Kaitlin U. Campbell and Thomas O. Crist, and records from The Lost Ladybug Project (<http://www.lostladybug.org/>) were made available by Rebecca Smyth. This work was partially supported by a grant to CAB from the National Science Foundation DBI number 2045721; and a USDA NIFA Food, Agriculture, Natural Resources and Human Sciences Education and Literacy Initiative Postdoctoral Fellowship (2018-67012-28011) to KIP.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data and code (Bahlai & Perry, 2024) are available in Zenodo at <https://doi.org/10.5281/zenodo.11263088>. Ohio lady beetle records were compiled from the 25 institutions listed in Appendix S1: Table S1, with query details for obtaining these data in Appendix S1: Section S1.

ORCID

Kayla I. Perry  <https://orcid.org/0000-0001-9903-8057>

Christie A. Bahlai  <https://orcid.org/0000-0002-8937-8709>

Timothy J. Assal  <https://orcid.org/0000-0001-6342-2954>

Christopher B. Riley  <https://orcid.org/0000-0003-4274-1563>

Katherine J. Turo  <https://orcid.org/0000-0003-1578-4592>

Leo Taylor  <https://orcid.org/0000-0002-2367-6949>

Yvan A. Delgado de la flor  <https://orcid.org/0000-0002-8721-7303>

Mary M. Gardiner  <https://orcid.org/0000-0003-1796-6146>

REFERENCES

- Alyokhin, A., and G. Sewell. 2004. "Changes in a Lady Beetle Community Following the Establishment of Three Alien Species." *Biological Invasions* 6: 463–471.

- Angalet, G. W., J. M. Tropp, and A. N. Eggert. 1979. "*Coccinella septempunctata* in the United States: Recolonizations and Notes on Its Ecology." *Environmental Entomology* 8: 896–901.
- Bahlai, C., and K. I. Perry. 2024. "BahlaiLab/Ohio_ladybeetles: Ecological Applications Manuscript Final Code (V1.1)." Zenodo. <https://doi.org/10.5281/zenodo.11263088>.
- Bahlai, C. A., M. Colunga-Garcia, S. H. Gage, and D. A. Landis. 2013. "Long-Term Functional Dynamics of an Aphidophagous Coccinellid Community Remain Unchanged despite Repeated Invasions." *PLoS One* 8: e83407.
- Bahlai, C. A., M. Colunga-Garcia, S. H. Gage, and D. A. Landis. 2015. "The Role of Exotic Lady Beetles in the Decline of Native Lady Beetle Populations: Evidence From Long-Term Monitoring." *Biological Invasions* 17: 1005–24.
- Bahlai, C. A., C. Hart, M. T. Kavanaugh, J. D. White, R. W. Ruess, T. J. Brinkman, H. W. Ducklow, et al. 2021. "Cascading Effects: Insights From the U.S. Long Term Ecological Research Network." *Ecosphere* 12: e03430.
- Bahlai, C. A., and E. F. Zipkin. 2020. "The Dynamic Shift Detector: An Algorithm to Identify Changes in Parameter Values Governing Populations." *PLoS Computational Biology* 16: e1007542.
- Bartomeus, I., J. S. Ascher, J. Gibbs, B. N. Danforth, D. L. Wagner, S. M. Hedtke, and R. Winfree. 2013. "Historical Changes in Northeastern US Bee Pollinators Related to Shared Ecological Traits." *Proceedings of the National Academy of Sciences of the United States of America* 110: 4656–60.
- Baselga, A. 2010. "Partitioning the Turnover and Nestedness Components of Beta Diversity." *Global Ecology and Biogeography* 19: 134–143.
- Baselga, A., and C. D. L. Orme. 2012. "Betapart: An R Package for the Study of Beta Diversity." *Methods in Ecology and Evolution* 3: 808–812.
- Bell, K. O., and W. H. Whitcomb. 1964. "Field Studies on Egg Predators of the Bollworm, *Heliothis zea* (Boddie)." *The Florida Entomologist* 47: 171–180.
- Berkvens, N., J. Bonte, D. Berkvens, K. Deforce, L. Tirry, and P. De Clercq. 2008. "Pollen as an Alternative Food for *Harmonia axyridis*." *BioControl* 53: 201–210.
- Berkvens, N., C. Landuyt, K. Deforce, D. Berkvens, L. Tirry, and P. De Clercq. 2010. "Alternative Foods for the Multicoloured Asian Lady Beetle *Harmonia axyridis* (Coleoptera: Coccinellidae)." *European Journal of Entomology* 107: 189–195.
- Boakes, E. H., P. J. K. McGowan, R. A. Fuller, D. Chang-qing, N. E. Clark, K. O'Connor, and G. M. Mace. 2010. "Distorted Views of Biodiversity: Spatial and Temporal Bias in Species Occurrence Data." *PLoS Biology* 8: e1000385.
- Brooks, D. R., J. E. Bater, S. J. Clark, D. T. Monteith, C. Andrews, S. J. Corbett, D. A. Beaumont, and J. W. Chapman. 2012. "Large Carabid Beetle Declines in a United Kingdom Monitoring Network Increases Evidence for a Widespread Loss in Insect Biodiversity." *Journal of Applied Ecology* 49: 1009–19.
- Brown, P. M. J., B. Ingels, A. Wheatley, E. L. Rhule, P. de Clercq, T. van Leeuwen, and A. Thomas. 2015. "Intraguild Predation by *Harmonia axyridis* (Coleoptera: Coccinellidae) on Native Insects in Europe: Molecular Detection From Field Samples." *Entomological Science* 18: 130–33.
- Brown, P. M. J., and H. E. Roy. 2018. "Native Ladybird Decline Caused by the Invasive Harlequin Ladybird *Harmonia axyridis*: Evidence From a Long-Term Field Study." *Insect Conservation and Diversity* 11: 230–39.

- Bubna, M. 1902. "Coleoptera of Cuyahoga County, Ohio." *Ohio Naturalist* 2: 193–97.
- Caltagirone, L. E., and R. L. Doutt. 1989. "The History of the Vedalia Beetle Importation to California and Its Impact on the Development of Biological Control." *Annual Review of Entomology* 34: 1–16.
- Cardoso, P., P. S. Barton, K. Birkhofer, F. Chichorro, C. Deacon, T. Fartmann, C. S. Fukushima, et al. 2020. "Scientists' Warning to Humanity on Insect Extinctions." *Biological Conservation* 242: 108426.
- Ceballos, G., P. R. Ehrlich, A. D. Barnosky, A. García, R. M. Pringle, and T. M. Palmer. 2015. "Accelerated Modern Human-Induced Species Losses: Entering the Sixth Mass Extinction." *Science Advances* 1: e1400253.
- Chen, I.-C., H.-J. Shiu, S. Benedick, J. D. Holloway, V. K. Chey, H. S. Barlow, J. K. Hill, and C. D. Thomas. 2009. "Elevation Increases in Moth Assemblages over 42 Years on a Tropical Mountain." *Proceedings of the National Academy of Sciences of the United States of America* 106: 1479–83.
- Colla, S. R., F. Gadallah, L. Richardson, D. Wagner, and L. Gall. 2012. "Assessing Declines of North American Bumble Bees (*Bombus* spp.) Using Museum Specimens." *Biodiversity and Conservation* 21: 3585–95.
- Colunga-Garcia, M., and S. H. Gage. 1998. "Arrival, Establishment, and Habitat Use of the Multicolored Asian Lady Beetle (Coleoptera: Coccinellidae) in a Michigan Landscape." *Environmental Entomology* 27: 1574–80.
- Conrad, K. F., M. S. Warren, R. Fox, M. S. Parsons, and I. P. Woiwod. 2006. "Rapid Declines of Common, Widespread British Moths Provide Evidence of an Insect Biodiversity Crisis." *Biological Conservation* 132: 279–291.
- Crossley, M. S., K. D. Burke, S. D. Schoville, and V. C. Radeloff. 2021. "Recent Collapse of Crop Belts and Declining Diversity of US Agriculture Since 1840." *Global Change Biology* 27: 151–164.
- Dewitz, J. 2019. "National Land Cover Database (NLCD) 2016 Products (ver. 2.0, July 2020): U.S. Geological Survey Data Release." <https://doi.org/10.5066/P96HHBIE>.
- Didham, R. K., Y. Basset, C. M. Collins, S. R. Leather, N. A. Littlewood, M. H. M. Menz, J. Müller, et al. 2020. "Interpreting Insect Declines: Seven Challenges and a Way Forward." *Insect Conservation and Diversity* 13: 103–114.
- Didham, R. K., J. M. Tylianakis, N. J. Gemmell, T. A. Rand, and R. M. Ewers. 2007. "Interactive Effects of Habitat Modification and Species Invasion on Native Species Decline." *Trends in Ecology & Evolution* 22: 489–496.
- Diepenbrock, L. M., K. Fothergill, K. V. Tindall, J. E. Losey, R. R. Smyth, and D. L. Finke. 2016. "The Influence of Exotic Lady Beetle (Coleoptera: Coccinellidae) Establishment on the Species Composition of the Native Lady Beetle Community in Missouri." *Environmental Entomology* 45: 855–864.
- Dirzo, R., and P. H. Raven. 2003. "Global State of Biodiversity and Loss." *Annual Review of Environment and Resources* 28: 137–167.
- Dixon, A. F. G., and A. E. Dixon. 2000. *Insect Predator-Prey Dynamics: Ladybird Beetles and Biological Control*. Cambridge: Cambridge University Press.
- Dumas, B. A., W. P. Boyer, and W. H. Whitcomb. 1964. "Effect of Various Factors on Surveys of Predaceous Insects in Soybeans." *Journal of the Kansas Entomological Society* 37: 192–201.
- Dury, C. 1879. "List of the Coleoptera Observed in the Vicinity of Cincinnati." *The Journal of the Cincinnati Society of Natural History* 2: 162–178.
- Elliott, N. C., R. Kieckhefer, and W. Kauffman. 1996. "Effects of an Invading Coccinellid on Native Coccinellids in an Agricultural Landscape." *Oecologia* 105: 537–544.
- Entomological Collections Network (ENC). 2020. "Collections and Archives." <https://ecnweb.net/resources/collections/>.
- Erickson, K. D., and A. B. Smith. 2021. "Accounting for Imperfect Detection in Data From Museums and Herbaria When Modeling Species Distributions: Combining and Contrasting Data-Level Versus Model-Level Bias Correction." *Ecography* 44: 1341–52. <https://doi.org/10.1111/ecog.05679>.
- Evans, E. W. 2009. "Lady Beetles as Predators of Insects Other than Hemiptera." *Biological Control* 51: 255–267.
- Fox, R., T. H. Oliver, C. Harrower, M. S. Parsons, C. D. Thomas, and D. B. Roy. 2014. "Long-Term Changes to the Frequency of Occurrence of British Moths Are Consistent With Opposing and Synergistic Effects of Climate and Land-Use Changes." *Journal of Applied Ecology* 51: 949–957.
- Gagnon, A.-È., G. E. Heimpel, and J. Brodeur. 2011. "The Ubiquity of Intraguild Predation among Predatory Arthropods." *PLoS One* 6: e28061.
- Gardiner, M. M., L. L. Allee, P. M. J. Brown, J. E. Losey, H. E. Roy, and R. R. Smyth. 2012. "Lessons From Lady Beetles: Accuracy of Monitoring Data from US and UK Citizen-Science Programs." *Frontiers in Ecology and the Environment* 10: 471–76.
- Gardiner, M. M., D. A. Landis, C. Gratton, N. Schmidt, M. O'Neal, E. Mueller, J. Chacon, G. E. Heimpel, and C. D. DiFonzo. 2009. "Landscape Composition Influences Patterns of Native and Exotic Lady Beetle Abundance." *Diversity and Distributions* 15: 554–564.
- Gardiner, M. M., K. I. Perry, C. B. Riley, K. J. Turo, Y. A. Delgado de la flor, and F. S. Sivakoff. 2021. "Community Science Data Suggests that Urbanization and Forest Habitat Loss Threaten Aphidophagous Native Lady Beetles." *Ecology and Evolution* 11: 2761–74.
- Goodarzy, K., and D. W. Davis. 1958. "Natural Enemies of the Spotted Alfalfa Aphid in Utah." *Journal of Economic Entomology* 51: 612–16.
- Gordon, R. D. 1985. "The Coccinellidae (Coleoptera) of America North of Mexico." *Journal of the New York Entomological Society* 93: 912.
- Greze, A. A., T. A. Rand, T. Zaviezo, and F. Castillo-Serey. 2013. "Land Use Intensification Differentially Benefits Alien Over Native Predators in Agricultural Landscape Mosaics." *Diversity and Distributions* 19: 749–759.
- Greze, A. A., T. Zaviezo, M. M. Gardiner, and A. J. Alaniz. 2019. "Urbanization Filters Coccinellids Composition and Functional Trait Distributions in Greenspaces Across Greater Santiago, Chile." *Urban Forestry & Urban Greening* 38: 337–345.
- Grixti, J. C., L. T. Wong, S. A. Cameron, and C. Favret. 2009. "Decline of Bumble Bees (*Bombus*) in the North American Midwest." *Biological Conservation* 142: 75–84.
- Guzman, L. M., S. A. Johnson, A. O. Mooers, and L. K. M'Gonigle. 2021. "Using Historical Data to Estimate Bumble Bee Occurrence: Variable Trends Across Species Provide Little Support for Community-Level Declines." *Biological Conservation* 257: 109141.

- Habel, J. C., A. Segerer, W. Ulrich, O. Torchik, W. W. Weisser, and T. Schmitt. 2016. "Butterfly Community Shifts over Two Centuries." *Conservation Biology* 30: 754–762.
- Hallmann, C. A., M. Sorg, E. Jongejans, H. Siepel, N. Hofland, H. Schwan, W. Stenmans, et al. 2017. "More than 75 Percent Decline Over 27 Years in Total Flying Insect Biomass in Protected Areas." *PLoS One* 12: e0185809.
- Harmon, J. P., E. Stephens, and J. Losey. 2007. "The Decline of Native Coccinellids (Coleoptera: Coccinellidae) in the United States and Canada." *Journal of Insect Conservation* 11: 85–94.
- Hijmans, R. J. 2020. "raster: Geographic Data Analysis and Modeling." R package Version 3.5-2. <https://cran.r-project.org/web/packages/raster/index.html>.
- Hodek, I., and A. Honěk. 2009. "Scale Insects, Mealybugs, Whiteflies and Psyllids (Hemiptera, Sternorrhyncha) as Prey of Ladybirds." *Biological Control* 51: 232–243.
- Hodek, I., A. Honek, and H. F. Van Emden. 2012. *Ecology and Behaviour of the Ladybird Beetles (Coccinellidae)*. Chichester: John Wiley & Sons.
- Homburg, K., C. Drees, E. Boutaud, D. Nolte, W. Schuett, P. Zumstein, E. von Ruschkowski, and T. Assmann. 2019. "Where Have All the Beetles Gone? Long-Term Study Reveals Carabid Species Decline in a Nature Reserve in Northern Germany." *Insect Conservation and Diversity* 12: 268–277.
- Honěk, A., A. F. G. Dixon, A. O. Soares, J. Skuhrovec, and Z. Martinkova. 2017. "Spatial and Temporal Changes in the Abundance and Composition of Ladybird (Coleoptera: Coccinellidae) Communities." *Current Opinion in Insect Science* 20: 61–67.
- Honěk, A., Z. Martinkova, P. Kindlmann, O. M. C. C. Ameixa, and A. F. G. Dixon. 2014. "Long-Term Trends in the Composition of Aphidophagous Coccinellid Communities in Central Europe." *Insect Conservation and Diversity* 7: 55–63.
- Jeppsson, T., A. Lindhe, U. Gärdenfors, and P. Forslund. 2010. "The Use of Historical Collections to Estimate Population Trends: A Case Study Using Swedish Longhorn Beetles (Coleoptera: Cerambycidae)." *Biological Conservation* 143: 1940–50.
- Johnson, K. G., S. J. Brooks, P. B. Fenberg, A. G. Glover, K. E. James, A. M. Lister, E. Michel, et al. 2011. "Climate Change and Biosphere Response: Unlocking the Collections Vault." *Bioscience* 61: 147–153.
- Katsanis, A., D. Babendreier, W. Nentwig, and M. Kenis. 2013. "Intraguild Predation Between the Invasive Ladybird *Harmonia axyridis* and Non-target European Coccinellid Species." *BioControl* 58: 73–83.
- Kharouba, H. M., J. M. M. Lewthwaite, R. Guralnick, J. T. Kerr, and M. Vellend. 2019. "Using Insect Natural History Collections to Study Global Change Impacts: Challenges and Opportunities." *Philosophical Transactions of the Royal Society B: Biological Sciences* 374: 20170405.
- Koch, R. L. 2003. "The Multicolored Asian Lady Beetle, *Harmonia axyridis*: A Review of Its Biology, Uses in Biological Control, and Non-target Impacts." *Journal of Insect Science* 3: 32.
- Li, H., B. Li, G. L. Lövei, T. J. Kring, and J. J. Obrycki. 2021. "Interactions Among Native and Non-native Predatory Coccinellidae Influence Biological Control and Biodiversity." *Annals of the Entomological Society of America* 114: 119–136.
- Lister, A. M. 2011. "Natural History Collections as Sources of Long-Term Datasets." *Trends in Ecology & Evolution* 26: 153–54.
- Long, E. Y., and D. L. Finke. 2014. "Contribution of Predator Identity to the Suppression of Herbivores by a Diverse Predator Assemblage." *Environmental Entomology* 43: 569–576.
- Losey, J. E., J. E. Perlman, and E. R. Hoebeke. 2007. "Citizen Scientist Rediscovered Rare Nine-Spotted Lady Beetle, *Coccinella novemnotata*, in Eastern North America." *Journal of Insect Conservation* 11: 415–17.
- Losey, J. E., and M. Vaughan. 2006. "The Economic Value of Ecological Services Provided by Insects." *Bioscience* 56: 311–323.
- Lövei, G. L. 1997. "Global Change Through Invasion." *Nature* 388: 627–28.
- Mace, G. M., K. Norris, and A. H. Fitter. 2012. "Biodiversity and Ecosystem Services: A Multilayered Relationship." *Trends in Ecology and Evolution* 27: 24–31.
- Maes, D., and H. Van Dyck. 2001. "Butterfly Diversity Loss in Flanders (North Belgium): Europe's Worst Case Scenario?" *Biological Conservation* 99: 263–276.
- Majerus, M. E. 2016. *A Natural History of Ladybird Beetles*. Cambridge: Cambridge University Press.
- Mayer, M., and D. C. Allen. 1983. "*Chilocorus stigma* (Coleoptera: Coccinellidae) and Other Predators of Beech Scale in Central New York." Proceedings, IUFRO Beech Bark Disease Working Party Conference; 1982 September 26–October 8; Hamden, CT. USDA Forest Service, Northeastern Forest Experiment Station. Gen. Tech. Rep. WO-37. US Department of Agriculture, Forest Service: 89–98. pp. 89–98.
- McClure, M. S. 1986. "Role of Predators in Regulation of Endemic Populations of *Matsucoccus matsumurae* (Homoptera: Margarodidae) in Japan." *Environmental Entomology* 15: 976–983.
- Meineke, E. K., and B. H. Daru. 2021. "Bias Assessments to Expand Research Harnessing Biological Collections." *Trends in Ecology & Evolution* 36: 1071–82.
- Meineke, E. K., T. J. Davies, B. H. Daru, and C. C. Davis. 2019. "Biological Collections for Understanding Biodiversity in the Anthropocene." *Philosophical Transactions of the Royal Society B: Biological Sciences* 374: 20170386.
- Michaud, J. P. 2001. "Numerical Response of *Olla v-nigrum* (Coleoptera: Coccinellidae) to Infestations of Asian Citrus Psyllid, (Hemiptera: Psyllidae) in Florida." *The Florida Entomologist* 84: 608–612.
- Montgomery, G. A., R. R. Dunn, R. Fox, E. Jongejans, S. R. Leather, M. E. Saunders, C. R. Shortall, M. W. Tingley, and D. L. Wagner. 2020. "Is the Insect Apocalypse Upon Us? How to Find Out." *Biological Conservation* 241: 108327.
- Newbold, T., L. N. Hudson, S. L. L. Hill, S. Contu, I. Lysenko, R. A. Senior, L. Börger, et al. 2015. "Global Effects of Land Use on Local Terrestrial Biodiversity." *Nature* 520: 45–50.
- Oatman, E. R., E. F. Legner, and R. F. Brooks. 1964. "An Ecological Study of Arthropod Populations on Apple in Northeastern Wisconsin: Insect Species Present." *Journal of Economic Entomology* 57: 978–983.
- Obrycki, J. J., and T. J. Kring. 1998. "Predaceous Coccinellidae in Biological Control." *Annual Review of Entomology* 43: 295–321.
- Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, R. B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, and H. Wagner. 2011. "vegan: Community Ecology Package." R Package Version 2.5-6. <https://CRAN.R-project.org/package=vegan>.

- Ortiz-Martínez, S., K. Staudacher, V. Baumgartner, M. Traugott, and B. Lavandero. 2020. "Intraguild Predation Is Independent of Landscape Context and Does Not Affect the Temporal Dynamics of Aphids in Cereal Fields." *Journal of Pest Science* 93: 235–249.
- Parker, D. M., K. J. Turo, Y. A. Delgado de la flor, and M. M. Gardiner. 2020. "Landscape Context Influences the Abundance and Richness of Native Lady Beetles Occupying Urban Vacant Land." *Urban Ecosystems* 23: 1299–1310.
- Pebesma, E. J. 2018. "Simple Features for R: Standardized Support for Spatial Vector Data." *The R Journal* 10: 439.
- Pell, J. K., J. Baverstock, H. E. Roy, R. L. Ware, and M. E. N. Majerus. 2008. "Intraguild Predation Involving *Harmonia axyridis*: A Review of Current Knowledge and Future Perspectives." *BioControl* 53: 147–168.
- Pimentel, D., and A. G. Wheeler, Jr. 1973. "Species and Diversity of Arthropods in the Alfalfa Community." *Environmental Entomology* 2: 659–668.
- Potts, S. G., J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W. E. Kunin. 2010. "Global Pollinator Declines: Trends, Impacts and Drivers." *Trends in Ecology & Evolution* 25: 345–353.
- Putman, W. L. 1964. "Occurrence and Food of Some Coccinellids (Coleoptera) in Ontario Peach Orchards." *The Canadian Entomologist* 96: 1149–55.
- R Core Team. 2020. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Ricci, C., L. Ponti, and A. Pires. 2005. "Migratory Flight and Pre-Diapause Feeding of *Coccinella septempunctata* (Coleoptera) Adults in Agricultural and Mountain Ecosystems of Central Italy." *European Journal of Entomology* 102: 531–38.
- Ries, L., E. F. Zipkin, and R. P. Guralnick. 2019. "Tracking Trends in Monarch Abundance Over the 20th Century Is Currently Impossible Using Museum Records." *Proceedings of the National Academy of Sciences of the United States of America* 116: 13745–48.
- Rondoni, G., I. Borges, J. Collatz, E. Conti, A. C. Costamagna, F. Dumont, E. W. Evans, et al. 2021. "Exotic Ladybirds for Biological Control of Herbivorous Insects – A Review." *Entomologia Experimentalis et Applicata* 169: 6–27.
- Ross, N. 2019. "GAMs in R: Interactive Course." <https://noamross.github.io/gams-in-r/course/>.
- Roy, H. E., T. Adriaens, N. J. B. Isaac, M. Kenis, T. Onkelinx, G. S. Martin, P. M. J. Brown, et al. 2012. "Invasive Alien Predator Causes Rapid Declines of Native European Ladybirds." *Diversity and Distributions* 18: 717–725.
- Roy, H. E., P. M. J. Brown, T. Adriaens, N. Berkvens, I. Borges, S. Clusella-Trullas, R. F. Comont, et al. 2016. "The Harlequin Ladybird, *Harmonia axyridis*: Global Perspectives on Invasion History and Ecology." *Biological Invasions* 18: 997–1044.
- Rusch, A., R. Chaplin-Kramer, M. M. Gardiner, V. Hawro, J. Holland, D. Landis, C. Thies, et al. 2016. "Agricultural Landscape Simplification Reduces Natural Pest Control: A Quantitative Synthesis." *Agriculture, Ecosystems & Environment* 221: 198–204.
- Sánchez-Bayo, F., and K. A. G. Wyckhuys. 2019. "Worldwide Decline of the Entomofauna: A Review of Its Drivers." *Biological Conservation* 232: 8–27.
- Saunders, M. E., J. K. Janes, and J. C. O'Hanlon. 2020. "Moving On From the Insect Apocalypse Narrative: Engaging With Evidence-Based Insect Conservation." *Bioscience* 70: 80–89.
- Schowalter, T. D., M. Pandey, S. J. Presley, M. R. Willig, and J. K. Zimmerman. 2021. "Arthropods Are Not Declining but Are Responsive to Disturbance in the Luquillo Experimental Forest, Puerto Rico." *Proceedings of the National Academy of Sciences of the United States of America* 118: e2002556117.
- Schuch, S., K. Wesche, and M. Schaefer. 2012. "Long-Term Decline in the Abundance of Leafhoppers and Planthoppers (Auchenorrhyncha) in Central European Protected Dry Grasslands." *Biological Conservation* 149: 75–83.
- Seibold, S., M. M. Gossner, N. K. Simons, N. Blüthgen, J. Müller, D. Ambarli, C. Ammer, et al. 2019. "Arthropod Decline in Grasslands and Forests Is Associated With Landscape-Level Drivers." *Nature* 574: 671–74.
- Shaffer, H. B., R. N. Fisher, and C. Davidson. 1998. "The Role of Natural History Collections in Documenting Species Declines." *Trends in Ecology & Evolution* 13(1): 27–30.
- Shortall, C. R., A. Moore, E. Smith, M. J. Hall, I. P. Woiwod, and R. Harrington. 2009. "Long-Term Changes in the Abundance of Flying Insects." *Insect Conservation and Diversity* 2: 251–260.
- Simmons, B. I., A. Balmford, A. J. Bladon, A. P. Christie, A. De Palma, L. V. Dicks, J. Gallego-Zamorano, et al. 2019. "Worldwide Insect Declines: An Important Message, but Interpret With Caution." *Ecology and Evolution* 9: 3678–80.
- Smith, B. C. 1971. "Effects of Various Factors on the Local Distribution and Density of Coccinellid Adults on Corn (Coleoptera: Coccinellidae)." *The Canadian Entomologist* 103: 1115–20.
- Smith, C. A., and M. M. Gardiner. 2013. "Biodiversity Loss Following the Introduction of Exotic Competitors: Does Intraguild Predation Explain the Decline of Native Lady Beetles?" *PLoS One* 8: e84448.
- Snyder, W. E. 2009. "Coccinellids in Diverse Communities: Which Niche Fits?" *Biological Control* 51: 323–335.
- Snyder, W. E., G. M. Clevenger, and S. D. Eigenbrode. 2004. "Intraguild Predation and Successful Invasion by Introduced Ladybird Beetles." *Oecologia* 140: 559–565.
- Snyder, W. E., and E. W. Evans. 2006. "Ecological Effects of Invasive Arthropod Generalist Predators." *Annual Review of Ecology, Evolution, and Systematics* 37: 95–122.
- Sohl, T., R. Reker, M. Bouchard, K. Sayler, J. Dornbierer, S. Wika, R. Quenzer, and A. Friesz. 2016. "Modeled Historical Land Use and Land Cover for the Conterminous United States." *Journal of Land Use Science* 11: 476–499.
- Sohl, T., R. Reker, M. Bouchard, K. Sayler, J. Dornbierer, S. Wika, R. Quenzer, and A. Friesz. 2018. "Modeled Historical Land Use and Land Cover for the Conterminous United States: 1938–1992: U.S. Geological Survey Data Release." <https://doi.org/10.5066/F7KK99RR>.
- Staines, C. L. 2008. "Coccinellidae or Ladybird Beetles (Insecta: Coleoptera) of Plummers Island, Maryland." *Bulletin of the Biological Society of Washington* 15: 149–150.
- Staines, C., Jr., M. Rothschild, and R. Trumbule. 1990. "A Survey of the Coccinellidae (Coleoptera) Associated with Nursery Stock in Maryland." *Proceedings of the Entomological Society of Washington* 92: 310–13.
- Steffens, W. P., and R. Lumen. 2015. "Decline in Relative Abundance of *Hippodamia convergens* (Coleoptera: Coccinellidae) in Fall Shoreline Aggregations on Western Lake Superior." *Great Lakes Entomologist* 48: 8.
- Suarez, A. V., and N. D. Tsutsui. 2004. "The Value of Museum Collections for Research and Society." *Bioscience* 54: 66–74.

- Thomas, A. P., J. Trotman, A. Wheatley, A. Aebi, R. Zindel, and P. M. J. Brown. 2013. "Predation of Native Coccinellids by the Invasive Alien *Harmonia axyridis* (Coleoptera: Coccinellidae): Detection in Britain by PCR-Based Gut Analysis." *Insect Conservation and Diversity* 6: 20–27.
- Thomas, C. D., T. H. Jones, and S. E. Hartley. 2019. "Insectageddon": A Call for More Robust Data and Rigorous Analyses." *Global Change Biology* 25: 1891–92.
- Turnipseed, R., T. A. Ugine, and J. E. Losey. 2015. "Egg Predation by the Introduced Lady Beetle, *Coccinella septempunctata* (Coleoptera: Coccinellidae), Lowers Mortality but Raises Relative Risk for the Native Lady Beetle, *Coccinella novemnotata*." *PLoS One* 10(6): e0118493. <https://doi.org/10.1371/journal.pone.0118493>.
- Turnock, W. J., I. L. Wise, and F. O. Matheson. 2003. "Abundance of some Native Coccinellines (Coleoptera: Coccinellidae) Before and After the Appearance of *Coccinella septempunctata*." *The Canadian Entomologist* 135: 391–404.
- Ugine, T. A., E. Hoki, and J. E. Losey. 2018. "Interactions of *Coccinella novemnotata* (Coleoptera: Coccinellidae) and *Coccinella septempunctata* (Coleoptera: Coccinellidae) while Foraging for Aphids." *Environmental Entomology* 47: 87–92. <https://doi.org/10.1093/ee/nvx198>.
- Vitousek, P. M., H. A. Mooney, J. Lubchenco, and J. M. Melillo. 1997. "Human Domination of Earth's Ecosystems." *Science* 277: 494–99.
- Wagner, D. L. 2020. "Insect Declines in the Anthropocene." *Annual Review of Entomology* 65: 457–480.
- Wagner, D. L., E. M. Grames, M. L. Forister, M. R. Berenbaum, and D. Stopak. 2021. "Insect Decline in the Anthropocene: Death by a Thousand Cuts." *Proceedings of the National Academy of Sciences of the United States of America* 118: e2023989118.
- Warren, M. S., D. Maes, C. A. M. van Swaay, P. Goffart, H. Van Dyck, N. A. D. Bourn, I. Wynhoff, D. Hoare, and S. Ellis. 2021. "The Decline of Butterflies in Europe: Problems, Significance, and Possible Solutions." *Proceedings of the National Academy of Sciences of the United States of America* 118: e2002551117.
- Weber, D. C., and J. G. Lundgren. 2009. "Assessing the Trophic Ecology of the Coccinellidae: Their Roles as Predators and as Prey." *Biological Control* 51: 199–214.
- Wheeler, A., Jr., and E. Hoebeke. 1995. "*Coccinella novemnotata* in Northeastern North America: Historical Occurrence and Current Status (Coleoptera: Coccinellidae)." *Entomological Society of Washington* 97(3): 701–716.
- Wickham, J., S. V. Stehman, D. G. Sorenson, L. Gass, and J. A. Dewitz. 2021. "Thematic Accuracy Assessment of the NLCD 2016 Land Cover for the Conterminous United States." *Remote Sensing of Environment* 257: 112357.
- Wilcove, D. S., D. Rothstein, D. Jason, A. Phillips, and E. Losos. 1998. "Quantifying Threats to Imperiled Species in the United States." *Bioscience* 48: 607–615.
- Winker, K. 2004. "Natural History Museums in a Postbiodiversity Era." *Bioscience* 54: 455–59.
- Woltz, M. J., and D. A. Landis. 2014. "Coccinellid Response to Landscape Composition and Configuration." *Agricultural and Forest Entomology* 16: 341–49.
- Wood, S. N. 2017. *Generalized Additive Models: An Introduction with R*, 2nd ed. New York: Chapman and Hall/CRC.
- Yang, L. H., and C. Gratton. 2014. "Insects as Drivers of Ecosystem Processes." *Current Opinion in Insect Science* 2: 26–32.
- Zaviezo, T., A. A. Grez, C. F. Estades, and A. Perez. 2006. "Effects of Habitat Loss, Habitat Fragmentation, and Isolation on the Density, Species Richness, and Distribution of Ladybeetles in Manipulated Alfalfa Landscapes." *Ecological Entomology* 31: 646–656.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Perry, Kayla I., Christie A. Bahlai, Timothy J. Assal, Christopher B. Riley, Katherine J. Turo, Leo Taylor, James Radl, Yvan A. Delgado de la flor, Frances S. Sivakoff, and Mary M. Gardiner. 2024. "Landscape Change and Alien Invasions Drive Shifts in Native Lady Beetle Communities over a Century." *Ecological Applications* 34(7): e3024. <https://doi.org/10.1002/eap.3024>