

# The $P = W$ conjecture for $\mathrm{GL}_n$

By DAVESH MAULIK and JUNLIANG SHEN

## Abstract

We prove the  $P = W$  conjecture for  $\mathrm{GL}_n$  for all ranks  $n$  and curves of arbitrary genus  $g \geq 2$ . The proof combines a strong perversity result on tautological classes with the curious Hard Lefschetz theorem of Mellit. For the perversity statement, we apply the vanishing cycles constructions in our earlier work to global Springer theory in the sense of Yun, and prove a parabolic support theorem.

## Contents

0. Introduction	529
1. Perverse filtrations and vanishing cycles	532
2. Strong perversity for Chern classes	536
3. Global Springer theory	542
4. Parabolic support theorem	546
References	553

## 0. Introduction

Throughout, we work over the complex numbers  $\mathbb{C}$ .

0.1. *The  $P = W$  conjecture.* The purpose of this paper is to present a proof of the  $P = W$  conjecture by de Cataldo–Hausel–Migliorini [8] for arbitrary rank  $n$  and genus  $g \geq 2$ .

Let  $C$  be a non-singular irreducible projective curve of genus  $g \geq 2$ . For two coprime integers  $n \in \mathbb{Z}_{\geq 1}$  and  $d \in \mathbb{Z}$ , there are two moduli spaces  $M_{\mathrm{Dol}}$  and  $M_{\mathrm{B}}$ , called the Dolbeault and the Betti moduli spaces, attached to  $C, n$ , and  $d$ .

The Dolbeault moduli space parametrizes stable Higgs bundles  $(\mathcal{E}, \theta)$  on  $C$ , where  $\mathcal{E}$  is a vector bundle on  $C$  of rank  $n$  and degree  $d$ ,  $\theta : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_C^1$

---

Keywords: Higgs bundles, character variety, nonabelian Hodge theory, perverse filtration, weight filtration

AMS Classification: Primary: 14H60, 14F45, 14D20.

© 2024 Department of Mathematics, Princeton University.

is a Higgs field, and stability is defined with respect to the slope  $\mu(\mathcal{E}, \theta) = \deg(\mathcal{E})/\text{rk}(\mathcal{E})$ . The moduli space  $M_{\text{Dol}}$  admits the structure of a completely integrable system

$$h : M_{\text{Dol}} \rightarrow A := \bigoplus_{i=1}^n H^0(C, \Omega_C^1)^{\otimes i}, \quad (\mathcal{E}, \theta) \mapsto \text{char.polynomial}(\theta),$$

which is referred to as the Hitchin system [20], [21]. The Hitchin map  $h$  is surjective and proper; it is also Lagrangian with respect to the canonical holomorphic symplectic form on  $M_{\text{Dol}}$  induced by the hyper-Kähler metric. The *perverse filtration* is an increasing filtration

$$P_0 H^*(M_{\text{Dol}}, \mathbb{Q}) \subset P_1 H^*(M_{\text{Dol}}, \mathbb{Q}) \subset \cdots \subset H^*(M_{\text{Dol}}, \mathbb{Q})$$

on the (singular) cohomology of  $M_{\text{Dol}}$  governed by the topology of the Hitchin system  $h$ ; see [Section 1.1](#) for a brief review.

The Betti moduli space  $M_B$  is the (twisted) character variety associated with  $\text{GL}_n(\mathbb{C})$  and degree  $d$ . It parametrizes isomorphism classes of irreducible local systems

$$\rho : \pi_1(C \setminus \{p\}) \rightarrow \text{GL}_n(\mathbb{C}),$$

where  $\rho$  sends a loop around a chosen point  $p$  to  $e^{\frac{2\pi\sqrt{-1}d}{n} \text{Id}_n}$ . Concretely, we have

$$M_B := \left\{ a_k, b_k \in \text{GL}_n(\mathbb{C}), k = 1, 2, \dots, g : \prod_{j=1}^g [a_j, b_j] = e^{\frac{2\pi\sqrt{-1}d}{n} \text{Id}_n} \right\} // \text{GL}_n(\mathbb{C}).$$

It is an affine variety whose mixed Hodge structure admits a non-trivial weight filtration

$$W_0 H^*(M_B, \mathbb{Q}) \subset W_1 H^*(M_B, \mathbb{Q}) \subset \cdots \subset H^*(M_B, \mathbb{Q}).$$

Non-abelian Hodge theory [34], [35] gives a diffeomorphism between the two very different algebraic varieties  $M_{\text{Dol}}$  and  $M_B$ , which canonically identifies their cohomology:

$$(1) \quad H^*(M_{\text{Dol}}, \mathbb{Q}) = H^*(M_B, \mathbb{Q}).$$

The  $P = W$  conjecture by de Cataldo–Hausel–Migliorini [8] refines the identification (1); it predicts that the perverse filtration associated with the Hitchin system is matched with the (double-indexed) weight filtration associated with the Betti moduli space. This establishes a surprising connection between topology of Hitchin systems and Hodge theory of character varieties.

**CONJECTURE 0.1** (The  $P = W$  conjecture for  $\text{GL}_n$  [8]). *For any  $k, m \in \mathbb{Z}_{\geq 0}$ , we have*

$$P_k H^m(M_{\text{Dol}}, \mathbb{Q}) = W_{2k} H^m(M_B, \mathbb{Q}) = W_{2k+1} H^m(M_B, \mathbb{Q}).$$

[Conjecture 0.1](#) has previously been proven for  $n = 2$  and arbitrary genus  $g \geq 2$  by de Cataldo–Hausel–Migliorini [8], and recently for arbitrary rank  $n$  and genus 2 by de Cataldo–Maulik–Shen [11]. The compatibility between the  $P = W$  conjecture and Galois conjugation on the Betti side was proven in [13], which implies that  $P = W$  does not depend on the degree  $d$  as long as it is coprime to  $n$ . We refer to [8, §1] and the paragraphs following [11, Th. 0.2] for discussions concerning connections between [Conjecture 0.1](#) and other directions. In particular, by [3] and [27, §9.3], [Conjecture 0.1](#) implies the correspondence between Gopakumar–Vafa invariants and Pandharipande–Thomas invariants [27, Conj. 3.13] for the local Calabi–Yau 3-fold  $T^*C \times \mathbb{C}$  in any curve class  $n[C]$ .

The main result of this paper is a full proof of [Conjecture 0.1](#).

**THEOREM 0.2.** *[Conjecture 0.1](#) holds.*

[Conjecture 0.1](#) has several variants which, to the best of our knowledge, are still open. These include the version formulated for possibly singular moduli spaces, intersection cohomology, and general reductive groups [8], [10], [15], and the version formulated for moduli stacks [5]. There also exist parabolic versions of [Conjecture 0.1](#), and we expect that our argument applies to these settings using parabolic variants of the ingredients here [29], [31]. We refer to [4], [17], [37], [41] and references therein for the  $P = W$  phenomenon in other settings.

For the case of type  $A_{n-1}$  ( $\mathrm{GL}_n, \mathrm{PGL}_n, \mathrm{SL}_n$ ) and a degree  $d$  coprime to  $n$ , [Conjecture 0.1](#) (i.e., the  $P = W$  conjecture for  $\mathrm{GL}_n$ ) is equivalent to the  $P = W$  conjecture for  $\mathrm{PGL}_n$ ; see the paragraph after [11, Th. 0.2]. The case of  $\mathrm{SL}_n$  is more subtle — it is closely related to the endoscopic decomposition of the  $\mathrm{SL}_n$  Hitchin moduli spaces. A systematic discussion on this aspect can be found in [24, §5]. In the special case when  $n = p$  is a prime number, [12, Th. 0.2] implies that the  $P = W$  conjecture for  $\mathrm{GL}_p, \mathrm{SL}_p$ , and  $\mathrm{PGL}_p$  are all equivalent. In particular, we have the following immediate consequence of [Theorem 0.2](#), which generalizes the result of [8] for  $\mathrm{SL}_2$ .

**THEOREM 0.3.** *The  $P = W$  conjecture holds for a curve  $C$  of any genus at least two and  $\mathrm{SL}_p$  with  $p$  a prime number.*

**0.2. Idea of the proof.** Our proof of [Conjecture 0.1](#) has four major steps.

*Step 1: Strong perversity of Chern classes.* Using work of Markman [23], Shende [33], and Mellit [29], we may reduce the  $P = W$  conjecture to a statement about the interaction between Chern classes of the universal family and the perverse filtration. This allows us to reduce the  $P = W$  conjecture to a sheaf-theoretic statement which concerns strong perversity of Chern classes.

*Step 2: Vanishing cycle techniques.* We apply the formalism of vanishing cycles to reduce the sheaf-theoretic formulation of Step 1 for the Hitchin system

to the case of twisted Hitchin systems associated with meromorphic Higgs bundles. Our motivation is from the key observation by Ngô [30] and Chaudouard–Laumon [2], that the decomposition theorem for such twisted Hitchin systems is more manageable.

Steps 1 and 2 are carried out in [Sections 1 and 2](#).

*Step 3: Global Springer theory.* The global Springer theory of Yun [39], [40], [38] produces rich symmetries for certain Hitchin moduli spaces; this proves the sheaf-theoretic formulation of Step 1 over the *elliptic locus*. For our purpose, we need a version of global Springer theory for the stable locus over the total Hitchin base. This is described in [Section 3](#).

*Step 4: A support theorem.* Lastly, in order to extend part of Yun’s symmetries induced by Chern classes from the elliptic locus to the total Hitchin base, we prove a support theorem parallel to [2] for certain parabolic Hitchin moduli spaces. This is completed in [Section 4](#).

The idea of combining vanishing cycle functors and support theorems was also applied in [24] to give a proof of the topological mirror symmetry conjecture of Hausel–Thaddeus [18], [16].

**0.3. Acknowledgements.** We would like to thank Mark Andrea de Cataldo, Bhargav Bhatt, Ben Davison, Jochen Heinloth, and Max Lieblich for various discussions. We are especially grateful to Zhiwei Yun for explaining his thesis to us, both in his office and at the playground. We also thank the anonymous referee for careful reading and useful suggestions on the exposition. J.S. was supported by the NSF grants DMS-2134315 and DMS-2301474.

## 1. Perverse filtrations and vanishing cycles

In this section, we introduce the notion of *strong perversity* and show its compatibility with vanishing cycle functors ([Proposition 1.5](#)). This plays a crucial role in [Section 2](#) in lifting the  $P = W$  conjecture sheaf-theoretically and reduce it to the twisted case.

**1.1. Perverse filtrations.** Let  $f : X \rightarrow Y$  be a proper morphism between irreducible non-singular quasi-projective varieties (or Deligne–Mumford stacks) with  $\dim X = a$  and  $\dim Y = b$ . Let  $r$  be the defect of semismallness of  $f$ :

$$r := \dim X \times_Y X - \dim X.$$

In particular, we have  $r = a - b$  when  $f$  has equi-dimensional fibers. The perverse filtration

$$P_0 H^m(X, \mathbb{Q}) \subset P_1 H^m(X, \mathbb{Q}) \subset \cdots \subset H^m(X, \mathbb{Q})$$

is an increasing filtration on the cohomology of  $X$  governed by the topology of the morphism  $f$ ; it is defined to be

$$P_i H^m(X, \mathbb{Q}) := \mathrm{Im} \left\{ H^{m-a+r}(Y, {}^{\mathbf{P}}\tau_{\leq i}(Rf_* \mathbb{Q}_X[a-r])) \rightarrow H^m(X, \mathbb{Q}) \right\},$$

where  ${}^{\mathbf{P}}\tau_{\leq *}$  is the perverse truncation functor [1].

LEMMA 1.1. *If  $f$  has equi-dimensional fibers, the perverse filtration on  $H^m(X, \mathbb{Q})$  terminates at  $P_m H^m(X, \mathbb{Q})$ , i.e.,*

$$P_m H^m(X, \mathbb{Q}) = H^m(X, \mathbb{Q}).$$

In particular, we have  $1 \in P_0 H^0(X, \mathbb{Q})$ .

*Proof.* By definition and the decomposition theorem [1], the dimensions of the graded pieces of the perverse filtration are given by

$$(2) \quad \dim \mathrm{Gr}_i^P H^m(X, \mathbb{Q}) = \dim H^{m-i-(a-r)}(Y, {}^{\mathbf{P}}\mathcal{H}^i(Rf_* \mathbb{Q}_X[a-r])).$$

Since a perverse sheaf, as a complex of constructible sheaves, is concentrated in degrees  $[-b, 0]$ , it only has non-trivial cohomology in degrees  $\geq -b$ . In particular, the right-hand side of (2) is non-trivial only if

$$m - i - (a - r) \geq -b.$$

Since  $r = a - b$ , this inequality is equivalent to  $m \geq i$ .  $\square$

A cohomology class  $\gamma \in H^l(X, \mathbb{Q})$  can be viewed as a morphism  $\gamma : \mathbb{Q}_X \rightarrow \mathbb{Q}_X[l]$ , which naturally induces

$$(3) \quad \gamma : Rf_* \mathbb{Q}_X \rightarrow Rf_* \mathbb{Q}_X[l]$$

after pushing forward along  $f$ . For an integer  $c \geq 0$ , we say that  $\gamma \in H^l(X, \mathbb{Q})$  has *strong perversity*  $c$  with respect to  $f$  if its induced morphism (3) satisfies

$$(4) \quad \gamma({}^{\mathbf{P}}\tau_{\leq i} Rf_* \mathbb{Q}_X) \subset {}^{\mathbf{P}}\tau_{\leq i+(c-l)}(Rf_* \mathbb{Q}_X[l]) \quad \forall i;$$

more precisely, condition (4) says that the composition

$${}^{\mathbf{P}}\tau_{\leq i} Rf_* \mathbb{Q}_X \hookrightarrow Rf_* \mathbb{Q}_X \xrightarrow{\gamma} Rf_* \mathbb{Q}_X[l]$$

factors through  ${}^{\mathbf{P}}\tau_{\leq i+(c-l)}(Rf_* \mathbb{Q}_X[l]) \hookrightarrow Rf_* \mathbb{Q}_X[l]$  for any  $i$ . Notice that  $\gamma$  automatically has strong perversity  $l$ , so this condition is interesting only when  $c < l$ . Combining Lemma 1.1 and the following lemma, we see that if  $f$  has equi-dimensional fibers and  $\gamma$  has strong perversity  $c$ , then  $\gamma \in P_c H^*(X, \mathbb{Q})$ .

LEMMA 1.2. *If  $\gamma \in H^l(X, \mathbb{Q})$  has strong perversity  $c$  with respect to  $f$ , then taking cup-product with  $\gamma$  satisfies*

$$\gamma \cup - : H^m(X, \mathbb{Q}) \rightarrow H^{m+l}(X, \mathbb{Q}), \quad P_i H^m(X, \mathbb{Q}) \mapsto P_{i+c} H^{m+l}(X, \mathbb{Q}).$$

*Proof.* This follows from taking global cohomology for (4) and noticing that

$$\begin{aligned} H^{m-a+r}(Y, {}^{\mathbf{p}}\tau_{\leq i+(c-l)}(Rf_*\mathbb{Q}_X[l+a-r])) \\ = H^{(m-a+r)+l}(Y, {}^{\mathbf{p}}\tau_{\leq i+c}(Rf_*\mathbb{Q}_X[a-r])). \quad \square \end{aligned}$$

In general, the perverse filtration  $P_{\bullet}H^*(X, \mathbb{Q})$  may not be multiplicative, i.e., for  $\gamma_j \in P_{c_j}H^*(X, \mathbb{Q})$  ( $j = 1, 2, \dots, s$ ), it may not be true that

$$\gamma_1 \cup \gamma_2 \cup \dots \cup \gamma_s \in P_{c_1+\dots+c_s}H^*(X, \mathbb{Q});$$

see [6, Exer. 5.6.8]. In fact, it was proven in [11, Th. 0.6] that [Conjecture 0.1](#) is equivalent to the multiplicativity of the perverse filtration associated with the Hitchin system. The following easy observation illustrates the advantage of considering strong perversity in view of the multiplicativity issue.

**LEMMA 1.3.** *If the class  $\gamma_j \in H^{l_j}(X, \mathbb{Q})$  ( $j = 1, 2, \dots, s$ ) has strong perversity  $c_j$  with respect to  $f$ , then the cup product*

$$\gamma_1 \cup \gamma_2 \cup \dots \cup \gamma_s \in H^{l_1+\dots+l_s}(X, \mathbb{Q})$$

*has strong perversity  $\sum_j c_j$ .*

**1.2. Vanishing cycles.** Throughout [Section 1.2](#), we let  $g : X \rightarrow \mathbb{A}^1$  be a morphism such that  $X$  is non-singular and irreducible with  $X_0 = g^{-1}(0)$  the closed fiber over  $0 \in \mathbb{A}^1$ . We consider the vanishing cycle functor

$$\varphi_g : D_c^b(X) \rightarrow D_c^b(X_0)$$

which preserves the perverse  $t$ -structures,

$$\varphi_g : \text{Perv}(X) \rightarrow \text{Perv}(X_0).$$

Here  $\text{Perv}(-)$  stands for the abelian category of perverse sheaves. We denote by

$$\varphi_g := \varphi_g(\text{IC}_X) = \varphi_g(\mathbb{Q}_X[\dim X]) \in \text{Perv}(X_0)$$

the perverse sheaf of vanishing cycles. We use  $X' \subset X$  to denote the support of the vanishing cycle complex  $\varphi_g$  so that  $\varphi_g \in \text{Perv}(X')$ .

Recall that for any bounded constructible object  $\mathcal{K} \in D_c^b(X)$ , a class  $\gamma \in H^l(X, \mathbb{Q})$  induces a morphism

$$\gamma : \mathcal{K} \rightarrow \mathcal{K}[l]$$

via taking the tensor product with  $\gamma : \mathbb{Q}_X \rightarrow \mathbb{Q}_X[l]$ . The following lemma shows the compatibility between the vanishing cycle functor and restriction of cohomology classes.

LEMMA 1.4. *With the same notation as above, let  $i : X' \hookrightarrow X$  be the closed embedding. The morphism*

$$i^* \gamma : \varphi_g \rightarrow \varphi_g[l] \in D_c^b(X')$$

*induced by the class  $i^* \gamma$  (applied to the object  $\mathcal{K} = \varphi_g$ ) coincides with*

$$\varphi_g(\gamma) : \varphi_g \rightarrow \varphi_g[l] \in D_c^b(X')$$

*obtained by applying the functor  $\varphi_g$  to  $\gamma : \mathbb{Q}_X \rightarrow \mathbb{Q}_X[l]$ .*

*Proof.* Let  $\iota : X_0 \hookrightarrow X$  be the closed embedding of the closed fiber over 0. By [22, Def. 8.6.2], the vanishing cycle functor can be written as

$$(5) \quad \varphi_g(-) = \iota^* R\mathcal{H}\mathrm{om}(\mathcal{C}, -) \in D_c^b(X_0)$$

with  $\mathcal{C}$  a fixed complex of sheaves. The morphism obtained by applying  $R\mathcal{H}\mathrm{om}(\mathcal{C}, -)$  to  $\gamma : \mathbb{Q}_X \rightarrow \mathbb{Q}_X[l]$  is equivalent to the morphism induced by applying  $\gamma$  to  $R\mathcal{H}\mathrm{om}(\mathcal{C}, \mathbb{Q}_X)$ . Similarly, the functor  $\iota^*$  sends the morphism induced by  $\gamma$  to the morphism induced by  $\iota^* \gamma$ .

Therefore, if we denote by  $\iota' : X' \hookrightarrow X_0$  the closed embedding and view  $\varphi_g$  as a perverse sheaf on  $X'$ , we have an equivalent morphism

$$(6) \quad \varphi_g(\gamma) = \iota^* \gamma : \iota'_* \varphi_g \rightarrow \iota'_* \varphi_g[l] \in D_c^b(X_0).$$

Finally, after applying  $\iota'^* : D_c^b(X_0) \rightarrow D_c^b(X')$  to (6) and noticing  $\iota'^* \iota'_* = \mathrm{id}$ , we obtain that the class  $i^* \gamma = \iota'^* \iota^* \gamma$  induces

$$\varphi_g(\gamma) : \varphi_g \rightarrow \varphi_g[l] \in D_c^b(X').$$

This completes the proof.  $\square$

PROPOSITION 1.5. *Let  $g : X \rightarrow \mathbb{A}^1$  and  $X'$  be as above. Assume that  $X'$  is non-singular and*

$$(7) \quad \varphi_g \simeq \mathrm{IC}_{X'} = \mathbb{Q}_{X'}[\dim X'] \in \mathrm{Perv}(X').$$

*Assume further that we have the commutative diagram*

$$\begin{array}{ccc} X' & \xrightarrow{i} & X \\ \downarrow f' & & \downarrow f \\ Y' & \longrightarrow & Y \end{array}$$

*such that  $f$  is proper and  $g = f \circ \nu$  with  $\nu : Y \rightarrow \mathbb{A}^1$ . Then if a class  $\gamma \in H^l(X, \mathbb{Q})$  has strong perversity  $c$  with respect to  $f$ , its restriction  $i^* \gamma \in H^l(X', \mathbb{Q})$  has strong perversity  $c$  with respect to  $f'$ .*

*Proof.* By definition, the morphism

$$(8) \quad \gamma : Rf_* \mathbb{Q}_X \rightarrow Rf_* \mathbb{Q}_X[l] \in D_c^b(Y)$$

induced by  $\gamma$  satisfies

$$(9) \quad \gamma({}^{\mathbf{P}}\tau_{\leq i}Rf_*\mathbb{Q}_X) \subset {}^{\mathbf{P}}\tau_{\leq i+(c-l)}(Rf_*\mathbb{Q}_X[l]) \quad \forall i.$$

Now we apply the vanishing cycle functor  $\varphi_\nu$  to (8). On the one hand, we have the base change  $Rf'_*\circ\varphi_g \simeq \varphi_\nu\circ Rf_*$  and the fact that the vanishing cycle functor preserves the perverse  $t$ -structures. Therefore (9) implies that the morphism

$$\varphi_g(\gamma) : Rf'_*\varphi_g \rightarrow Rf'_*\varphi_g[l] \in D_c^b(Y')$$

satisfies

$$\varphi_g(\gamma)({}^{\mathbf{P}}\tau_{\leq i}Rf'_*\varphi_g) \subset {}^{\mathbf{P}}\tau_{\leq i+(c-l)}(Rf'_*\varphi_g[l]) \quad \forall i.$$

On the other hand, using the isomorphism (7) and Lemma 1.4, the above equation means precisely that the morphism  $i^*\gamma : \mathbb{Q}_{X'} \rightarrow \mathbb{Q}_{X'}[l]$  satisfies

$$(i^*\gamma)({}^{\mathbf{P}}\tau_{\leq i}Rf'_*\mathbb{Q}_{X'}) \subset {}^{\mathbf{P}}\tau_{\leq i+(c-l)}(Rf'_*\mathbb{Q}_{X'}[l]) \quad \forall i;$$

that is, the class  $i^*\gamma$  has strong perversity  $c$  with respect to  $f'$ .  $\square$

## 2. Strong perversity for Chern classes

In this section, we fix the rank  $n$  and the degree  $d$  with  $(n, d) = 1$ . In Theorem 2.6, we rephrase and then enhance Conjecture 0.1 to a statement involving  $\mathcal{L}$ -twisted Hitchin systems and strong perversity of Chern classes. It will be proven in Sections 3 and 4.

**2.1. Tautological classes.** As discussed in [11, §0.3], the  $P = W$  conjecture for  $\mathrm{GL}_n$  can be reduced to a statement involving tautological classes on  $M_{\mathrm{Dol}}$  and the perverse filtration associated with  $h : M_{\mathrm{Dol}} \rightarrow A$ , without reference to the Betti moduli space  $M_B$ . In this subsection, we recall this reduction step.

For convenience, we work with the  $\mathrm{PGL}_n$  Dolbeault moduli space to avoid normalization of a universal family as in [11]; we refer to [33] for a detailed discussion concerning the formulation of the  $P = W$  conjecture in terms of tautological classes for the  $\mathrm{PGL}_n$  Dolbeault moduli space.

Fix  $\mathcal{N} \in \mathrm{Pic}^d(C)$ . Let  $\widetilde{M}_{\mathrm{Dol}}$  be the moduli stack of stable Higgs bundles  $(\mathcal{E}, \theta)$  with  $\det(\mathcal{E}) \simeq \mathcal{N}$  and  $\mathrm{trace}(\theta) = 0$ , rigidified with respect to the generic  $\mu_n$ -stabilizer; this is the same as taking its coarse moduli space. We refer to this (non-singular) variety as the  $\mathrm{SL}_n$  Dolbeault moduli space of degree  $d$ . The finite group  $\Gamma := \mathrm{Pic}^0(C)[n]$  acts naturally on  $\widetilde{M}_{\mathrm{Dol}}$  via tensor product. The  $\mathrm{PGL}_n$  Dolbeault moduli space of degree  $d$  is recovered by taking the quotient stack

$$(10) \quad \widehat{M}_{\mathrm{Dol}} := \widetilde{M}_{\mathrm{Dol}}/\Gamma$$

which is a non-singular Deligne–Mumford stack. The  $\mathrm{PGL}_n$  Hitchin system

$$\widehat{h} : \widehat{M}_{\mathrm{Dol}} \rightarrow \widehat{A} := \bigoplus_{i=2}^n H^0(C, \Omega_C^1 \otimes^i)$$

is induced by the Hitchin map associated with  $\widetilde{M}_{\mathrm{Dol}}$  as the  $\Gamma$ -action is fiberwise with respect to  $h$ . Analogous to the  $\mathrm{GL}_n$  case, we have the perverse filtration  $P_* H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q})$  associated with  $\widehat{h}$ . The universal  $\mathrm{PGL}_n$ -bundle  $\mathcal{U}$  on  $C \times \widehat{M}_{\mathrm{Dol}}$  induces Chern characters

$$\mathrm{ch}_k(\mathcal{U}) \in H^{2k}(C \times \widehat{M}_{\mathrm{Dol}}, \mathbb{Q}), \quad k \geq 2.$$

The tautological classes  $c_k(\gamma)$  are defined to be

$$c_k(\gamma) := \int_{\gamma} \mathrm{ch}_k(\mathcal{U}) = q_{M*}(q_C^* \gamma \cup \mathrm{ch}_k(\mathcal{U})) \in H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q}), \quad \gamma \in H^*(C, \mathbb{Q}),$$

where  $q_{(-)}$  are the projections from  $C \times \widehat{M}_{\mathrm{Dol}}$ .

Now we consider the  $\mathrm{PGL}_n$  Betti moduli space of degree  $d$  with the isomorphism on cohomology provided by non-abelian Hodge theory (cf. [8, Th. 1.2.4]):

$$(11) \quad H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q}) = H^*(\widehat{M}_B, \mathbb{Q}).$$

Define the Hodge sub-vector space

$${}^k \mathrm{Hdg}^m(\widehat{M}_B) := W_{2k} H^m(M_B, \mathbb{Q}) \cap F^k H^m(M_B, \mathbb{C}) \subset H^m(\widehat{M}_B, \mathbb{Q}).$$

The following theorem, collecting results of Markman and Shende, provides a complete description of  $H^*(\widehat{M}_B, \mathbb{Q})$  in terms of the Chern classes  $\mathrm{ch}_k(\mathcal{U})$  and the weight filtration.

**THEOREM 2.1** ([23, 33]). *We use the same notation as above.*

- (i) *The tautological classes  $c_k(\gamma) \in H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q})$  generate  $H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q})$  as a  $\mathbb{Q}$ -algebra.*
- (ii) *The class  $c_k(\gamma)$ , passing through the non-abelian Hodge correspondence (11), lies in  ${}^k \mathrm{Hdg}^*(\widehat{M}_B)$ . In particular, we have a canonical decomposition*

$$H^*(\widehat{M}_B, \mathbb{Q}) = \bigoplus_{m,k} {}^k \mathrm{Hdg}^m(\widehat{M}_B).$$

*Proof.* The first part was proven in [23], and the second part was proven in [33].  $\square$

**Theorem 2.1** (ii) yields immediately that

$$W_{2k} H^m(\widehat{M}_B, \mathbb{Q}) = W_{2k+1} H^m(\widehat{M}_B, \mathbb{Q}).$$

Moreover, by **Theorem 2.1**, the  $P = W$  conjecture implies that each class  $\prod_{i=1}^s c_{k_i}(\gamma_i)$  lies in the perverse piece  $P_{\sum_i k_i} H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q})$ ; the latter is in fact equivalent to the  $P = W$  conjecture. Indeed, suppose we have

$$(12) \quad \prod_{i=1}^s c_{k_i}(\gamma_i) \in P_{\sum_i k_i} H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q})$$

for any product of tautological classes. Then we know that  $W_{2k}H^*(\widehat{M}_B, \mathbb{Q}) \subset P_kH^*(\widehat{M}_{\text{Dol}}, \mathbb{Q})$ . The curious hard Lefschetz theorem proven by Mellit [29] forces the two filtrations  $P_\bullet$  and  $W_{2\bullet}$  to coincide as long as one contains the other. This was mentioned in the last paragraph of [29, §1]; we include its proof here for the reader's convenience.

LEMMA 2.2. *We denote by  $V$  the  $\mathbb{Q}$ -vector space (11) with the perverse and the weight filtrations  $P_\bullet$  and  $W_{2\bullet}$ . If  $W_{2k} \subset P_k$  for all  $k$ , then  $W_{2k} = P_k$  for all  $k$ .*

*Proof.* Assume  $r = \dim \widehat{M}$ ; both filtrations terminate at the  $r$ -th pieces, i.e.,  $W_{2r} = P_r = V$ . We first show that

$$W_0 = P_0, \quad W_{2(r-1)} = P_{r-1}.$$

In fact, since  $W_\bullet \subset P_\bullet$ , we have

$$\dim W_0 \leq \dim P_0 = \dim V / P_{r-1} \leq \dim V / W_{2(r-1)};$$

moreover, by the curious hard Lefschetz theorem, each inequality has to be an equality. So our claim follows.

We proceed by applying the same argument to  $W_1, P_1$  and  $W_{2(r-2)}, P_{r-1}$ . The lemma follows by a simple induction.  $\square$

In conclusion, we have reduced the  $P = W$  conjecture to the following:

CONJECTURE 2.3 (Equivalent version of  $P = W$ ). *Condition (12) holds for all products of tautological classes.*

## 2.2. Strong perversity for Chern classes for $\mathcal{L}$ -twisted Hitchin systems.

For our purposes, it is important to consider Dolbeault moduli spaces of Higgs bundles, twisted by an effective line bundle  $\mathcal{L}$  (i.e.,  $H^0(C, \mathcal{L}) \neq 0$ ). These moduli spaces have already appeared in [2], [19], [18], [39], [40]; we review the construction here briefly.

Set  $\Omega_{\mathcal{L}}$  to be the line bundle  $\Omega_C^1 \otimes \mathcal{L}$  on the curve  $C$ . We denote by  $M_{\text{Dol}}^{\mathcal{L}}$  the moduli space of stable twisted Higgs bundles

$$(\mathcal{E}, \theta), \quad \theta : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_{\mathcal{L}}, \quad \text{rk}(\mathcal{E}) = n, \quad \deg(\mathcal{E}) = d,$$

with respect to the slope stability condition. The corresponding Hitchin map

$$h^{\mathcal{L}} : M_{\text{Dol}}^{\mathcal{L}} \rightarrow A^{\mathcal{L}} := \bigoplus_{i=1}^n H^0(C, \Omega_{\mathcal{L}}^{\otimes i}), \quad (\mathcal{E}, \theta) \mapsto \text{char.polynomial}(\theta),$$

is still proper as in the untwisted case; but it fails to be a Lagrangian fibration when  $\deg(\mathcal{L}) > 0$ . The  $\mathcal{L}$ -twisted  $\text{SL}_n$  and  $\text{PGL}_n$  Dolbeault moduli spaces  $\widetilde{M}_{\text{Dol}}^{\mathcal{L}}$  and  $\widehat{M}_{\text{Dol}}^{\mathcal{L}}$  can be constructed similarly. The moduli space

$$\widetilde{M}_{\text{Dol}}^{\mathcal{L}} := \{(\mathcal{E}, \theta) \in M_{\text{Dol}}^{\mathcal{L}} \mid \det(\mathcal{E}) \simeq \mathcal{N} \in \text{Pic}^d(C), \text{trace}(\theta) = 0\}$$

admits a Hitchin map

$$\check{h}^{\mathcal{L}} : \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}} \rightarrow \widehat{A}^{\mathcal{L}} := \bigoplus_{i=2}^n H^0(C, \Omega_{\mathcal{L}}^{\otimes i})$$

and a fiberwise  $\Gamma = \mathrm{Pic}^0(C)[n]$  action by tensor product. Taking the  $\Gamma$ -quotient recovers the  $\mathcal{L}$ -twisted  $\mathrm{PGL}_n$  Hitchin map

$$\widehat{h}^{\mathcal{L}} : \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}} = \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}} / \Gamma \rightarrow \widehat{A}^{\mathcal{L}}.$$

An observation in [24, §4] is that, for a fixed closed point  $p \in C$ , the  $\mathcal{L}$ -twisted and  $\mathcal{L}(p)$ -twisted  $\mathrm{SL}_n$  Dolbeault moduli spaces can be related via critical loci and vanishing cycles, which we recall in the following.

By viewing a  $\mathcal{L}$ -twisted Higgs bundle naturally as a  $\mathcal{L}(p)$ -twisted Higgs bundle, we have the natural embedding  $i : \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}} \hookrightarrow \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}(p)}$  which induces the commutative diagram

$$(13) \quad \begin{array}{ccc} \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}} & \xhookrightarrow{i} & \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}(p)} \\ \downarrow \check{h}^{\mathcal{L}} & & \downarrow \check{h}^{\mathcal{L}(p)} \\ \widehat{A}^{\mathcal{L}} & \hookrightarrow & \widehat{A}^{\mathcal{L}(p)}. \end{array}$$

We recall the following theorem from [24]:

**THEOREM 2.4** ([24, Th. 4.5]). *There exists a regular function  $g : \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}(p)} \rightarrow \mathbb{A}^1$  factorized as  $g = \nu \circ \check{h}^{\mathcal{L}(p)}$  with  $\nu : \widehat{A}^{\mathcal{L}(p)} \rightarrow \mathbb{A}^1$  such that*

$$\varphi_g \simeq \mathrm{IC}_{\widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}}} = \mathbb{Q}_{\widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}}} [\dim \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}}].$$

Since the embedding  $i : \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}} \hookrightarrow \widetilde{M}_{\mathrm{Dol}}^{\mathcal{L}(p)}$  is  $\Gamma$ -equivariant, taking  $\Gamma$ -quotients in the diagram (13) yields the following commutative diagram:

$$(14) \quad \begin{array}{ccc} \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}} & \xhookrightarrow{\widehat{i}} & \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}(p)} \\ \downarrow \widehat{h}^{\mathcal{L}} & & \downarrow \widehat{h}^{\mathcal{L}(p)} \\ \widehat{A}^{\mathcal{L}} & \hookrightarrow & \widehat{A}^{\mathcal{L}(p)}. \end{array}$$

**PROPOSITION 2.5.** *The vanishing cycle complex  $\varphi_{\widehat{g}}$  associated with the regular function*

$$\widehat{g} := \nu \circ \widehat{h}^{\mathcal{L}(p)} : \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}(p)} \rightarrow \mathbb{A}^1$$

*satisfies*

$$\varphi_{\widehat{g}} \simeq \mathrm{IC}_{\widehat{M}_{\mathrm{Dol}}^{\mathcal{L}}} = \mathbb{Q}_{\widehat{M}_{\mathrm{Dol}}^{\mathcal{L}}} [\dim \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}}].$$

*Proof.* We reduce [Proposition 2.5](#) to [Theorem 2.4](#). Consider the  $\Gamma$ -quotient map  $r : \widetilde{M}_{\text{Dol}}^{\mathcal{L}(p)} \rightarrow \widehat{M}_{\text{Dol}}^{\mathcal{L}(p)}$ . The direct image  $r_* \mathbb{Q}_{\widetilde{M}_{\text{Dol}}^{\mathcal{L}(p)}}$  admits a natural  $\Gamma$ -equivariant structure, whose invariant part recovers

$$(15) \quad \left( r_* \mathbb{Q}_{\widetilde{M}_{\text{Dol}}^{\mathcal{L}(p)}} \right)^\Gamma \simeq \mathbb{Q}_{\widehat{M}_{\text{Dol}}^{\mathcal{L}(p)}}.$$

Since  $g = \widehat{g} \circ r$ , we have

$$\begin{aligned} \varphi_{\widehat{g}} &= \varphi_{\widehat{g}} \left( \mathbb{Q}_{\widehat{M}_{\text{Dol}}^{\mathcal{L}(p)}} \left[ \dim \widehat{M}_{\text{Dol}}^{\mathcal{L}(p)} \right] \right) \\ &\simeq \varphi_{\widehat{g}} \left( r_* \mathbb{Q}_{\widetilde{M}_{\text{Dol}}^{\mathcal{L}(p)}} \left[ \dim \widetilde{M}_{\text{Dol}}^{\mathcal{L}(p)} \right] \right)^\Gamma \\ &\simeq (r_* \varphi_g)^\Gamma \\ &\simeq \mathbb{Q}_{\widehat{M}_{\text{Dol}}^{\mathcal{L}}} \left[ \dim \widehat{M}_{\text{Dol}}^{\mathcal{L}} \right]. \end{aligned}$$

Here the first equation follows by definition, the second uses (15), the third follows from the base change, and the last is given by [Theorem 2.4](#).  $\square$

Now we formulate a sheaf-theoretic enhancement of [Conjecture 2.3](#).

**THEOREM 2.6.** *There exists an effective line bundle  $\mathcal{L}$  such that the class*

$$\text{ch}_k(\mathcal{U}^{\mathcal{L}}) \in H^{2k}(C \times \widehat{M}_{\text{Dol}}^{\mathcal{L}}, \mathbb{Q})$$

*has strong perversity  $k$  with respect to*

$$\mathbf{h}^{\mathcal{L}} := \text{id} \times \widehat{h}^{\mathcal{L}} : C \times \widehat{M}_{\text{Dol}}^{\mathcal{L}} \rightarrow C \times \widehat{A}^{\mathcal{L}}.$$

*Here  $\mathcal{U}^{\mathcal{L}}$  is the universal  $\text{PGL}_n$ -bundle over  $C \times \widehat{M}_{\text{Dol}}^{\mathcal{L}}$ .*

Notice that the perversity bound here is stronger than the trivial one of  $2k$ . In the rest of this section we prove that [Theorem 2.6](#) indeed implies [Conjecture 2.3](#); therefore it implies [Conjecture 0.1](#). We prove [Theorem 2.6](#) in [Sections 3 and 4](#).

**2.3. Theorem 2.6 implies Conjecture 2.3.** We start with the following claim:

**CLAIM.** *For a point  $p \in C$ , if the class*

$$\text{ch}_k(\mathcal{U}^{\mathcal{L}(p)}) \in H^{2k}(C \times \widehat{M}_{\text{Dol}}^{\mathcal{L}(p)}, \mathbb{Q})$$

*has strong perversity  $k$  with respect to  $\mathbf{h}^{\mathcal{L}(p)} : C \times \widehat{M}_{\text{Dol}}^{\mathcal{L}(p)} \rightarrow C \times \widehat{A}^{\mathcal{L}(p)}$ , then the class*

$$\text{ch}_k(\mathcal{U}^{\mathcal{L}}) \in H^{2k}(C \times \widehat{M}_{\text{Dol}}^{\mathcal{L}}, \mathbb{Q})$$

*has strong perversity  $k$  with respect to  $\mathbf{h}^{\mathcal{L}} : C \times \widehat{M}_{\text{Dol}}^{\mathcal{L}} \rightarrow C \times \widehat{A}^{\mathcal{L}}$ .*

*Proof of claim.* We consider the commutative diagram obtained from (14):

$$(16) \quad \begin{array}{ccc} C \times \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}} & \xhookrightarrow{\mathbf{i}} & C \times \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}(p)} \\ \downarrow \mathbf{h}^{\mathcal{L}} & & \downarrow \mathbf{h}^{\mathcal{L}(p)} \\ C \times \widehat{A}^{\mathcal{L}} & \longrightarrow & C \times \widehat{A}^{\mathcal{L}(p)}. \end{array}$$

Here  $\mathbf{i} = \mathrm{id} \times \widehat{i}$ . By definition, we have  $\mathbf{i}^* \mathcal{U}^{\mathcal{L}(p)} = \mathcal{U}^{\mathcal{L}}$ . Therefore

$$\mathbf{i}^* \mathrm{ch}_k(\mathcal{U}^{\mathcal{L}(p)}) = \mathrm{ch}_k(\mathcal{U}^{\mathcal{L}}).$$

We know from [Proposition 2.5](#) that the vanishing cycle complex associated with the regular function

$$\mathbf{g} := \widehat{g} \circ \mathrm{pr}_M : C \times \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}(p)} \rightarrow \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}(p)} \rightarrow \mathbb{A}^1$$

satisfies

$$\varphi_{\mathbf{g}} \simeq \mathrm{IC}_{C \times \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}}} = \mathbb{Q}_{C \times \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}}} \left[ \dim(C \times \widehat{M}_{\mathrm{Dol}}^{\mathcal{L}}) \right].$$

Our claim then follows from [Proposition 1.5](#) (applied to the diagram (16), the class  $\mathrm{ch}_k(\mathcal{U}^{\mathcal{L}(p)})$ , and the map  $\mathbf{g}$ ).  $\square$

As a consequence of the claim, the statement of [Theorem 2.6](#) holds for  $\mathcal{L} \simeq \mathcal{O}_C$ ; i.e., the class

$$\mathrm{ch}_k(\mathcal{U}) \in H^{2k}(C \times \widehat{M}_{\mathrm{Dol}}, \mathbb{Q})$$

has strong perversity  $k$  with respect to

$$\mathbf{h} := \mathrm{id} \times \widehat{h} : C \times \widehat{M}_{\mathrm{Dol}} \rightarrow C \times \widehat{A}.$$

Finally, to deduce the cohomological statement, we note that since  $\mathbf{h}$  is the identity map restricting to  $C$ , its induced perverse filtration can be described as

$$(17) \quad P_k H^*(C \times \widehat{M}_{\mathrm{Dol}}, \mathbb{Q}) = H^*(C, \mathbb{Q}) \otimes P_k H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q}).$$

Choose a homogeneous basis of  $H^*(C, \mathbb{Q})$ ,

$$\Sigma = \{\sigma_0, \sigma_1, \dots, \sigma_{2g+2}\},$$

with Poincaré-dual basis  $\{\sigma_0^\vee, \dots, \sigma_{2g+2}^\vee\}$ .

We may express  $\mathrm{ch}_k(\mathcal{U})$  as

$$(18) \quad \mathrm{ch}_k(\mathcal{U}) = \sum_{\sigma \in \Sigma} \sigma^\vee \otimes c_k(\sigma) \in H^*(C, \mathbb{Q}) \otimes H^*(\widehat{M}_{\mathrm{Dol}}, \mathbb{Q}).$$

Since  $\mathrm{ch}_k(\mathcal{U})$  has strong perversity  $k$ , [Lemma 1.2](#) implies that

$$\mathrm{ch}_k(\mathcal{U}) \cup - : P_s H^*(C \times \widehat{M}_{\mathrm{Dol}}, \mathbb{Q}) \rightarrow P_{s+k} H^*(C \times \widehat{M}_{\mathrm{Dol}}, \mathbb{Q}).$$

Applying this operator to  $\sigma \otimes P_s H^*(\widehat{M}_{\text{Dol}}, \mathbb{Q}) \subset H^*(C \times \widehat{M}_{\text{Dol}}, \mathbb{Q})$  with  $\sigma \in \Sigma$ , we obtain from (17) and (18) that

$$c_k(\sigma) \cup - : P_s H^*(\widehat{M}_{\text{Dol}}, \mathbb{Q}) \rightarrow P_{s+k} H^*(\widehat{M}_{\text{Dol}}, \mathbb{Q}).$$

In particular, for any class  $\gamma \in H^*(C, \mathbb{Q})$ , we have

$$c_k(\gamma) : P_s H^*(\widehat{M}_{\text{Dol}}, \mathbb{Q}) \rightarrow P_{s+k} H^*(\widehat{M}_{\text{Dol}}, \mathbb{Q}),$$

which further yields

$$\prod_i c_{k_1}(\gamma_i) = \left( \prod_i c_{k_1}(\gamma_i) \right) \cup 1 \in P_{\sum_i k_i} H^*(\widehat{M}_{\text{Dol}}, \mathbb{Q}).$$

Here we used  $1 \in P_0 H^0(\widehat{M}_{\text{Dol}}, \mathbb{Q})$  in the last equation, which is given by Lemma 1.1.

This completes the proof that Theorem 2.6 implies Conjecture 2.3.  $\square$

### 3. Global Springer theory

In this section, we review Yun's global Springer theory [39], [40], [38] and use it to deduce Theorem 2.6, assuming a support theorem (Theorem 3.2). Global Springer theory was previously used to study perverse filtrations for affine Springer fibers in terms of Chern classes [31], [32]. In our setting, we require a partial extension of Yun's results from the elliptic locus to the entire Hitchin base.

**3.1. Notation.** We fix  $\mathcal{L}$  to be an effective line bundle of sufficiently large degree ( $\deg(\mathcal{L}) > 2g$  is enough for us). Since from now on we only concern the  $\mathcal{L}$ -twisted moduli spaces, we will use  $M$ ,  $\widetilde{M}$ , and  $\widehat{M}$  to denote the  $\mathcal{L}$ -twisted  $\text{GL}_n$ ,  $\text{SL}_n$ , and  $\text{PGL}_n$  Dolbeault moduli spaces  $M_{\text{Dol}}^{\mathcal{L}}$ ,  $\widetilde{M}_{\text{Dol}}^{\mathcal{L}}$ , and  $\widehat{M}_{\text{Dol}}^{\mathcal{L}}$  respectively. For the same reason, we will uniformly use the term *Higgs bundles* to call  $\mathcal{L}$ -twisted Higgs bundles.

From now on we let  $G = \text{PGL}_n$ . We let  $\mathfrak{g}$  be its Lie algebra,  $B \subset G$  the Borel subgroup induced by upper triangular matrices with  $\mathfrak{b}$  the Lie algebra,  $T \subset B$  the maximal torus given by diagonal matrices, and  $W \simeq \mathfrak{S}_n$  the Weyl group. We denote by  $\mathbb{X}^*(T)$  the character group of  $T$ ; it is isomorphic to  $\mathbb{Z}^{n-1}$  as an abelian group.

**3.2. Parabolic moduli stacks.** Let  $\widehat{\mathfrak{M}}$  be the moduli stack of  $G$ -Higgs bundles on  $C$ ; we do not impose any stability condition on  $\widehat{\mathfrak{M}}$  so that it is only a (singular) algebraic stack. The stable locus of  $\widehat{\mathfrak{M}}$  is a non-singular Deligne–Mumford substack

$$\widehat{M} \hookrightarrow \widehat{\mathfrak{M}}.$$

Yun's global Springer theory [39] constructs an algebraic stack  $\widehat{\mathfrak{M}}^{\mathrm{par}}$  over  $C \times \widehat{\mathfrak{M}}$ ,

$$(19) \quad \pi : \widehat{\mathfrak{M}}^{\mathrm{par}} \rightarrow C \times \widehat{\mathfrak{M}},$$

which is a global analog of the Grothendieck simultaneous resolution. There are two equivalent constructions of  $\widehat{\mathfrak{M}}^{\mathrm{par}}$  given in [39, §2.1].

The first is to construct  $\widehat{\mathfrak{M}}^{\mathrm{par}}$  as the moduli stack of parabolic Higgs bundles, which are quadruples  $(x, \mathcal{E}, \theta, \mathcal{E}_x^B)$ , with  $(\mathcal{E}, \theta)$  a  $G$ -Higgs bundle,  $x \in C$  a closed point, and  $\mathcal{E}_x^B$  a  $B$ -reduction of  $\mathcal{E}$  at  $x$ , satisfying the constraint that  $\theta$  is compatible with  $\mathcal{E}_x^B$ ; see [39, Def. 2.1.1]. Then the morphism  $\pi$  is given by forgetting the  $B$ -reduction:

$$\pi(x, \mathcal{E}, \theta, \mathcal{E}_x^B) = (x, (\mathcal{E}, \theta)) \in C \times \widehat{\mathfrak{M}}.$$

The second construction is via the Grothendieck simultaneous resolution  $\pi_G : [\mathfrak{b}/B] \rightarrow [\mathfrak{g}/G]$ . More precisely, let  $\rho_{\mathcal{L}}$  be the  $\mathbb{G}_m$ -torsor over  $C$  associated with the line bundle  $\Omega_{\mathcal{L}}$ . Denote by  $[\mathfrak{g}/G]_{\mathcal{L}}$  (resp.  $[\mathfrak{b}/B]_{\mathcal{L}}$ ) the family of  $[\mathfrak{g}/G]$  (resp.  $[\mathfrak{b}/B]$ ) over the curve  $C$  twisted by the torsor  $\rho_{\mathcal{L}}$ . We have a tautological evaluation map

$$(20) \quad \begin{array}{ccc} C \times \widehat{\mathfrak{M}} & \xrightarrow{\mathrm{ev}} & [\mathfrak{g}/G]_{\mathcal{L}} \\ & \searrow & \swarrow \\ & C & \end{array}$$

which is a natural  $C$ -morphism; after base change to a closed point  $x \in C$ , the map  $\mathrm{ev}$  sends a  $G$ -Higgs bundle to the evaluation of its Higgs field at  $x$ . The morphism (19) is then induced by the base change of the Grothendieck simultaneous resolution along the evaluation map (see [39, Lemma 2.1.2]):

$$\begin{array}{ccc} \widehat{\mathfrak{M}}^{\mathrm{par}} & \xrightarrow{\mathrm{ev}^p} & [\mathfrak{b}/B]_{\mathcal{L}} \\ \downarrow \pi & & \downarrow \pi_G \\ C \times \widehat{\mathfrak{M}} & \xrightarrow{\mathrm{ev}} & [\mathfrak{g}/G]_{\mathcal{L}}. \end{array}$$

The parabolic Hitchin system

$$\widehat{h}^{\mathrm{par}} : \widehat{\mathfrak{M}}^{\mathrm{par}} \rightarrow C \times \widehat{A}$$

is the composition of  $\pi : \widehat{\mathfrak{M}}^{\mathrm{par}} \rightarrow C \times \widehat{\mathfrak{M}}$  and the morphism  $\mathbf{h} : C \times \widehat{\mathfrak{M}} \rightarrow C \times \widehat{A}$  induced by the standard (stacky) Hitchin map  $\widehat{h} : \widehat{\mathfrak{M}} \rightarrow \widehat{A}$ .<sup>1</sup>

In general the moduli stack  $\widehat{\mathfrak{M}}^{\mathrm{par}}$  is singular, and the parabolic Hitchin map is not proper. In the next section we will impose a stability condition and then restrict to the stable locus.

<sup>1</sup>Recall that we always use the  $\mathcal{L}$ -twisted version.

3.3. *Stable loci.* We define the *stable locus* of the moduli of parabolic  $G$ -Higgs bundles to be

$$\widehat{M}^{\text{par}} := (C \times \widehat{M}) \times_{C \times \widehat{\mathfrak{M}}} \widehat{\mathfrak{M}}^{\text{par}};$$

equivalently, it fits into the Cartesian diagrams

$$(21) \quad \begin{array}{ccccc} \widehat{M}^{\text{par}} & \longrightarrow & \widehat{\mathfrak{M}}^{\text{par}} & \xrightarrow{\text{ev}^p} & [\mathfrak{b}/B]_{\mathcal{L}} \\ \downarrow \pi & & \downarrow \pi & & \downarrow \pi_G \\ C \times \widehat{M} & \hookrightarrow & C \times \widehat{\mathfrak{M}} & \xrightarrow{\text{ev}} & [\mathfrak{g}/G]_{\mathcal{L}}. \end{array}$$

We use the same notation as for the stacky case to denote the parabolic Hitchin map restricted to the stable locus:

$$(22) \quad \widehat{h}^{\text{par}} : \widehat{M}^{\text{par}} \rightarrow C \times \widehat{A}.$$

PROPOSITION 3.1.

- (i) *The moduli stack  $\widehat{M}^{\text{par}}$  is non-singular and Deligne–Mumford; and*
- (ii) *the parabolic Hitchin map (22) is proper.*

*Proof.* The Deligne–Mumford part of (i) follows from the proof of [39, Prop. 2.5.1 (3)]. Indeed, the left vertical arrow in the diagram (21) is the pullback of  $\pi_G$ , therefore it is schematic and of finite type. This implies that  $\widehat{M}^{\text{par}}$  is Deligne–Mumford since  $C \times \widehat{M}$  is Deligne–Mumford.

To prove the smoothness part of (i), we use the evaluation map (20). Recall that [24, Prop. 4.1] (which was proven via deformation theory) shows that the  $C$ -morphism  $\text{ev}$  is smooth after restricting over the stable locus  $C \times \widehat{M}$ . Hence by the Cartesian diagrams of (21), the evaluation map

$$\text{ev} : \widehat{M}^{\text{par}} \rightarrow [\mathfrak{b}/B]_{\mathcal{L}}$$

is also smooth. Since the target is a non-singular algebraic stack, so is the source.

(ii) follows directly from (21) and that  $\widehat{h} : \widehat{M} \rightarrow \widehat{A}$  is proper.  $\square$

As a consequence of Proposition 3.1, the direct image complex

$$R\widehat{h}_*^{\text{par}} \mathbb{Q}_{\widehat{M}^{\text{par}}} \in D_c^b(C \times \widehat{A})$$

satisfies the decomposition theorem [1].

THEOREM 3.2 (Support theorem for parabolic Hitchin map). *The decomposition for the parabolic Hitchin map  $\widehat{h}^{\text{par}}$  has full support; i.e., any non-trivial simple perverse summand of*

$$\mathfrak{p}\mathcal{H}^i(R\widehat{h}_*^{\text{par}} \mathbb{Q}_{\widehat{M}^{\text{par}}}) \quad \forall i \in \mathbb{Z}$$

*has support  $C \times \widehat{A}$ .*

We will postpone the proof of [Theorem 3.2](#) to [Section 4](#).

**3.4. Proof of Theorem 2.6.** We first prove [Theorem 2.6](#) assuming [Theorem 3.2](#).

There are three ingredients of global Springer theory

$$\pi : \widehat{M}^{\mathrm{par}} \rightarrow C \times \widehat{M}$$

which are important in the proof of [Theorem 2.6](#). We summarize them as follows.

(A) (*Splitting of the universal  $G$ -bundle*). As explained in the second bullet point of [39, Construction 6.1.4], each Chern root of  $\pi^* \mathcal{U}$  is of the form  $c_1(L(\xi))$ , where  $\xi$  is an element in  $\mathbb{X}^*(T)$  and  $L(\xi)$  is the tautological line bundle on  $\widehat{M}^{\mathrm{par}}$  associated with  $\xi$ .

More precisely,  $\pi^* \mathcal{U}$  is the universal  $G$ -bundle on  $\widehat{M}^{\mathrm{par}}$  whose fiber over  $(x, \mathcal{E}, \theta, \mathcal{E}_x^B)$  is  $\mathcal{E}_x$ ; the  $B$ -reduction  $\mathcal{E}_x^B$  yields a  $T$ -torsor over  $\widehat{M}^{\mathrm{par}}$  which induces for each  $\xi \in \mathbb{X}^*(T)$  a line bundle  $L(\xi)$ .

(B) (*Strong perversity for  $c_1(L(\xi))$* ). By [40, Lemma 3.2.3], there exists a Zariski dense open subset of  $C \times \widehat{A}$  over which the operator

$$c_1(L(\xi)) : R\widehat{h}_*^{\mathrm{par}} \mathbb{Q}_{\widehat{M}^{\mathrm{par}}} \rightarrow R\widehat{h}_*^{\mathrm{par}} \mathbb{Q}_{\widehat{M}^{\mathrm{par}}} [2]$$

has strong perversity 1 for any  $\xi \in \mathbb{X}^*(T)$ . Since  $c_1(L(\xi))$  automatically has strong perversity 2, showing it has strong perversity 1 is equivalent to showing that the induced morphism of perverse cohomology sheaves

$${}^p \mathcal{H}^i \left( R\widehat{h}_*^{\mathrm{par}} \mathbb{Q}_{\widehat{M}^{\mathrm{par}}} \right) \rightarrow {}^p \mathcal{H}^i \left( R\widehat{h}_*^{\mathrm{par}} \mathbb{Q}_{\widehat{M}^{\mathrm{par}}} [2] \right)$$

vanishes for each  $i$ . Once we have [Theorem 3.2](#), these sheaves have full support over the entire base, so this vanishing (and thus strong perversity 1) extends over the total base  $C \times \widehat{A}$  as well. In fact, [40, Lemma 3.2.3] was proven in exactly this way using a support theorem [40, §4.6.2] for the elliptic locus.

(C) (*Springer's Weyl group action*). By the Cartesian diagrams (21), we may pullback Springer's sheaf-theoretic Weyl group action from the Grothendieck simultaneous resolution; in particular, the object  $R\pi_* \mathbb{Q}_{\widehat{M}^{\mathrm{par}}}$  admits a canonical  $W$ -action whose invariant part recovers the trivial local system

$$(R\pi_* \mathbb{Q}_{\widehat{M}^{\mathrm{par}}})^W = \mathbb{Q}_{C \times \widehat{M}}.$$

By taking global cohomology we have

$$H^*(\widehat{M}^{\mathrm{par}}, \mathbb{Q}) = H^*(C \times \widehat{M}, \mathbb{Q}) \oplus (\text{variant part of } W).$$

Now we prove [Theorem 2.6](#). We first prove its parabolic version.

**CLAIM.** *The class*

$$\mathrm{ch}_k(\pi^* \mathcal{U}) \in H^{2k}(\widehat{M}^{\mathrm{par}}, \mathbb{Q})$$

*has strong perversity  $k$  with respect to the parabolic Hitchin system (22).*

*Proof of claim.* By (A), the Chern character  $\text{ch}_k(\pi^* \mathcal{U})$  can be expressed in terms of  $c_1(L(\xi))$ . Hence the claim follows from [Lemma 1.3](#) and the strong perversity of  $c_1(L(\xi))$  given by (B).  $\square$

Next, we reduce [Theorem 2.6](#) to the claim above via the following lemma.

**LEMMA 3.3.** *For a class  $\gamma \in H^l(C \times \widehat{M}, \mathbb{Q})$ , if  $\pi^* \gamma \in H^l(\widehat{M}^{\text{par}}, \mathbb{Q})$  has strong perversity  $k$  with respect to  $\widehat{h}^{\text{par}} : \widehat{M}^{\text{par}} \rightarrow C \times \widehat{A}$ , then  $\gamma$  has strong perversity  $k$  with respect to  $\mathbf{h} : C \times \widehat{M} \rightarrow C \times \widehat{A}$ .*

*Proof.* This is a consequence of (C). We write the class  $\gamma$  as a map

$$(23) \quad \gamma : \mathbb{Q}_{C \times \widehat{M}} \rightarrow \mathbb{Q}_{C \times \widehat{M}}[l],$$

whose pullback

$$(24) \quad \pi^* \gamma : \mathbb{Q}_{\widehat{M}^{\text{par}}} \rightarrow \mathbb{Q}_{\widehat{M}^{\text{par}}}[l]$$

recovers the class  $\pi^* \gamma \in H^l(\widehat{M}^{\text{par}}, \mathbb{Q})$ . By the projection formula  $\pi_* \pi^* \gamma = |W| \cdot \gamma$ , we may recover the morphism (23) from (24) by derived pushing forward to  $C \times \widehat{M}$  and taking the  $W$ -invariant part.

Now by the assumption we know that the action of  $\pi^* \gamma$  on the object  $R\widehat{h}_*^{\text{par}} \mathbb{Q}_{\widehat{M}^{\text{par}}}$  satisfies

$$(25) \quad \pi^* \gamma : {}^{\mathbf{p}} \tau_{\leq i} R\widehat{h}_*^{\text{par}} \mathbb{Q}_{\widehat{M}^{\text{par}}} \rightarrow {}^{\mathbf{p}} \tau_{\leq i+(k-l)} (R\widehat{h}_*^{\text{par}} \mathbb{Q}_{\widehat{M}^{\text{par}}}[l]).$$

Since we have

$$R\widehat{h}_*^{\text{par}} \mathbb{Q}_{\widehat{M}^{\text{par}}} = R\mathbf{h}_* (R\pi_* \mathbb{Q}_{\widehat{M}^{\text{par}}}),$$

the ingredient (C) produces a natural  $W$ -action on this object whose invariant part recovers  $R\mathbf{h}_* \mathbb{Q}_{C \times \widehat{M}}$ ; the operator

$$(26) \quad \gamma : R\mathbf{h}_* \mathbb{Q}_{C \times \widehat{M}} \rightarrow R\mathbf{h}_* \mathbb{Q}_{C \times \widehat{M}}[l]$$

is then recovered from the  $W$ -invariant part of the action of  $\pi^* \gamma$  on  $R\widehat{h}_*^{\text{par}} \mathbb{Q}_{\widehat{M}^{\text{par}}}$ . In particular, the desired property concerning (26) follows from the  $W$ -invariant part of (25).  $\square$

Thus we have completed the proof of [Theorem 2.6](#).  $\square$

#### 4. Parabolic support theorem

In the last section, we prove [Theorem 3.2](#). This is the parabolic version of the Chaudouard–Laumon support theorem [2]. For the proof, we ultimately reduce it to a relative dimension bound which we establish in [Section 4.4](#).

4.1. *Review of support theorems.* We start with a review of support theorems for Hitchin systems.

For a proper morphism  $f : X \rightarrow Y$  with  $X, Y$  non-singular Deligne–Mumford stacks, the decomposition theorem [1] yields

$$Rf_* \mathbb{Q}_X \simeq \bigoplus_i {}^p\mathcal{H}^i(Rf_* \mathbb{Q}_X)[-i]$$

with  ${}^p\mathcal{H}^i(Rf_* \mathbb{Q}_X)$  semisimple perverse sheaves on  $Y$ ; we say that a closed  $Z \subset Y$  is a support of  $f$  if it is a support of a simple summand of some  ${}^p\mathcal{H}^i(Rf_* \mathbb{Q}_X)$ . A particularly interesting case is that  $f$  has full support, that is,  $Y$  is the only support of  $f$ ; in this case the cohomology of any closed fiber of  $f$  is governed by the non-singular fibers.

The study of supports for Hitchin systems was initiated by B. C. Ngô and is crucial in his proof of the fundamental lemma of the Langlands program [30]. He determines all the supports for the Hitchin system (including the  $\mathcal{L}$ -twisted cases) after restricting to the elliptic locus of the Hitchin base; that is the subset formed by integral spectral curves.

After Ngô’s work, Chaudouard–Laumon [2] observed that, if we consider the moduli space of  $\mathcal{L}$ -twisted  $\mathrm{GL}_n$  stable Higgs bundles with  $\deg(\mathcal{L}) > 0$ , then Ngô’s support theorem can be extended to the total Hitchin base; in particular they showed that the  $\mathcal{L}$ -twisted  $\mathrm{GL}_n$  Hitchin system has full support. Chaudouard–Laumon’s idea was extended to the  $\mathrm{SL}_n$  case [7], the endoscopic moduli spaces [24], and singular cases involving strictly semistable Higgs bundles [26], [25]. See also [9], [28] concerning the supports over the open subset of reduced spectral curves for the untwisted (i.e.,  $\mathcal{L} = 0$ ) Hitchin system.

The idea of Chaudouard–Laumon [2] is to show that each support of the  $\mathcal{L}$ -twisted Hitchin system has generic point lying in the elliptic locus; this is achieved by combining two constraints: (I) the support inequality for a weak abelian fibration, and (II)  $\delta$ -regularity for spectral curves. We describe them in more detail.

- (I) (*Support inequality*). The Hitchin system admits the structure of a weak abelian fibration; that is, there is a commutative group scheme  $P$  over the Hitchin base acting on the moduli space which satisfies certain properties. Then an argument generalizing the Goresky–MacPherson inequality leads to a codimension estimate for the supports. More concretely, it says that any support  $Z$  has codimension bounded above by the  $\delta$ -function of the group  $P$ . This part was already carried out in Ngô [30, §7]; see [26, §1] for a summary.
- (II) ( *$\delta$ -regularity*). In the case of type  $A$  and for the stable locus, the group scheme is obtained from the multi-degree 0 relative Picard (or, for  $\mathrm{SL}_n$ , Prym) variety associated with the family of spectral curves. Then we

have the Severi inequality, referred to as  *$\delta$ -regularity*, for the spectral curves. We refer to [7, Th. 5.4.4] for the precise statement.

Using (I) and (II), one can deduce that no support is allowed to have a generic point lying outside the open subset of integral curves; this was explained in [2, §11] for  $GL_n$ , in [7, §6.2] for  $SL_n$ , and in [26, §4.5] for a general complete linear system in a del Pezzo surface. The argument is to combine the two inequalities above to deduce a numerical contradiction if there is a support that appears outside the elliptic locus.

**4.2. Parabolic Hitchin systems.** Now we focus on the  $\mathcal{L}$ -twisted parabolic Hitchin system (22). Let  $\widehat{A}^{\text{ell}} \subset \widehat{A}$  be the open subset parametrizing integral spectral curves (the elliptic locus). Following Ngô's method, Yun [40], [38] proved a parabolic support theorem over  $C \times \widehat{A}^{\text{ell}}$  and determined all the supports of (22) in  $C \times \widehat{A}^{\text{ell}}$ . More precisely, by [38, §2] any strict subset of  $C \times \widehat{A}^{\text{ell}}$ , which is a support of  $\widehat{h}^{\text{par}}|_{C \times \widehat{A}^{\text{ell}}}$ , is a component of the *endoscopic loci* governed by the endoscopic theory of  $G$ . As we only consider the special case  $G = \text{PGL}_n$ , there are no non-trivial endoscopic loci and the restricted Hitchin map  $\widehat{h}^{\text{par}}|_{C \times \widehat{A}^{\text{ell}}}$  has full support.

To prove [Theorem 3.2](#), it suffices to show that there is no support of (22) whose generic point is lying outside  $C \times \widehat{A}^{\text{ell}}$ .

Since stability for a  $\text{PGL}_n$  Higgs bundle is described by its corresponding vector bundle (see (10)), it is more convenient to work with the  $SL_n$  moduli spaces. We consider the following Cartesian diagram:

$$(27) \quad \begin{array}{ccc} \widetilde{M}^{\text{par}} & \xrightarrow{(-)/\Gamma} & \widehat{M}^{\text{par}} \\ \downarrow \pi' & & \downarrow \pi \\ C \times \widetilde{M} & \xrightarrow{(-)/\Gamma} & C \times \widehat{M}. \end{array}$$

Here the horizontal maps are given by the natural quotient maps by  $\Gamma = \text{Pic}^0(C)[n]$ . To describe the map  $\pi'$  on the left side of the diagram (see [39, Exam. 2.2.5]), we recall that  $\widetilde{M}$  parametrizes traceless Higgs bundles with fixed determinant

$$(\mathcal{E}, \theta), \quad \theta : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_{\mathcal{L}}, \quad \text{rk}(\mathcal{E}) = n, \quad \det(\mathcal{E}) \simeq \mathcal{N}, \quad \text{trace}(\theta) = 0$$

with respect to slope stability. Similarly  $\widetilde{M}^{\text{par}}$  parametrizes

$$(x, \mathcal{E} = \mathcal{E}_0 \supset \mathcal{E}_1 \supset \cdots \supset \mathcal{E}_n = \mathcal{E}_0(-x), \theta),$$

where  $x \in C$ ,  $(\mathcal{E}, \theta) \in \widetilde{M}$ , each  $\mathcal{E}_i$  in the flag is of rank  $n$  with  $\mathcal{E}_i/\mathcal{E}_{i+1}$  a length 1 skyscraper sheaf supported at  $x$ , the Higgs field preserves the flag  $\theta(\mathcal{E}_i) \subset \mathcal{E}_i \otimes \Omega_{\mathcal{L}}$ , and the map  $\pi'$  in the diagram is the forgetful map. We note that the stability condition for a parabolic Higgs bundle is determined by the stability of the underlying Higgs bundle.

Following [39, Exam. 2.2.5] we may also describe  $\breve{M}^{\mathrm{par}}$  via spectral curves. If we present a point in  $\breve{M}$  as a 1-dimensional sheaf  $\mathcal{F}_\alpha$  supported on a spectral curve  $C_\alpha \subset \mathrm{Tot}(\Omega_{\mathcal{L}})$  with  $p_\alpha : C_\alpha \rightarrow C$  the projection, then for any  $x \in C$ , a closed point in the fiber  $\pi'^{-1}(x, \mathcal{F}_\alpha)$  is represented by

$$\mathcal{F}_\alpha = \mathcal{F}_0 \supset \mathcal{F}_1 \supset \cdots \supset \mathcal{F}_n = \mathcal{F}_0 \otimes p_\alpha^* \mathcal{O}_C(-x), \quad \mathrm{length}(\mathcal{F}_i/\mathcal{F}_{i+1}) = 1;$$

see [39, (2.4)].

Now we consider the  $\mathrm{SL}_n$  parabolic Hitchin system

$$(28) \quad \check{h}^{\mathrm{par}} : \breve{M}^{\mathrm{par}} \rightarrow C \times \breve{M} \rightarrow C \times \widehat{A}.$$

From the diagram (27), it recovers the  $\mathrm{PGL}_n$  Hitchin system  $\widehat{h}^{\mathrm{par}} : \widehat{M}^{\mathrm{par}} \rightarrow C \times \widehat{A}$  by taking the quotient of the source by  $\Gamma$ . In particular, we have

$$R\widehat{h}_*^{\mathrm{par}} \mathbb{Q}_{\widehat{M}^{\mathrm{par}}} = (R\check{h}_*^{\mathrm{par}} \mathbb{Q}_{\breve{M}^{\mathrm{par}}})^\Gamma \in D_c^b(C \times \widehat{A}).$$

Hence in order to prove [Theorem 3.2](#), it suffices to show that there is no support of (28) with generic point lying outside  $C \times \widehat{A}^{\mathrm{ell}}$ . In the next two sections, we adapt the strategy of Chaudouard–Laumon [2] and de Cataldo [7] (for the  $\mathrm{SL}_n$ -version) to this parabolic setting, and verify the parabolic analogs of (I) and (II) of [Section 4.1](#).

**4.3. Weak abelian fibrations and  $\delta$ -regularity.** A general approach for proving support theorems was given in [26, Th. 1.8]. We apply it here for the parabolic Hitchin system (28). In this section, we explain how existing results allow us to reduce [Theorem 3.2](#) to a relative dimension bound (30). Then in the next section we prove this bound.

For the reader's convenience, we first recall some of the necessary ingredients. As mentioned in (I) of [Section 4.1](#), the notion of *weak abelian fibration* introduced by Ngô [30] plays a key role. A weak abelian fibration is a triple  $(\mathcal{P}, \mathcal{M}, \mathcal{A})$  with morphisms

$$f : \mathcal{M} \rightarrow \mathcal{A}, \quad g : \mathcal{P} \rightarrow \mathcal{A},$$

such that all  $\mathcal{P}, \mathcal{M}, \mathcal{A}$  are non-singular,  $f$  is proper,  $g : \mathcal{P} \rightarrow \mathcal{A}$  is a smooth group scheme, and  $\mathcal{P}$  acts on  $\mathcal{M}$  relatively over  $\mathcal{A}$ ; they satisfy the following conditions:

- (i) every closed fiber of the  $\mathcal{P} \rightarrow \mathcal{A}$  has dimension  $= \dim \mathcal{M} - \dim \mathcal{A}$ ;
- (ii) the action of  $\mathcal{P}$  on  $\mathcal{M}$  has affine stabilizers; and
- (iii) the Tate module associated with  $\mathcal{P} \rightarrow \mathcal{A}$  is polarizable.

We refer to [26, §1.1] for more details.

Now we show that (28) is naturally enhanced into a weak abelian fibration. We first construct the  $(C \times \widehat{A})$ -group scheme  $P$  which acts on  $\breve{M}^{\mathrm{par}}$ . Recall that in [7] de Cataldo showed that  $\breve{M}$  admits a weak abelian fibration structure  $(\check{P}, \breve{M}, \widehat{A})$ . Here the  $\widehat{A}$ -group scheme  $\check{P}$  which acts on  $\breve{M}$  is given by the identity

component of the relative Prym variety associated with the spectral curves [7, (43)]. For a spectral curve  $p_\alpha : C_\alpha \rightarrow C$  with a Higgs bundle given by  $\mathcal{F}_\alpha$  supported on  $C_\alpha$ , the Prym action is induced by tensor product

$$\mathcal{Q} \cdot \mathcal{F}_\alpha = \mathcal{Q} \otimes \mathcal{F}_\alpha, \quad \mathcal{Q} \in \text{Prym}^0(C_\alpha/C) = \check{P}_{[C_\alpha]}, \quad [C_\alpha] \in \widehat{A}.$$

We denote by  $P$  the  $(C \times \widehat{A})$ -group scheme obtained by the pullback of  $\check{P}$ . The  $P$ -action on  $C \times \widetilde{M}$  can be lifted to  $\widetilde{M}^{\text{par}}$ :

$$\begin{aligned} \mathcal{Q} \cdot (x, \mathcal{F}_\alpha = \mathcal{F}_0 \supset \mathcal{F}_1 \supset \cdots \supset \mathcal{F}_n) \\ = (x, \mathcal{Q} \otimes \mathcal{F}_\alpha = \mathcal{Q} \otimes \mathcal{F}_0 \supset \mathcal{Q} \otimes \mathcal{F}_1 \supset \cdots \supset \mathcal{Q} \otimes \mathcal{F}_n), \end{aligned}$$

since the stability condition does not rely on the flag.

**PROPOSITION 4.1.** *The triple  $(P, \widetilde{M}^{\text{par}}, C \times \widehat{A})$  forms a weak abelian fibration; i.e., it satisfies (i), (ii), and (iii) above.*

*Proof.* (i) only concerns the dimension of  $\widetilde{M}$  which is clear from (21), and (iii) only depends on the group scheme  $P$  which follows from the corresponding property for  $\check{P}$  proved in [7, Th. 4.7.2]. To prove (ii): since the  $P$ -action on  $\widetilde{M}^{\text{par}}$  is a lifting of the  $P$ -action on  $C \times \widetilde{M}$ , and the latter has affine stabilizers by [7], we obtain that the  $P$ -stabilizer for any point  $z \in \widetilde{M}^{\text{par}}$  is contained in the (affine)  $P$ -stabilizer for  $\pi(z) \in C \times \widetilde{M}$ . This proves (ii).  $\square$

Next, we consider the  $\delta$ -function on the base  $C \times \widehat{A}$ . The group scheme  $P$  endows  $C \times \widehat{A}$  with an upper semi-continuous function

$$\delta : C \times \widehat{A} \rightarrow \mathbb{N}$$

calculating the dimension of the affine part of the commutative group scheme given by each closed fiber. For a closed subset  $Z \subset C \times \widehat{A}$ , we define  $\delta_Z$  to be the minimal value of the function  $\delta$  on  $Z$ .

We say that the weak abelian fibration  $(P, \widetilde{M}^{\text{par}}, C \times \widehat{A})$  given by [Proposition 4.1](#) satisfies the *support inequality* ((I) of [Section 4.1](#)) if the inequality

$$(29) \quad \text{codim}_{C \times \widehat{A}} Z \leq \delta_Z$$

holds for any irreducible support  $Z \subset C \times \widehat{A}$  associated with (28). Furthermore, by [26, Th. 1.8], the support inequality (29) follows from the relative dimension bound

$$(30) \quad \tau_{>2\mathbf{d}} (R\check{h}_*^{\text{par}} \mathbb{Q}_{\widetilde{M}^{\text{par}}}) = 0, \quad \mathbf{d} := \dim \widetilde{M}^{\text{par}} - \dim(C \times \widehat{A})$$

with  $\tau_{>*}$  the standard truncation functor.

Once we have (30), we can combine the support inequality (29) with  $\delta$ -regularity of  $\check{P}$  [7, Cor. 5.4.4] to prove [Theorem 3.2](#) as follows. If  $Z \subset C \times \widehat{A}$

is an irreducible support of  $\check{h}^{\mathrm{par}}$ , the projection  $W = \mathrm{pr}_{\widehat{A}}(Z) \subset \widehat{A}$  also satisfies the support inequality

$$\mathrm{codim}_{\widehat{A}} W \leq \mathrm{codim}_{C \times \widehat{A}} Z \leq \delta_Z = \delta_W.$$

Here, in the second inequality we used the support inequality (29) for  $\check{h}^{\mathrm{par}}$ , and in the last equality we used that the  $\delta$ -function on  $C \times \widehat{A}$  is pulled back from  $\widehat{A}$ . The argument of [7, §6.2] then shows that the generic point of  $W$  lies in  $\widehat{A}^{\mathrm{ell}}$ . Hence the generic point of  $Z$  lies in  $C \times \widehat{A}^{\mathrm{ell}}$ , as desired.

In conclusion, this reduces [Theorem 3.2](#) to (30).

**4.4. Proof of the relative dimension bound (30).** We need to show that each closed fiber of the morphism

$$\check{h}^{\mathrm{par}} : \check{M}^{\mathrm{par}} \rightarrow C \times \widehat{A}$$

has dimension

$$\mathbf{d} = \dim \check{M}^{\mathrm{par}} - \dim(C \times \widehat{A}) = \frac{n(n-1)}{2} \deg(\mathcal{L}) + (n^2 - 1)(g-1).$$

Here the last equation is obtained by the dimension formula of [7, Prop. 2.4.6] directly:

$$\mathbf{d} = \dim \check{M} - \dim \widehat{A} = \frac{n(n-1)}{2} \deg(\mathcal{L}) + (n^2 - 1)(g-1).$$

First, we note that the morphism  $\check{h}^{\mathrm{par}}$  is surjective since the usual Hitchin map  $\check{h} : \check{M} \rightarrow \widehat{A}$  is surjective. Furthermore, as  $\check{h}^{\mathrm{par}}$  is equivariant with respect to the scaling  $\mathbb{G}_m$ -action on the Higgs fields, it suffices to bound from above the dimension of the fiber over  $(x, 0) \in C \times \widehat{A}$  for each point  $x \in C$  by  $\mathbf{d}$ , since upper semicontinuity will force all other fibers to have the same dimension upper-bound.<sup>2</sup> We fix the point  $p$  from now on.

Using the assumption  $\deg(\mathcal{L}) > 2g$ , we may express  $\mathcal{L}$  as  $\mathcal{O}_C(D)$  with  $D = x_0 + x_1 + \cdots + x_t$  an effective reduced divisor containing  $x = x_0$ . In particular, a Higgs bundle  $(\mathcal{E}, \theta : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_C)$  can be viewed as a meromorphic (un-twisted) Higgs bundles with at most simple poles along  $D$ :

$$(\mathcal{E}, \theta), \quad \theta : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_C^1(D).$$

We will control the fiber dimension using an analogous bound for the nilpotent cone for *strongly parabolic* Higgs bundles for the pair  $(C, D)$ .

Recall that given  $D$  as above, a strongly parabolic Higgs bundle consists of an  $\mathrm{SL}_n$  Higgs bundle of degree  $d$ ,

$$(\mathcal{E}, \theta), \quad \theta : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_C^1(D), \quad \mathrm{rk}(\mathcal{E}) = n, \quad \det(\mathcal{E}) \simeq \mathcal{N}, \quad \mathrm{trace}(\theta) = 0,$$

---

<sup>2</sup>Furthermore, we actually show that each fiber has the same dimension. But the dimension bound is enough for our purpose.

along with a flag at each point  $x_i \in D$ ,

$$\mathcal{E}|_{x_i} = F_{x_i,0} \supset F_{x_i,1} \supset F_{x_i,2} \supset \cdots \supset F_{x_i,n} = 0, \quad \dim F_{x_i,j}/F_{x_i,j+1} = 1,$$

such that the Higgs field satisfies  $\theta(F_{x_i,j}) \subset F_{x_i,j+1}$ .<sup>3</sup>

Let  $\widetilde{M}^{\text{spar}}(D)$  denote the moduli space of strongly parabolic Higgs bundles associated with the pair  $(C, D)$ , such that the underlying twisted Higgs bundle is stable. Let

$$A(D) = \bigoplus_{i=2}^n H^0(C, \Omega_{\mathcal{L}}^{\otimes i}(-D)) \subset \widehat{A}$$

denote the linear subspace consisting of sections which vanish along the points of  $D$ . There is a Hitchin map for the strongly parabolic Higgs moduli space

$$\check{h}^{\text{spar}} : \widetilde{M}^{\text{spar}}(D) \rightarrow A(D)$$

which is proper, surjective, and Lagrangian with respect to a natural holomorphic symplectic form [14]. In particular, the dimension of the zero fiber is given by

$$\begin{aligned} \dim \widetilde{M}^{\text{spar}}(D)_0 &= \dim A(D) = \sum_{j=2}^n ((j-1)(\deg(\Omega_L)) + g - 1) \\ &= \frac{n(n-1)}{2} \deg(\mathcal{L}) + (n^2 - 1)(g - 1). \end{aligned}$$

See also [36, Th. 6.9] for a direct proof of the above dimension formula.<sup>4</sup>

We now consider the following diagram relating the two types of parabolic Higgs moduli spaces  $\widetilde{M}^{\text{spar}}(D)$  and  $\widetilde{M}^{\text{par}}$ :

$$(31) \quad \begin{array}{ccccc} \widetilde{M}^{\text{spar}}(D)_0 & \hookrightarrow & \widetilde{M}^{\text{spar}}(D) & & \\ \downarrow q_0 & & \downarrow q & & \\ \widetilde{M}^{\text{par}}_{(x,0)} & \hookrightarrow & \widetilde{M}^{\text{par}}|_{x \times A(D)} & \hookrightarrow & \widetilde{M}^{\text{par}} \\ \downarrow & & \downarrow & & \downarrow \check{h}^{\text{par}} \\ \{(x,0)\} & \hookrightarrow & x \times A(D) & \hookrightarrow & C \times \widehat{A}, \end{array}$$

where all squares are Cartesian.

We first claim that the natural map  $q$  (sending a strong parabolic Higgs bundle to a parabolic Higgs bundle) is surjective. Indeed, given a point  $z \in$

<sup>3</sup>One may compare this with the (non-strong) parabolic condition  $\theta(F_{x,j}) \subset F_{x,j}$  we used earlier in this paper for the global Springer theory.

<sup>4</sup>The formula above has  $g$  less than the dimension formula obtained in [36, Th. 6.9], since we consider the  $\text{SL}_n$  case where we fixed the determinant on  $C$ .

$\widetilde{M}^{\mathrm{par}}$  lying over  $C \times A(D)$ , choosing a point in its preimage  $q^{-1}(z)$  only consists of fixing a flag at each point of  $x_i$  with  $i > 0$ , preserved by the Higgs field; since the characteristic polynomial already has zero roots at all points of  $D$ , the Higgs field is automatically strongly parabolic.

By base change, this implies that  $q_0$  is surjective as well, so we get the desired dimension upper bound

$$\dim \widetilde{M}_{(x,0)}^{\mathrm{par}} \leq \dim \widetilde{M}^{\mathrm{spar}}(D)_0 = \frac{n(n-1)}{2} \deg(\mathcal{L}) + (n^2 - 1)(g-1),$$

which completes the proof.  $\square$

*Remark 4.2.* Combining [Sections 3](#) and [4](#) proves that [Theorem 2.6](#) holds for a line bundle  $\mathcal{L}$  of sufficiently large degree. Then the argument of [Section 2.3](#) implies that it actually holds for any effective  $\mathcal{L}$ .

## References

- [1] A. A. BEĬLINSON, J. BERNSTEIN, and P. DELIGNE, Faisceaux pervers, in *Analysis and Topology on Singular Spaces, I* (Luminy, 1981), *Astérisque* **100**, Soc. Math. France, Paris, 1982, pp. 5–171. [MR 0751966](#). [Zbl 0536.14011](#). Available at [http://numdam.org/item/AST\\_1982\\_100\\_1\\_0/](http://numdam.org/item/AST_1982_100_1_0/).
- [2] P.-H. CHAUDOUARD and G. LAUMON, Un théorème du support pour la fibration de Hitchin, *Ann. Inst. Fourier (Grenoble)* **66** no. 2 (2016), 711–727. [MR 3477888](#). [Zbl 1375.14069](#). <https://doi.org/10.5802/aif.3023>.
- [3] W.-Y. CHUANG, D.-E. DIACONESCU, and G. PAN, BPS states and the  $P = W$  conjecture, in *Moduli Spaces, London Math. Soc. Lecture Note Ser.* **411**, Cambridge Univ. Press, Cambridge, 2014, pp. 132–150. [MR 3221294](#). [Zbl 1320.14057](#). <https://doi.org/10.1017/CBO9781107279544.003>.
- [4] Z. DANCSO, M. MCBREEN, and V. SHENDE, Deletion-contraction triangles for Hausel–Proudfoot varieties, *J. Eur. Math. Soc.* **26** no. 7 (2024), 2565–2653. [MR 4756571](#). [Zbl 7875318](#). <https://doi.org/10.4171/JEMS/1369>.
- [5] B. DAVISON, Nonabelian Hodge theory for stacks and a stacky  $P = W$  conjecture, *Adv. Math.* **415** (2023), Paper No. 108889, 61. [MR 4543450](#). [Zbl 1516.14026](#). <https://doi.org/10.1016/j.aim.2023.108889>.
- [6] M. A. DE CATALDO, Perverse sheaves and the topology of algebraic varieties, in *Geometry of Moduli Spaces and Representation Theory, IAS/Park City Math. Ser.* **24**, Amer. Math. Soc., Providence, RI, 2017, pp. 1–58. [MR 3752459](#). [Zbl 1439.14060](#).
- [7] M. A. DE CATALDO, A support theorem for the Hitchin fibration: the case of  $\mathrm{SL}_n$ , *Compos. Math.* **153** no. 6 (2017), 1316–1347. [MR 3705257](#). [Zbl 1453.14029](#). <https://doi.org/10.1112/S0010437X17007096>.
- [8] M. A. DE CATALDO, T. HAUSEL, and L. MIGLIORINI, Topology of Hitchin systems and Hodge theory of character varieties: the case  $A_1$ , *Ann. of Math.* (2) **175** no. 3 (2012), 1329–1407. [MR 2912707](#). [Zbl 1375.14047](#). <https://doi.org/10.4007/annals.2012.175.3.7>.

- [9] M. A. DE CATALDO, J. HEINLOTH, and L. MIGLIORINI, A support theorem for the Hitchin fibration: the case of  $GL_n$  and  $K_C$ , *J. Reine Angew. Math.* **780** (2021), 41–77. [MR 4333982](#). [Zbl 1476.14068](#). <https://doi.org/10.1515/crelle-2021-0045>.
- [10] M. A. DE CATALDO and D. MAULIK, The perverse filtration for the Hitchin fibration is locally constant, *Pure Appl. Math. Q.* **16** no. 5 (2020), 1441–1464. [MR 4221002](#). [Zbl 1469.14018](#). <https://doi.org/10.4310/PAMQ.2020.v16.n5.a4>.
- [11] M. A. DE CATALDO, D. MAULIK, and J. SHEN, Hitchin fibrations, abelian surfaces, and the  $P = W$  conjecture, *J. Amer. Math. Soc.* **35** no. 3 (2022), 911–953. [MR 4433080](#). [Zbl 1523.14061](#). <https://doi.org/10.1090/jams/989>.
- [12] M. A. DE CATALDO, D. MAULIK, and J. SHEN, On the  $P = W$  conjecture for  $SL_n$ , *Selecta Math. (N.S.)* **28** no. 5 (2022), Paper No. 90, 21. [MR 4496523](#). [Zbl 1528.14038](#). <https://doi.org/10.1007/s00029-022-00803-0>.
- [13] M. A. DE CATALDO, D. MAULIK, J. SHEN, and S. ZHANG, Cohomology of the moduli of Higgs bundles via positive characteristic, 2023, published online first 16 December 2023, 23 pp. [https://doi.org/10.4171/JEMS/1393](#).
- [14] G. FALTINGS, Stable  $G$ -bundles and projective connections, *J. Algebraic Geom.* **2** no. 3 (1993), 507–568. [MR 1211997](#). [Zbl 0790.14019](#).
- [15] C. FELISETTI and M. MAURI,  $P = W$  conjectures for character varieties with symplectic resolution, *J. Éc. Polytech. Math.* **9** (2022), 853–905. [MR 4436035](#). [Zbl 1499.14087](#). <https://doi.org/10.5802/jep.196>.
- [16] M. GROECHENIG, D. WYSS, and P. ZIEGLER, Mirror symmetry for moduli spaces of Higgs bundles via  $p$ -adic integration, *Invent. Math.* **221** no. 2 (2020), 505–596. [MR 4121158](#). [Zbl 1451.14123](#). <https://doi.org/10.1007/s00222-020-00957-8>.
- [17] A. HARDER, Z. LI, J. SHEN, and Q. YIN,  $P = W$  for Lagrangian fibrations and degenerations of hyper-Kähler manifolds, *Forum Math. Sigma* **9** (2021), Paper No. e50, 6. [MR 4277887](#). [Zbl 1467.14101](#). <https://doi.org/10.1017/fms.2021.31>.
- [18] T. HAUSEL and M. THADDEUS, Mirror symmetry, Langlands duality, and the Hitchin system, *Invent. Math.* **153** no. 1 (2003), 197–229. [MR 1990670](#). [Zbl 1043.14011](#). <https://doi.org/10.1007/s00222-003-0286-7>.
- [19] T. HAUSEL and M. THADDEUS, Relations in the cohomology ring of the moduli space of rank 2 Higgs bundles, *J. Amer. Math. Soc.* **16** no. 2 (2003), 303–329. [MR 1949162](#). [Zbl 1015.14018](#). <https://doi.org/10.1090/S0894-0347-02-00417-4>.
- [20] N. J. HITCHIN, The self-duality equations on a Riemann surface, *Proc. London Math. Soc.* (3) **55** no. 1 (1987), 59–126. [MR 0887284](#). [Zbl 0634.53045](#). <https://doi.org/10.1112/plms/s3-55.1.59>.
- [21] N. HITCHIN, Stable bundles and integrable systems, *Duke Math. J.* **54** no. 1 (1987), 91–114. [MR 0885778](#). [Zbl 0627.14024](#). <https://doi.org/10.1215/S0012-7094-87-05408-1>.
- [22] M. KASHIWARA and P. SCHAPIRA, *Sheaves on Manifolds*, Grundlehren Math. Wissen. **292**, Springer-Verlag, Berlin, 1990, with a chapter in French by Christian Houzel. [MR 1074006](#). [Zbl 0709.18001](#). <https://doi.org/10.1007/978-3-662-02661-8>.

- [23] E. MARKMAN, Generators of the cohomology ring of moduli spaces of sheaves on symplectic surfaces, *J. Reine Angew. Math.* **544** (2002), 61–82. [MR 1887889](#). [Zbl 0988.14019](#). <https://doi.org/10.1515/crll.2002.028>.
- [24] D. MAULIK and J. SHEN, Endoscopic decompositions and the Hausel-Thaddeus conjecture, *Forum Math. Pi* **9** (2021), Paper No. e8, 49 pp. [MR 4304301](#). [Zbl 1478.14054](#). <https://doi.org/10.1017/fmp.2021.7>.
- [25] D. MAULIK and J. SHEN, On the intersection cohomology of the moduli of  $\mathrm{SL}_n$ -Higgs bundles on a curve, *J. Topol.* **15** no. 3 (2022), 1034–1057. [MR 4442682](#). [Zbl 1521.14025](#). <https://doi.org/10.1112/topo.12250>.
- [26] D. MAULIK and J. SHEN, Cohomological  $\chi$ -independence for moduli of one-dimensional sheaves and moduli of Higgs bundles, *Geom. Topol.* **27** no. 4 (2023), 1539–1586. [MR 4602420](#). [Zbl 1514.14038](#). <https://doi.org/10.2140/gt.2023.27.1539>.
- [27] D. MAULIK and Y. TODA, Gopakumar-Vafa invariants via vanishing cycles, *Invent. Math.* **213** no. 3 (2018), 1017–1097. [MR 3842061](#). [Zbl 1400.14141](#). <https://doi.org/10.1007/s00222-018-0800-6>.
- [28] M. MAURI and L. MIGLIORINI, Hodge-to-singular correspondence for reduced curve, *J. Eur. Math. Soc.* (2024), published online first 23 July 2024. <https://doi.org/10.4171/JEMS/1508>.
- [29] A. MELLIT, Cell decompositions of character varieties, 2019. [arXiv 1905.10685](#).
- [30] B. C. NGÔ, Le lemme fondamental pour les algèbres de Lie, *Publ. Math. Inst. Hautes Études Sci.* no. 111 (2010), 1–169. [MR 2653248](#). [Zbl 1200.22011](#). <https://doi.org/10.1007/s10240-010-0026-7>.
- [31] A. OBLOMKOV and Z. YUN, Geometric representations of graded and rational Cherednik algebras, *Adv. Math.* **292** (2016), 601–706. [MR 3464031](#). [Zbl 1403.20007](#). <https://doi.org/10.1016/j.aim.2016.01.015>.
- [32] A. OBLOMKOV and Z. YUN, The cohomology ring of certain compactified Jacobians, 2017. [arXiv 1710.05391](#).
- [33] V. SHENDE, The weights of the tautological classes of character varieties, *Int. Math. Res. Not. IMRN* no. 22 (2017), 6832–6840. [MR 3737322](#). [Zbl 1405.14037](#). <https://doi.org/10.1093/imrn/rnv363>.
- [34] C. T. SIMPSON, Higgs bundles and local systems, *Inst. Hautes Études Sci. Publ. Math.* no. 75 (1992), 5–95. [MR 1179076](#). [Zbl 0814.32003](#). Available at [http://www.numdam.org/item?id=PMIHES\\_1992\\_75\\_5\\_0](http://www.numdam.org/item?id=PMIHES_1992_75_5_0).
- [35] C. T. SIMPSON, Moduli of representations of the fundamental group of a smooth projective variety. II, *Inst. Hautes Études Sci. Publ. Math.* no. 80 (1995), 5–79. [MR 1320603](#). [Zbl 0891.14006](#). Available at [http://www.numdam.org/item?id=PMIHES\\_1994\\_80\\_5\\_0](http://www.numdam.org/item?id=PMIHES_1994_80_5_0).
- [36] X. SU, B. WANG, and X. WEN, Parabolic Hitchin maps and their generic fibers, *Math. Z.* **301** no. 1 (2022), 343–372. [MR 4405653](#). [Zbl 1496.14016](#). <https://doi.org/10.1007/s00209-021-02896-3>.
- [37] S. SZABÓ, Perversity equals weight for Painlevé spaces, *Adv. Math.* **383** (2021), Paper No. 107667, 45 pp. [MR 4232540](#). [Zbl 1470.14068](#). <https://doi.org/10.1016/j.aim.2021.107667>.

- [38] Z. YUN, Towards a global Springer theory III: Endoscopy and Langlands duality, 2009. [arXiv 0904.3372](https://arxiv.org/abs/0904.3372).
- [39] Z. YUN, Global Springer theory, *Adv. Math.* **228** no. 1 (2011), 266–328. [MR 2822234](https://doi.org/10.1016/j.aim.2011.05.012). [Zbl 1230.14048](https://doi.org/10.1016/j.aim.2011.05.012). <https://doi.org/10.1016/j.aim.2011.05.012>.
- [40] Z. YUN, Langlands duality and global Springer theory, *Compos. Math.* **148** no. 3 (2012), 835–867. [MR 2925401](https://doi.org/10.1112/S0010437X11007433). [Zbl 1279.14018](https://doi.org/10.1112/S0010437X11007433). <https://doi.org/10.1112/S0010437X11007433>.
- [41] Z. ZHANG, The  $P = W$  identity for cluster varieties, *Math. Res. Lett.* **28** no. 3 (2021), 925–944. [MR 4270279](https://doi.org/10.4310/MRL.2021.v28.n3.a13). [Zbl 1462.13024](https://doi.org/10.4310/MRL.2021.v28.n3.a13). <https://doi.org/10.4310/MRL.2021.v28.n3.a13>.

(Received: September 12, 2022)

(Revised: January 4, 2024)

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA

*E-mail:* [maulik@mit.edu](mailto:maulik@mit.edu)

YALE UNIVERSITY, NEW HAVEN, CT

*E-mail:* [junliang.shen@yale.edu](mailto:junliang.shen@yale.edu)