

POLYNOMIAL SUPERPOTENTIAL FOR GRASSMANNIAN $\mathrm{Gr}(k, n)$ FROM A LIMIT OF VERTEX FUNCTION

ANDREY SMIRNOV* AND ALEXANDER VARCHENKO[◦]

Department of Mathematics, University of North Carolina at Chapel Hill
Chapel Hill, NC 27599-3250, USA

Key words: Superpotentials; Vertex Functions; J -functions; Landau-Ginzburg model.

2020 Mathematics Subject Classification: 14G33 (11D79, 32G34, 33C05, 33E30)

ABSTRACT. In this note we discuss an integral representation for the vertex function of the cotangent bundle over the Grassmannian, $X = T^* \mathrm{Gr}(k, n)$. This integral representation can be used to compute the $\hbar \rightarrow \infty$ limit of the vertex function, where \hbar denotes the equivariant parameter of a torus acting on X by dilating the cotangent fibers. We show that in this limit the integral turns into the standard mirror integral representation for the A -series of the Grassmannian $\mathrm{Gr}(k, n)$ with the Laurent polynomial Landau-Ginzburg superpotential of Eguchi, Hori and Xiong. We also observe some Dwork type congruences for the coefficients of the A -series.

1. INTRODUCTION

1.1. The vertex functions have been introduced in [Oko17] as generating functions counting rational quasimaps to Nakajima varieties. In this respect, a vertex function is a “quasimap” analog of Givental’s J -function in quantum cohomology. In this paper we consider the the cohomological vertex function for the cotangent bundle over the Grassmannian $X = T^* \mathrm{Gr}(k, n)$. By definition, this function is a power series in the quantum parameter z with coefficients in the equivariant cohomology:

$$\mathrm{Vertex}(z) \in H_T^\bullet(X)[[z]]$$

where T is a torus acting on X , see Section 2. Let $V(z)$ denote the coefficient of the fundamental class in the vertex function

$$V(z) := \langle \mathrm{Vertex}(z), [X] \rangle$$

where $\langle -, - \rangle$ stands for the standard pairing in the equivariant cohomology. The power series $V(z)$ is the analog of the so-called A -series in quantum cohomology. The coefficients of $V(z)$ depend non-trivially on the equivariant parameter \hbar , which corresponds to the torus acting on X by dilating the cotangent fibers. In this note we describe the following result (Theorem 4.1):

Theorem 1.1. *In the non-equivariant specialization one has the following limit:*

$$(1.1) \quad \lim_{\hbar \rightarrow \infty} V(z/\hbar^n) = \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint e^{\frac{1}{\epsilon} S(x, z)} \bigwedge_{i,j} \frac{dx_{i,j}}{x_{i,j}}$$

*E-mail: asmirnov@email.unc.edu

[◦]E-mail: anv@email.unc.edu

where $S(x, z)$ denotes a Laurent polynomial in $k(n - k)$ variables $x = (x_{i,j})$ given by (4.1).

The integral in the right-hand side of (1.1) denotes the constant term of the integrand, see Section 4.2 for the definition. The Laurent polynomial $S(x, z)$ appearing in the limit above is the well-known version of the *superpotential* for $\mathrm{Gr}(k, n)$. It first appeared in the paper [EHX97] by Eguchi-Hori-Xiong and has since been revisited and generalized by many authors in the extensive literature on quantum cohomology of Grassmannians. A partial listing of references can be found in [BCKS98, Cas20, GGI16, KoSt10, LaTe17, MaRi20]. Therefore, the expression on the right-hand side of equation (1.1) corresponds to the widely used integral representation for the A -series of the Grassmannian $\mathrm{Gr}(k, n)$. A closed combinatorial formula for the coefficients of this series is also known, it was first conjectured in [BCKS98] and later proved in [MaRi20], see Corollary 4.8 in [MaRi20] (our z is their q).

Informally speaking formula (1.1) means that in the limit $\hbar \rightarrow \infty$ the cotangent directions of X do not contribute to the quasimap partition function and the vertex function of $T^* \mathrm{Gr}(k, n)$ degenerates to the J -function of $\mathrm{Gr}(k, n)$. The idea that $\hbar \rightarrow \infty$ bridges the vertex functions with the J -functions is not new, see [TV22] or Section 5 of [KPSZ21]. However, the derivation of the Laurent superpotential of [EHX97] from the limit of $V(z)$ has not been documented well. The goal of this letter is to fill up this gap in the existing literature. The main technical tool which allows us to compute the limit is the integral representation for $V(z)$ obtained previously by the authors in Theorem 3.2 of [SmV23].

1.2. As an illustration, let us consider the statement of Theorem 1.1 in the simplest case.

Example. Let $X = T^*\mathbb{P}(\mathbb{C}^2)$. There is a two-dimensional torus $\mathbf{A} = (\mathbb{C}^\times)^2$ with equivariant parameters u_1, u_2 naturally acting on \mathbb{C}^2 by dilating the coordinate subspaces. There is also a one-dimensional torus \mathbb{C}_\hbar^\times acting on X by dilation of the cotangent fibers, we denote the corresponding character by \hbar . Finally, there is a one-dimensional torus $\mathbb{C}_\epsilon^\times$ with equivariant parameter ϵ which acts on the source of the quasimaps $\mathbb{P}^1 \dashrightarrow X$. By definition, the vertex function is a power series with coefficients in equivariant cohomology [Oko17]:

$$\mathrm{Vertex}(z) \in H_{\mathbf{A} \times \mathbb{C}_\hbar^\times \times \mathbb{C}_\epsilon^\times}^\bullet(X)_{loc}[[z]]$$

where the subscript loc denotes localization with respect to $\mathbb{C}_\epsilon^\times$. In the basis of $H_{\mathbf{A} \times \mathbb{C}_\hbar^\times \times \mathbb{C}_\epsilon^\times}^\bullet(X)_{loc}$ given by the classes of the torus fixed points $[p_1], [p_2] \in X^\mathbf{A}$ (which correspond to the coordinate lines in \mathbb{C}^2) we have the following closed formulas in terms of the Gauss hypergeometric functions:

$$(1.2) \quad \begin{aligned} \langle \mathrm{Vertex}(z), [p_1] \rangle &= {}_2F_1\left(\frac{\hbar}{\epsilon}, \frac{u_2 - u_1 + \hbar}{\epsilon}; \frac{u_2 - u_1 + \epsilon}{\epsilon}; z\right), \\ \langle \mathrm{Vertex}(z), [p_2] \rangle &= {}_2F_1\left(\frac{\hbar}{\epsilon}, \frac{u_1 - u_2 + \hbar}{\epsilon}; \frac{u_1 - u_2 + \epsilon}{\epsilon}; z\right). \end{aligned}$$

In the *non-equivariant* limit $u_1 = u_2 = 0$, corresponding to “turning off” the action of the torus \mathbf{A} , the above functions coincide and give the coefficient of the vertex function at the fundamental class $[X]$:

$$(1.3) \quad \langle \mathrm{Vertex}(z), [X] \rangle = {}_2F_1\left(\frac{\hbar}{\epsilon}, \frac{\hbar}{\epsilon}; 1; z\right)$$

We denote this coefficient by $V(z)$. Explicitly we have

$$V(z) = \sum_{d=0}^{\infty} \frac{(\hbar)_d^2}{(d!)^2 \epsilon^{2d}} z^d, \quad \text{where } (\hbar)_d = \hbar(\hbar + \epsilon)(\hbar + 2\epsilon) \dots (\hbar + (d-1)\epsilon).$$

Since $\lim_{\hbar \rightarrow \infty} (\hbar)_d / \hbar^d = 1$, we obtain:

$$\lim_{\hbar \rightarrow \infty} V(z/\hbar^2) = \sum_{d=0}^{\infty} \frac{z^d}{(d!)^2 \epsilon^{2d}}.$$

Let $S(x, z) = x + z/x$, then

$$\oint \frac{dx}{x} S(x, z)^d := [S(x, z)^d]_0 = \begin{cases} \frac{(d)!}{(d/2)!(d/2)!} z^d, & d \text{ is even} \\ 0, & d \text{ is odd} \end{cases}$$

where $[S(x, z)^d]_0$ denotes the constant term in x of the Laurent polynomial $S(x, z)^d$. Combining all these together, we can write

$$\lim_{\hbar \rightarrow \infty} V(z/\hbar^2) = \sum_{d=0}^{\infty} \frac{1}{d!} \oint \frac{dx}{x} S(x, z)^d = \oint \frac{dx}{x} e^{\frac{S(x, z)}{\epsilon}}.$$

This formula is the statement of Theorem 1.1 in this case. A more straightforward way to compute this limit is to note that the hypergeometric function (1.3) has the integral representation:

$$(1.4) \quad V(z) = \oint_{|x|=\epsilon} \frac{dx}{x} \left(1 - x\right)^{-\frac{\hbar}{\epsilon}} \left(1 - \frac{z}{x}\right)^{-\frac{\hbar}{\epsilon}}$$

where ϵ is any positive real number such that $|z| < \epsilon < 1$. We note that the change of variables $z \rightarrow z/\hbar^2$, $x \rightarrow x/\hbar$ together with change of the contour $\epsilon \rightarrow \epsilon/\hbar$ does not affect this condition for large $|\hbar|$. Thus, for large $|\hbar|$ we have:

$$V(z/\hbar^2) = \oint_{|x|=\epsilon} \frac{dx}{x} \left(1 - \frac{x}{\hbar}\right)^{-\frac{\hbar}{\epsilon}} \left(1 - \frac{z}{x\hbar}\right)^{-\frac{\hbar}{\epsilon}}$$

which allows one to compute the limit using elementary tools:

$$\lim_{\hbar \rightarrow \infty} \left(1 - \frac{x}{\hbar}\right)^{-\frac{\hbar}{\epsilon}} \left(1 - \frac{z}{x\hbar}\right)^{-\frac{\hbar}{\epsilon}} = e^{\frac{S(x, z)}{\epsilon}}$$

1.3. Exposition of the material. In Section 2 we recall a combinatorial formula for the vertex functions generalizing (1.2) to the case of $X = T^* \text{Gr}(k, n)$. In Section 3 we describe the analog of the integral representation (1.4) for this case. In Section 4 we use this integral representation to compute the $\hbar \rightarrow \infty$ limit of $V(z)$ similarly to the example above.

In our previous paper [SmV23], we show that certain truncations of $V(z)$ with parameters specialized to \mathbb{Q}_p satisfy some Dwork type congruence relations. In Section 5 we show that a similar structure exists in the limit $\hbar \rightarrow \infty$.

Acknowledgements. We thank Thomas Lam for very useful comments. Work of A. Smirnov is partially supported by NSF grant DMS - 2054527 and by the RSF under grant 19-11-00062. Work of A. Varchenko is partially supported by NSF grant DMS - 1954266.

2. THE VERTEX FUNCTION OF $T^* \mathrm{Gr}(k, n)$

2.1. For $X = T^* \mathrm{Gr}(k, n)$ we consider the following explicit power series:

$$(2.1) \quad \langle \mathrm{Vertex}(z), [1, \dots, k] \rangle := \sum_{d=0}^{\infty} c_d(u_1, \dots, u_n, \hbar) z^d$$

with the coefficients $c_d(u_1, \dots, u_n, \hbar) \in \mathbb{Q}(u_1, \dots, u_n, \hbar, \epsilon)$ given by:

$$(2.2) \quad c_d(u_1, \dots, u_n, \hbar) = \sum_{\substack{d_1, \dots, d_k: \\ d_1 + \dots + d_k = d}} \left(\prod_{i,j=1}^k \frac{(\epsilon - u_i + u_j)_{d_i - d_j}}{(\hbar - u_i + u_j)_{d_i - d_j}} \right) \left(\prod_{j=1}^n \prod_{i=1}^k \frac{(\hbar + u_j - u_i)_{d_i}}{(\epsilon + u_j - u_i)_{d_i}} \right),$$

where $(x)_d$ denotes the Pochhammer symbol with step ϵ :

$$(x)_d = \begin{cases} x(x + \epsilon) \dots (x + (d-1)\epsilon), & d > 0 \\ 1, & d = 0 \\ \frac{1}{(x - \epsilon)(x - 2\epsilon) \dots (x + d\epsilon)}, & d < 0 \end{cases}$$

The degree d coefficient of this series counts (equivariantly) the number of degree d rational curves in X . More precisely, it is given by the equivariant integral

$$(2.3) \quad c_d(u_1, \dots, u_n, \hbar) = \int_{[\mathrm{QM}_d(X, \infty)]^{\mathrm{vir}}} \omega^{\mathrm{vir}}$$

over the virtual fundamental class on moduli space $\mathrm{QM}_d(X, \infty)$ of quasimaps from \mathbb{P}^1 to X , which send $\infty \in \mathbb{P}^1$ to a prescribed torus fixed point $[1, \dots, k] \in X$, see Section 7.2 of [Oko17] for definitions. Using the equivariant localization, the integral (2.3) reduces to the sum over the torus fixed points on $\mathrm{QM}_d(X, \infty)$ which gives the sum (2.2). We refer to Section 4.5 of [PSZ16] where this computation is done in some details.

The parameters $u_1, \dots, u_n, \hbar, \epsilon$ are the equivariant parameters of the torus $T = (\mathbb{C}^\times)^n \times \mathbb{C}_\hbar^\times \times \mathbb{C}_\epsilon^\times$ acting on the moduli space $\mathrm{QM}_d(X, \infty)$ in the following way:

- $(\mathbb{C}^\times)^n$ acts on \mathbb{C}^n in a natural way, scaling the coordinates with weights u_1, \dots, u_n .
- The set of torus fixed points $X^{(\mathbb{C}^\times)^n}$ corresponds to k -subspaces in \mathbb{C}^n spanned by any set of k coordinate lines. The fixed point $[1, \dots, k] \in X^{(\mathbb{C}^\times)^n}$ corresponds to the k -subspace spanned by the first k coordinate lines.
- \mathbb{C}_\hbar^\times acts on X by scaling the cotangent fibers with weight \hbar .
- $\mathbb{C}_\epsilon^\times$ acts on the source of the quasimaps $C \cong \mathbb{P}^1$ fixing the points $0, \infty \in \mathbb{P}^1$. The parameter ϵ denotes the corresponding weight of the tangent space $T_0 C$.

The full vertex function is a power series with coefficients in equivariant cohomology:

$$\mathrm{Vertex}(z) \in H_T^\bullet(X)_{\mathrm{loc}}[[z]]$$

where loc denotes the equivariant localization with respect to torus $\mathbb{C}_\epsilon^\times$. Using the equivariant localization, we can expand $\mathrm{Vertex}(z)$ in the basis of $H_{T \times \mathbb{C}_\epsilon^\times}^\bullet(X)_{\mathrm{loc}}$ given by the classes of

torus fixed points. The power series (2.1) gives the coefficient $\text{Vertex}(z)$ at the “first” torus fixed point $[1, \dots, k]$. Other coefficients have the same structure and can be obtained from (2.1) by permutations of parameters u_i .

2.2. In this paper we consider the specialization of the equivariant parameters:

$$(2.4) \quad u_1 = 0, \dots, u_n = 0$$

which corresponds to non-equivariant limit when the action of the torus $(\mathbb{C}^\times)^n$ is “turned off”. The coefficients of $\text{Vertex}(z)$ at the torus fixed points all reduce to the same function (simply because without $(\mathbb{C}^\times)^n$ -action these points are indistinguishable) which corresponds to the coefficient of the vertex function at the fundamental class:

$$(2.5) \quad V(z) := \left\langle \text{Vertex}(z), [X] \right\rangle \Big|_{u_1=0, \dots, u_n=0}$$

Thus, $V(z)$ can be obtained by specializing the coefficients of the power series (2.1) at (2.4). We note that this specialization is non-trivial: already in the case of $T^* \text{Gr}(2, 4)$ the terms in the sum (2.2) have poles at $u_i = u_j$. The total sum (2.2) is, however, non-singular since the vertex function is an integral equivariant cohomology class (we recall that only $\mathbb{C}_\epsilon^\times$ -localization is required to define it).

3. INTEGRAL REPRESENTATION OF $V(z)$

3.1. In this section we describe an integral representation for the function (2.5)

$$V(z) = \int_\gamma \Phi(x, z) dx$$

which has its origin in 3D-mirror symmetry, we refer to Section 3 of [SmV23] for more details.

3.2. Assume that $n \geq 2k$. Let v_i , $i = 1, \dots, n-1$ be integers defined by:

$$v_i = \begin{cases} i, & i < k, \\ k, & k \leq i \leq n-k, \\ n-i, & n-k < i, \end{cases}$$

We denote by $\omega = \hbar/\epsilon$ and define the *superpotential* function:

$$(3.1) \quad \Phi(x, z) = \left(\prod_{i=1}^{n-1} \prod_{j=1}^{v_i} x_{i,j} \right)^{-1+\omega} \left(\prod_{m=1}^{v_m} \prod_{1 \leq i < j \leq v_m} (x_{m,j} - x_{m,i}) \right)^{2\omega} \times \left(\prod_{i=1}^{n-2} \prod_{a=1}^{v_i} \prod_{b=1}^{v_{i+1}} (x_{i,a} - x_{i+1,b}) \right)^{-\omega} \left(\prod_{i=1}^k (z_1 - x_{k,i})(z_2 - x_{n-k,i}) \right)^{-\omega}.$$

We note that this function is an example of the *master functions* in the theory of integral representations of the trigonometric Knizhnik-Zamolodchikov equations. In particular, (3.1) corresponds to the KZ equation associated with the weight subspace of weight $[1, \dots, 1]$ in the tensor product the k -th and $(n-k)$ -th fundamental representations of \mathfrak{gl}_n , see [SV91, MV02].

3.3. The dimension vector \mathbf{v}_i and the variables $x_{i,j}$ have a convenient combinatorial visualization. Let us consider a $k \times (n - k)$ rectangle rotated counterclockwise by 45° , see Fig. 1. Note that in this picture the number of boxes in i -th vertical column is exactly v_i . In this way, we may assign the variables $x_{i,j}$ to the boxes in this picture. We will order them as in Fig. 1. Note that the total number of variables $x_{i,j}$ equals to $\dim \mathrm{Gr}(k, n) = k(n - k)$. To a




FIGURE 1. Set of variables $x_{i,j}$ for $k = 4$ and $n = 8$.

box (i, j) in the Fig. 1 we assign a weight

$$(3.2) \quad m_{i,j} = (|i - k| + 2j - 1) \in \mathbb{N}$$

This function ranges from $m_{k,1} = 1$ to $m_{n-k,k} = n - 1$. The definition of $m_{i,j}$ is clear from the Fig. 2. We have a partial ordering on the boxes (i, j) corresponding to:

$$(3.3) \quad m_{k,1} < m_{k-1,1} = m_{k+1,1} < \cdots < m_{n-k,k}$$

For a small real number $0 < \varepsilon \ll 1$ let us define the torus by the following equations:


$$(3.4) \quad \gamma_{k,n} \subset \mathbb{C}^{k(n-k)}, \quad |x_{i,j}| = m_{i,j} \varepsilon$$

where i, j run through all possible values.

Proposition 3.1 ([SmV23]). *Assume that $|z_1| < \varepsilon$ and $(n - 1)\varepsilon < |z_2|$, then the superpotential (3.1) has a single-valued branch on the torus $\gamma_{k,n}$, which is distinguished in the proof and which will be used in the paper.*

Proof. Let us denote

$$(3.5) \quad L(x_{i,a}, x_{j,b}) = \begin{cases} (1 - x_{i,a}/x_{j,b})^{-\omega}, & m_{i,a} < m_{j,b}, \quad i \neq j \\ (x_{j,b}/x_{i,a} - 1)^{-\omega}, & m_{i,a} > m_{j,b}, \quad i \neq j. \end{cases}$$

FIGURE 2. The values of the weight function $m_{i,j}$

Each of these ratios $x_{i,a}/x_{j,b}$, $x_{j,b}/x_{i,a}$ restricted to $\gamma_{k,n}$ has absolute value less than 1. We replace $(1 - x_{i,a}/x_{j,b})^{-\omega}$ on $\gamma_{k,n}$ with $\sum_{m=0}^{\infty} \binom{-\omega}{m} (-x_{i,a}/x_{j,b})^m$ and replace $(x_{j,b}/x_{i,a} - 1)^{-\omega}$ with $e^{-\pi\sqrt{-1}\omega} \sum_{m=0}^{\infty} \binom{-\omega}{m} (-x_{j,b}/x_{i,a})^m$.

Next, we denote $L(z_1, x_{k,a}) = (1 - z_1/x_{k,a})^{-\omega}$ and $L(z_2, x_{k,a}) = (1 - x_{n-k,a}/z_2)^{-\omega}$. On $\gamma_{k,n}$ we have $|x_{k,i}| \geq \epsilon$, and $|x_{n-k,i}| \leq |x_{n-k,k}| = n\epsilon$, therefore $|z_1/x_{k,i}| < 1$ and $|x_{n-k,i}/z_2| < 1$. We replace on $\gamma_{k,n}$ the factor $(1 - z_1/x_{k,a})^{-\omega}$ with $\sum_{m=0}^{\infty} \binom{-\omega}{m} (-z_1/x_{k,a})^m$ and the factor $(1 - x_{n-k,a}/z_2)^{-\omega}$ with $\sum_{m=0}^{\infty} \binom{-\omega}{m} (-x_{n-k,a}/z_2)^m$.

Finally, we denote $\Delta(x_{m,i}, x_{m,j}) = (1 - x_{m,i}/x_{m,j})^{2\omega}$ for $1 \leq i < j \leq v_m$. On $\gamma_{k,n}$ we have $|x_{m,i}/x_{m,j}| < 1$. We replace on $\gamma_{k,n}$ the factor $\Delta(x_{m,i}, x_{m,j})$ with $\sum_{m=0}^{\infty} \binom{2\omega}{m} (-x_{m,i}/x_{m,j})^m$.

In these notations we have:

(3.6)

$$\Phi(x, z) = \frac{\left(\prod_{i=1}^{n-1} \prod_{a < b} \Delta(x_{i,a}, x_{i,b}) \right) \left(\prod_{i=1}^{n-2} \prod_{a=1}^{v_i} \prod_{b=1}^{v_{i+1}} L(x_{i,a}, x_{i+1,b}) \right) \left(\prod_{i=1}^k L(z_1, x_{k,i}) L(z_2, x_{n-k,i}) \right)}{\prod_{i=1}^{n-1} \prod_{j=1}^{v_i} x_{i,j}},$$

and for each factor a single-valued branch is chosen by replacing that factor with the corresponding power series. The product of those power series distinguishes a single-valued branch of $\Phi(x, z)$ on $\gamma_{k,n}$. \square

Example. For $X = T^* \mathrm{Gr}(2, 4)$ we have

$$\begin{aligned}\Phi(x, z) &= (x_{11}x_{21}x_{22}x_{31})^{-1} \\ &\times (1 - x_{21}/x_{22})^{2\omega} ((x_{21}/x_{11} - 1)(1 - x_{21}/x_{31})(z_1/x_{21} - 1)(1 - x_{21}/z_2))^{-\omega} \\ &\times ((1 - x_{11}/x_{22})(x_{31}/x_{22} - 1)(z_1/x_{22} - 1)(1 - x_{22}/z_2))^{-\omega}.\end{aligned}$$

From the previous proposition, the integral of $\Phi(x, z)$ over $\gamma_{k,n}$ is an analytic function of $z = z_1/z_2$ in the disc $|z| < \epsilon$.

Theorem 3.2 ([SmV23]). *The function (2.5) has the following integral representation*

$$(3.7) \quad V(z) = \frac{\alpha}{(2\pi\sqrt{-1})^{k(n-k)}} \oint_{\gamma_{k,n}} \Phi(x, z) \bigwedge_{i,j} dx_{i,j}$$

where $\Phi(x, z)$ is the branch of superpotential function (3.1) on the torus $\gamma_{k,n}$ chosen in Proposition 3.1, and $\alpha = e^{\pi\sqrt{-1}N\omega}$ is a normalization constant where N is the number of factors in (3.6) having the form $(x_{j,b}/x_{i,a} - 1)^{-\omega}$.

Definition 3.3. Let $\gamma'_{k,m}$ be another contour defined by $|x_{i,j}| = R_{i,j}$ for $R_{i,j} \in \mathbb{R}$ such that $|z_1| < R_{1,1}, R_{n-k,k} < |z_2|$ and the conditions

$$(3.8) \quad m_{i,j} < m_{a,b} \implies R_{i,j} < R_{a,b}$$

are satisfied for all pairs of indices (i, j) and (a, b) . Then, we say that $\gamma'_{k,m}$ is homologous to $\gamma_{k,m}$ and write $\gamma'_{k,m} \sim \gamma_{k,m}$.

Note that (3.7) remains invariant if we replace $\gamma_{k,m}$ by a homologous $\gamma'_{k,m}$. This is simply because the evaluation of the integral over $\gamma'_{k,n}$ is again reduces to computing the residues at $x_{i,j} = 0$, and the residues are computed in the same order as for $\gamma_{k,n}$. This implies that the result remains the same.

3.4. Relation to 3D-mirror symmetry. Let us explain the origin of the superpotential function (3.1). The factors of (3.1) correspond to the edges of the quiver which describes the 3D-mirror variety $X^!$. For $X = T^* \mathrm{Gr}(k, n)$ the quiver of $X^!$ is given in Sections 3.2-3.3 of [SmV23], the correspondence between the factors of (3.1) and the edges of this quiver is also explained there.

For general Nakajima quiver varieties, the superpotential function (3.1) is constructed by the same procedure if the quiver for the 3D-mirror variety $X^!$ is known. For the Nakajima quiver varieties of type A , which include cotangent bundles over partial flag varieties as special cases, a conjectural description of the 3D-mirrors was given by physicists. It is explained for instance in [GK13]. We expect that the results of this note and of [SmV23] have straightforward generalizations to those cases.

The 3D-mirror symmetry conjecture is formulated on the level of K-theory rather than cohomology. Recall, that the quantum difference equations [OS22] are the K-theoretic generalizations of quantum differential equations in quantum cohomology. The 3D-mirror symmetry conjecture claims that the quantum difference equations for X and $X^!$ are equivalent. The K-theoretic vertex functions of X and $X^!$ provide two different bases of solutions to the this common system of q -difference equations. For cotangent bundles over Grassmannians

this conjecture was proved by Dinkins in [Din20] and for full flag varieties in [Din22]. For the hypertoric varieties this result is obtained in [SZ22].

An alternative definition of 3D-mirror symmetry postulates the equality of the elliptic stable envelopes [AO21] of X and $X^!$. This idea was first proposed in [Oko18] and later examined for various cases of X in [RSV19, RSVZ21, RW20, SZ22]. It was shown in [KoSm22, Ko] that the elliptic stable envelope of X determines the corresponding quantum difference equation of X and vice versa. This established an equivalence between the two definitions of 3D-mirror symmetry.

Theorem 1.1 says that the mirror description of the J -function for $\mathrm{Gr}(k, n)$ arises as a double limit of 3D-mirror symmetry. In the first limit one considers the cohomological limit of the K-theoretic vertex functions for $T^* \mathrm{Gr}(k, n)$. In this limit, the 3D-mirror symmetry description of these functions [Din20] degenerates to the integral representation (3.7). In the second limit $\hbar \rightarrow \infty$ we obtain Theorem 1.1.

4. THE LIMIT $\hbar \rightarrow \infty$

4.1. Polynomial superpotential. Let Γ be an oriented graph, with vertices given by boxes inside the $k \times (n-k)$ Young diagram, plus two extra vertices corresponding to z_1 and z_2 , see Fig.3. The edges of the graph are defined as follows: every two adjacent boxes are connected by an edge. Each edge is oriented in the direction of decrease of weight function $m_{i,j}$, which is defined by (3.2). Two additional edges are from $x_{k,1}$ to z_1 and from z_2 to $x_{n-k,k}$, Fig.3. Given an edge e of Γ we denote by $h(e)$ and $t(e)$ the corresponding head and tail. We define the following Laurent polynomial:

$$(4.1) \quad S(x, z) = \sum_{e \in \text{edges}(\Gamma)} \frac{x_{h(e)}}{x_{t(e)}}.$$

Example. For $k = 1$ we obtain:

$$(4.2) \quad S(x, z) = \frac{z_1}{x_{1,1}} + \frac{x_{1,1}}{x_{2,1}} + \frac{x_{2,1}}{x_{3,1}} + \cdots + \frac{x_{n-2,1}}{x_{n-1,1}} + \frac{x_{n-1,1}}{z_2}.$$

Substituting $z_1 = q, z_2 = 1$, and introducing new variables by

$$x_{1,1} = a_1 a_2 \dots a_{n-1}, \quad x_{2,1} = a_1 a_2 \dots a_{n-2}, \quad \dots, \quad x_{n-1,1} = a_1.$$

We arrive at the standard Givental's superpotential of projective space

$$S(x) = a_1 + a_2 + \cdots + a_{n-1} + \frac{q}{a_1 \dots a_{n-1}}.$$

Example. For $X = T^* \mathrm{Gr}(2, 4)$ we obtain

$$(4.3) \quad S(x, z) = \frac{z_1}{x_{2,1}} + \frac{x_{2,1}}{x_{1,1}} + \frac{x_{2,1}}{x_{3,1}} + \frac{x_{3,1}}{x_{2,2}} + \frac{x_{1,1}}{x_{2,2}} + \frac{x_{2,2}}{z_2}.$$

4.2. Exponential integral. For $S(x, z)$ defined by (4.1) we consider the power series:

$$(4.4) \quad \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint e^{\frac{1}{\epsilon} S(x, z)} \bigwedge_{i,j} \frac{dx_{i,j}}{x_{i,j}} := \sum_{d=0}^{\infty} \frac{[S(x, z)^d]_0}{d! \epsilon^d}$$

where $[S(x, z)^d]_0$ denotes the constant term of the Laurent polynomial $S(x, z)^d$ in the variables $x = (x_{i,j})$. We assume that $[S(x, z)^0]_0 = 1$. From the structure of the superpotential




FIGURE 3. The graph Γ associated with $X = T^* \text{Gr}(k, n)$.

(4.1) it is easy to see that $[S(x, z)^d]_0$ is a monomial in $z = z_1/z_2$ and thus (4.4) is a power series in z .

Example. For $X = T^*\mathbb{P}^{n-1}$, the superpotential is given by (4.2). In this case, elementary computations shows that $[S(x, z)^d]_0$ is non-vanishing only if the degree d is of the form $d = nm$ for some $m \in \mathbb{N}$. In this case we have

$$[S(x, z)^{mn}]_0 = \frac{(z_1/z_2)^m (nm)!}{(m!)^n}.$$

We thus conclude that

$$\oint e^{\frac{1}{\epsilon} S(x, z)} \bigwedge_{i,j} \frac{dx_{i,j}}{x_{i,j}} = \sum_{m=0}^{\infty} \frac{z^m}{(m!)^n \epsilon^{mn}}.$$

Example. For $X = T^* \text{Gr}(2, 4)$, the superpotential is given by (4.3). In this case $[S(x, z)^d]_0$ is non-vanishing only if $d = 4m$, in which case

$$[S(x, z)^{4m}]_0 = \frac{(2m)!(4m)!}{(m!)^6} \frac{z^m}{\epsilon^{4m}}.$$

Thus, we obtain:

$$\oint e^{\frac{1}{\epsilon}S(x, z)} \bigwedge_{i,j} \frac{dx_{i,j}}{x_{i,j}} = \sum_{m=0}^{\infty} \frac{(2m)!}{(m!)^6 \epsilon^{4m}} z^m.$$

4.3. The vertex function in the $\hbar \rightarrow \infty$ limit.

Theorem 4.1. *Let $V(z)$ be the function (2.5), then:*

$$\lim_{\hbar \rightarrow \infty} V(z/\hbar^n) = \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint e^{\frac{1}{\epsilon}S(x, z)} \bigwedge_{i,j} \frac{dx_{i,j}}{x_{i,j}}$$

where the integral is defined by (4.4) and $S(x, z)$ is the polynomial superpotential (4.1).

Proof. By Theorem 3.2 we have:

$$V(z) = \frac{\alpha}{(2\pi\sqrt{-1})^{k(n-k)}} \oint_{\gamma_{k,n}} \Phi(x, z) \bigwedge_{i,j} dx_{i,j}$$

where the contour $\gamma_{k,n}$ is defined by (3.4) and $\Phi(x, z)$ is the branch of the superpotential (3.6) distinguished by Proposition 3.1. It will be convenient to define:

$$(4.5) \quad \tilde{L}(x_{i,a}, x_{j,b}) = \begin{cases} (1 - x_{i,a}/x_{j,b})^{-\omega}, & m_{i,a} < m_{j,b}, \\ (1 - x_{j,b}/x_{i,a})^{-\omega}, & m_{i,a} > m_{j,b} \end{cases}$$

which differ from (3.5) by a factor

$$\tilde{L}(x_{i,a}, x_{j,b}) = \begin{cases} L(x_{i,a}, x_{j,b}), & m_{i,a} < m_{j,b}, \\ e^{-\pi\sqrt{-1}\hbar/\epsilon} L(x_{i,a}, x_{j,b}), & m_{i,a} > m_{j,b}. \end{cases}$$

Recall that $\alpha = e^{\pi\sqrt{-1}N\hbar/\epsilon}$ where N is the total number of factors in $\Phi(x, z)$ for which $\tilde{L}(x_{i,a}, x_{j,b})/L(x_{i,a}, x_{j,b}) = e^{-\pi\sqrt{-1}\hbar/\epsilon}$. Thus, in these notations we have

$$(4.6) \quad V(z) = \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint_{\gamma_{k,n}} \tilde{\Phi}(x, z) \bigwedge_{i,j} \frac{dx_{i,j}}{x_{i,j}}$$

where

$$\tilde{\Phi}(x, z) = \left(\prod_{i=1}^{n-1} \prod_{a < b} \Delta(x_{i,a}, x_{i,b}) \right) \left(\prod_{i=1}^{n-2} \prod_{a=1}^{v_i} \prod_{b=1}^{v_{i+1}} \tilde{L}(x_{i,a}, x_{i+1,b}) \right) \left(\prod_{i=1}^k \tilde{L}(z_1, x_{k,i}) L(z_2, x_{n-k,i}) \right).$$

In this integral we rescale the variables by: $z_1 \rightarrow z_1$, $z_2 \rightarrow z_2 \hbar^n$, since in our notations $z = z_1/z_2$, this is equivalent to substitution $z \rightarrow z/\hbar^n$ in the left-hand side of (4.6). Let $\gamma'_{k,n}(\hbar)$ be the contour defined by the conditions

$$|x_{i,j}| = m_{i,j} \varepsilon |\hbar|^{m_{i,j}}.$$

Assuming that $|\hbar| > 1$ we have

$$m_{i,j} < m_{a,b} \implies m_{i,j} \varepsilon |\hbar|^{m_{i,j}} < m_{a,b} \varepsilon |\hbar|^{m_{a,b}}$$

for all pairs (i, j) and (a, b) . By assumption of Proposition 3.1, we have $|z_1| < |x_{1,1}|$ and $|x_{n-k,k}| < |z_2 \hbar^n|$ on $\gamma'_{k,n}(\hbar)$. Therefore $\gamma'_{k,n}(\hbar) \sim \gamma_{k,n}$ in the sense of Definition 3.3. Thus

$$(4.7) \quad V(z/\hbar^n) = \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint_{\gamma'_{k,n}(\hbar)} \tilde{\Phi}(x, z) \bigwedge_{i,j} \frac{dx_{i,j}}{x_{i,j}}$$

Now, in this integral we change the variables of integration by $x_{i,j} = y_{i,j}\hbar^{m_{i,j}}$. The contour $\gamma'_{k,n}(\hbar)$ in the variables $y_{i,j}$ is given by $|y_{i,j}| = m_{i,j}\varepsilon$, i.e., in the coordinates $y_{i,j}$ we integrate over the original contour $\gamma_{k,n}$. Overall we obtain:

$$(4.8) \quad V(z/\hbar^n) = \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint_{\gamma_{k,n}} \tilde{\Phi}(y, z) \bigwedge_{i,j} \frac{dy_{i,j}}{y_{i,j}}$$

where

$$(4.9) \quad \begin{aligned} \tilde{\Phi}(y, z) = & \left(\prod_{i=1}^{n-1} \prod_{a < b} \Delta(y_{i,a}\hbar^{m_{i,a}}, y_{i,b}\hbar^{m_{i,b}}) \right) \left(\prod_{i=1}^{n-2} \prod_{a=1}^{v_i} \prod_{b=1}^{v_{i+1}} \tilde{L}(y_{i,a}\hbar^{m_{i,a}}, y_{i+1,b}\hbar^{m_{i+1,b}}) \right) \times \\ & \times \left(\prod_{i=1}^k \tilde{L}(z_1, y_{k,i}\hbar^{m_{k,i}}) \tilde{L}(z_2\hbar^n, y_{n-k,i}\hbar^{m_{n-k,i}}) \right). \end{aligned}$$

We have:

$$\tilde{L}(y_{i,a}\hbar^{m_{i,a}}, y_{i+1,b}\hbar^{m_{i+1,b}}) = \begin{cases} \left(1 - (y_{i,a}/y_{i+1,b})/(\hbar^{m_{i+1,b}-m_{i,a}})\right)^{-\hbar/\epsilon}, & m_{i,a} < m_{i+1,b}, \\ \left(1 - (y_{i+1,b}/y_{i,a})/(\hbar^{m_{i,a}-m_{i+1,b}})\right)^{-\hbar/\epsilon}, & m_{i,a} > m_{i+1,b}. \end{cases}$$

Note that the powers of \hbar appearing in these factors are positive integers. Thus, we compute

$$\lim_{\hbar \rightarrow \infty} L(y_{i,a}\hbar^{m_{i,a}}, y_{i+1,b}\hbar^{m_{i+1,b}}) = \begin{cases} e^{\frac{1}{\epsilon} \frac{y_{i,a}}{y_{i+1,b}}}, & m_{i,a} = m_{i+1,b} - 1, \\ e^{\frac{1}{\epsilon} \frac{y_{i+1,b}}{y_{i,a}}}, & m_{i,a} = m_{i+1,b} + 1, \\ 1, & \text{otherwise.} \end{cases}$$

We also have

$$\tilde{L}(z_1, y_{k,i}\hbar^{m_{k,i}}) = \left(1 - (z_1/y_{k,i})/\hbar^{m_{k,i}}\right)^{-\hbar/\epsilon},$$

$$\tilde{L}(z_2\hbar^n, y_{n-k,i}\hbar^{m_{n-k,i}}) = \left(1 - y_{n-k,i}\hbar^{m_{n-k,i}-n}/z_2\right)^{-\hbar/\epsilon}.$$

with $m_{k,i} = 2i - 1$ and $m_{n-k,i} = n - 2k + 2i - 1$, therefore

$$\lim_{\hbar \rightarrow \infty} \tilde{L}(z_1, y_{k,i}\hbar^{m_{k,i}}) = \begin{cases} e^{\frac{1}{\epsilon} \frac{z_1}{y_{k,1}}}, & i = 1, \\ 1 & i \neq 1 \end{cases}$$

and

$$\lim_{\hbar \rightarrow \infty} \tilde{L}(z_2\hbar^n, y_{n-k,i}\hbar^{m_{n-k,i}}) = \begin{cases} e^{\frac{1}{\epsilon} \frac{y_{n-k,k}}{z_2}}, & i = k, \\ 1 & i \neq k \end{cases}$$

Finally

$$\Delta(y_{i,a}\hbar^{m_{i,a}}, y_{i,b}\hbar^{m_{i,b}}) = (1 - y_{i,a}/y_{i,b}/(\hbar^{m_{i,b}-m_{i,a}}))^{2\hbar/\epsilon}$$

and since $m_{i,b} - m_{i,a} \geq 2$ for $b > a$ we have

$$\lim_{\hbar \rightarrow \infty} \Delta(y_{i,a}\hbar^{m_{i,a}}, y_{i,b}\hbar^{m_{i,b}}) = 1.$$

In summary, the limit $\hbar \rightarrow \infty$ of a factor in (4.9) is non-trivial only if it corresponds to an edge of the graph Γ and:

$$(4.10) \quad \lim_{\hbar \rightarrow \infty} \tilde{\Phi}(y, z) = \prod_{e \in \text{edges}(\Gamma)} e^{\frac{1}{\epsilon} \frac{y_{h(e)}}{y_{t(e)}}} = e^{\frac{S(y, z)}{\epsilon}}$$

Finally, the point-wise limit (4.10) on the compact set $\gamma_{k,n}$ is uniform, the limit commutes with the integration and from (4.8) we obtain

$$\begin{aligned} \lim_{\hbar \rightarrow \infty} V(z/\hbar^n) &= \lim_{\hbar \rightarrow \infty} \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint_{\gamma_{k,n}} \tilde{\Phi}(y, z) \bigwedge_{i,j} \frac{dy_{i,j}}{y_{i,j}} \\ &= \frac{1}{(2\pi\sqrt{-1})^{k(n-k)}} \oint_{\gamma_{k,n}} e^{\frac{S(y, z)}{\epsilon}} \bigwedge_{i,j} \frac{dy_{i,j}}{y_{i,j}}, \end{aligned}$$

which completes the proof. \square

5. DWORK CONGRUENCES

Let p be a prime number. Consider the power series

$$(5.1) \quad F(z) = \sum_{d=0}^{\infty} [S(x, z)^d]_0 \in \mathbb{Z}[[z]].$$

Note that the coefficients of this series differ from the coefficients of the A -series in (4.4) by a factor $d!e^d$, i.e., these two series are related by the Borel integral transform.

Let us consider a system of polynomial truncations of the function $F(z)$:

$$F_s(z) = \sum_{d=0}^{p^s-1} [S(x, z)^d]_0 \in \mathbb{Z}[z], \quad s = 0, 1, 2, \dots$$

where as before we assume that $[S(x, z)^0]_0 = 1$.

Theorem 5.1. *We have the following congruences:*

$$\frac{F(z)}{F(z^p)} \equiv \frac{F_s(z)}{F_{s-1}(z^p)} \pmod{p^s}, \quad s = 1, 2, \dots$$

In particular, the polynomials $F_s(z)$ satisfy the Dwork type congruences:

$$\frac{F_{s+1}(z)}{F_s(z^p)} \equiv \frac{F_s(z)}{F_{s-1}(z^p)} \pmod{p^s}, \quad s = 1, 2, \dots$$

Proof. The proof of these congruences is based on the properties of the Newton polytope

$$\Delta(k, n) = \mathbf{N}(S(x, z)) \subset \mathbb{R}^{k(n-k)}$$

of the Laurent polynomial (4.1) in the variables $x = (x_{i,j})$. Let $f_{i,j}$ with $i = 1, \dots, k$, $j = 1, \dots, n-k$ denote the standard basis in $\mathbb{R}^{k(n-k)}$. The vectors $f_{i,j}$ correspond to the boxes of the $k \times (n-k)$ -diagram in Fig. 3. From (4.1), we see that $\Delta(k, n)$ is the convex hull of the vectors:

$$\begin{aligned} f_{1,1}, \quad & f_{i,j+1} - f_{i-1,j+1}, \quad i = 2, \dots, k, \quad j = 0, \dots, n-k-1, \\ -f_{k,n-k}, \quad & f_{i,j+1} - f_{i,j}, \quad i = 1, \dots, k, \quad j = 1, \dots, n-k-1. \end{aligned}$$

This polytope has been considered in many publications, in particular it is known to be *reflexive* see Theorem 3.1.3 in [BCKS98]. We recall that the origin $(0, \dots, 0)$ is the only integral point in a reflexive polytope, see for instance [Nill05] for an overview.

Now the proof of Theorem 5.1 follows from Theorem 1.1 in [MeV16], after simple modifications. Let $S(x, 1)$ denote the superpotential (4.1) with $z_1 = z_2 = 1$. Clearly, this Laurent polynomial has the same Newton polytope $\mathbf{N}(S(z, 1)) = \mathbf{N}(S(x, z)) = \Delta(k, n)$.

Consider

$$\mathbf{M}(\xi) = \sum_{d=0}^{\infty} [S(x, 1)^d]_0 \xi^d, \quad \mathbf{M}_s(\xi) = \sum_{d=0}^{p^s-1} [S(x, 1)^d]_0 \xi^d.$$

Since $(0, \dots, 0)$ is the only interior point of $\mathbf{N}(S(z, 1))$ we may apply Theorem 1.1 in [MeV16] and conclude that these functions satisfy the congruences:

$$(5.2) \quad \frac{\mathbf{M}(\xi)}{\mathbf{M}(\xi^p)} \equiv \frac{\mathbf{M}_s(\xi)}{\mathbf{M}_{s-1}(\xi^p)}, \quad \frac{\mathbf{M}_{s+1}(\xi)}{\mathbf{M}_s(\xi^p)} \equiv \frac{\mathbf{M}_s(\xi)}{\mathbf{M}_{s-1}(\xi^p)} \pmod{p^s}, \quad s = 1, 2, \dots.$$

From the structure of the superpotential $S(x, z)$ it is clear that

$$[S(x, z)^d]_0 = [S(x, 1)^d]_0 z^{\frac{d}{n}}.$$

In particular, this coefficient equals zero unless n divides d . From this we find that

$$\mathbf{F}(z) = \sum_{d=0}^{\infty} [S(x, 1)^d]_0 z^{\frac{d}{n}} = \mathbf{M}(z^{\frac{1}{n}})$$

and similarly $\mathbf{F}_s(z) = \mathbf{M}_s(z^{\frac{1}{n}})$. Now theorem 5.1 follows from (5.2) after the substitution $\xi \rightarrow z^{\frac{1}{n}}$. \square

For further discussion of Dwork type congruences for vertex functions and solutions of KZ equations we refer to [SmV23, Var22, SV19, VZ21].

Remark. Theorem 5.1 implies an infinite factorization:

$$\mathbf{F}(z) = \prod_{i=0}^{\infty} \frac{\mathbf{F}_s(z^{p^i})}{\mathbf{F}_{s-1}(z^{p^{i+1}})} \pmod{p^s},$$

cf. Theorem 5.3 in [SmV23].

REFERENCES

- [AO21] Mina Aganagic and Andrei Okounkov, *Elliptic stable envelopes*, J. Amer. Math. Soc., **34** (2021), n.1, 79–133
- [BCKS98] V. Batyrev, I. Ciocan-Fontanine, B. Kim, and D. van Straten, *Conifold transitions and mirror symmetry for Calabi-Yau complete intersections in Grassmannians*, Nuclear Physics B **514** (1998), 640–666
- [Cas20] Marco Castronovo, *Fukaya category of Grassmannians: Rectangles*, Adv. Math., **372**, <https://doi.org/10.1016/j.aim.2020.107287>.
- [Dan22] Ivan Danilenko, *Quantum differential equation for slices of the affine Grassmannian*, [arXiv:2210.17061](https://arxiv.org/abs/2210.17061), 1–57
- [Din20] Hunter Dinkins, *Symplectic Duality of $T^* \text{Gr}(k, n)$* , Math. Res. Lett., **29**, **3**, (2022), <https://dx.doi.org/10.4310/MRL.2022.v29.n3.a3>
- [Din22] *3d mirror symmetry of the cotangent bundle of the full flag variety*, Lett. in Math. Phys. **112**, **100**, (2022)
- [Dwo69] B. Dwork, *p -adic cycles*, Publ. Math. de l’IHÉS **37** (1969), 27–115
- [EHX97] Tohru Eguchi, Kentaro Hori, Chuan-Sheng Xiong, *Gravitational Quantum Cohomology*, Int.J.Mod.Phys. A**12**, (1997), 1743–1782
- [GK13] Davide Gaiotto, Peter Koroteev *On three dimensional quiver gauge theories and integrability*, JHEP, **126** (2013).
- [Giv96] A. Givental, *Equivariant Gromov-Witten invariants*, Internat. Math. Res. Notices (1996), no. 13, 613–663
- [GGI16] Sergey Galkin, Vasily Golyshev, Hiroshi Iritani, *Gamma classes and quantum cohomology of Fano manifolds: Gamma conjecture*, Duke Math. J. **165**(11), 2005–2077, (2016).
- [KoSm22] Yakov Kononov, Andrey Smirnov, *Pursuing Quantum Difference Equations II: 3D mirror symmetry*, IMRN, (2022), <https://doi.org/10.1093/imrn/rnac196>
- [Ko] Iakov Kononov, *Elliptic Stable Envelopes and 3D Mirror Symmetry*, PhD Thesis, Columbia University.
- [KoSt10] Christian Korff, Catharina Stroppel, *The $\widehat{\mathfrak{sl}}(n)_k$ -WZNW fusion ring: A combinatorial construction and a realisation as quotient of quantum cohomology*. Adv. in Math., **225**, **1**, 200–268, (2010).
- [KPSZ21] Peter Koroteev, Petr Pushkar, Andrey Smirnov, Anton Zeitlin, *Quantum K-theory of Quiver Varieties and Many-Body Systems*, Sel. Math. New Ser. **27**, **87** (2021)
- [LaTe17] Thomas Lam, Nicolas Templier, *The mirror conjecture for minuscule flag varieties*, [arXiv:1705.00758](https://arxiv.org/abs/1705.00758), (2017).
- [MaRi20] R. J. Marsh and K. Rietsch, *The B-model connection and mirror symmetry for Grassmannians*, Adv. Math., 366:107027, 131, 2020
- [MO19] D. Maulik, A. Okounkov, *Quantum groups and quantum cohomology*, Astérisque, t. 408, Société Mathématique de France, 2019, 1–277; <https://doi.org/10.24033/ast.1074>
- [MV02] Y. Markov and A. Varchenko, *Hypergeometric solutions of trigonometric KZ equations satisfy dynamical difference equations*, Adv. Math., 166(1):100–147, 2002
- [MeVl16] A. Mellit, M. Vlasenko, *Dwork’s congruences for the constant terms of powers of a Laurent polynomial*, Int. J. Number Theory **12** (2016), no. 2, 313–321

- [Nill05] B. Nill, *Reflexive Polytopes - Combinatorics and Convex Geometry*, <https://personales.unican.es/santosf/anogia05/slides/Nill-anogia05.pdf>
- [Oko17] Andrei Okounkov, *Lectures on K-theoretic computations in enumerative geometry*, volume 24 of IAS/Park City Math. Ser., pages 251–380. Amer. Math. Soc., Providence, RI, 2017.
- [Oko18] Andrei Okounkov, *Enumerative symplectic duality*, MSRI workshop “Structures in Enumerative Geometry” (the talk is accessible from MSRI web-page)
- [OS22] Andrei Okounkov, Andrey Smirnov, *Quantum difference equation for Nakajima varieties*, *Invent. Math.* **229**, 1203–1299 (2022).
- [PSZ16] Petr Pushkar, Andrey Smirnov, and Anton Zeitlin, *Baxter Operator from quantum K-theory*, *Advances in Mathematics*, 360, 12 2016.
- [RSVZ19] R. Rimanyi, A. Smirnov, A. Varchenko, Z. Zhou, *Three-dimensional mirror self-symmetry of the cotangent bundle of the full flag variety* SIGMA, 15: **093**, **22**, (2019)
- [RSVZ21] R. Rimanyi, A. Smirnov, A. Varchenko, Z. Zhou, *3D Mirror Symmetry and Elliptic Stable Envelopes*, IMRN, (2021), <https://doi.org/10.1093/imrn/rnaa389>
- [RW20] Richárd Rimányi, Andrzej Weber, *Elliptic classes of Schubert varieties via Bott–Samelson resolution*, *J. of Topol.*, (2020), <https://doi.org/10.1112/topo.12152>
- [SmV23] Andrey Smirnov, Alexander Varchenko, *The p -adic approximations of vertex functions via 3D-mirror symmetry*, arXiv:2302.03092
- [SV91] V. Schechtman, A. Varchenko, *Arrangements of hyperplanes and Lie algebra homology*, *Invent. Math.* **106** (1991), 139–194
- [SV19] V. Schechtman, A. Varchenko, *Solutions of KZ differential equations modulo p* , *Ramanujan J.* **48** (2019), no. 3, 655–683; <https://doi.org/10.1007/s11139-018-0068-x>, arXiv:1707.02615
- [SZ22] Andrey Smirnov, Zijun Zhou, *3d mirror symmetry and quantum K-theory of hypertoric varieties*, *Adv. in Math.* **395**, (2022) <https://doi.org/10.1016/j.aim.2021.108081>
- [TV22] V. Tarasov, A. Varchenko, *Landau–Ginzburg mirror, quantum differential equations and q KZ difference equations for a partial flag variety*, *Journal of Geometry and Physics*, Vol. **184**, Available online 23 November 2022, 104711, 1–58, <https://doi.org/10.1016/j.geomphys.2022.104711>
- [Var22] A. Varchenko, *Dwork-type congruences and p -adic KZ connection*, arXiv:2205.01479, 1–26
- [Vl18] M. Vlasenko, *Higher Hasse–Witt matrices*, *Indagationes Mathematicae*, **29** (2018) 1411–1424
- [VZ21] A. Varchenko and W. Zudilin, *Ghosts and congruences for p^s -approximations of hypergeometric periods*, arXiv:2107.08548, 1–29