

Statistical mechanics of cell aggregates: explaining the phase transition and paradoxical piezoelectric behavior of soft biological tissues

Pratik Khandagale¹, Hao Lin², Liping Liu^{2, 3*}, and Pradeep Sharma^{4, †}

¹*Department of Mechanical Engineering, University of Houston, Houston, Texas 77204, USA*

²*Department of Mechanical and Aerospace Engineering, Rutgers University, New Jersey 08854, USA*

³*Department of Mathematics, Rutgers University, New Jersey 08854, USA*

⁴*Departments of Mechanical Engineering, Physics, and the Materials Science and Engineering Program, University of Houston, Houston, Texas 77204, USA*

June 2, 2025

Abstract

Piezoelectricity in biological soft tissues is a controversial issue with differing opinions. While there is compelling experimental evidence to suggest a piezoelectric-like response in tissues such as the aortic wall (among others), there are equally compelling experiments that argue against this notion. In addition, the lack of a polar structure in the underlying components of most soft biological tissues supports the latter. In this paper, we address the collective behavior of cells within a two-dimensional cell aggregate from the viewpoint of statistical mechanics. Our starting point is the simplest form of energy for cell behavior that only includes known observable facts e.g. the electrical Maxwell stress or electrostriction, resting potential across cell membranes, elasticity, and we explicitly *exclude* any possibility of electromechanical coupling reminiscent of piezoelectricity at the cellular level. We coarse-grain our cellular aggregate to obtain its emergent mechanical, physical, and electromechanical properties. Our findings indicate that the fluctuation of cellular strain (\mathbf{E}) plays a similar role as the absolute temperature in a conventional atomistic-level statistical model. The coarse-grained effective free energy reveals several intriguing features of the collective behavior of cell aggregates, such as solid-fluid phase transitions and a distinct piezoelectric-like coupling, even though it is completely absent at the microscale. Closed-form formulas are obtained for key electromechanical properties, including stiffness, effective resting potential, critical \mathbf{E}^2 -temperature (or fluctuation) for solid-fluid phase transitions, and apparent piezoelectric coupling in terms of fluctuation and electric potential regulated by active cellular processes.

Keywords: statistical mechanics, cell aggregate, fluctuations, solid-to-fluid phase transition, electro-mechanical coupling

*liu.liping@rutgers.edu

†psharma@central.uh.edu

1 Introduction

Cell aggregates or tissues are composed of individual cells, intricately organized with notable variability in their shapes, sizes, and spatial arrangements. Each constituent cell is structurally complex, featuring a bilipid membrane interspersed with ion channels and molecular pumps. Enclosed within this membrane are cytoskeleton, cytoplasm, and various organelles. The hierarchical structures and complex interplay of biophysical processes within cells give rise to intriguing electromechanical properties of single cells and cell aggregates. In particular, the bilipid membrane and cytoskeleton provide shear resistance and maintain the structural integrity of a cell [1]; the combined activities of numerous ion channels and molecular pumps induce a long-term, steady-state electric potential difference across the cell membrane [2]. The propagation of changes in transmembrane potentials mediates cellular signaling and regulates critical biophysical processes [3]. Conversely, mechanical deformations can regulate ion channels and pumps, inducing cellular bioelectricity within cell aggregates. These electromechanical couplings provide the fundamental physical mechanism underlying biological functions, including cellular adhesion, migration, differentiation, and mechanosensing [4–9].

Cells in an aggregate interact with each other both mechanically and electrically. The resulting electromechanical properties of cell aggregates or tissues are derived from the collective behavior of individual cells, leading to properties and functionalities that are significantly richer than those of a single cell. For example, cell aggregates reversibly transition between a rigid solid-like phase and a floppy fluid-like phase referred to as a solid-fluid phase transition, which plays a critical role in the biological function of living tissues [10–17]. This phase transition can be actively regulated by transmembrane potential [18–21].

Currently, several mechanisms have been proposed to explain the solid-fluid phase transition in cell aggregates. These include a topological transition which promotes tissue fluidization by relaxing elastic stresses over longer time scales [10, 22, 23]; the sliding of coherent cells within the aggregate, where external forces induce gradual rearrangements of cell conformations [21]; and transitions driven by the density of multicellular vertices (rosettes) and intracellular tensions [11]. Within a tissue, cellular deformation and electric potential exhibit considerable fluctuation and variability. The fluctuation in cellular shape and size can be readily quantified, providing valuable insight into the physiological stage of the tissue [24].

The true nature of the electromechanical behavior of soft biological tissues has attracted repeated controversy. Claims have been made regarding the piezoelectricity of these tissues c.f. Fukada and reference therein [25] and even the presence of ferroelectricity [26]. Convincing counter-arguments have been made against this by Lenz et. al. [27], who argue that the observable electromechanical coupling is due to Maxwell stress or electrostriction. In an interesting turn of events, in a more recent work, we turn full circle where Ikushima et. al. [28] highlight ultrasound experiments that *do* appear to indicate the presence of piezoelectricity in aortic wall tissue.

Existing approaches to modeling cells and their aggregates can be broadly divided into three categories: agent-based models, continuum models, and statistical models. The agent-based approach treats a single cell as a homogeneous, isotropic, elastic, and spherical body [29–32]. While this approach is valuable for studying the interaction of individual cells with each other and with their environment [33–35], its applicability to the macroscopic behavior of cell aggregates remains limited. Continuum models for cell aggregates [36–41], by contrast, are well-suited for modeling large-scale phenomena where aggregate properties vary smoothly over length scales spanning several cell diameters. These models effectively capture the macroscopic response of cell aggregates and can be extended to incorporate additional physical phenomena such as intercellular interactions and formation of cell aggregates [42]. For instance, a continuum description of cell motility,

driven by cell-cell and cell-ECM (extracellular matrix) interactions, was developed using a non-local interaction term to account for adhesion between cells and between cells and the ECM [43–45]. Despite these successes, continuum approaches fall short in capturing the inherent variability in cellular deformation and electric potential within tissues. In contrast, statistical models provide a robust framework for addressing these limitations [46–49]. By applying the framework of statistical mechanics to cell aggregates, we can effectively describe the mechanical and electrical fluctuations observed in tissues, bridging the gap between microscopic cellular properties and macroscopic tissue behavior. As discussed below, this approach also allows for a quantitative analysis of deformation and electric potential fluctuation, offering deeper insights into the emergent electromechanical properties of cell aggregates.

The method of statistical mechanics has been employed in a number of studies on cells and their aggregates. A recently developed theory [50, 51] addressed the *active* processes in cells to provide insights into cell size distribution. They extended that work to include electrical fields to establish the threshold electrical fields that can be sensed by a cell [52]. The work [10] based on non-equilibrium statistical mechanics and a 2D vertex model was able to capture mechanical instabilities in cell aggregates. The study in [53] for active matter (living cells) focused on steric and aligning interactions and interactions driven by shape changes. The impact of thermal fluctuations in polarization on the renormalization of the flexoelectric coupling coefficient of cell membranes has also been studied using statistical mechanics [54]. However, none of the existing models incorporate both the mechanical and electrical fluctuations and inter-cellular interactions for cell aggregates.

In this work, we propose an electromechanical model to describe both single cells and cell aggregates based on structural symmetries and derive its macroscopic continuum limit. Notably, our starting point for the microscopic model (at the cell level) does not include any piezoelectric behavior and simply takes cognizance of the fact that there is a resting potential across the cell and the presence of electrostriction (which is exhibited by all materials). Fluctuations in geometric parameters, such as cell area and aspect ratio, can be effectively quantified through the analysis of tissue micrographs. Notably, the fluctuation in strain — later referred to as the \mathbf{E}^2 -temperature due to its analogous role to absolute temperature in classical thermal physics — serves as a quantitative measure of the inherent system variability [55]. Our resulting coarse-grained electromechanical model for cell aggregates is particularly well-suited for applications to epithelial tissues, which consist of flattened, tightly packed cell layers. However, our insights are broader and transferrable to other tissue configurations. Based on the coarse-grained model, we investigate the macroscopic electromechanical behavior of cell aggregates, demonstrating that effective electromechanical coupling and solid-fluid phase transitions arise from the collective cellular interactions. These resolve the existing paradox in the literature. The emergent properties depend on the \mathbf{E}^2 -temperature. The quantitative dependence of effective resting potential, tissue stiffness, $T1$ -transition rate, viscosity, and critical point for solid-fluid phase transition on the \mathbf{E}^2 -temperature could be validated by experimental observations. Such comparisons open the door to understanding how fluctuation and variability contribute to tissue behavior in different biological contexts, making the \mathbf{E}^2 -temperature an experimentally testable concept for linking microscopic fluctuations with macroscopic emergent properties.

Central premise

Following the paradigm of statistical mechanics, we treat a cell aggregate as a canonical ensemble of many cells. We borrow the terminologies and methodology from statistical mechanics with the caveat that the fluctuation or variability of the system mainly arise from biological active processes (e.g. growing, diffusion, etc), defects, inhomogeneities, among others, instead of the actual




Figure 1: Cell aggregate and single cell. (a) A typical cell aggregate (plated Madin-Darby Canine Kidney cells, Courtesy of K. Irvine). (b) Geometrically, a cell aggregate is regarded as a tessellation of a plane by polygons. (c) Schematic of a single cell occupying a regular hexagonal domain ($n = 6$) and (d) a deformed hexagon, where \mathbf{c} , \mathbf{x}_j , \mathbf{y}_j is the position of the centroid of the cell, the j^{th} vertex of the cell in undeformed and deformed configuration, respectively.

thermal agitations. Therefore, the benchmark energy scale ($\tilde{k}_B T = 1/\beta$) used in this work is a phenomenological constant; the fluctuation or variability in cellular states arises not only from thermal agitations but also from many other passive and active processes that entail “equilibrium” of the system at the observation time-scale. The collection of cellular strains and electric potentials in a cell aggregate are assumed to describe the microstates of the system. For their collective properties in a cell aggregates, we first propose a single-cell Hamiltonian based on the underlying structural symmetries and experimental observations of their electromechanical responses. We employ the mean-field approximation, which allows for an analytical solution of the model and yields a closed-form expression for the effective free energy of cell aggregates. Based on this effective free energy, we present several experimentally testable predictions, including the macroscopic electromechanical properties, the solid-fluid phase transitions observed in cell aggregates, and their dependence on the \mathbf{E}^2 -temperature and average electric potential regulated by active processes.

2 An electromechanical model for cell aggregates

We are interested in the macroscopic electromechanical response of a cell aggregate as illustrated in Fig. 1(a). We first introduce independent state variables for describing physical and geometrical configurations of cells and their aggregates. Figure 1(a) shows a schematic of a typical amorphous cell aggregate. We idealize the cell aggregate as a tessellation of a plane by equal-area polygons as illustrated in Fig. 1(b). Denote by $\mathcal{C}_i \subset \mathbb{R}^2$ the polygon/domain occupied by the i th cell, A the area of the polygon \mathcal{C}_i , and \mathbf{c}_i the spatial positions of the centroids of cells. For simplicity, we assume that each cell precisely occupies the Voronoi cell associated with $\{\mathbf{c}_i : i = 1, \dots, N\}$ [56]. Though it is tempting to employ the centroids of cells as the fundamental state variables, a reasonable physical model or Hamiltonian for the tessellation in terms of $\{\mathbf{c}_i : i = 1, \dots, N\}$ must be interacting and permutation invariant, and hence not amenable to systematic perturbation methods. On the other hand, for cell aggregates illustrated in Fig. 1(a), we see that the shapes of cells and statistical

properties of aspect ratios are good indicator of different physiological stages of cell aggregates (or tissues) which, presumably, dictate the physical properties of cell aggregates as well [55, 57, 58]. Therefore, we choose the deformation or strain as one of the independent state variables for the cell aggregate.

Unlike crystalline solids for which there exists a natural reference configuration (i.e., crystalline lattice), at least locally, for defining the deformation and strain, a cell aggregate is amorphous whose deformation or strain, strictly speaking, cannot be defined for a single current configuration. Nevertheless, neglecting the compatibility requirement for neighboring cells, “strain” could be defined for each individual cell by a few alternative procedures. The essence of these procedures is such that the introduced strain tensor $\mathbf{E} \in \mathbb{R}_{\text{sym}}^{2 \times 2}$ is a measure of orientation and aspect ratio of individual cells and the underlying arrangement of cells. In particular, we require that the strain tensor $\mathbf{E} \in \mathbb{R}_{\text{sym}}^{2 \times 2}$ satisfy the following.

- (R1) The strain $\mathbf{E} = 0$ if the cell is a regular polygon.
- (R2) For small strains with $|\mathbf{E}| \ll 1$, the ratio of eigenvalues of strain tensor $\mathbf{E} \in \mathbb{R}_{\text{sym}}^{2 \times 2}$ measures the “aspect ratio” of the cell.
- (R3) For small strains with $|\mathbf{E}| \ll 1$, the eigenvectors of strain tensor $\mathbf{E} \in \mathbb{R}_{\text{sym}}^{2 \times 2}$ defines the “orientation” of the cell.

As a tessellation of a plane, cells in aggregates occupy polygonal domain \mathcal{C} with n vertices \mathbf{y}_j ($j = 1, \dots, n$). As illustrated in Fig. 1(a), typically $n = 4, 5, 6, 7$ from experiments. Suppose a cell occupies an irregular hexagon as in Fig. 1(d). We choose the regular hexagon of the same area in Fig. 1(c) as the reference undeformed configuration.

Below we introduce two procedures to identify the strain tensor \mathbf{E} associated with a single cell in the regime of small strain ($|\mathbf{E}| \ll 1$).

- (i) Let \mathbf{x}_j ($j = 1, \dots, n$) be vertices of the regular n -gon centered at the origin:

$$\mathbf{x}_j = r_0 \mathbf{m}_j, \quad \mathbf{m}_j = [\cos \frac{j2\pi}{n}, \sin \frac{j2\pi}{n}], \quad (1)$$

where r_0 ($A_0 = \frac{1}{2}nr_0^2 \sin(2\pi/n)$) is such that the area of the regular n -gon equals the area of the cell. By the method of least square, we can in general define a deformation gradient $\mathbf{F} \in \mathbb{R}^{2 \times 2}$ and a translational vector $\mathbf{u}_0 \in \mathbb{R}^2$ for the cell:

$$\min\{L(\mathbf{F}, \mathbf{u}_0) = \sum_{j=1}^n |\mathbf{y}_j - \mathbf{F}\mathbf{x}_j - \mathbf{u}_0|^2 : \mathbf{F} \in \mathbb{R}^{2 \times 2}, \mathbf{u}_0 \in \mathbb{R}^2\}.$$

Let $\mathbf{u}_i = \mathbf{y}_i - \mathbf{x}_i$ be the displacement of the i th vertex and $\mathbf{H} = \mathbf{F} - \mathbf{I}$ the unsymmetrized strain ($\mathbf{I} \in \mathbb{R}^{2 \times 2}$ is the identity matrix). We write $L(\mathbf{F}, \mathbf{u}_0)$ as

$$L(\mathbf{H}, \mathbf{u}_0) = \sum_{i=1}^n \left[|\mathbf{u}_i - \mathbf{u}_0|^2 - 2(\mathbf{u}_i - \mathbf{u}_0) \cdot \mathbf{H}\mathbf{x}_i + \mathbf{x}_i \cdot \mathbf{H}^T \mathbf{H}\mathbf{x}_i \right].$$

By the first-order necessary conditions we have

$$\begin{aligned} 0 &= \frac{\partial L}{2\partial \mathbf{u}_0} = \sum_{i=1}^n (-\mathbf{u}_i + \mathbf{H}\mathbf{x}_i) + n\mathbf{u}_0, \\ 0 &= \frac{\partial L}{2\partial \mathbf{H}} = -\sum_{i=1}^n (\mathbf{u}_i - \mathbf{u}_0) \otimes \mathbf{x}_i + \mathbf{H} \sum_{i=1}^n \mathbf{x}_i \otimes \mathbf{x}_i, \end{aligned}$$

which imply that

$$\mathbf{u}_0 = \frac{1}{n} \sum_{j=1}^n \mathbf{u}_j, \quad \mathbf{F} = \mathbf{I} + \frac{1}{r_0} \sum_{j=1}^n \mathbf{u}_j \otimes \mathbf{m}_j.$$

From classical continuum mechanics, the linearized symmetric strain for the cell can be written as

$$\mathbf{E} = \frac{1}{2}(\mathbf{F} + \mathbf{F}^T) - \mathbf{I} = \frac{1}{2r_0} \sum_{j=1}^n (\mathbf{u}_j \otimes \mathbf{m}_j + \mathbf{m}_j \otimes \mathbf{u}_j). \quad (2)$$

(ii) A second procedure to determine the strain tensor is through the normalized second moment of the current position vector. For a polygon \mathcal{C} illustrated in Fig. 1(d), without loss of generality we assume the centroid is at the origin: $\mathbf{c} = \int_{\mathcal{C}} \mathbf{y} da = 0$. Then the normalized second moment tensor is given by

$$\mathbf{B} = \frac{1}{J_n} \int_{\mathcal{C}} \mathbf{y} \otimes \mathbf{y} da, \quad (3)$$

where $J_n = \frac{nr_0^4}{48}(4\sin(2\pi/n) + \sin(2\pi/n))$ is the second moment tensor of a regular n -gon of the same area (Cf. Fig. 1(c)). Suppose that the current shape \mathcal{C} is obtained by deforming the regular n -gon \mathcal{C}_0 with vertices given by \mathbf{x}_j in (1), i.e., $\mathbf{y}_j = \mathbf{F}\mathbf{x}_j$. The second normalized moment tensor is given by

$$\mathbf{B} = \frac{1}{J_n} \int_{\mathcal{C}_0} (\mathbf{F}\mathbf{x}) \otimes (\mathbf{F}\mathbf{x}) da = \mathbf{F}\mathbf{F}^\top.$$

Then the strain tensor may be defined as

$$\mathbf{E} = (\mathbf{B})^{1/2} - \mathbf{I}, \quad (4)$$

which clearly fulfills the previously mentioned requirements (R1)-(R3).

We remark that strains defined by (2) and (4) coincide to the leading order for small strains with $|\mathbf{E}| \ll 1$ and both of them have been used to quantify experimental and numerical results [55, 57]. Also, for hexagons the shape index [57] or perimeter (p)-to-sqrt of area (\sqrt{A}) ratio can be expressed as a function of strain \mathbf{E} :

$$\hat{p} = \frac{p}{\sqrt{A}} \approx \frac{\sqrt{2}(6 + \frac{9}{16}|\mathbf{E}|^2)}{3^{3/4}} \approx 3.72 + 0.35|\mathbf{E}|^2. \quad (5)$$

Strain from rearrangement of cells. As for crystalline solids, significant strain can be generated by structural rearrangement of cells from the viewpoint of centroid positions. Such a transformation strain cannot be characterized by the two definitions ((2) and (4)) of strain tensor. As illustrated in Fig. 2, in a rearrangement process called T1-transition commonly observed in the development of epithelial tissues, neighboring cells exchange their positions through a reorganization of intercellular junctions without significant changes in the shape of individual cells. Though the final configuration in Fig. 2(c) in T1-transition is a deformed configuration relative to the initial configuration in Fig. 2(a), the strain calculated by either (2) or (4) vanishes for every cell. To remedy this issue, we observe that, as a cluster of cells, the final configuration in Fig. 2(c) can be obtained by compressing (and stretching) the initial configuration in Fig. 2(a) in y -direction (in x -direction). The strain associated with this deformation process may be identified as

$$\mathbf{E}_{T1} = \frac{\gamma^*}{\sqrt{2}} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (6)$$




Figure 2: T1-transition in cell aggregates. Cell aggregates undergo rearrangement in which neighboring cells exchange their positions through a reorganization of intercellular junctions without significant changes in the shape of individual cells and transition from (a) initial configuration to (b) intermediate configuration to (c) final configuration.

where the T1-transition strain $\gamma^* \sim 1$ is beyond the regime of small strains and will be determined later from empirical data.

To summarize, we note that both definitions of the strain tensor, (2) and (4), encounter difficulty in consistently capturing the local structural rearrangements of cells. This limitation arises from the inherent complexity of cellular configurations. In this work, we take a pragmatic approach and treat the strain tensor \mathbf{E} as an ‘‘order parameter’’ as in the framework of Landau’s phenomenological theory [59, 60]. In this context, the precise mathematical definition of strain tensor \mathbf{E} beyond the regime of small strain is secondary, as long as it qualitatively reflects the key geometric characteristics — namely, the shape changes and structural arrangement of the underlying cells. By adopting this perspective, the strain tensor serves as a convenient and effective state variable for quantifying the geometry and arrangement of cells in a cell aggregate and can be confidently computed by either (2) or (4) in the regime of small strain.

In addition to elasticity, we will consider electric interactions between cells. Cell aggregates, in spite of adhesion between them, in general, are immersed in an extracellular matrix (ECM) whose electric potential may be assigned as the ground potential. Ionic channels, i.e., specialized protein structures embedded in the cell membrane, allow selective passage of ions (e.g., Na^+ , K^+ , Ca^{2+} , Cl^- , etc) across the membrane. This selective transport ensures and maintains a resting membrane potential ξ^* , which is essential for numerous biological functions [2]. For simplicity, we assume the interior of each cell is of equipotential. The actual potential of the i^{th} -cell, denoted by ξ_i , could be significantly different from the resting potential ξ^* due to electromechanical coupling and other passive or active transport processes in cell aggregates.

In summary, for a phenomenological model of cell aggregates we assume that the state of the cell aggregates are completely described by the strain tensor and electric potential of cells:

$$(\mathbf{E}_i, \xi_i) \in \mathcal{D} := \{(\mathbf{E}, \xi) : (\mathbf{E}, \xi) \in \mathbb{R}_{\text{sym}}^{2 \times 2} \times \mathbb{R}, \text{Tr}\mathbf{E} = 0\} \quad (i = 1, \dots, N), \quad (7)$$

where the constraint $\text{Tr}\mathbf{E} = 0$ arises from the incompressibility of cells. For brevity and future convenience, denote by s , \bar{s} , s' the collection of state variables of N -cells, the average, and their

deviations:

$$\begin{aligned} s &= \{\mathbf{E}_i, \xi_i\}_{i=1}^N \in \mathcal{D}^N, \\ \bar{s} &= (\bar{\mathbf{E}}, \bar{\xi}) = \frac{1}{N} \sum_{i=1}^N (\mathbf{E}_i, \xi_i) \in \mathcal{D}, \\ s' &= s - \bar{s} \in \mathcal{D}^N. \end{aligned} \quad (8)$$

Energetics. In line with standard practices in physical modeling, we formulate the electromechanical model for cell aggregates via an energy function. As illustrated in Fig. 1 (d), we consider a single cell from the aggregate with strain-electric potential $(\mathbf{E}, \xi) \in \mathcal{D}$ and propose that the cellular energy is given by

$$\Psi(\mathbf{E}, \xi) = \left(\frac{\mu}{\gamma^* 2} |\mathbf{E}|^2 (|\mathbf{E}|^2 - \gamma^*)^2 + \frac{\epsilon}{2\lambda^2} (\xi - \xi^*)^2 + K(\xi - \xi^*) |\mathbf{E}|^2 \right) A, \quad (9)$$

where A is the area of the cell, and $\mu, \gamma^* (\sim 1), \epsilon, \lambda > 0$ and ξ^*, K are constant model parameters. Physically, the energy terms in (9) are motivated from the following consideration. The first term characterizes the strain energy; the specific expression is the minimal form that is isotropic, thermodynamically stable, and admits the regular hexagonal tessellation (Cf. Fig. 2(a)) and its equivalent configuration after $T1$ -transition (Cf. Fig. 2(c)) as the strain-energy minimizing states. In particular, the constant μ may be recognized as the shear modulus (for small strain). The second term $\frac{\epsilon}{2\lambda^2} (\xi - \xi^*)^2$ is the electric field energy with ξ^* being the resting potential (at the absence of electromechanical coupling, i.e., $K = 0$), ϵ the dielectric constant, and λ a length scale comparable to the thickness of cell membrane. The electric field energy penalizes the deviation of interior cellular potential from the resting potential. The last term $K(\xi - \xi^*) |\mathbf{E}|^2$ is for capturing the coupling between electrical field and mechanical strain. We remark that the proposed single-cell energy (9) represents the minimal isotropic electromechanical model that achieves the following: (i) it effectively reproduces the observed $T1$ -transitions in cell aggregates, featuring with the characteristic rearrangements during tissue remodeling, and (ii) it incorporates nontrivial electromechanical coupling, allowing for a meaningful interplay between mechanical deformations and electrical responses. This balance of simplicity and functionality makes the proposed model (9) a robust foundation for exploring the electromechanical properties of tissues.

We next consider cell aggregates that are *nonuniformly* deformed or charged. Let \mathcal{I}_i be the index set that contains all neighboring cells in contact with the i th cell. In account of nonuniformity, we postulate that the *interaction energy* for the i^{th} cell is:

$$H_{int}(s) = \frac{1}{2} \frac{A}{l^2} J^{\text{elast}} \sum_{j \in \mathcal{I}_i} |\mathbf{E}_i - \mathbf{E}_j|^2 + \frac{1}{2} \frac{A}{l^2} J^{\text{elect}} \sum_{j \in \mathcal{I}_i} |\xi_i - \xi_j|^2, \quad (10)$$

where the length scale l is comparable to the overall size of a cell, and constant $J^{\text{elast}} > 0$ (resp. $J^{\text{elect}} > 0$) measures the increase of energy when the strain (resp. electric potential) of a cell is different from that of neighboring cells. In the context of the Ising model, J^{elast} and J^{elect} are referred to as the exchange constants. Combining (9) with (10), we identify the Hamiltonian of an N -cell system as

$$H(s) = \sum_{i=1}^N \Psi(\mathbf{E}_i, \xi_i) + \sum_{i=1}^N H_{int}(\mathbf{E}_i, \xi_i), \quad (11)$$

which serves as the foundation for our statistical mechanics model.

We will follow the standard procedure of statistical mechanics to analyze the cell aggregate. In the classical statistical and thermal physics, the critical concept of (absolute) temperature reflects the fluctuation of energy associated with microstates of the system, whereas entropy measures the number of accessible microstates of the system. In thermal equilibrium, the probability distribution over all admissible microstates of the system is dictated by the Second Law, which can then be used to determine macroscopic properties and thermodynamic relations of the system. To apply this approach to the electromechanical model (11) of a cell aggregate, we start from the following hypotheses [61].

(H1) The variability or fluctuations in state variables (\mathbf{E}, ξ) are macroscopically homogeneous within the system (a cell aggregate) and can still be characterized by a single benchmark energy scale $\tilde{k}_B T = 1/\beta$.

We remark that unlike the classical systems such as ideal gases, the stochasticity in cell aggregates has many contributions including but not limited to thermal agitations, passive and active biological processes (e.g., growing, diffusion, etc.), among others. Therefore, the benchmark energy scale $\tilde{k}_B T = 1/\beta$ should be understood as a phenomenological parameter instead of the actual thermal energy scale that originates from the oscillations of constituent molecules or atoms. Nevertheless, for physiological relevance we can still interpret T as the absolute temperature of the system. Meanwhile, \tilde{k}_B should be regarded as an empirical constant in parallel to the classical Boltzmann's constant. Similar ideas have been used in the literature to describe novel phenomena in a range of physical systems, including granular media [62] and plasticity in polycrystalline materials [61].

(H2) In quasi-static processes, the cell aggregate may exchange energy with the environment and stay in instantaneous equilibrium with a constant and uniform temperature T , and hence follows the Boltzmann distribution:

$$p(s) \propto e^{-\beta H(s)}, \quad (12)$$

where $s = \{\mathbf{E}_i, \xi_i\}_{i=1}^N \in \mathcal{D}^N$ represents an admissible microstate of the cell aggregate and $p(s)$ is the probability of the aggregate in the microstate s .

Based on these hypotheses, we consider a canonical ensemble of the system at a constant temperature T and upscale from the microscopic model described by the Hamiltonian (11) to a macroscopic coarse-grained model. Denote the collection of all microstates with prescribed average $(\bar{\mathbf{E}}, \bar{\xi})$ by

$$\mathcal{D}_{(\bar{\mathbf{E}}, \bar{\xi})}^N = \{s \in \mathcal{D}^N : \frac{1}{N} \sum_{i=1}^N (\mathbf{E}_i, \xi_i) = (\bar{\mathbf{E}}, \bar{\xi})\}. \quad (13)$$

For a coarse-grained macroscopic model, we aim to calculate the effective free energy (per cell) with a prescribed average:

$$F^{\text{eff}}(\bar{\mathbf{E}}, \bar{\xi}; \beta) = -\frac{1}{N\beta} \log \mathbf{Z}(\bar{\mathbf{E}}, \bar{\xi}; \beta), \quad (14)$$

where, by (12), the partition function with prescribed average is identified as

$$Z(\bar{\mathbf{E}}, \bar{\xi}; \beta) = \int_{\mathcal{D}_{(\bar{\mathbf{E}}, \bar{\xi})}^N} \exp(-\beta \mathbf{H}(\mathbf{s})) \, d\mathbf{s}. \quad (15)$$

We remark that the effective free energy defined in (14) is such that the partition function without prescribed average is given by

$$\begin{aligned} Z &:= \int_{\mathcal{D}^N} e^{-\beta H(s)} ds = \int_{\mathcal{D}} \left[\int_{\mathcal{D}_{(\bar{\mathbf{E}}, \bar{\xi})}^N} e^{-\beta H(s)} ds \right] d\bar{s} = \int_{\mathcal{D}} Z(\bar{\mathbf{E}}, \bar{\xi}; \beta) d\bar{s} \\ &= \int_{\mathcal{D}} e^{-N\beta F^{\text{eff}}(\bar{\mathbf{E}}, \bar{\xi}; \beta)} d\bar{s}, \end{aligned} \quad (16)$$

where the last equality justifies the terminology of “effective” free energy that governs the macroscopic coarse-grained properties of cell aggregates.

3 Effective free energy by mean-field approximation

For an N -cell aggregate in equilibrium, presumably cells would stay at the resting potential ξ^* and tessellate the plane by regular hexagons to minimize the total energy. However, the presence of passive and active noises, size dispersion, topological defects, and thermal agitations give rise to variability and randomness in cell aggregates. From the postulated Boltzmann distribution (12), we can (i) identify the temperature or energy scale $1/\beta = \tilde{k}_B T$ from the *fluctuations* of states of cell aggregates, and (ii) predict the macroscopic measurable effective material properties (e.g., shear modulus and electromechanical coupling coefficients) and how they depend on the fluctuations or, equivalently, the energy scale $\tilde{k}_B T$. For these purposes, we need to evaluate the partition function (15) and the effective free energy (14) for some prescribed average/macrosopic strain and potential $(\bar{\mathbf{E}}, \bar{\xi})$.

3.1 Mean-field approximation

As for a standard Ising model, an exact solution to (15) is still an open problem for nontrivial interactions (encoded by the index set \mathcal{I}_i). A widely accepted approximation method is the so-called *the mean-field theory*, which simplifies the problem by neglecting the contribution of interaction energy from fluctuations. More precisely, we rewrite the interaction energy of cell aggregates in (11) as

$$\begin{aligned} \sum_{i=1}^N H_{\text{int}}(\mathbf{E}_i, \xi_i) &= \sum_{i=1}^N \left[\frac{1}{2} \frac{A}{l^2} J^{\text{elast}} \sum_{j \in \mathcal{I}_i} |\mathbf{E}_i - \mathbf{E}_j|^2 + \frac{1}{2} \frac{A}{l^2} J^{\text{elect}} \sum_{j \in \mathcal{I}_i} |\xi_i - \xi_j|^2 \right] \\ &= \sum_{i=1}^N \left[\frac{1}{2} \frac{A}{l^2} J^{\text{elast}} \sum_{j \in \mathcal{I}_i} |(\mathbf{E}_i - \bar{\mathbf{E}}) - (\mathbf{E}_j - \bar{\mathbf{E}})|^2 + \frac{1}{2} \frac{\mathbf{A}}{l^2} \mathbf{J}^{\text{elect}} \sum_{j \in \mathcal{I}_i} |(\xi_i - \bar{\xi}) - (\xi_j - \bar{\xi})|^2 \right] \\ &\approx q \frac{A}{l^2} J^{\text{elast}} \sum_{i=1}^N |\mathbf{E}_i - \bar{\mathbf{E}}|^2 + \mathbf{q} \frac{\mathbf{A}}{l^2} \mathbf{J}^{\text{elect}} \sum_{i=1}^N |\xi_i - \bar{\xi}|^2, \end{aligned} \quad (17)$$

where q is the number of neighboring cells (i.e. the size of the index set \mathcal{I}_i) which is assumed to be independent of i . Denote by

$$\mathbf{E}'_i = \mathbf{E}_i - \bar{\mathbf{E}} \quad \text{and} \quad \xi'_i = \xi_i - \bar{\xi}$$

the deviation from the average strain and average electric potential, respectively. In general, a Hamiltonian can be decomposed into three parts: the first depends only on the averaged or coarse-grained state variables, the second only on the deviations or high-frequency microscopic

fluctuations, and the third on both. For instance, within the mean-field approximation (17) the Hamiltonian (11) of the cell aggregate can be written as

$$H(s) = H_0(\bar{s}) + H_1(s') + H_2(s'; \bar{s}), \quad (18)$$

where, by direct calculation, we find that

$$\begin{aligned} H_0(\bar{s}) &= N\Psi(\bar{\mathbf{E}}, \bar{\xi}), \quad \mathbf{H}_1(s') = \sum_{i=1}^N \left(\Psi(\mathbf{E}'_i, \xi'_i) + \frac{\mathbf{q}\mathbf{J}^{\text{elast}}\mathbf{A}}{\mathbf{l}^2} |\mathbf{E}'_i|^2 + \frac{\mathbf{q}\mathbf{J}^{\text{elec}}\mathbf{A}}{\mathbf{l}^2} |\xi'_i|^2 \right), \\ H_2(s'; \bar{s}) &= \sum_{i=1}^N (\Psi(\mathbf{E}'_i + \bar{\mathbf{E}}, \xi'_i + \bar{\xi}) - \Psi(\mathbf{E}'_i, \xi'_i) - \Psi(\bar{\mathbf{E}}, \bar{\xi})). \end{aligned} \quad (19)$$

We remark that the Hamiltonian (18), albeit anharmonic, is *non-interacting* because of the mean-field approximation. The associated partition function (15) and free energy (14) can be approximately evaluated by perturbation methods. To this end, we insert (18) and (19) into (14) and obtain

$$\begin{aligned} F^{\text{eff}}(\bar{\mathbf{E}}, \bar{\xi}; \beta) &= \Psi(\bar{\mathbf{E}}, \bar{\xi}) - \frac{1}{N\beta} \log \left[\int_{\mathcal{D}_{(0,0)}^N} e^{-\beta\mathbf{H}_2(s'; \bar{s})} e^{-\beta\mathbf{H}_1(s')} d\mathbf{s}' \right] \\ &=: \Psi(\bar{\mathbf{E}}, \bar{\xi}) + \tilde{F}(\bar{\mathbf{E}}, \bar{\xi}; \beta). \end{aligned} \quad (20)$$

Further, upon neglecting the constraint in the integration domain, i.e., replacing $\mathcal{D}_{(0,0)}^N$ by \mathcal{D}^N , we can write \tilde{F} defined above as

$$\begin{aligned} \tilde{F}(\bar{\mathbf{E}}, \bar{\xi}; \beta) &= -\frac{1}{\beta} \log \left[\int_{\mathcal{D}} e^{-\beta\tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi})} d(\mathbf{E}', \xi') \right], \\ \tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi}) &= \Psi(\mathbf{E}' + \bar{\mathbf{E}}, \xi' + \bar{\xi}) - \Psi(\bar{\mathbf{E}}, \bar{\xi}) + \frac{\mathbf{q}\mathbf{J}^{\text{elast}}\mathbf{A}}{\mathbf{l}^2} |\mathbf{E}'|^2 + \frac{\mathbf{q}\mathbf{J}^{\text{elec}}\mathbf{A}}{\mathbf{l}^2} |\xi'|^2. \end{aligned} \quad (21)$$

Two remarks are worth noting here. First, the replacement of $\mathcal{D}_{(0,0)}^N$ by \mathcal{D}^N can be justified *a posteriori*. As will be shown shortly, the expected value of the deviations in our subsequent calculations does satisfy $\langle \mathbf{E}' \rangle = \langle \xi' \rangle = 0$. Second, the reduction of the integral over the high-dimensional space \mathcal{D}^N to \mathcal{D} is possible due to the mean-field approximation (18), which accounts for interactions between neighboring cells by mean-fields.

3.2 Variational perturbation method

We now evaluate the effective free energy (F^{eff}), i.e., the integral (21), using the variational perturbation method. However, the presence of anharmonic terms like $|\mathbf{E}'|^4$ and $|\mathbf{E}'|^6$ in Ψ (Cf. (9)) defies an exact closed-form solution to (21). One approach to making progress is through the standard perturbation method. To this end, we choose a comparison energy function $\Psi_c = \Psi_c(\mathbf{E}', \xi')$ and denote by

$$\rho_c(\mathbf{E}', \xi') = \frac{1}{Z_c} e^{-\beta\Psi_c(\mathbf{E}', \xi')} \quad \left(Z_c = \int_{\mathcal{D}} e^{-\beta\Psi_c(\mathbf{E}', \xi')} d\mathbf{E}' d\xi' \right), \quad (22)$$

the probability distribution function (PDF) associated with the comparison energy function Ψ_c . The statistical average using PDF (22) is defined as

$$\langle \cdot \rangle_c := \int_{\mathcal{D}} (\cdot) \rho_c(\mathbf{E}', \xi') d\mathbf{E}' d\xi'. \quad (23)$$

Then the Bogoliubov inequality [63] asserts that for any energy function of \mathbf{E}' and ξ' , e.g., the function $\tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi})$ defined in (21), we have

$$\tilde{F}(\bar{\mathbf{E}}, \bar{\xi}; \beta) \lesssim \tilde{\mathbf{F}}^{\mathbf{U}}(\bar{\mathbf{E}}, \bar{\xi}; \beta) = \mathbf{F}_{\mathbf{c}} + \langle \tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi}) - \Psi_{\mathbf{c}}(\mathbf{E}', \xi') \rangle_{\mathbf{c}}, \quad (24)$$

where $F_c = -\frac{1}{\beta} \log Z_c$ is a constant independent of coarse-grained state variables $(\bar{\mathbf{E}}, \bar{\xi})$.

We now analytically evaluate the right-hand-side of (24) by choosing a quadratic comparison energy function:

$$\Psi_c(\mathbf{E}', \xi') = A \left(\mu_c |\mathbf{E}'|^2 + \eta_c \xi'^2 + \kappa_c \xi' (E'_{11} + E'_{12}) \right), \quad (25)$$

where $\mu_c, \eta_c > 0$ and $\kappa_c \in \mathbb{R}$ are some constants that guarantee positivity of (25) and will be fixed later. For the best upper-bound estimate of F^{eff} , by (20) we may set

$$F^{\text{eff}}(\bar{\mathbf{E}}, \bar{\xi}; \beta) = \Psi(\bar{\mathbf{E}}, \bar{\xi}) + \min\{\tilde{\mathbf{F}}^{\mathbf{U}}(\bar{\mathbf{E}}, \bar{\xi}; \beta) : \mu_{\mathbf{c}}, \eta_{\mathbf{c}} > \mathbf{0}, \kappa_{\mathbf{c}} \in \mathbb{R}\}, \quad (26)$$

where the minimization problem is over all reasonable Gaussian distribution (22) or positive harmonic comparison energy function $\Psi_c(\mathbf{E}', \xi')$.

In this work, we consider a planar model in which the fluctuating strain \mathbf{E}' , average strain $\bar{\mathbf{E}}$ can be written as

$$\mathbf{E}' = \begin{bmatrix} E'_{11} & E'_{12} \\ E'_{12} & -E'_{11} \end{bmatrix}, \quad \bar{\mathbf{E}} = \begin{bmatrix} \bar{E}_{11} & \bar{E}_{12} \\ \bar{E}_{12} & -\bar{E}_{11} \end{bmatrix}. \quad (27)$$

The quadratic nature of the comparison energy turns integrals in (23) into Gaussian-type integrals which can be analytically evaluated. In particular, we are interested in evaluating the $(\bar{\mathbf{E}}, \bar{\xi})$ -dependent terms on the right-hand side of (24), i.e., $\langle \tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi}) \rangle_{\mathbf{c}}$. From (9) and (21)₂, we find that

$$\begin{aligned} \frac{1}{A} \tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi}) &= (\mu + \frac{\mathbf{qJ}_{\text{elast}}}{\mathbf{l}^2} + \mathbf{K}(\bar{\xi} - \xi^*)) |\mathbf{E}'|^2 + (\frac{\epsilon}{2\lambda^2} + \frac{\mathbf{qJ}_{\text{elec}}}{\mathbf{l}^2}) |\xi'|^2 + 2\mathbf{K}\xi'(\bar{\mathbf{E}} \cdot \mathbf{E}') \\ &- \frac{4\mu}{\gamma^*} |\bar{\mathbf{E}}|^2 |\mathbf{E}'|^2 - \frac{2\mu}{\gamma^*} |\mathbf{E}'|^4 + \frac{\mu}{\gamma^*} |\mathbf{E}'|^6 - \frac{8\mu}{\gamma^*} (\bar{\mathbf{E}} \cdot \mathbf{E}')^2 + \frac{3\mu}{\gamma^*} |\bar{\mathbf{E}}|^2 |\mathbf{E}'|^4 \\ &+ \frac{12\mu}{\gamma^*} (\bar{\mathbf{E}} \cdot \mathbf{E}')^2 (|\mathbf{E}'|^2 + |\bar{\mathbf{E}}|^2) + \frac{3\mu}{\gamma^*} |\bar{\mathbf{E}}|^4 |\mathbf{E}'|^2 + \text{OT}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi}), \end{aligned} \quad (28)$$

where $\text{OT}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi})$ represents terms that are odd in (\mathbf{E}', ξ') , and hence will not contribute to $\langle \tilde{\Psi} \rangle_{\mathbf{c}}$. By tedious but straightforward calculations, we find that

$$\begin{aligned} \frac{1}{A} \langle \tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi}) \rangle_{\mathbf{c}} &= \mu_0 + 2K \langle \xi' \mathbf{E}' \rangle_c \cdot \bar{\mathbf{E}} + \mathbf{K} \langle |\mathbf{E}'|^2 \rangle_c (\bar{\xi} - \xi^*) \\ &+ \mu_2 |\bar{\mathbf{E}}|^2 + \bar{\mathbf{E}} \cdot \mathbb{C} \bar{\mathbf{E}} + \mu_4 |\bar{\mathbf{E}}|^4 + \frac{12\mu |\bar{\mathbf{E}}|^2}{\gamma^*} \bar{\mathbf{E}} \cdot \hat{\mathbb{C}} \bar{\mathbf{E}}, \end{aligned} \quad (29)$$

where μ_0 includes all terms that are independent of $(\bar{\mathbf{E}}, \bar{\xi})$, and

$$\begin{aligned} \mu_2 &= -\frac{4\mu}{\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c + \frac{3\mu}{\gamma^*} \langle |\mathbf{E}'|^4 \rangle_c, \\ \mathbb{C} &= \left\langle -\frac{8\mu}{\gamma^*} \mathbf{E}' \otimes \mathbf{E}' + \frac{12\mu |\mathbf{E}'|^2}{\gamma^*} \mathbf{E}' \otimes \mathbf{E}' \right\rangle_c, \\ \hat{\mathbb{C}} &= \langle \mathbf{E}' \otimes \mathbf{E}' \rangle_c, \quad \mu_4 = \frac{3\mu}{\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c, \end{aligned} \quad (30)$$

and

$$\begin{aligned}
\langle |\xi'|^2 \rangle_c &= \frac{4\mu_c}{k_0}, \quad k_0 := 2\beta A (4\eta_c\mu_c - \kappa_c^2), \\
\langle \xi' \mathbf{E}' \rangle_c &= -\frac{\kappa_c}{k_0} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \\
\langle |\mathbf{E}'|^2 \rangle_c &= 2 [\langle E_{11}'^2 \rangle_c + \langle E_{12}'^2 \rangle_c] = \frac{8\eta_c\mu_c - \kappa_c^2}{\mu_c k_0}.
\end{aligned} \tag{31}$$

Further, we find that the fourth-order tensor $\hat{\mathbb{C}}$ satisfies that

$$\langle \mathbf{Q} \mathbf{E}' \mathbf{Q}^\top \otimes \mathbf{Q} \mathbf{E}' \mathbf{Q}^\top \rangle_c = \int_{\mathcal{D}} \mathbf{E}' \otimes \mathbf{E}' \rho_c(\mathbf{E}', \xi') d\mathbf{E}' d\xi' = \langle \mathbf{E}' \otimes \mathbf{E}' \rangle_c \quad \forall \mathbf{Q} \in \text{SO}(2),$$

meaning that the fourth-order tensor $\hat{\mathbb{C}}$ is isotropic and such that

$$\bar{\mathbf{E}} \cdot \hat{\mathbb{C}} \bar{\mathbf{E}} = \frac{1}{2} \langle |\mathbf{E}'|^2 \rangle_c |\bar{\mathbf{E}}|^2. \tag{32}$$

Inserting (30)-(32) into (29), we obtain the closed-form upper bound estimate for the effective free energy $F^{\text{eff}}(\bar{\mathbf{E}}, \bar{\xi}; \beta)$ defined in (26) as

$$\begin{aligned}
\frac{1}{A} F^{\text{eff}}(\bar{\mathbf{E}}, \bar{\xi}) &\approx \mu_0 + \Psi(\bar{\mathbf{E}}, \bar{\xi}) + \frac{1}{A} \langle \tilde{\Psi}(\mathbf{E}', \xi'; \bar{\mathbf{E}}, \bar{\xi}) \rangle_c \\
&\approx \mu_2^{\text{eff}} |\bar{\mathbf{E}}|^2 + \mu_4^{\text{eff}} |\bar{\mathbf{E}}|^4 + \frac{\mu}{\gamma^* 2} |\bar{\mathbf{E}}|^6 + \mathbf{Q}^{\text{eff}}(\bar{\xi} - \xi^*) + \frac{\epsilon}{2\lambda^2} (\bar{\xi} - \xi^*)^2 + \mathbf{K}(\bar{\xi} - \xi^*) |\bar{\mathbf{E}}|^2,
\end{aligned} \tag{33}$$

where higher-order terms beyond $O(|\bar{\mathbf{E}}|^6, (\bar{\xi} - \xi^*)^2, \frac{1}{\beta})$ and immaterial constants are neglected, and the relevant effective properties are given by

$$\begin{aligned}
\mu_2^{\text{eff}} &= \mu(1 - \frac{8}{\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c), \quad Q^{\text{eff}} = K \langle |\mathbf{E}'|^2 \rangle_c, \\
\mu_4^{\text{eff}} &= \frac{2\mu}{\gamma^*} \left(-1 + \frac{9}{2\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c \right).
\end{aligned} \tag{34}$$

We remark that the effective free energy function, as defined in (33), governs the macroscopic electromechanical properties of cell aggregates.

4 Results and discussion

In this section, we explore biophysical implications of the coarse-grained model (33), with a particular focus on understanding how fluctuations impact the electromechanical behavior of cell aggregates. To facilitate meaningful comparisons with experimental results and numerical simulations, we select relevant model parameters in (9) such that their values lie within the physiological range. Specifically, the resting potential is chosen as $\xi^* = -60 \text{ mV}$ [12, 64], the range of change in electric potential of a single cell as $(\bar{\xi} - \xi^*) \in [-200 \text{ mV}, 200 \text{ mV}]$ [13, 16, 17, 64], the thickness of cell membrane as $\lambda = 5 \text{ nm}$ [65], shear modulus as $\mu = 200 \text{ Pa}$ [66–69], the relative permittivity of a single cell as $\epsilon_r = 20$ [65], and the electromechanical coupling constant for a single cell is chosen as $K \sim 0.1 \frac{\mu}{|\xi^*|}$. For simplicity, we focus on scenarios of simple shear with

$$\bar{\mathbf{E}} = \begin{bmatrix} 0 & \gamma \\ \gamma & 0 \end{bmatrix} \quad \text{and} \quad |\bar{\mathbf{E}}|^2 = 2\gamma^2.$$

Upon neglecting an $(\bar{\mathbf{E}}, \bar{\xi})$ -independent constant, we rewrite the effective free energy function (33) as

$$\frac{1}{A} F^{\text{eff}}(\gamma, \bar{\xi}) = (\mu_2^{\text{eff}} + K(\bar{\xi} - \xi^*))2\gamma^2 + 4\mu_4^{\text{eff}}\gamma^4 + \frac{8\mu}{\gamma^{*2}}\gamma^6 + \frac{\epsilon}{2\lambda^2}(\bar{\xi} - \xi^* + \frac{Q^{\text{eff}}\lambda^2}{\epsilon})^2. \quad (35)$$

Characterization of relevant energy scale. As demonstrated in [55, 57], the *fluctuation* $\langle |\mathbf{E}'|^2 \rangle_c$ can be directly obtained from experimental and numerical images of tissues. By (31), we see that the relevant energy scale

$$\tilde{k}_B T = \frac{1}{\beta} \propto \langle |\mathbf{E}'|^2 \rangle_c. \quad (36)$$

Aside from a proportionality constant, the fluctuation $\langle |\mathbf{E}'|^2 \rangle_c$ plays exactly the same role as the temperature in the classical statistical mechanics and is referred to as the \mathbf{E}^2 -temperature in [55]. **\mathbf{E}^2 -temperature characterizes stochasticity (or fluctuations) in cellular strains contributed by complex active and passive biological processes.** Therefore, \mathbf{E}^2 -temperature should be understood as some kind of empirical parameter or effective temperature [61, 70–72] instead of the actual thermal temperature that originates from the vibration of atoms in classical thermodynamics. Subsequently, we study how macroscopic electromechanical properties of the cell aggregate depend on the \mathbf{E}^2 -temperature (or fluctuation).

Renormalized resting potential. At the absence of external mechanical loading, the equilibrium state of the cell aggregate is determined by the principle of minimum free energy:

$$\begin{aligned} \min_{(\gamma, \bar{\xi})} F^{\text{eff}}(\gamma, \bar{\xi}) \quad &\Rightarrow \\ \frac{\partial}{\partial \gamma} F^{\text{eff}}(\gamma, \bar{\xi}) = 0 \quad &\text{and} \quad \frac{\partial}{\partial \bar{\xi}} F^{\text{eff}}(\gamma, \bar{\xi}) = 0. \end{aligned} \quad (37)$$

It is easy to see that a local small-strain minimizer is given by

$$(\gamma, \bar{\xi}) = (0, \xi^{*\text{eff}}) \quad \text{and} \quad \xi^{*\text{eff}} = \xi^* - \frac{Q^{\text{eff}}\lambda^2}{\epsilon} = \xi^* - \frac{K\lambda^2}{\epsilon} \langle |\mathbf{E}'|^2 \rangle_c. \quad (38)$$

Depending on the sign of K in the single-cell Hamiltonian (9), the \mathbf{E}^2 -temperature $\langle |\mathbf{E}'|^2 \rangle_c$ effectively lowers (if $K > 0$) or increases (if $K < 0$) the resting potential. We remark that this effective resting potential $\xi^{*\text{eff}}$ does not account for active processes of living cells. Biological processes, including ionic transport through channels and pumps, metabolic activities, and other cellular interactions, can significantly alter the electrical state of the cell.

Solid-to-fluid phase transition. It has been widely shown that cell aggregates may undergo a phase transition from a rigid solid-like phase to a floppy fluid-like phase, depending on the degree of “deviations” from the regular hexagonal tessellation [16, 17, 55, 57]. In the current setting, we interpret this transition as the critical point such that the small strain local minimizer (38) loses its stability, meaning that

$$\frac{\partial^2}{\partial \gamma^2} F^{\text{eff}}(\gamma, \bar{\xi}) \bigg|_{(0, \xi^{*\text{eff}})} = 0 \quad \Rightarrow \quad \mu_2^{\text{eff}} + K(\xi^{*\text{eff}} - \xi^*) = 0. \quad (39)$$

By (34) and (38), we find the *critical \mathbf{E}^2 -temperature* to be the one at which the cell aggregate at resting condition (i.e., no external shear stress), transits from a solid-like phase to a fluid-like phase as

$$\langle |\mathbf{E}'|^2 \rangle_{c,cr} = \frac{\gamma^*}{8} \left(1 + \frac{K^2 \lambda^2 \gamma^*}{8\mu\epsilon} \right)^{-1}. \quad (40)$$




Figure 3: Phase diagram for solid-to-fluid phase transition in cell aggregates. For chosen numerical values in this work, the range of non-dimensional x -axis corresponds to the range of change in electric potential as $(\bar{\xi} - \xi^*) \in [-200\text{mV}, 200\text{mV}]$ consistent with the physiological range; the y -axis represents the \mathbf{E}^2 -temperature $\langle |\mathbf{E}'|^2 \rangle_c$.

Bi *et al.* (2016 [73]) show that the solid-fluid phase transition occurs at the critical shape index $\hat{p}_{cr} = 3.81$ at the absence of electromechanical coupling. By (5) we may fix $\gamma^* \approx 2.0$ for consistency.

Moreover, living cells are capable of regulating their transmembrane potentials. At the presence of electromechanical coupling, tissues can actively regulate their stiffness and solid-to-fluid transition by manipulating their electric potentials. This capability allows them to adaptively respond to varying environmental conditions and physiological demands. Suppose that the average electric potential of a cell aggregate is fixed at $\bar{\xi}$ via certain active biophysical process. By (35) we find that the apparent shear modulus as

$$\mu^{\text{apparent}} = \frac{1}{4A} \frac{\partial^2}{\partial \gamma^2} F^{\text{eff}}(\gamma, \bar{\xi}) \Big|_{\gamma=0} = \mu \left(1 - \frac{8}{\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c \right) + K(\bar{\xi} - \xi^*). \quad (41)$$

The critical \mathbf{E}^2 -temperature (or fluctuation) for solid-fluid phase transition at this prescribed potential $\bar{\xi}$ is identified as that at which the apparent shear modulus μ^{apparent} of the cell aggregate vanishes:

$$\langle |\mathbf{E}'|^2 \rangle_c = \frac{\gamma^*}{8} \left(1 + \frac{K}{\mu} (\bar{\xi} - \xi^*) \right). \quad (42)$$

Figure 3 shows the phase diagram for solid-to-fluid phase transition governed by the coarse-grained model (35). The black solid line shows the phase boundary. The region below and above the phase boundary corresponds to the rigid solid-like phase and floppy fluid-like phase, respectively. At a fixed applied electric potential $\bar{\xi}$, an increase in \mathbf{E}^2 -temperature $\langle |\mathbf{E}'|^2 \rangle_c$ would transform the cell aggregate from a rigid solid-like phase to a floppy fluid-like phase. Similarly, at a prescribed \mathbf{E}^2 -temperature, an increase in electric potential $\bar{\xi}$ would transform the cell aggregate from a fluid-like phase to a solid-like phase.

Energy barrier. The effective free energy function (35) is anharmonic and admits multiple critical points in γ for a prescribed average potential $\bar{\xi}$. Graphically, as γ increases from 0, we observe that the first critical point ($\gamma = 0$) and the third critical point (γ_{T1}) are locally stable while the second critical point is unstable, reflecting the $T1$ -transition illustrated in Fig. 2. To find the energy barrier between the two stable equilibria, we consider the equations for critical points:

$$\frac{\partial}{\partial \gamma} F^{\text{eff}}(\gamma, \bar{\xi}) = 0 \quad \Rightarrow \quad \gamma = 0, \gamma^{\text{unstable}}, \gamma_{T1}.$$

By (35) we find that the unstable small-strain solution to the above equation is approximately given by

$$\gamma^{\text{unstable}} \approx \sqrt{\frac{(\mu_2^{\text{eff}} + K(\bar{\xi} - \xi^*))^2}{-4\mu_4^{\text{eff}}}}.$$

Therefore, by (35) the energy barrier between the stable equilibria $\gamma = 0$ and $\gamma = \gamma_{T1}$ is given by

$$\begin{aligned} U^{\text{barrier}} &= F^{\text{eff}}(\gamma, \bar{\xi}) \Big|_{\gamma=\gamma^{\text{unstable}}} - F^{\text{eff}}(\gamma, \bar{\xi}) \Big|_{\gamma=0} \approx A \frac{(\mu_2^{\text{eff}} + K(\bar{\xi} - \xi^*))^2}{-4\mu_4^{\text{eff}}} \\ &= \frac{\mu\gamma^* A}{8} \cdot \frac{\left(1 - \frac{8}{\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c + \frac{K}{\mu}(\bar{\xi} - \xi^*)\right)^2}{1 - \frac{9}{2\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c}. \end{aligned}$$

The energy barrier can be used to quantitatively assess the rate of $T1$ -transition. Following the classical Kramer's model [74], we may postulate the rate k of $T1$ -transition per unit volume as

$$k \propto \exp(-\beta U^{\text{barrier}}) = \exp\left(-\theta \frac{(\mu^{\text{apparent}}/\mu)^2}{\langle |\mathbf{E}'|^2 \rangle_c (1 - \frac{9}{2\gamma^*} \langle |\mathbf{E}'|^2 \rangle_c)}\right), \quad (43)$$

where the positive dimensionless constant $\theta = \frac{\mu\gamma^*(8\eta_c\mu_c - \kappa_c^2)}{16\mu_c(4\eta_c\mu_c - \kappa_c^2)} > 0$ follows from the last equality in (31). We remark that (43) is consistent with the physical interpretation of solid-to-fluid phase transitions determined by (41). In particular, the transition rate $k \ll 1$ if the fluctuation $\langle |\mathbf{E}'|^2 \rangle_c \ll 1$ whereas $k \sim 1$ if $\mu^{\text{apparent}} = 0$. Moreover, the rate of transition may be identified as the self-diffusivity which, by the Stokes-Einstein's relation, implies that the *viscosity* η of the cell aggregates should satisfy

$$\eta \propto \frac{1}{k} \propto \exp(\beta U^{\text{barrier}}).$$

Electromechanical responses. Finally, we consider the electromechanical properties of the cell aggregate in the solid phase. Suppose that the cell aggregate is under the application of external shear stress σ^e . By (35), we observe that the external shear stress can alter the average cross-membrane potential $\bar{\xi}$. Conversely, changes of the average transmembrane potential $\bar{\xi}$ can influence the overall shear strain of the cell aggregate. These interactions highlight a two-way coupling between mechanical and electrical responses within the system. Furthermore, both responses are significantly modulated by \mathbf{E}^2 -temperature (or fluctuations) of the system. For a fixed applied shear stress σ^e , average potential $\bar{\xi}$, and \mathbf{E}^2 -temperature, we obtain the equilibrium shear strain γ^{eq} by the principle of minimum free energy:

$$\gamma^{eq} = \text{argmin}_{\gamma} \left(\frac{1}{A} F^{\text{eff}}(\gamma, \bar{\xi}) - 2\sigma^e \gamma \right). \quad (44)$$

That is, γ^{eq} is the solution of the following nonlinear algebraic equation (Cf. (35)):

$$\left(\left(\mu_2^{\text{eff}} + K(\bar{\xi} - \xi^*) \right) 4\gamma + 16\mu_4^{\text{eff}}\gamma^3 + \frac{48\mu}{\gamma^{*2}}\gamma^5 - 2\sigma^e \right) = 0. \quad (45)$$

In general, the above nonlinear equation admits multiple solutions. We are only interested in the “small-strain” solution, which we numerically compute using MATLAB.




Figure 4: Macroscopic non-linear elastic response of cell aggregate for known fluctuation and applied electric potential $\bar{\xi} = \xi^{*\text{eff}}$.

We first consider the dependence of strain on the external stress under different \mathbf{E}^2 -temperature $\langle |\mathbf{E}'|^2 \rangle_c$. As an example, we set the average electric potential the same as the effective resting potential $\xi^{*\text{eff}}$ in (38): $\bar{\xi} = \xi^{*\text{eff}}$. Figure 4 shows the stress-strain relation of cell aggregate at different \mathbf{E}^2 -temperatures (or fluctuations) below the critical \mathbf{E}^2 -temperature in (40). We observe that the macroscopic elastic response of the cell aggregate is generally nonlinear and matches the typical range for cell aggregates [68]. We also notice that an increase in \mathbf{E}^2 -temperature (or fluctuation) softens the macroscopic elastic response, consistent with the effect of thermal temperature for classical crystalline solids.

Next, we consider the dependence of strain on the average cellular potential. Figure 5 shows the equilibrium shear strain γ^{eq} plotted against normalized applied electric potential $\frac{K(\bar{\xi} - \xi^*)}{\mu}$ for $\sigma^e = 0.02\mu$ and different \mathbf{E}^2 -temperatures $\langle |\mathbf{E}'|^2 \rangle_c$. We observe that as the average electric potential increases, the equilibrium shear strain decreases nonlinearly, showing a nonlinear piezoelectric response for cell aggregates. For a given average electric potential, the equilibrium shear strain is higher for higher \mathbf{E}^2 -temperature. We also observe that the changes in equilibrium shear strain due to changes in average electric potential are more significant at higher \mathbf{E}^2 -temperature. Physically, this implies that a softer cell aggregate possesses an enhanced electro-mechanical response, making its shape changes more significant and more sensitive to the changes in transmembrane electric potential. This observation is consistent with the solid-to-fluid phase transition for cell aggregate (Cf. Fig. 3) that for a given average electric potential, an increase in \mathbf{E}^2 -temperature (or fluctuation) brings a solid cell aggregate closer to the solid-to-fluid phase transition point.




Figure 5: Macroscopic inverse piezoelectric-like response of cell aggregate. Equilibrium shear strain γ^{eq} is plotted against normalized applied electric potential along x -axis. For chosen numerical values in this work, the range of non-dimensional x -axis in the plot corresponds to the range of change in electric potential as $(\bar{\xi} - \xi^*) \in [-200\text{mV}, 200\text{mV}]$ consistent with the physiological range.

5 Concluding remarks

In this work we first established the concept of strain tensor that characterizes the shape and orientation of individual cells and local arrangement of cells in amorphous cell aggregates. Based on the underlying structural symmetry and $T1$ -transition experimentally observed in tissues, we proposed an electromechanical model for single cells and their interactions. For collective behavior of cell aggregates, we employed the approach of statistical mechanics and achieved a coarse-grained model for macroscopic electromechanical properties of cell aggregates. Fluctuation in cellular strains can be precisely quantified, which is referred as the \mathbf{E}^2 -temperature for its analogous role as the absolute temperature in a conventional statistical model. The closed-form expression of the effective free energy enabled us to predict the macroscopic electromechanical properties of the cell aggregate. In particular, we derived the renormalized mechanical and electromechanical coupling coefficients and highlight how the average electric potential and \mathbf{E}^2 -temperature can independently regulate the electromechanical properties and solid-fluid phase transitions in cell aggregates. Our work resolves the controversy of whether soft biological tissues are piezoelectric. The answer is simple. They are not intrinsically piezoelectric since they lack an underlying polar structure however, due to the electrostriction (or Maxwell stress) behavior of a single cell, and the resting potential, the aggregate or collective behavior of cells can mimic piezoelectric-like behavior.

There are several possible future directions: (i) The developed theory for *living* cell aggregates could be extended to explicitly account for the active processes (e.g., effects of active protein forces) by using the principles of non-equilibrium statistical mechanics [50, 75, 76]. (ii) The derived effective free-energy may be used to explore the rich instability and bifurcation behavior in soft tissues c.f. [77, 78]. (iii) We have ignored the electromechanical coupling mechanism of flexoelectricity [6] or rather its possible emergence at the tissue scale. Prior work appears to indicate that flexoelectricity, at least at the cellular level, is implicated in a variety of biophysical phenomena such as the hearing

mechanism c.f. [79–81], and may also have consequences for energy extraction at the coarser level c.f. [82]. An approach similar to outlined in this work may be used to address this mechanism. (iv) The developed theory can be augmented to include the non-local inter-cellular interactions [83–87] by going beyond the nearest-neighbor interaction assumption used in this work.

Acknowledgment. L.L. gratefully acknowledges the support of NSF DMS-2306254 and Simons travel grant. P.S. gratefully acknowledges the support of the Hugh Roy and Lillie Cranz Cullen Distinguished University Chair from the University of Houston.

References

- [1] SA Safran, Nir Gov, A Nicolas, Ulrich S Schwarz, and Tsvi Tlusty. “Physics of cell elasticity, shape and adhesion”. In: *Physica A: Statistical Mechanics and its Applications* 352.1 (2005), pp. 171–201.
- [2] Bertil Hille. *Ion Channels of Excitable Membranes*. 3rd. Sunderland, MA: Sinauer Associates, 2001. ISBN: 0-87893-321-2.
- [3] Michael Levin. “Molecular bioelectricity: how endogenous voltage potentials control cell behavior and instruct pattern regulation in vivo”. In: *Molecular biology of the cell* 25.24 (2014), pp. 3835–3850.
- [4] Ido Nitsan, Stavit Drori, Yair E Lewis, Shlomi Cohen, and Shelly Tzlil. “Mechanical communication in cardiac cell synchronized beating”. In: *Nature Physics* 12.5 (2016), pp. 472–477.
- [5] Alexander D Bershadsky, Nathalie Q Balaban, and Benjamin Geiger. “Adhesion-dependent cell mechanosensitivity”. In: *Annual review of cell and developmental biology* 19.1 (2003), pp. 677–695.
- [6] Mehdi Torbati, Kosar Mozaffari, Liping Liu, and Pradeep Sharma. “Coupling of mechanical deformation and electromagnetic fields in biological cells”. In: *Reviews of Modern Physics* 94.2 (2022), p. 025003.
- [7] Thomas E Angelini, Edouard Hannezo, Xavier Trepat, Jeffrey J Fredberg, and David A Weitz. “Cell migration driven by cooperative substrate deformation patterns”. In: *Physical review letters* 104.16 (2010), p. 168104.
- [8] Xin Tang, Piyush Bajaj, Rashid Bashir, and Taher A Saif. “How far cardiac cells can see each other mechanically”. In: *Soft Matter* 7.13 (2011), pp. 6151–6158.
- [9] Shelly Tzlil and David A Tirrell. “Strain propagation in artificial extracellular matrix proteins can accelerate cell spreading and polarization”. In: *Soft Matter* 9.23 (2013), pp. 5602–5608.
- [10] Doron Grossman and Jean-Francois Joanny. “Instabilities and geometry of growing tissues”. In: *Physical Review Letters* 129.4 (2022), p. 048102.
- [11] Le Yan and Dapeng Bi. “Multicellular rosettes drive fluid-solid transition in epithelial tissues”. In: *Physical Review X* 9.1 (2019), p. 011029.
- [12] Thomas Heimburg and Andrew D Jackson. “On soliton propagation in biomembranes and nerves”. In: *Proceedings of the National Academy of Sciences* 102.28 (2005), pp. 9790–9795.
- [13] VF Antonov, E Yu Smirnova, and EV Shevchenko. “Electric field increases the phase transition temperature in the bilayer membrane of phosphatidic acid”. In: *Chemistry and physics of lipids* 52.3-4 (1990), pp. 251–257.

- [14] RMJ Cotterill. “Field effects on lipid membrane melting”. In: *Physica Scripta* 18.3 (1978), p. 191.
- [15] Tea Mužić, Fatma Tounsi, Søren B Madsen, Denis Pollakowski, Manfred Konrad, and Thomas Heimburg. “Melting transitions in biomembranes”. In: *Biochimica et Biophysica Acta (BBA)-Biomembranes* 1861.11 (2019), p. 183026.
- [16] Thomas Heimburg. “The capacitance and electromechanical coupling of lipid membranes close to transitions: the effect of electrostriction”. In: *Biophysical journal* 103.5 (2012), pp. 918–929.
- [17] Lars D Mosgaard, Karis A Zecchi, and Thomas Heimburg. “Mechano-capacitive properties of polarized membranes”. In: *Soft Matter* 11.40 (2015), pp. 7899–7910.
- [18] Daniel A Beysens, Gabor Forgacs, and James A Glazier. “Cell sorting is analogous to phase ordering in fluids”. In: *Proceedings of the National Academy of Sciences* 97.17 (2000), pp. 9467–9471.
- [19] MS Steinberg. “Liquid behavior of embryonic tissues”. In: *Cell behavior* (1982).
- [20] Gabor Forgacs, Ramsey A Foty, Yinon Shafrir, and Malcolm S Steinberg. “Viscoelastic properties of living embryonic tissues: a quantitative study”. In: *Biophysical journal* 74.5 (1998), pp. 2227–2234.
- [21] HM Phillips, MS Steinberg, and BH Lipton. “Embryonic tissues as elasticoviscous liquids”. In: *Developmental Biology* 59.2 (1977), pp. 124–134.
- [22] Edouard Hannezo, Jacques Prost, and Jean-Francois Joanny. “Theory of epithelial sheet morphology in three dimensions”. In: *Proceedings of the National Academy of Sciences* 111.1 (2014), pp. 27–32.
- [23] Nebojsa Murisic, Vincent Hakim, Ioannis G Kevrekidis, Stanislav Y Shvartsman, and Basile Audoly. “From discrete to continuum models of three-dimensional deformations in epithelial sheets”. In: *Biophysical journal* 109.1 (2015), pp. 154–163.
- [24] Javier Cervera, Jose Antonio Manzanares, Salvador Mafe, and Michael Levin. “Synchronization of bioelectric oscillations in networks of nonexcitable cells: from single-cell to multicellular states”. In: *The Journal of Physical Chemistry B* 123.18 (2019), pp. 3924–3934.
- [25] Eiichi Fukada. “Piezoelectricity of biopolymers”. In: *Biorheology* 32.6 (1995), pp. 593–609.
- [26] Yuanming Liu, Hong-Ling Cai, Matthew Zelisko, Yunjie Wang, Jinglan Sun, Fei Yan, Feiyue Ma, Peiqi Wang, Qian Nataly Chen, Hairong Zheng, et al. “Ferroelectric switching of elastin”. In: *Proceedings of the National Academy of Sciences* 111.27 (2014), E2780–E2786.
- [27] Thomas Lenz, Regina Hummel, Ilias Katsouras, Wilhelm A Groen, Marlies Nijemeisland, Robert Ruemmler, Michael KE Schäfer, and Dago M De Leeuw. “Ferroelectricity and piezoelectricity in soft biological tissue: Porcine aortic walls revisited”. In: *Applied Physics Letters* 111.13 (2017).
- [28] Kenji Ikushima, Takashi Kumamoto, Kenshiro Ito, and Yamato Anzai. “Electric polarization of soft biological tissues induced by ultrasound waves”. In: *Physical review letters* 123.23 (2019), p. 238101.
- [29] Dirk Drasdo and Stefan Höhme. “A single-cell-based model of tumor growth in vitro: monolayers and spheroids”. In: *Physical biology* 2.3 (2005), p. 133.
- [30] Juliette Griffié, Ruby Peters, and Dylan M Owen. “An agent-based model of molecular aggregation at the cell membrane”. In: *Plos one* 15.2 (2020), e0226825.

- [31] Alexander RA Anderson, Mark AJ Chaplain, E Luke Newman, Robert JC Steele, and Alastair M Thompson. “Mathematical modelling of tumour invasion and metastasis”. In: *Computational and mathematical methods in medicine* 2.2 (2000), pp. 129–154.
- [32] M Block, E Schöll, and D Drasdo. “Classifying the expansion kinetics and critical surface dynamics of growing cell populations”. In: *Physical review letters* 99.24 (2007), p. 248101.
- [33] Joana Moreira and Andreas Deutsch. “Cellular automaton models of tumor development: a critical review”. In: *Advances in Complex Systems* 5.02n03 (2002), pp. 247–267.
- [34] Mark S Alber, Maria A Kiskowski, James A Glazier, and Yi Jiang. “On cellular automaton approaches to modeling biological cells”. In: *Mathematical systems theory in biology, communications, computation, and finance*. Springer. 2003, pp. 1–39.
- [35] Tomás Alarcón, Helen M Byrne, and Philip K Maini. “A mathematical model of the effects of hypoxia on the cell-cycle of normal and cancer cells”. In: *Journal of theoretical biology* 229.3 (2004), pp. 395–411.
- [36] Andrew C Callan-Jones and Frank Jülicher. “Hydrodynamics of active permeating gels”. In: *New Journal of Physics* 13.9 (2011), p. 093027.
- [37] Jonas Ranft, Markus Basan, Jens Elgeti, Jean-François Joanny, Jacques Prost, and Frank Jülicher. “Fluidization of tissues by cell division and apoptosis”. In: *Proceedings of the National Academy of Sciences* 107.49 (2010), pp. 20863–20868.
- [38] Jacques Prost, Frank Jülicher, and Jean-François Joanny. “Active gel physics”. In: *Nature physics* 11.2 (2015), pp. 111–117.
- [39] Michael H Köpf and Len M Pismen. “A continuum model of epithelial spreading”. In: *Soft matter* 9.14 (2013), pp. 3727–3734.
- [40] Julia C Arciero, Qi Mi, Maria F Branca, David J Hackam, and David Swigon. “Continuum model of collective cell migration in wound healing and colony expansion”. In: *Biophysical journal* 100.3 (2011), pp. 535–543.
- [41] Shuji Ishihara, Philippe Marcq, and Kaoru Sugimura. “From cells to tissue: A continuum model of epithelial mechanics”. In: *Physical Review E* 96.2 (2017), p. 022418.
- [42] Soheil Firooz, Stefan Kaessmair, Vasily Zaburdaev, Ali Javili, and Paul Steinmann. “On continuum modeling of cell aggregation phenomena”. In: *Journal of the Mechanics and Physics of Solids* 167 (2022), p. 105004.
- [43] Nicola J Armstrong, Kevin J Painter, and Jonathan A Sherratt. “A continuum approach to modelling cell–cell adhesion”. In: *Journal of theoretical biology* 243.1 (2006), pp. 98–113.
- [44] Alf Gerisch and Mark AJ Chaplain. “Mathematical modelling of cancer cell invasion of tissue: local and non-local models and the effect of adhesion”. In: *Journal of Theoretical Biology* 250.4 (2008), pp. 684–704.
- [45] Ignacio Ramis-Conde, Mark AJ Chaplain, and Alexander RA Anderson. “Mathematical modelling of cancer cell invasion of tissue”. In: *Mathematical and Computer Modelling* 47.5-6 (2008), pp. 533–545.
- [46] Frederik W Wiegel and Alan S Perelson. “Statistical mechanics of red blood cell aggregation: the distribution of rouleaux in thermal equilibrium”. In: *Journal of statistical physics* 29 (1982), pp. 813–848.
- [47] Andrew E Teschendorff and Andrew P Feinberg. “Statistical mechanics meets single-cell biology”. In: *Nature Reviews Genetics* 22.7 (2021), pp. 459–476.

[48] Nadia Loy and Luigi Preziosi. “A Statistical Mechanics Approach to Describe Cell Reorientation Under Stretch”. In: *Bulletin of Mathematical Biology* 85.7 (2023), p. 60.

[49] Christoffer Åberg, Juan A Varela, Laurence W Fitzpatrick, and Kenneth A Dawson. “Spatial and structural metrics for living cells inspired by statistical mechanics”. In: *Scientific reports* 6.1 (2016), p. 34457.

[50] Yashashree Kulkarni. “Fluctuations of active membranes with nonlinear curvature elasticity”. In: *Journal of the Mechanics and Physics of Solids* 173 (2023), p. 105240.

[51] Sreekanth Ramesh and Yashashree Kulkarni. “Statistical mechanics of active vesicles and the size distribution paradox”. In: *Journal of the Mechanics and Physics of Solids* 191 (2024), p. 105749.

[52] Anand Mathew and Yashashree Kulkarni. “Active matter as the underpinning agency for extraordinary sensitivity of biological membranes to electric fields”. In: *arXiv preprint arXiv:2412.16319* (2024).

[53] Étienne Fodor and M Cristina Marchetti. “The statistical physics of active matter: From self-catalytic colloids to living cells”. In: *Physica A: Statistical Mechanics and its Applications* 504 (2018), pp. 106–120.

[54] LP Liu and P Sharma. “Flexoelectricity and thermal fluctuations of lipid bilayer membranes: Renormalization of flexoelectric, dielectric, and elastic properties”. In: *Physical Review E* 87.3 (2013), p. 032715.

[55] R Li, C Ibar, Z Zhou, S Moazzeni, AN Norris, KD Irvine, L Liu, and H Lin. “ \mathbf{E}^2 and Gamma Distributions in Polygonal Networks”. In: *Physical Review Research* 3.4 (2021), p. L042001.

[56] Martin Bock, Ajay K. Tyagi, Jan-Ulrich Kreft, and Wolfgang Alt. “Generalized Voronoi Tessellation as a Model of Two-dimensional Cell Tissue Dynamics”. In: *Bulletin of Mathematical Biology* 72 (2010), pp. 1696–1731.

[57] L Atia, D Bi, Y Sharma, JA Mitchel, B Gweon, SA Koehler, SJ DeCamp, B Lan, JH Kim, R Hirsch, AF Pegoraro, KH Lee, JR Starr, DA Weitz, AC Martin, J-A Park, JP Butler, and JJ Fredberg. “Geometrical constraints during epithelial jamming”. In: *Nature Physics* 14 (2018), p. 613.

[58] R Li, S Moazzeni, L Liu, and H Lin. “Micro and Macroscopic Stress-Strain Relations in Disordered Tessellated Networks”. In: *Physical Review Letters* 130.18 (2023), p. 188201.

[59] L. D. Landau. “On the Theory of Phase Transitions”. In: *Phy. Z. Sow.* 11 (1937), p. 26.

[60] L. D. Landau and E. M. Lifshitz. *Statistical Physics (3rd ed.)* Pergamon Press, 1999.

[61] Pratik Khandagale, Liping Liu, and Pradeep Sharma. “Statistical mechanics of plasticity: Elucidating anomalous size-effects and emergent fractional nonlocal continuum behavior”. In: *Journal of the Mechanics and Physics of Solids* 191 (2024), p. 105747.

[62] S. Ogawa, A. Umemura, and N. Oshima. “On the equations of fully fluidized granular materials”. In: *Journal of Applied Mathematics and Physics (ZAMP)* 31 (1980), pp. 483–493. DOI: [10.1007/BF01590859](https://doi.org/10.1007/BF01590859).

[63] N. N. Jr. Bogolyubov. “On model dynamical systems in statistical mechanics”. In: *Physica* 32 (1966), pp. 933–944.

[64] Arthur Beyder and Frederick Sachs. “Electromechanical coupling in the membranes of Shaker-transfected HEK cells”. In: *Proceedings of the National Academy of Sciences* 106.16 (2009), pp. 6626–6631.

[65] S Krichen, Liping Liu, and P Sharma. “Biological cell as a soft magnetoelectric material: Elucidating the physical mechanisms underpinning the detection of magnetic fields by animals”. In: *Physical Review E* 96.4 (2017), p. 042404.

[66] Ilya Levental, Penelope C Georges, and Paul A Janmey. “Soft biological materials and their impact on cell function”. In: *Soft Matter* 3.3 (2007), pp. 299–306.

[67] Shinichiro Hiratsuka, Yusuke Mizutani, Masahiro Tsuchiya, Koichi Kawahara, Hiroshi Tukamoto, and Takaharu Okajima. “The number distribution of complex shear modulus of single cells measured by atomic force microscopy”. In: *Ultramicroscopy* 109.8 (2009), pp. 937–941.

[68] Junxiang Huang, James O Cochran, Suzanne M Fielding, M Cristina Marchetti, and Dapeng Bi. “Shear-driven solidification and nonlinear elasticity in epithelial tissues”. In: *Physical Review Letters* 128.17 (2022), p. 178001.

[69] Dapeng Bi, JH Lopez, Jennifer M Schwarz, and M Lisa Manning. “A density-independent rigidity transition in biological tissues”. In: *Nature Physics* 11.12 (2015), pp. 1074–1079.

[70] Edwin T Jaynes. “Information theory and statistical mechanics”. In: *Physical review* 106.4 (1957), p. 620.

[71] Sam F Edwards and RBS Oakeshott. “Theory of powders”. In: *Physica A: Statistical Mechanics and its Applications* 157.3 (1989), pp. 1080–1090.

[72] Alexander S Balankin and Leonardo Flores-Cano. “Edwards’s statistical mechanics of crumpling networks in crushed self-avoiding sheets with finite bending rigidity”. In: *Physical Review E* 91.3 (2015), p. 032109.

[73] Dapeng Bi, Xingbo Yang, M Cristina Marchetti, and M Lisa Manning. “Motility-driven glass and jamming transitions in biological tissues”. In: *Physical Review X* 6.2 (2016), p. 021011.

[74] H. A. Kramers. “Brownian motion in a field of force and the diffusion model of chemical reactions”. In: *Physica* 7 (1940), pp. 284–304.

[75] Travis Leadbetter, Prashant K Purohit, and Celia Reina. “A statistical mechanics framework for constructing nonequilibrium thermodynamic models”. In: *PNAS Nexus* 2.12 (2023), pgad417.

[76] Shenglin Huang, Ian R Graham, Robert A Riddleman, Paulo E Arratia, Steve Fitzgerald, and Celia Reina. “Predicting the unobserved: A statistical mechanics framework for non-equilibrium material response with quantified uncertainty”. In: *Journal of the Mechanics and Physics of Solids* 161 (2022), p. 104779.

[77] Shengyou Yang and Pradeep Sharma. “A tutorial on the stability and bifurcation analysis of the electromechanical behaviour of soft materials”. In: *Applied Mechanics Reviews* 75.4 (2023), p. 044801.

[78] Shengyou Yang, Xuanhe Zhao, and Pradeep Sharma. “Revisiting the instability and bifurcation behavior of soft dielectrics”. In: *Journal of Applied Mechanics* 84.3 (2017), p. 031008.

[79] Kathryn D Breneman, William E Brownell, and Richard D Rabbitt. “Hair cell bundles: flexoelectric motors of the inner ear”. In: *PLoS One* 4.4 (2009), e5201.

[80] Kosar Mozaffari, Fatemeh Ahmadpoor, Qian Deng, and Pradeep Sharma. “A minimal physics-based model for musical perception”. In: *Proceedings of the National Academy of Sciences* 120.5 (2023), e2216146120.

- [81] Qian Deng, Fatemeh Ahmadpoor, William E Brownell, and Pradeep Sharma. “The collusion of flexoelectricity and Hopf bifurcation in the hearing mechanism”. In: *Journal of the Mechanics and Physics of Solids* 130 (2019), pp. 245–261.
- [82] Binglei Wang, Shengyou Yang, and Pradeep Sharma. “Flexoelectricity as a universal mechanism for energy harvesting from crumpling of thin sheets”. In: *Physical Review B* 100.3 (2019), p. 035438.
- [83] Jason Marshall and Kaushik Dayal. “Atomistic-to-continuum multiscale modeling with long-range electrostatic interactions in ionic solids”. In: *Journal of the Mechanics and Physics of Solids* 62 (2014), pp. 137–162.
- [84] Pratik Khandagale, Timothy Breitzman, Carmel Majidi, and Kaushik Dayal. “Statistical field theory for nonlinear elasticity of polymer networks with excluded volume interactions”. In: *Physical Review E* 107.6 (2023), p. 064501.
- [85] VI Abkevich, AM Gutin, and EI Shakhnovich. “Impact of local and non-local interactions on thermodynamics and kinetics of protein folding”. In: *Journal of molecular biology* 252.4 (1995), pp. 460–471.
- [86] Roger H French, V Adrian Parsegian, Rudolf Podgornik, Rick F Rajter, Anand Jagota, Jian Luo, Dilip Asthagiri, Manoj K Chaudhury, Yet-ming Chiang, Steve Granick, et al. “Long range interactions in nanoscale science”. In: *Reviews of Modern Physics* 82.2 (2010), p. 1887.
- [87] Alessandro Campa, Thierry Dauxois, and Stefano Ruffo. “Statistical mechanics and dynamics of solvable models with long-range interactions”. In: *Physics Reports* 480.3-6 (2009), pp. 57–159.