



Contents lists available at ScienceDirect

# European Journal of Combinatorics

journal homepage: <https://www.sciencedirect.com/journal/european-journal-of-combinatorics>



## Linear extensions and continued fractions

Swee Hong Chan <sup>a</sup>, Igor Pak <sup>b</sup>



<sup>a</sup> Department of Mathematics, Rutgers University, Piscataway, NJ 08854, United States of America

<sup>b</sup> Department of Mathematics, UCLA, Los Angeles, CA 90095, United States of America

### ARTICLE INFO

#### Article history:

Received 16 April 2024

Accepted 11 June 2024

Available online 2 July 2024

### ABSTRACT

We introduce several new constructions of finite posets with the number of linear extensions given by generalized continued fractions. We apply our results to the problem of the minimum number of elements needed for a poset with a given number of linear extensions.

© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

## 1. Introduction

### 1.1. Foreword

Continued fractions go back to antiquity [5] and are surprisingly versatile. They appear across mathematics, from number theory [3,28] to analysis [18,21], from cluster algebras [7] to discrete geometry [20] to signal processing [31]. In combinatorics, they famously enumerate partitions [2], lattice paths [12] (see also [13,14,27]), permutations [11,36], and perfect matchings [42] (see also [32,38]).

Curiously, the applications go in both directions: the asymptotics of combinatorial sequences can be derived from analytic properties of continued functions, while combinatorial interpretations imply positivity properties. This paper explores connections between linear extensions of finite posets and continued fractions, and their asymptotic applications to counting.

Note that we utilize standard terminologies in order theory and continued fraction theory, and we include detailed definitions in Section 2 for the reader's convenience and clarity.

E-mail addresses: [sc2518@rutgers.edu](mailto:sc2518@rutgers.edu) (S.H. Chan), [pak@math.ucla.edu](mailto:pak@math.ucla.edu) (I. Pak).

<https://doi.org/10.1016/j.ejc.2024.104018>

0195-6698/© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

## 1.2. Linear extensions

Let  $P = (X, \prec)$  be a poset with  $|X| = n$  elements. Denote  $[n] := \{1, \dots, n\}$ . A *linear extension* of  $P$  is a bijection  $f : X \rightarrow [n]$ , such that  $f(x) < f(y)$  for all  $x \prec y$ . Let  $\mathcal{E}(P)$  be the set of linear extensions of  $P$ , and denote  $e(P) := |\mathcal{E}(P)|$ . Clearly,  $1 \leq e(P) \leq n!$  See [10] for a detailed recent survey.

Denote by  $\mu(n)$  the minimum number of elements in a poset with  $n$  linear extensions. See [35, A160371] for the numerical data (see also [35, A281723]). For example,  $\mu(5) = 4$  since  $e(Z_4) = 5$ , where  $Z_4$  is a *zigzag poset* on 4 elements (with an  $N$ -shaped comparability graph).

The asymptotics of  $\{\mu(n)\}$  remains an important open problem. Clearly,  $\mu(n) \leq n$  since for the parallel sum of chains we have:  $e(C_{n-1} \oplus C_1) = n$ . In a different direction,  $\mu(n) = \Omega(\log n / \log \log n)$  since  $e(P) \leq n!$ : The first nontrivial upper bound  $\mu(n) = O(\sqrt{n})$  was found by Tenner [40]. Most recently, this bound was greatly improved:

**Theorem 1.1** (Kravitz–Sah [23, Thm 1.1]). *We have:  $\mu(n) = O(\log n \log \log n)$ .*

The authors use a simple but surprising connection to continued fractions, the starting point of this paper (see below). They state the following:

**Conjecture 1.2** ([23, Conj. 7.3]). *We have:  $\mu(n) = O(\log n)$ .*

In this paper, we are mostly interested in the combinatorial aspects of the connection between linear extensions and continued fractions, suggesting new technical tools towards the conjecture.

## 1.3. Simple continued fractions

Let  $\mathbb{N} := \{0, 1, 2, \dots\}$  and  $\mathbb{P} := \{1, 2, \dots\}$ . A *simple continued fraction* (CF) is defined as follows:

$$[b_0, b_1, b_2, \dots, b_m] := b_0 + \cfrac{1}{b_1 + \cfrac{1}{b_2 + \cfrac{1}{\ddots + \cfrac{1}{b_m}}}}, \quad (1.1)$$

where integers  $b_0 \geq 0, b_1, \dots, b_{m-1} \geq 1$ , and  $b_m \geq 2$  for  $m \geq 1$ . Integers  $b_i$  are called *quotients*. The sum of these quotients  $S(b_0, \dots, b_m) := b_0 + \dots + b_m$  is called the *weight* of  $[b_0, \dots, b_m]$ . Recall that for every  $\alpha \in \mathbb{Q}_{\geq 0}$  there is a unique simple continued fraction  $[b_0, b_1, b_2, \dots, b_m] = \alpha$ , and we write  $s(\alpha) := S(b_0, b_1, b_2, \dots, b_m)$  in this case. Note that  $s(\alpha) = s(\alpha^{-1})$ .

In the terminology of [43] (see also [22, §4.5.3]), the weight  $s\left(\frac{c}{d}\right)$  is the number of steps of the *subtraction algorithm*, the original (classical) version of the Euclidean algorithm for finding the greatest common divisor that uses only subtractions instead of divisions. The following result is the key to the proof of [Theorem 1.1](#).

**Theorem 1.3** (Larcher [24], See Also [23, Thm 1.2]). *For every integer  $d \geq 1$ , there exists an integer  $1 \leq c < d$ ,  $\gcd(c, d) = 1$ , such that*

$$s\left(\frac{c}{d}\right) \leq C \frac{d}{\phi(d)} \log d \log \log d, \quad (1.2)$$

where  $\phi(n)$  is Euler's totient function, and  $C > 0$  is a universal constant.

See Section 5.3 for more on the theorem. Now, Kravitz and Sah observed that [Conjecture 1.2](#) follows from the following conjectural extension of [Theorem 1.3](#).

**Conjecture 1.4** ([23, Conj. 7.2]). *For every prime  $d$ , there is an integer  $1 \leq c < d$ , such that*

$$s\left(\frac{c}{d}\right) \leq C \log d, \quad (1.3)$$

where  $C > 0$  is a universal constant.

Note that in a CF (1.1) for  $\frac{c}{d}$ , the number of quotients is  $m = O(\log d)$ . Thus, [Conjecture 1.4](#) follows from the celebrated *Zaremba's conjecture* (see also [Section 5.5](#)):

**Conjecture 1.5** (Zaremba [44, p. 76]). *For every integer  $d \geq 1$ , there is an integer  $1 \leq c < d$ , such that  $c/d = [0, b_1, \dots, b_m]$  and  $b_1, \dots, b_m \leq A$ , where  $A > 0$  is a universal constant.*

#### 1.4. From continued fractions to linear extensions

In poset  $P = (X, \prec)$ , an *antichain* is a subset of pairwise independent elements. The *width* of a poset is the size of the maximal antichain. An element  $x \in X$  is *minimal*, if for every  $y \in X$  we have either  $x \preccurlyeq y$  or  $x \parallel y$ . Denote by  $\min(P)$  the set of all minimal elements in  $P$ . Denote by  $P - x$  the poset obtained by removing the element  $x$ .

**Theorem 1.6** (See [23, Prop. 4.1]). *For all integers  $1 \leq c < d$  with  $\gcd(c, d) = 1$ , there is a poset  $P = (X, \prec)$  of width two, such that  $|X| = s\left(\frac{c}{d}\right)$ ,  $e(P) = d$  and  $e(P - x) = c$  for some minimal element  $x \in \min(P)$ .*

The proof of the theorem uses two simple transformations of posets  $(P, x) \rightarrow (P', x')$  and  $(P'', x'')$ , such that for  $e(P) = d$ ,  $e(P - x) = c$  the new posets satisfy  $e(P') = e(P'') = c + d$ ,  $e(P' - x') = c$ ,  $e(P'' - x'') = d - c$ . In [Section 3](#) we modify and generalize this construction.

Before we proceed to generalizations, consider

$$\mathcal{T}(k) := \{e(P) : P = (X, \prec), |X| \leq k\},$$

so that  $\mu(n) = \min\{k : n \in \mathcal{T}(k)\}$ . Open Problems 7.5 and 7.6 in [23] ask about the asymptotics of  $|\mathcal{T}(k)|$ , and of the largest  $L = L_c(k)$  such that  $|\mathcal{T}(k) \cap \{1, \dots, L\}| > cL$ . We have the following direct application of [Theorem 1.6](#) (not noticed in [23]), which gives partial answers to both open problems:

**Corollary 1.7.** *We have:  $|\mathcal{T}(k)| = \exp \mathcal{Q}(k)$ . Moreover, there is a constant  $C > 1$ , such that*

$$\frac{1}{C^k} |\mathcal{T}(k) \cap \{1, 2, \dots, \lfloor C^k \rfloor\}| \rightarrow 1 \quad \text{as } k \rightarrow \infty. \quad (1.4)$$

**Proof.** Recall the following remarkable result of Bourgain and Kontorovich [4] (see also [Section 5.5](#)), giving an asymptotic version of Zaremba's [Conjecture 1.5](#):  $\gamma(n) \rightarrow 1$  as  $n \rightarrow \infty$ , where  $\gamma(n)$  denotes the proportion of  $d \in \{1, \dots, n\}$ , such that  $c/d$  has all quotients  $\leq 50$  for some  $1 \leq c < d$ ,  $\gcd(c, d) = 1$ . Since  $s\left(\frac{c}{d}\right) = O(\log d)$  for such fractions, by [Theorem 1.6](#) we obtain the result.  $\square$

#### 1.5. Relative version

Let  $P = (X, \prec)$  and let  $x \in X$ . Following [9], consider the *relative number of linear extensions*:

$$\rho(P, x) := \frac{e(P)}{e(P - x)}.$$

It follows from [Theorem 1.6](#), that every rational number  $\alpha \geq 1$  is equal to  $\rho(P, x)$  for some poset  $P$  and element  $x \in X$ .

For  $d \geq c \geq 1$ , let  $\nu(c, d)$  denote the minimal number of elements in a poset  $P = (X, \prec)$ , such that  $\rho(P, x) = \frac{d}{c}$  for some  $x \in X$ . The following upper bound can be viewed as a relative version of [Theorem 1.1](#).

**Theorem 1.8.** *For all  $d \geq 3c$ , we have:*

$$\nu(c, d) \leq \frac{d}{c} + O(\log d \log \log d). \quad (1.5)$$

In [9, Prop. 8.8], we showed an asymptotically matching lower bound:

$$v(c, d) \geq \frac{d}{c}. \quad (1.6)$$

The motivation of **Theorem 1.8** comes from the approach in [8], where we studied relative versions of several counting functions (domino tilings, spanning trees, etc.) The proof of **Theorem 1.8** is based on the approach in [9, §8.2]. It would be interesting to see if the condition  $d \geq 3c$  can be weakened to  $d \geq (1 + \varepsilon)c$  or even dropped. Additionally, by analogy with the Kravitz–Sah **Conjecture 1.2**, we conjecture that (1.5) can be improved to

$$v(c, d) \leq \frac{d}{c} + O(\log d). \quad (1.7)$$

In a different direction, one can ask about the smallest size poset with  $e(P) = d$  and  $e(P - x) = c$ , since the construction in the proof can result in an integer multiple of both.

The key part of the proof of **Theorem 1.8** is the following tail estimate for the weight of random continued fractions:

**Theorem 1.9** (Rukavishnikova [30]). *There is a universal constant  $C > 0$ , such that*

$$\frac{1}{d} \# \left\{ c \in [d] : \left| s\left(\frac{c}{d}\right) - \frac{12}{\pi^2} \log d \log \log d \right| > (\log d)(\log \log d)^{2/3} \right\} < \frac{C}{(\log \log d)^{1/3}}. \quad (1.8)$$

Here we are stating a special case of the main theorem in [30] which suffices for our purposes.

### 1.6. Generalized continued fractions

Let  $m \geq 0$ ,  $a_1, \dots, a_m \in \mathbb{P}$ ,  $b_0, \dots, b_m \in \mathbb{P}$ . A *generalized continued fraction* (GCF) is defined as

$$[a_1, \dots, a_m; b_0, \dots, b_m] := b_0 + \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots + \frac{a_m}{b_m}}}}. \quad (1.9)$$

Note that when  $a_1 = \dots = a_m = 1$  we get a simple continued fraction. We define the *weight* of GCFs as follows:

$$G(a_1, \dots, a_m; b_0, \dots, b_m) := (b_0 + \dots + b_m) - (a_1 + \dots + a_m) + m,$$

and note that  $G(1, \dots, 1; b_0, \dots, b_m) = S(b_0, \dots, b_m)$ . Observe that a rational number can have many presentations as a GCF, some of which can have weight smaller than the weight of the corresponding CFs. For example,

$$\frac{20}{7} = 2 + \frac{1}{1 + \frac{1}{6}} = 2 + \frac{2}{2 + \frac{1}{3}},$$

so  $S\left(\frac{20}{7}\right) = S(2, 1, 6) = 9$  and  $G(2, 1; 2, 2, 3) = 6$ .

A generalized continued fraction (1.9) is called *balanced* if

$$b_i \geq a_i + a_{i+1} - 1 \quad \text{for all } 0 \leq i \leq m, \quad (1.10)$$

where by convention we assume that  $a_0 = a_{m+1} = 1$ . Clearly, every simple continued fraction of  $\alpha \in \mathbb{Q}_{\geq 1}$  is balanced. The following is the GCF analogue of **Theorem 1.6**.

**Theorem 1.10.** *Let  $m \geq 0$ ,  $a_1, \dots, a_m \in \mathbb{P}$ ,  $b_0, \dots, b_m \in \mathbb{P}$  be integers satisfying (1.10). Then there exists a poset  $P = (X, \prec)$  of width at most three, and a minimal element  $x \in \min(P)$ , such that  $|X| = G(a_1, \dots, a_m; b_0, \dots, b_m)$ , and*

$$[a_1, \dots, a_m; b_0, \dots, b_m] = \rho(P, x),$$

where  $[a_1, \dots, a_m; b_0, \dots, b_m]$  is a balanced GCF defined in (1.9).

For  $\alpha \in \mathbb{Q}_{\geq 1}$ , define

$$g(\alpha) := \min \{ G(a_1, \dots, a_m; b_0, \dots, b_m) : [a_1, \dots, a_m; b_0, \dots, b_m] = \alpha \},$$

where the minimum is over all balanced GCF (1.10) such that all partial fractions  $\frac{c_i}{D_i}$  are reduced, i.e.  $\gcd(c_i, D_i) = 1$  for all  $1 \leq i \leq m$  (see the definition in Section 2.2). For example, if  $a_1 = \dots = a_m = r$  for some integer  $r \geq 1$ , and integers  $b_1, \dots, b_m$  are coprime to  $r$ , then all partial fractions  $\frac{c_i}{D_i}$  are reduced. In particular, this condition automatically holds for all simple CFs. From above, we have  $g(\alpha) \leq s(\alpha)$ . Thus, the following conjecture is a natural weakening of Conjecture 1.4.

**Conjecture 1.11.** For every prime  $d$ , there is an integer  $1 \leq c < d$ , such that

$$g\left(\frac{d}{c}\right) \leq C \log d, \quad (1.11)$$

where  $C > 0$  is a universal constant.

From Theorem 1.10, we have:

**Proposition 1.12.** Conjecture 1.11 implies Conjecture 1.2.

### 1.7. Rational GCFs

We call a continued fraction of the form (1.9) *rational* if  $a_i \in \mathbb{Q}_{\geq 1}$ . A rational generalized continued fraction (RGCF) is called *balanced* if it is of the form

$$b_0 + \alpha_1 + \frac{\alpha_1}{s(\alpha_1) - 1 + b_1 + \alpha_2 + \frac{\alpha_2}{s(\alpha_2) - 1 + b_2 + \alpha_3 + \frac{\alpha_3}{\ddots + \frac{\alpha_m}{s(\alpha_m) - 1 + b_m}}}}, \quad (1.12)$$

where  $\alpha_1, \dots, \alpha_m \in \mathbb{Q}_{\geq 1}$  and  $b_0, \dots, b_m \in \mathbb{N}$  s.t.  $b_m \geq 1$ . We use  $[\alpha_1, \dots, \alpha_m; b_0, \dots, b_m]$  to denote this RGCF.

Note that for  $\alpha_1, \dots, \alpha_m \in \mathbb{P}$ , this is a balanced GCF, since the inequalities (1.10) are automatically satisfied. Denote by

$$R(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m) := b_0 + \dots + b_m + s(\alpha_1) + \dots + s(\alpha_m)$$

the *weight* of (1.12). For example, take  $m = 1, \alpha_1 = \frac{3}{2}, b_0 = 1, b_1 = 3$ . Then

$$s\left(\frac{3}{2}\right) = 3, \quad \left[\frac{3}{2}; 1, 3\right] = 1 + \frac{3}{2} + \frac{\frac{3}{2}}{s\left(\frac{3}{2}\right) - 1 + 3} = \frac{14}{5} \quad \text{and} \quad R\left(\frac{3}{2}; 1, 3\right) = 1 + 3 + s\left(\frac{3}{2}\right) = 7.$$

The following result is a variation of Theorem 1.10 to RGCF:

**Theorem 1.13.** Let  $m \geq 0, \alpha_1, \dots, \alpha_m \in \mathbb{Q}_{\geq 1}$  and  $b_0, \dots, b_m \in \mathbb{P}$ . Then there exists a poset  $P = (X, \prec)$  of width at most three, and a minimal element  $x \in \min(P)$ , such that  $|X| = R(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m)$ , and

$$[\alpha_1, \dots, \alpha_m; b_0, \dots, b_m] = \rho(P, x),$$

where  $[\alpha_1, \dots, \alpha_m; b_0, \dots, b_m]$  is a balanced RGCF defined in (1.12).

For  $\beta \in \mathbb{Q}_{\geq 1}$ , define

$$r(\beta) := \min \{ R(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m) : [\alpha_1, \dots, \alpha_m; b_0, \dots, b_m] = \beta \},$$

where the minimum is over all RGCF (1.12) such that all partial fractions  $\frac{c_i}{D_i}$  are reduced (see the definition in Section 2.2). From above,  $r(\alpha) \leq s(\alpha)$ . Thus, the following conjecture is a natural weakening of both Conjectures 1.4 and 1.11.

**Conjecture 1.14.** For every prime  $d$ , there is an integer  $1 \leq c < d$ , such that

$$r\left(\frac{d}{c}\right) \leq C \log d, \quad (1.13)$$

where  $C > 0$  is a universal constant.

To motivate the conjecture, note that  $r(\beta)$  can be much smaller than  $s(\beta)$ . Take, for example,  $m = 1$ ,  $\alpha_1 = \frac{13}{7}$  and  $\beta = \frac{173}{56}$ . We have:

$$\alpha_1 = 1 + \frac{1}{1 + \frac{1}{6}}, \quad s(\alpha_1) = 8, \quad \beta = 3 + \frac{1}{11 + \frac{1}{5}} = 1 + \alpha_1 + \frac{\alpha_1}{s(\alpha_1) - 1 + 1} = [\alpha_1; 1, 1].$$

Thus,  $s(\beta) = g(\beta) = 19$  while  $r(\beta) \leq R(\alpha_1; 1, 1) = 10$  in this case. Again, by Theorem 1.13 we have:

**Proposition 1.15.** Conjecture 1.14 implies Conjecture 1.2.

### 1.8. Paper structure

We recall poset theoretic definitions and notation in Section 2. Recursive constructions of posets are studied in Section 3. We present the proofs in Section 4. We conclude with final remarks and open problems in Section 5.

## 2. Basic definitions and notation

### 2.1. Posets

For a poset  $P = (X, \prec)$  and a subset  $Y \subset X$ , denote by  $P_Y = (Y, \prec)$  a subposet of  $P$ . We use  $(P - z)$  to denote a subposet  $P_{X-z}$ , where  $z \in X$ . Element  $x \in X$  is *minimal* in  $P$ , if there exists no element  $y \in X - x$  such that  $y \prec x$ . Denote by  $\min(P)$  the set of minimal elements in  $P$ .

In a poset  $P = (X, \prec)$ , elements  $x, y \in X$  are called *incomparable* if  $x \not\prec y$  and  $y \not\prec x$ . We write  $x \parallel y$  in this case. An *antichain* is a subset  $A \subset X$  of pairwise incomparable elements. The *width* of poset  $P = (X, \prec)$ , denoted  $\text{width}(P)$ , is the size of a maximal antichain. A *chain* is a subset  $C \subset X$  of pairwise comparable elements. Denote by  $A_n$  and  $C_n$  the antichain and the chain with  $n$  elements, respectively.

A *dual poset* is a poset  $P^* = (X, \prec^*)$ , where  $x \prec^* y$  if and only if  $y \prec x$ . A *parallel sum*  $P \oplus Q$  of posets  $P = (X, \prec)$  and  $Q = (Y, \prec')$  is a poset  $(X \cup Y, \prec^\diamond)$ , where the relation  $\prec^\diamond$  coincides with  $\prec$  and  $\prec'$  on  $X$  and  $Y$ , and  $x \parallel y$  for all  $x \in X, y \in Y$ . A *linear sum*  $P \otimes Q$  of posets  $P = (X, \prec)$  and  $Q = (Y, \prec')$  is a poset  $(X \cup Y, \prec^\diamond)$ , where the relation  $\prec^\diamond$  coincides with  $\prec$  and  $\prec'$  on  $X$  and  $Y$ , and  $x \prec^\diamond y$  for all  $x \in X, y \in Y$ .

Note that  $e(P^*) = e(P)$ ,  $e(P \otimes Q) = e(P)e(Q)$  and  $e(P \oplus Q) = \binom{n+n'}{n} e(P)e(Q)$ , where  $|X| = n$  and  $|Y| = n'$ . We refer to [39, Ch. 3] for an accessible introduction, and to surveys [6, 10, 41] for further definitions and standard results.

### 2.2. Continued fractions

Consider a GCF  $[a_1, \dots, a_m; b_0, \dots, b_m]$  given by (1.9). Recursively define  $C_i := C_i(a_1, \dots, a_m; b_0, \dots, b_m)$  and  $D_i := D_i(a_1, \dots, a_m; b_0, \dots, b_m)$ ,  $0 \leq i \leq m$ , as follows:

$$\begin{aligned} C_m &:= b_m, & D_m &:= 1, \\ D_i &:= C_{i+1}, & C_i &:= b_i D_i + a_{i+1} D_{i+1}. \end{aligned}$$

It is easy to see by induction that

$$[a_{i+1}, a_{i+2}, \dots, a_m; b_i, b_{i+1}, \dots, b_m] = \frac{C_i}{D_i}.$$

These are called *partial fractions* or *tails of continued fractions*. Note that for simple CFs we have  $\gcd(C_i, D_i) = 1$ , but this does not always hold for GCFs.

Similarly, consider a RGCF  $[\alpha_1, \dots, \alpha_m; b_0, \dots, b_m]$  given by (1.12). Let  $\alpha_i = c_i/d_i$  where  $\gcd(c_i, d_i) = 1$ ,  $1 \leq i \leq m$ . Recursively define  $C_i := C_i(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m)$  and  $D_i := D_i(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m)$  as follows:

$$\begin{aligned} C_m &:= b_m, \quad D_m := 1, \\ D_i &:= d_{i+1} (C_{i+1} + (s(\alpha_{i+1}) - 1)D_{i+1}), \\ C_i &:= b_i D_i + c_{i+1} (C_{i+1} + s(\alpha_{i+1})D_{i+1}). \end{aligned}$$

It is easy to see by induction that

$$[\alpha_{i+1}, \alpha_{i+2}, \dots, \alpha_m; b_i, b_{i+1}, \dots, b_m] = \frac{C_i}{D_i}.$$

The ratios  $\frac{C_i}{D_i}$  are called *partial fractions* in this case.

### 3. Recursive constructions

#### 3.1. Hybrid sums

Let  $P = (X, \prec)$  and  $Q = (Y, \prec')$  be posets on  $|X| = n$  and  $|Y| = n'$  elements. Fix  $x \in \min(P)$ . The *hybrid sum*  $Q \otimes_x P$  is the poset  $R = (X \cup Y, \prec^\diamond)$  given by the relations

$$\begin{aligned} u \prec^\diamond u' &\quad \text{for every } u \prec u', \quad u, u' \in X, \\ v \prec^\diamond v' &\quad \text{for every } v \prec' v', \quad v, v' \in Y, \\ v \prec^\diamond u &\quad \text{for every } u \in X - x, \quad v \in Y. \end{aligned}$$

Note that  $x$  is incomparable to  $Y$  in  $R$ , and thus  $x \in \min(R)$ .

We have:

$$e(Q \otimes_x P) = e(Q)e(P) + e(Q \oplus x)e(P - x) - e(Q)e(P - x).$$

Indeed, the term  $e(Q)e(P)$  counts linear extensions  $f \in \mathcal{E}(R)$  for which  $f(x) \geq n' + 1$ . Similarly, the term  $e(Q \oplus x)e(P - x)$  counts  $f \in \mathcal{E}(R)$  for which  $f(x) \leq n' + 1$ . Finally, the term  $e(Q)e(P - x)$  counts  $f \in \mathcal{E}(R)$  for which  $f(x) = n' + 1$ . Because  $e(Q \oplus x) = (n' + 1) \cdot e(Q)$ , we then have

$$e(Q \otimes_x P) = e(Q)e(P) + n' \cdot e(Q)e(P - x). \quad (3.1)$$

It then follows that for all  $y \in \min(Q)$ , we have

$$e(R - y) = e((Q - y) \otimes_x P) = e(Q - y)e(P) + (n' - 1) \cdot e(Q - y)e(P - x). \quad (3.2)$$

Since  $(Q \otimes_x P) - x = Q \otimes (P - x)$ , we also have:

$$e((Q \otimes_x P) - x) = e(Q)e(P - x). \quad (3.3)$$

Finally, note that

$$\text{width}(Q \otimes_x P) \leq \max\{\text{width}(P) - 1, \text{width}(Q)\} + 1. \quad (3.4)$$

**Remark 3.1.** Hybrid sum is a special case of the *quasi-series composition* defined similarly in [15] for general subsets of minimal elements. Also, when  $Y = \{y\}$ , we have  $R = \{y\} \oplus_{y,x} P$ , where  $\oplus_{y,x}$  is the direct sum operation defined in [23, §2].

#### 3.2. Properties of hybrid sums

We now use hybrid sums to construct posets for which the numbers of linear extensions satisfy recurrence relations emulating continued fractions.

**Lemma 3.2.** Let  $P = (X, \prec)$  and  $Q = (Y, \prec')$  be posets on  $n = |X|$  and  $n' = |Y|$  elements, and let  $x \in \min(P)$ ,  $y \in \min(Q)$ . Let  $R = (Z, \prec^\diamond)$  be a poset and let  $z \in \min(R)$  be given by

$$R := Q \otimes_x P, \quad z := y.$$

Then we have

$$\begin{aligned} e(R) &= e(Q)(e(P) + n' \cdot e(P - x)), \\ e(R - z) &= e(Q - y)(e(P) + (n' - 1) \cdot e(P - x)), \\ |Z| &= n + n', \\ \text{width}(R) &\leq \max\{\text{width}(P), \text{width}(Q) + 1\}. \end{aligned}$$

Additionally, we have:

$$\rho(R, z) = \rho(Q, y) \left( 1 + \frac{1}{n' - 1 + \rho(P, x)} \right).$$

**Proof.** The first four conclusions follows from (3.1), (3.2), (3.4). We conclude that

$$\begin{aligned} \rho(R, z) &= \frac{e(Q)}{e(Q - y)} \cdot \frac{e(P) + n' \cdot e(P - x)}{e(P) + (n' - 1) \cdot e(P - x)} \\ &= \rho(Q, y) \left( 1 + \frac{e(P - x)}{e(P) + (n' - 1) \cdot e(P - x)} \right) \\ &= \rho(Q, y) \left( 1 + \frac{1}{\rho(P, x) + (n' - 1)} \right), \end{aligned}$$

as desired.  $\square$

**Lemma 3.3.** Let  $P = (X, \prec)$  be a poset on  $|X| = n$  elements, let  $x \in \min(P)$ , and let  $b \geq 0$ . Let  $R = (Z, \prec^\diamond)$  be a poset and let  $z \in \min(R)$  be given by

$$R := C_b \otimes_x P, \quad z := x,$$

where  $C_b$  is a chain of  $b$  elements. Then we have

$$\begin{aligned} e(R) &= e(P) + b \cdot e(P - x), \\ e(R - z) &= e(P - x), \\ |Z| &= n + b, \\ \text{width}(R) &\leq \max\{\text{width}(P), 2\}. \end{aligned}$$

Additionally, we have:

$$\rho(R, z) = b + \rho(P, x).$$

**Proof.** The first four conclusions follows from (3.1), (3.3), and (3.4). We conclude that

$$\rho(R, z) = \frac{e(P) + b \cdot e(P - x)}{e(P - x)} = \rho(P, x) + b,$$

as desired.  $\square$

By combining the two lemmas above, we get the following:

**Lemma 3.4.** Let  $P = (X, \prec)$  and  $Q = (Y, \prec')$  be posets on  $n = |X|$  and  $n' = |Y|$  elements, and let  $x \in \min(P)$ ,  $y \in \min(Q)$ . Fix  $b \geq 0$ . Let  $R = (Z, \prec^\diamond)$  be a poset and let  $z \in \min(R)$  be given by

$$R := C_b \otimes_y (Q \otimes_x P), \quad z := y.$$

where  $C_b$  is a chain of  $b$  elements. Then we have

$$\begin{aligned} e(R - z) &= e(Q - y)(e(P) + (n' - 1)e(P - x)), \\ e(R) &= b \cdot e(R - z) + e(Q)[e(P) + n' \cdot e(P - x)], \\ |Z| &= n + n' + b, \\ \text{width}(R) &\leq \max\{\text{width}(P), \text{width}(Q) + 1, 2\}. \end{aligned}$$

Additionally, we have:

$$\rho(R, z) = b + \rho(Q, y) \left( 1 + \frac{1}{n' - 1 + \rho(P, x)} \right).$$

**Proof.** This follows from first applying [Lemma 3.2](#) then applying [Lemma 3.3](#).  $\square$

**Lemma 3.5.** Let  $P = (X, \prec)$  be a poset on  $|X| = n$  elements, let  $x \in \min(P)$ , and let  $b \geq a \geq 0$ . Let  $R = (Z, \prec^\diamond)$  be a poset and let  $z \in \min(R)$  be given by

$$R := C_{b-a} \otimes_y ((y \oplus C_{a-1}) \otimes_x P), \quad z := y,$$

where  $C_b$  is a chain of  $b$  elements, and  $y$  is an element not contained in  $P$ . Then we have

$$\begin{aligned} e(R - z) &= e(P) + (a - 1) \cdot e(P - x), \\ e(R) &= (b - a) \cdot e(R - z) + a \cdot [e(P) + a \cdot e(P - x)] = b \cdot e(R - z) + a \cdot e(P - x), \\ |Z| &= n + b, \\ \text{width}(R) &\leq \max\{\text{width}(P), 3\}. \end{aligned}$$

Additionally, we have:

$$\rho(R, z) = b + \frac{a}{a - 1 + \rho(P, x)}.$$

**Proof.** Let  $Q = (Y, \prec') := y \oplus C_{a-1}$ : Note that

$$e(Q) = a, \quad e(Q - y) = 1, \quad |Y| = a, \quad \text{and} \quad \text{width}(Q) = 2.$$

The lemma now follows from substituting  $b \leftarrow (b - a)$  into [Lemma 3.4](#).  $\square$

### 3.3. A flip-flop construction

We will need the following variation on the hybrid sum construction to prove [Theorem 1.8](#).

Let  $P = (X, \prec)$  and  $Q = (Y, \prec')$  be posets on  $n = |X|$  and  $n' = |Y|$  elements, and let  $x \in \min(P)$ ,  $y \in \min(Q)$ . The *flip-flop poset*  $R = (Z, \prec^\diamond)$  is defined by

$$Z := (X - x) \cup (Y - y) \cup \{z, v\},$$

where  $z, v$  are new elements. The partial order  $\prec^\diamond$  is defined by

$$\begin{aligned} p \prec^\diamond p' &\quad \text{for every } p, p' \in X - x \text{ s.t. } p \succ p', \\ q \prec^\diamond q' &\quad \text{for every } q, q' \in Y - y \text{ s.t. } q \prec' q', \\ p \prec^\diamond z &\quad \text{for every } p \in X - x \text{ s.t. } x \prec p, \\ z \prec^\diamond q &\quad \text{for every } q \in Y - y \text{ s.t. } y \prec' q, \\ p \prec^\diamond v \prec^\diamond q &\quad \text{for every } p \in X - x, \quad q \in Y - y, \\ &\quad \text{and} \quad z \parallel_{\prec^\diamond} v. \end{aligned}$$

**Lemma 3.6.** *The flip-flop poset  $R = (Z, \prec^\diamond)$  satisfies*

$$\begin{aligned} e(R) &= e(P)e(Q - y) + e(P - x)e(Q), \\ e(R - z) &= e(P - x)e(Q - y), \\ |Z| &= n + n', \\ \text{width}(R) &\leq \text{width}(P) + \text{width}(Q). \end{aligned}$$

Additionally, we have:

$$\rho(R, z) = \rho(P, x) + \rho(Q, y).$$

We warn the reader that the element  $z$  is not necessarily a minimal element of  $R$ , so this construction cannot be easily iterated.

**Proof.** We have:

$$e(R) = e(P) \cdot e(Q - y) + e(P - x) \cdot e(Q).$$

Indeed, the factor  $e(P) \cdot e(Q - y)$  counts linear extensions  $f \in \mathcal{E}(R)$  for which  $f(u) < f(v)$ , while the factor  $e(P - x) e(Q)$  counts linear extensions  $f \in \mathcal{E}(R)$  for which  $f(u) > f(v)$ . Also note that

$$e(R - z) = e(P - x) \cdot e(Q - y),$$

because  $(R - z)$  is isomorphic to the linear sum  $(P - x) \oslash \{v\} \oslash (Q - y)$ . Finally, note that

$$\text{width}(R) \leq \text{width}(P) + \text{width}(Q),$$

$$|Z| = |X| + |Y|,$$

by construction. This completes the proof.  $\square$

## 4. Proofs

### 4.1. Proof of Theorem 1.10

We prove the claim by induction on  $m$ . First, let  $m = 0$ . Recall the notation in Section 2.2. Note that condition (1.10) becomes  $b_0 \geq 1$ , which holds by the assumption. Let  $P = (X, \prec) := \{x\} \oplus C_{b_0-1}$ . Then we have:

$$e(P) = b_0 = C_0(b_0) \quad \text{and} \quad e(P - x) = 1 = D_0(b_0).$$

We also have  $|X| = b_0$  and  $\text{width}(P) = 2$ , as desired.

Suppose now that the claim holds for  $m - 1$ . Let  $b'_1 := b_1 - a_1 + 1$ . The balanced assumptions (1.10) gives  $b'_1 \geq a_2$ . Thus, by the inductive assumption, there exist  $P' = (X', \prec')$  and  $x' \in \min(P')$ , such that

$$\begin{aligned} e(P' - x') &= D_0(a_2, \dots, a_m; b'_1, b_2, \dots, b_m) = D_1(a_1, \dots, a_m; b_0, b_1, \dots, b_m), \\ e(P') &= C_0(a_2, \dots, a_m; b'_1, b_2, \dots, b_m) \\ &= C_0(a_2, \dots, a_m; b_1, b_2, \dots, b_m) - (a_1 - 1) \cdot D_0(a_2, \dots, a_m; b_1, b_2, \dots, b_m) \\ &= C_1(a_1, \dots, a_m; b_0, \dots, b_m) - (a_1 - 1) \cdot D_1(a_1, \dots, a_m; b_0, \dots, b_m). \end{aligned}$$

Now, apply Lemma 3.5 to  $P \leftarrow P'$  with  $b \leftarrow b_0$  and  $a \leftarrow a_1$  and  $x \leftarrow x'$ . We obtain a poset  $R = (Z, \prec^\diamond)$  on  $|Z| = n$  elements, and  $z \in \min(R)$ , such that

$$\begin{aligned} e(R - z) &= e(P') + (a_1 - 1) \cdot e(P' - x') = C_1(a_1, \dots, a_m; b_0, \dots, b_m) \\ &= D_0(a_1, \dots, a_m; b_0, \dots, b_m), \\ e(R) &= b_0 \cdot e(R - z) + a_1 \cdot e(P' - x') \end{aligned}$$

$$\begin{aligned}
&= b_0 \cdot D_0(a_1, \dots, a_m; b_0, \dots, b_m) + a_1 \cdot D_1(a_1, \dots, a_m; b_0, \dots, b_m) \\
&= C_0(a_1, \dots, a_m; b_0, \dots, b_m).
\end{aligned}$$

Since induction assumption implies that  $|X'| = G(a_2, \dots, a_m; b'_1, b_2, \dots, b_m)$ , we also have

$$\begin{aligned}
n &= b_0 + |X'| = b_0 + b'_1 + \sum_{i=2}^m b_i - \sum_{i=2}^m a_i + m - 1 \\
&= \sum_{i=0}^m b_i - \sum_{i=1}^m a_i + m = G(a_1, \dots, a_m; b_0, \dots, b_m),
\end{aligned}$$

and  $\text{width}(R) \leq \max\{\text{width}(P'), 3\} \leq 3$ . Finally, we have:

$$\rho(R, z) = \frac{C_0(a_1, \dots, a_m; b_0, \dots, b_m)}{D_0(a_1, \dots, a_m; b_0, \dots, b_m)} = [a_1, \dots, a_m; b_0, \dots, b_m].$$

The theorem is now complete by substituting  $P \leftarrow R$  and  $x \leftarrow z$ .  $\square$

#### 4.2. Proof of Theorem 1.13

We prove the claim by induction on  $m$ . For  $m = 0$ , let  $P = (X, \prec) := x \oplus C_{b_0-1}$ . We have:

$$e(P) = b_0 = C_0(b_0) \quad \text{and} \quad e(P - x) = 1 = D_0(b_0).$$

We also have  $|X| = b_0$  and  $\text{width}(P) = 2$ , which proves the case  $m = 0$ .

We now suppose the claim is already proved for  $(m - 1)$ . By the induction assumption, there exists a poset  $P' = (X', \prec')$  and element  $x' \in \min(P')$ , such that

$$\begin{aligned}
e(P' - x') &= D_0(\alpha_2, \dots, \alpha_m; b_1, b_2, \dots, b_m) = D_1(\alpha_1, \dots, \alpha_m; b_0, b_1, \dots, b_m), \\
e(P') &= C_0(\alpha_2, \dots, \alpha_m; b_1, b_2, \dots, b_m) = C_1(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m).
\end{aligned}$$

Applying Theorem 1.6 to  $\alpha_1$ , there exists a poset  $Q = (Y, \prec')$  and  $y \in \min(Q)$  such that

$$e(Q) = c_1, \quad e(Q - y) = d_1, \quad |Y| = s(\alpha_1) \quad \text{and} \quad \text{width}(Q) \leq 2.$$

Now, apply Lemma 3.4 to posets  $P'$ ,  $Q$ , and element  $b_0$ . We obtain a poset  $P = (X, \prec)$  and  $x \in \min(P)$ , such that

$$\begin{aligned}
e(P - x) &= e(Q - y)[e(P') + (|Y| - 1)e(P' - x')], \\
&= d_1 \cdot [C_1(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m) + (s(\alpha_1) - 1) \cdot D_1(\alpha_1, \dots, \alpha_m; b_0, b_1, \dots, b_m)] \\
&= D_0(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m).
\end{aligned}$$

We also have

$$\begin{aligned}
e(P) &= b_0 \cdot e(P - x) + e(Q)[e(P') + |Y| \cdot e(P' - x')] \\
&= b_0 \cdot D_0(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m) \\
&\quad + c_1 \cdot [C_1(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m) + s(\alpha_1) \cdot D_1(\alpha_1, \dots, \alpha_m; b_0, b_1, \dots, b_m)] \\
&= C_0(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m)
\end{aligned}$$

and

$$\text{width}(P) \leq \max\{\text{width}(P'), \text{width}(Q) + 1, 2\} \leq 3.$$

Finally, we have

$$\begin{aligned}
|X| &= |X'| + |Y| + b_0 \\
&= (b_1 + \dots + b_m) + s(\alpha_2) + \dots + s(\alpha_m) + s(\alpha_1) + b_0 \\
&= R(\alpha_1, \dots, \alpha_m; b_0, \dots, b_m).
\end{aligned}$$

This completes the proof.  $\square$

### 4.3. Proof of [Proposition 1.12](#) and [1.15](#)

For [Proposition 1.12](#), recall from the introduction that [Conjecture 1.11](#) implies [Conjecture 1.2](#) for prime  $d$ . Indeed, by [Theorem 1.6](#) for a GCF  $[a_1, \dots, a_m; b_0, \dots, b_m] = \frac{d}{c}$ , we obtain a poset  $P = (X, \prec)$  and  $x \in X$  such that  $|X| = g\left(\frac{d}{c}\right) \leq C \log d$  and  $\frac{e(P)}{e(P-x)} = \frac{d}{c}$ . By the reduced condition on the definition of  $g$ , it then follows that  $e(P) = d$ , as desired.

To show that the first part of [Conjecture 1.11](#) suffices, let  $p_1^{m_1} \dots p_\ell^{m_\ell}$  be the prime factorization of  $d$ . For each prime  $p_i$ , let  $P_i = (X_i, \prec_i)$  be the corresponding poset with  $e(P_i) = p_i$  and  $|X_i| \leq C \log p_i$ . Define

$$P := \underbrace{P_1 \otimes \dots \otimes P_1}_{m_1 \text{ times}} \otimes \dots \otimes \underbrace{P_\ell \otimes \dots \otimes P_\ell}_{m_\ell \text{ times}}$$

be the linear sum of posets  $P_i$ . We have:

$$e(P) = \prod_{i=1}^{\ell} e(P_i)^{m_i} = d,$$

and

$$|X| = \sum_{i=1}^{\ell} m_i |X_i| \leq C \sum_{i=1}^{\ell} m_i \log p_i = C \log d.$$

This completes the proof of [Proposition 1.12](#). The proof of [Proposition 1.15](#) follows verbatim.  $\square$

### 4.4. Proof of [Theorem 1.8](#)

First, observe that there exists a constant  $C > 0$ , such that for all coprime integers  $a, b \leq d$  which satisfy  $C < b \leq a \leq 2b$ , there exists a positive integer  $\ell := \ell(a, b)$  such that  $1 \leq \ell < b$ , and

$$s\left(\frac{\ell}{b}\right) \leq 2 \log b \log \log b \quad \text{and} \quad s\left(\frac{a-\ell}{b}\right) \leq 2 \log b \log \log b. \quad (4.1)$$

Indeed, by [Theorem 1.9](#) and using  $\frac{12}{\pi^2} < 2$ , for random  $\ell \in \{1, \dots, b\}$ , the probability that each inequality fails  $\rightarrow 0$  as  $b \rightarrow \infty$ . Taking  $C$  large enough so that each probability is  $< \frac{1}{2}$  proves the claim.

Let  $a, b$  be given by

$$a := c + d - \left\lfloor \frac{d}{c} \right\rfloor c, \quad b := c,$$

so that  $b \leq a \leq 2b$  and  $b \leq d$ . From above, there exists  $1 \leq \ell \leq b$ , such that (4.1) holds. Let

$$\alpha := 1 + \frac{\ell}{b} \quad \text{and} \quad \beta := \left\lfloor \frac{d}{c} \right\rfloor - 2 + \frac{a-\ell}{b}.$$

It follows from the construction that  $\alpha + \beta = \frac{d}{c}$ : Since  $\frac{d}{c} \geq 3$ , we have  $\alpha, \beta \geq 1$ .

Applying [Theorem 1.10](#) to simple continued fractions, we obtain a poset  $P = (X, \prec)$  and element  $x \in \min(P)$ , such that

$$\rho(P, x) = \alpha \quad \text{and} \quad |X| = 1 + s\left(\frac{\ell}{b}\right).$$

Similarly, we obtain a poset  $Q = (Y, \prec')$  and element  $y \in \min(Q)$ , such that

$$\rho(Q, y) = \beta \quad \text{and} \quad |Y| = \left\lfloor \frac{d}{c} \right\rfloor - 2 + s\left(\frac{a-\ell}{b}\right).$$

By [Lemma 3.6](#), there exists a poset  $R = (Z, \prec^\diamond)$  and element  $z \in Z$ , such that

$$\rho(R, z) = \rho(P, x) + \rho(Q, y) = \frac{d}{c},$$

and

$$|Z| = |X| + |Y| = \left\lfloor \frac{d}{c} \right\rfloor - 1 + s\left(\frac{\ell}{b}\right) + s\left(\frac{a-\ell}{b}\right) \leq \frac{d}{c} + O(\log d \log \log d).$$

This completes the proof.  $\square$

## 5. Final remarks and open problems

### 5.1.

The nature of connections between counting combinatorial objects and continued fractions described in Section 1.1 is clear and easy to explain: when objects are decomposed into smaller objects, they often have simple recurrences of the type described in Section 2.2. Fundamentally, this is the same reason why the generating functions are so powerful in combinatorial enumeration, see e.g. [14,39]. And yet, every time such a connection is found it is an unexpected delight, stemming both from the sheer elegance of continued fractions as well as the power of technical tools developed for them. While we tend to be swayed by the latter arguments, we appreciate the former sentiments.

### 5.2.

The upper bound in Larcher's [Theorem 1.3](#) was sharpened by Rukavishnikova [30] to  $O(\log d \log \log d)$ . Since  $\frac{d}{\phi(d)}$  can be as large as  $C \log \log d$ , see e.g. [16, Thm 328], this is a significant asymptotic improvement. This result was further sharpened by Aistleitner, Borda and Hauke [1, Cor. 2], who proved that for all  $d \geq 3$  there exist  $1 \leq c < d$ , such that

$$s\left(\frac{c}{d}\right) \leq \frac{12}{\pi^2} \log d \log \log d + O(\log d). \quad (5.1)$$

Note that we are using only prime  $d$  for our applications, which is why we postponed this recent result. We note in passing that the authors of [23] stated [Conjecture 1.4](#) in the generality of all  $d$ ; while plausible this remains out of reach with the existing technology. They were unaware of the earlier work and rediscovered [Theorem 1.3](#).<sup>1</sup>

### 5.3.

The asymptotics in the upper bound (5.1) cannot be easily improved by probabilistic arguments. This follows from a version on the tail estimates (1.8) given in [29]. A stronger result was proved in [1, Thm 1], which implies that for all  $C < 0$ ,  $d \geq 3$ , and  $\varepsilon = \varepsilon(d) > (\log d)^C$ , for the  $(1 - \varepsilon)$  fraction of  $c \in \{1, \dots, d\}$  with  $\gcd(c, d) = 1$ , we have:

$$\left| s\left(\frac{c}{d}\right) - \frac{12}{\pi^2} \log d \log \log d \right| = O\left(\frac{\log d}{\varepsilon}\right). \quad (5.2)$$

Of course, this does not preclude the outlying small values predicted by Zaremba's conjecture. In fact, as was pointed out in [1], the distribution of  $s\left(\frac{c}{d}\right)$  is heavy-tailed and has a large mean:

$$\frac{1}{\phi(d)} \sum_c s\left(\frac{c}{d}\right) = \frac{6}{\pi^2} (\log d)^2 + O((\log d)(\log \log d)^2), \quad (5.3)$$

where the summation is over all  $c \in \{1, \dots, d\}$  such that  $\gcd(c, d) = 1$ . This was proved independently in [25,26,43].

<sup>1</sup> Personal communication.

## 5.4.

It was pointed out by Kravitz and Sah (see Remark 5.31 in [8]), that the numerator  $c$  in [Theorem 1.6](#) can be found in probabilistic polynomial time  $\text{poly}(\log d)$ . Tail estimates (5.2) give a simpler (and faster) probabilistic algorithm: pick a random  $c$ , check if  $\gcd(c, d) = 1$ , compute a simple CF (1.1), repeat if  $s(\frac{c}{d}) > 2 \log d \log \log d$ . It is an interesting open problem if this can be done deterministically. More broadly, is there a deterministic polynomial time construction of a poset with exactly  $n$  linear extensions? So far, the only deterministic construction we know of is by Tenner [40], which is exponential in  $(\log n)$ .

## 5.5.

Zaremba's [Conjecture 1.5](#) is often stated with  $A = 5$  or even  $A = 4$  for all sufficiently large integers. It is known to hold for integers of the form  $2^m 3^n$ , for other families of powers of small primes and sufficiently large powers of all primes, where the constant  $A$  can depend on the prime, see [34]. We refer to [3, §6.2] for an elegant presentation of the  $2^m$  case. Of course, the Kravitz-Sah [Conjecture 1.2](#) is trivial in this case. Note that the constant 50 in the Bourgain-Kontorovich theorem that was used in the proof of [Corollary 1.7](#), has been improved to 5 in [17]. See [19] for further extensions, and [33, §7] for an overview.

## 5.6.

It would be interesting to find an elementary proof of the first part of [Corollary 1.7](#). The result is especially surprising given that the bound is obtained on a relatively small family of posets of width two. On the other hand, we know of no nontrivial bound for the much larger family of height two posets (cf. [37]).

## 5.7.

In [8, Conj. 5.17], we conjecture that all but finitely many integers are the numbers of linear extensions of posets of height two. We also observe (Prop. 5.18, *ibid.*), that this would imply [Conjecture 1.2](#) with a sharp  $\Theta(\frac{\log n}{\log \log n})$  asymptotics.

## Acknowledgements

We are grateful to Christoph Aistleitner, Milan Haiman, Oleg Karpenkov, Alex Kontorovich, Noah Kravitz, Greta Panova, Ashwin Sah, Rolf Schiffler, Ilya Shkredov, Nikita Shulga and Alan Sokal for interesting discussions and helpful remarks. Special thanks to Maria Rukavishnikova for sending us [29]. Last but not least, we would like to thank the anonymous referees for valuable comments and suggestions. Both authors were partially supported by the United States National Science Foundation.

## References

- [1] Christoph Aistleitner, Bence Borda, Manuel Hauke, *On the distribution of partial quotients of reduced fractions with fixed denominator*, *Trans. Amer. Math. Soc.* 377 (2024) 1371–1408.
- [2] George E. Andrews, Bruce C. Berndt, *Ramanujan's Lost Notebook. Part I*, Springer, New York, 2005, p. 437.
- [3] Jonathan Borwein, Alf van der Poorten, Jeffrey Shallit, Wadim Zudilin, *Neverending Fractions*, Cambridge Univ. Press, Cambridge, UK, 2014, p. 212.
- [4] Jean Bourgain, Alex Kontorovich, *On Zaremba's conjecture*, *Ann. Math.* 180 (2014) 137–196.
- [5] Claude Brezinski, *History of Continued Fractions and Padé Approximants*, Springer, Berlin, 1991, p. 551.
- [6] Graham Brightwell, Douglas B. West, *West partially ordered sets*, in: *Handbook of Discrete and Combinatorial Mathematics*, CRC Press, Boca Raton, FL, 2000, pp. 717–752 (Ch. 11).
- [7] İlke Çanakçı, Ralf Schiffler, *Cluster algebras and continued fractions*, *Compos. Math.* 154 (2018) 565–593.
- [8] Swee Hong Chan, Igor Pak, *Computational complexity of counting coincidences*, *Theoret. Comput. Sci.* (in press). arXiv:2308.10214.

[9] Swee Hong Chan, Igor Pak, Equality cases of the Alexandrov–Fenchel inequality are not in the polynomial hierarchy, 2023, preprint (2023) 35 pp.; [arXiv:2309.05764](https://arxiv.org/abs/2309.05764).

[10] Swee Hong Chan, Igor Pak, Linear extensions of finite posets, 2023, preprint (2023) 55 pp.; [arXiv:2311.02743](https://arxiv.org/abs/2311.02743).

[11] Sergi Elizalde, Continued fractions for permutation statistics, *Discrete Math. Theor. Comput. Sci.* 19 (2) (2017) Paper No. 11 24pp.

[12] Philippe Flajolet, Combinatorial aspects of continued fractions, *Discrete Math.* 32 (1980) 125–161.

[13] Philippe Flajolet, Robert Sedgewick, *Analytic Combinatorics*, Cambridge Univ. Press, Cambridge, 2009.

[14] Ian P. Goulden, David M. Jackson, *Combinatorial Enumeration*, John Wiley, New York, 1983, 569pp.

[15] Michel Habib, Roland Jégo, *N*-Free posets as generalizations of series-parallel posets, *Discrete Appl. Math.* 12 (1985) 279–291.

[16] Godfrey H. Hardy, Edward M. Wright, *An Introduction to the Theory of Numbers*, sixth ed., revised, Oxford Univ. Press, Oxford, 2008, p. 621.

[17] ShinnYih Huang, An improvement to Zaremba’s conjecture, *Geom. Funct. Anal.* 25 (2015) 860–914.

[18] William B. Jones, Wolfgang J. Thron, *Continued Fractions. Analytic Theory and Applications*, Addison-Wesley, Reading, MA, 1980, p. 428.

[19] Igor D. Kan, A strengthening of the Bourgain–Kontorovich method: three new theorems, *Sb. Math.* 212 (2021) 921–964.

[20] Oleg Karpenkov, *Geometry of Continued Fractions*, Springer, Heidelberg, 2013, p. 405.

[21] Aleksandr Ya. Khinchin, *Continued Fractions*, Dover, Mineola, NY, 1997, p. 95.

[22] Donald E. Knuth, *The Art of Computer Programming. Vol. 2. Seminumerical Algorithms*, third ed., Addison-Wesley, Reading, MA, 1998, p. 762.

[23] Noah Kravitz, Ashwin Sah, Linear extension numbers of  $n$ -element posets, *Order* 38 (2021) 49–66.

[24] Gerhard Larcher, On the distribution of sequences connected with good lattice points, *Monatsh. Math.* 101 (1986) 135–150.

[25] Bernhard Liehl, Über die Teilnennen endlicher Kettenbrüche, *Arch. Math. (Basel)* 40 (1983) 139–147 (in German).

[26] Alexey A. Panov, Mean for the sum of elements over a class of finite continued fractions, *Math. Notes* 32 (1982) 781–785.

[27] Mathias Pétréolle, Alan Sokal, Bao-Xuan Zhu, Lattice paths and branched continued fractions, *Mem. Amer. Math. Soc.* 291 (1450) (2023) 154.

[28] Andrew M. Rockett, Peter Szűsz, *Continued Fractions*, World Sci., River Edge, NJ, 1992, p. 188.

[29] Maria G. Rukavishnikova, A probability estimate for the sum of incomplete partial quotients with fixed denominator, *Chebyshevskii Sb.* 7 (4) (2006) 113–121 (in Russian). available at [tinyurl.com/ytv33cct](https://tinyurl.com/ytv33cct).

[30] Maria G. Rukavishnikova, The law of large numbers for the sum of partial quotients of a rational number with a fixed denominator, *Math. Notes* 90 (2011) 418–430.

[31] Tomas Sauer, *Continued Fractions and Signal Processing*, Springer, Cham, 2021, p. 263.

[32] Rolf Schiffler, Snake graphs, perfect matchings and continued fractions, in: *Snapshots of Modern Mathematics from Oberwolfach*, No. 1, 2019, p. 10, available at [publications.mfo.de/handle/mfo/1405](https://publications.mfo.de/handle/mfo/1405).

[33] Ilya D. Shkredov, Non-commutative methods in additive combinatorics, *Russian Math. Surveys* 76 (2021) 1065–1122.

[34] Nikita Shulga, Radical bound for Zaremba’s conjecture, 2023, preprint (2023) 8 pp.; [arXiv:2310.09801](https://arxiv.org/abs/2310.09801).

[35] Neil J.A. Sloane, The online encyclopedia of integer sequences, [oeis.org](https://oeis.org).

[36] Alan D. Sokal, Jian Zeng, Some multivariate master polynomials for permutations, set partitions, and perfect matchings, and their continued fractions, *Adv. Appl. Math.* 138 (2022) Paper No. 102341 122pp.

[37] David Soukup, Complexity of sign imbalance, parity of linear extensions, and height 2 posets, 2023, preprint (2023) 9 pp.; [arXiv:2311.02203](https://arxiv.org/abs/2311.02203).

[38] Thomás Jung Spier, *Graph Continued Fractions* (Ph.D. thesis), IMPA, 2021, 94.

[39] Richard P. Stanley, *Enumerative Combinatorics*, Vol. 1 (second ed.) and Vol. 2, and 2, Cambridge Univ. Press, 2012 and 1999, 626pp. and 581 pp.

[40] Bridget E. Tenner, Optimizing linear extensions, *SIAM J. Discrete Math.* 23 (2009) 1450–1454.

[41] William T. Trotter, Partially ordered sets, in: *Handbook of Combinatorics*, Vol. 1, Elsevier, Amsterdam, 1995, 433–480.

[42] Xavier Gérard Viennot, A combinatorial theory for general orthogonal polynomials with extensions and applications, in: *Lecture Notes in Math.*, vol. 1171, Springer, Berlin, 1985, pp. 139–157.

[43] Andrew C. Yao, Donald E. Knuth, Analysis of the subtractive algorithm for greatest common divisors, *Proc. Natl. Acad. Sci. USA* 72 (1975) 4720–4722.

[44] Stanisław K. Zaremba, La méthode des bons treillis pour le calcul des intégrales multiples, in: *Applications of Number Theory to Numerical Analysis*, Academic Press, New York, 1972, pp. 39–119 (in French).