



Multi-linear forms, graphs, and L^p -improving measures in \mathbb{F}_q^d

Pablo Bhowmik, Alex Iosevich, Doowon Koh, and Thang Pham

Abstract. The purpose of this paper is to introduce and study the following graph-theoretic paradigm. Let

$$T_K f(x) = \int K(x, y) f(y) d\mu(y),$$

where $f : X \rightarrow \mathbb{R}$, X a set, finite or infinite, and K and μ denote a suitable kernel and a measure, respectively. Given a connected ordered graph G on n vertices, consider the multi-linear form

$$\Lambda_G(f_1, f_2, \dots, f_n) = \int_{x^1, \dots, x^n \in X} \prod_{(i, j) \in \mathcal{E}(G)} K(x^i, x^j) \prod_{l=1}^n f_l(x^l) d\mu(x^l),$$

where $\mathcal{E}(G)$ is the edge set of G . Define $\Lambda_G(p_1, \dots, p_n)$ as the smallest constant $C > 0$ such that the inequality

$$(0.1) \quad \Lambda_G(f_1, \dots, f_n) \leq C \prod_{i=1}^n \|f_i\|_{L^{p_i}(X, \mu)}$$

holds for all nonnegative real-valued functions f_i , $1 \leq i \leq n$, on X . The basic question is, how does the structure of G and the mapping properties of the operator T_K influence the sharp exponents in (0.1). In this paper, this question is investigated mainly in the case $X = \mathbb{F}_q^d$, the d -dimensional vector space over the field with q elements, $K(x^i, x^j)$ is the indicator function of the sphere evaluated at $x^i - x^j$, and connected graphs G with at most four vertices.

1 Introduction

One of the fundamental objects in harmonic analysis is the operator of the form

$$(1.1) \quad T_K f(x) = \int_{\mathbb{R}^d} K(x, y) f(y) dy,$$

Received by the editors February 15, 2023; revised November 18, 2023; accepted December 4, 2023.
 Published online on Cambridge Core December 27, 2023.

A. Iosevich and P. Bhowmik were supported in part by the National Science Foundation (Grant No. HDR TRIPODS-1934962) and the National Science Foundation (Grant No. DMS-2154232). D. Koh was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (Grant No. RS-2023-00249597). T. Pham was supported by the research project QG.23.03 of Vietnam National University, Hanoi.

AMS subject classification: 42B05.

Keywords: Multi-linear forms, graphs, finite fields, boundedness problems.



where $K : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a suitable kernel and f is a locally integrable function. See [16] and the references contained therein for a variety of manifestations of operators of this type and their bounds.

The purpose of this paper is to study operators from (1.1) in the context of vector spaces over finite fields. Let \mathbb{F}_q denote the finite field with q elements, and let \mathbb{F}_q^d be the d -dimensional vector space over this field. Let $K : \mathbb{F}_q^d \times \mathbb{F}_q^d \rightarrow \mathbb{C}$ be a suitable kernel, and define

$$T_K f(x) = \sum_{y \in \mathbb{F}_q^d} K(x, y) f(y).$$

Operators of this type have been studied before [4, 11–13]. In particular, the operator T_K with $K(x, y) = S_t(x - y)$, where S_t is the indicator function of the sphere

$$S_t = \{x \in \mathbb{F}_q^d : \|x\| = t\},$$

$\|x\| = x_1^2 + x_2^2 + \cdots + x_d^2$, comes up naturally in the study of the Erdős–Falconer distance problem in vector spaces over finite fields, namely the question of how large a subset $E \subset \mathbb{F}_q^d$ needs to be to ensure that if

$$\Delta(E) = \{\|x - y\| : x, y \in E\},$$

then $|\Delta(E)| \geq \frac{q}{2}$. Here and throughout, $|S|$, with S a finite set, denotes the number of elements in this set. See, for example, [3, 5, 8, 10, 15].

If one is interested in studying more complicated geometric objects than distances, an interesting modification of the spherical averaging operator needs to be made. Indeed, let $E \subset \mathbb{F}_q^d$, and suppose that we want to know how many equilateral triangles of side-length 1 it determines. The quantity that counts such triangles is given by

$$(1.2) \quad \sum_{x, y, z \in \mathbb{F}_q^d} K(x, y) K(x, z) K(y, z) E(x) E(y) E(z),$$

where $K(x, y) = S_1(x - y)$.

Let us interpret the quantity (1.2) in the following way. Let us view x, y, z as vertices, and let us view the presence of $K(x, y)$ as determining the edge connecting x and y , and so on. In this way, the quantity (1.2) is associated with the graph K_3 , the complete graph on three vertices (Figure 1b).

Another natural example is the following. Let $K(x, y) = S_1(x - y)$, and consider the quantity that counts rhombi of side-length 1, i.e.,

$$(1.3) \quad \sum_{x, y, z, w \in \mathbb{F}_q^d} K(x, y) K(y, z) K(z, w) K(w, x) E(x) E(y) E(z) E(w).$$

Arguing as above, we associate this form with the graph C_4 , the cycle on four vertices (Figure 1e).

In general, let K be a kernel function, and let G be a connected ordered graph on n vertices. Define

$$(1.4) \quad \Lambda_G(f_1, f_2, \dots, f_n) = \frac{1}{\mathcal{N}(G)} \sum_{x^1, \dots, x^n \in \mathbb{F}_q^d} \prod_{(i, j) \in \mathcal{E}(G)} K(x^i, x^j) \prod_{l=1}^n f_l(x^l),$$




Figure 1:

where $\mathcal{E}(G)$ is the edge set of G and $\mathcal{N}(G)$ is the normalizing factor defined as the number of distinct embeddings of G in \mathbb{F}_q^d . Notice that $\mathcal{N}(G)$ is the number of tuples $(x^1, \dots, x^n) \in (\mathbb{F}_q^d)^n$ such that $\prod_{(i,j) \in \mathcal{E}(G)} K(x^i, x^j) = 1$. We will call the operator Λ_G as the G form on \mathbb{F}_q^d .

We note in passing that the paradigm we just introduced extends readily to the setting of hypergraphs. If we replace our basic object, the linear operator T_K , by an m -linear operator M_K , the problem transforms to the setting where the edges dictated by the kernel K are replaced by hyperedges induced by the multi-linear kernel $K(x^1, \dots, x^{m+1})$. We shall address this formulation of the problem in the sequel.

The norm $\|f\|_p$, $1 \leq p < \infty$, is defined to be associated with normalizing counting measure on \mathbb{F}_q^d . More precisely, given a function f on \mathbb{F}_q^d , we define

$$\|f\|_p := \left(q^{-d} \sum_{x \in \mathbb{F}_q^d} |f(x)|^p \right)^{\frac{1}{p}} \quad (1 \leq p < \infty), \quad \text{and} \quad \|f\|_\infty := \max_{x \in \mathbb{F}_q^d} |f(x)|.$$

Definition 1.1 Let n and d be nonnegative integers. For each finite field \mathbb{F}_q , we consider a connected ordered graph G on n vertices in \mathbb{F}_q^d . For any numbers $1 \leq p_i \leq \infty$, $i = 1, \dots, n$, we define $\Lambda_G(p_1, \dots, p_n)$ as the smallest number such that the following inequality

$$(1.5) \quad \Lambda_G(f_1, \dots, f_n) \leq \Lambda_G(p_1, \dots, p_n) \prod_{i=1}^n \|f_i\|_{p_i}$$

holds for all nonnegative real-valued functions f_i , $1 \leq i \leq n$, on \mathbb{F}_q^d .

Notice that the graph G in the above definition is chosen based on the underlying finite field \mathbb{F}_q . Hence, the operator norm $\Lambda_G(p_1, \dots, p_n)$ may depend on q , the size of the underlying finite field \mathbb{F}_q so that it can grow with q . However, if there exists a constant C , independent of q , such that $\Lambda_G(p_1, \dots, p_n) \leq C$, then we will denote $\Lambda_G(p_1, \dots, p_n) \lesssim 1$.

The main purpose of this paper is to determine all numbers $1 \leq p_i \leq \infty$, $i = 1, 2, \dots, n$, such that the operator norm $\Lambda_G(p_1, \dots, p_n)$ is not allowed to grow with q , that is, $\Lambda_G(p_1, \dots, p_n) \lesssim 1$. We will refer to this problem as the boundedness problem for the operator Λ_G on \mathbb{F}_q^d .

For the remainder of this paper, the kernel function $K(x, y)$ is assumed to be $S_t(x - y)$ with $t \neq 0$. In addition, when the dimension d is 2, we assume that the number 3 in \mathbb{F}_q is a square number so that we can exclude the trivial case in which the shape of an equilateral triangle in \mathbb{F}_q^2 does not occur.

We shall mainly confine ourselves to the following connected graphs G with at most four vertices: K_2 (the graph with two vertices and one edge), K_3 (the cycle with three vertices and three edges), $K_3 + \text{tail}$ (a kite), P_2 (the path of length 2), P_3 (the path of length 3), C_4 (the cycle with four vertices and four edges), $C_4 + \text{diagonal}$, Y -shape (a space station). In particular, we have avoided the K_4 (the complete graph with four vertices) since there is no K_4 distance graph on \mathbb{F}_q^2 . However, it would be interesting to investigate the case when the graph G is a K_4 in higher dimensions, a graph with more than four vertices, or a disconnected graph. Despite the difficulties posed by this case, we anticipate that experts in this field will address advanced results in the near future.

When the graph G is the K_2 , the complete answer to the boundedness problem will be given in all dimensions. To deduce the result, we will invoke the spherical averaging estimates over finite fields (see Theorem 3.3).

When the number of the vertices of the graph G is 3 or 4, we will obtain reasonably good boundedness results in two dimensions. In particular, in the case when the degree of each vertex is at least 2 (K_3 , $C_4 + \text{a diagonal}$, and C_4), we shall prove sharp results (up to the endpoints) for the operators on \mathbb{F}_q^2 (see Theorems 4.7, 6.5, and 7.5). While the proofs for the graphs K_2 and K_3 use standard results in the literature, in other cases, a new approach will be introduced. We also note that there are several papers in the literature studying the distribution of the graphs P_2 , P_3 , and Y -shape in a large set (see [2, 9] for example); however, the techniques in those papers are not helpful for the question raised in this paper. For three and higher dimensions, the boundedness problem is not simple and we will address partial results.

It is very natural to ask whether or not one can prove a general theorem that addresses all connected graphs on n vertices. Unfortunately, such a result is beyond the scope of this paper. The main difficulties arise when the maximal degree is large or the edge set is dense, or if the graph contains a cycle or not. All of these issues will be illustrated in the proofs of our results.

We also study the boundedness relation between the operators associated with a graph G and its subgraph G' with n -vertices. Throughout the paper, we always assume that the graph G and its subgraph G' are connected ordered graphs with $|G| = |G'|$ in \mathbb{F}_q^d , and two vertices x, y in G is connected if $\|x - y\| = t \neq 0$.

In Theorem 5.5, we will see that any exponents $1 \leq p_1, p_2, p_3 \leq \infty$ with $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$ satisfy that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$. Notice that P_2 can be considered as a subgraph of K_3 , and the operators Λ_{P_2} and Λ_{K_3} are related to the graphs P_2 and K_3 , respectively. Hence, in view of Theorem 5.5, one may have a question that, “Compared to a graph G , does the operator associated with its subgraph yield less restricted mapping exponents?” More precisely, one may pose the following question.

Question 1.2 Suppose that G' is a subgraph of the graph G with n vertices in \mathbb{F}_q^d . Let $1 \leq p_i \leq \infty, 1 \leq i \leq n$. If $\Lambda_G(p_1, \dots, p_n) \lesssim 1$, is it true that $\Lambda_{G'}(p_1, \dots, p_n) \lesssim 1$?

Somewhat surprisingly, the answer turns out to be no! When $G = K_3$ and $G' = P_2$, the answer to Question 1.2 is positive as Theorem 5.5 shows. However, it turns out that there exist a graph G and its subgraph G' yielding a negative answer, although the answers are positive for the most graphs which we consider in this paper. For example, the answer to Question 1.2 is negative when G is the $C_4 +$ diagonal and G' is the C_4 (see Proposition 7.6).

Since the general answer to Question 1.2 is not always positive, we pose the following natural question.

Problem 1.3 Find general properties of the graph G and its subgraph G' which yield a positive answer to Question 1.2.

The main goal of this paper is to address a conjecture on this problem and to confirm it in two dimensions. To precisely state our conjecture on the problem, let us review the standard definition and notation for the minimal degree of a graph.

Definition 1.4 The Minimum Degree of a graph G , denoted by $\delta(G)$, is defined as the degree of the vertex with the least number of edges incident to it.

We propose the following conjecture which can be a solution of Problem 1.3.

Conjecture 1.5 Let G' be a subgraph of the graph G in $\mathbb{F}_q^d, d \geq 2$, with n vertices, and let $1 \leq p_i \leq \infty, 1 \leq i \leq n$. In addition, assume that

$$(1.6) \quad \min \{\delta(G), d\} > \delta(G').$$

Then, if $\Lambda_G(p_1, \dots, p_n) \lesssim 1$, we have $\Lambda_{G'}(p_1, \dots, p_n) \lesssim 1$.

Note that the condition (1.6) in Conjecture 1.5 is equivalent to the following:

$$(1.7) \quad (i) \delta(G) > \delta(G') \quad \text{and} \quad (ii) d > \delta(G').$$

We have some comments and further questions below, regarding the above conjecture and our main theorems which we will state and prove in the body of this paper.

- Our results in this paper confirm Conjecture 1.5, possibly up to endpoints, for all graphs G and their subgraphs G' on $n = 3, 4$ vertices in \mathbb{F}_q^2 (see Theorem 10.7). Also note that when $d = 2$ and $n = 3, 4$, the condition (1.6) is equivalent to the first condition (i) in (1.7) since $\delta(G) \leq 2$ (see the figures above).
- It will be shown that the conclusion of Conjecture 1.5 cannot be reversed at least for $n = 3, 4$ in two dimensions (see Remark 5.6 for $n = 3$, and see Remarks 8.9, 8.12, 9.11, and 10.5 for $n = 4$.)

It is worth investigating whether the key hypothesis (1.6) of Conjecture 1.5 can be relaxed.

- The conclusion of Conjecture 1.5 does not hold in general if the $>$ in the assumption (1.6) is replaced by \geq . To see this, consider $G = C_4 + \text{diagonal}$ and $G' = C_4$ on 4 vertices in \mathbb{F}_q^d , with $d = 2$. It is obvious that G' is a subgraph of G and $\min\{\delta(G), d\} = \delta(G') = 2$. However, Proposition 7.6(ii) implies that the conclusion of Conjecture 1.5 is not true.
- We are not sure what can we say about the conclusion of Conjecture 1.5 if the main hypothesis (1.6) of Conjecture 1.5 is relaxed by the second one of the conditions (1.7). To be precise, when (1.6) is replaced by the second statement of (1.7), that is, $d > \delta(G')$, we do not have a definitive answer even for $n = 4$ in \mathbb{F}_q^2 . For instance, let $G = Y\text{-shape}$ and $G' = K_3 + \text{a tail on } \mathbb{F}_q^2$. Then it is clear that $d = 2 > \delta(G') = 1$ and $1 = \delta(G) = \delta(G')$ and so this provides an example that does not satisfy the assumption (1.6) of Conjecture 1.5 but satisfy the second statement of (1.7). Unfortunately, in this paper, we have not found any inclusive boundedness relations between the operators corresponding to such graphs. In order to exclude this uncertain case, both conditions in (1.7) were taken as the hypothesis for Conjecture 1.5, namely the condition (1.6).

Notation:

- We denote $\Lambda_G(p_1, \dots, p_n) \lesssim 1$ if the inequality (1.5) holds true for all characteristic functions on \mathbb{F}_q^d .
- By \mathbb{F}_q^* , we mean the set of all nonzero elements in \mathbb{F}_q .
- For $t \in \mathbb{F}_q^*$, we denote by S_t^{n-1} the sphere of radius t centered at the origin in \mathbb{F}_q^n :

$$S_t^{n-1} := \{x \in \mathbb{F}_q^n : \|x\| = t\}.$$

Unless otherwise specified in this paper, d represents the general dimension of \mathbb{F}_q^d , $d \geq 2$. When $n = d$, we write S_t instead of S_t^{d-1} for simplicity.

- We identify the set S_t with its indicator function 1_{S_t} , namely, $S_t(x) = 1_{S_t}(x)$.
- We write δ_0 for the indicator function of the set of the zero vector in \mathbb{F}_q^d .
- For positive numbers $A, B > 0$, we write $A \lesssim B$ if $A \leq CB$ for some constant $C > 0$ independent of q , the size of the underlying finite field \mathbb{F}_q . The notation $A \sim B$ means that $A \lesssim B$ and $B \lesssim A$.

The rest of this paper is organized as follows: In Section 2, we recall known results on the spherical averaging operator, which functions as a fundamental tool to prove our theorems. Sections 3–10 are devoted to the presentation and proofs of our main

results associated with the graphs mentioned above. The Appendix contains some technical lemmas on the number of intersection points of two spheres in \mathbb{F}_q^d .

2 The spherical averaging problem

In the finite field setting, Carbery, Stones, and Wright [4] initially formulated and studied the averaging problem over the varieties defined by vector-valued polynomials. This problem for general varieties was studied by Chun-Yen Shen and the third-listed author [12]. Here, we introduce the standard results on the averaging problem over the spheres. We adopt the notation in [12].

Let dx be the normalizing counting measure on \mathbb{F}_q^d . For each nonzero t , we endow the sphere S_t with the normalizing surface measure $d\sigma_t$. We recall that

$$d\sigma_t(x) = \frac{q^d}{|S_t|} 1_{S_t}(x) dx$$

so that we can identify the measure $d\sigma_t$ with the function $\frac{q^d}{|S_t|} 1_{S_t}$ on \mathbb{F}_q^d .

The spherical averaging operator A_{S_t} is defined by

$$(2.1) \quad A_{S_t}f(x) = f * d\sigma_t(x) = \int_{S_t} f(x - y) d\sigma_t(y) = \frac{1}{|S_t|} \sum_{y \in S_t} f(x - y),$$

where f is a function on \mathbb{F}_q^d . By a change of variables, we also have

$$(2.2) \quad A_{S_t}f(x) = \frac{1}{|S_t|} \sum_{y \in \mathbb{F}_q^d} S_t(x - y) f(y).$$

For $1 \leq p, r \leq \infty$, we define $A_{S_t}(p \rightarrow r)$ to be the smallest number such that the averaging estimate

$$(2.3) \quad \|f * d\sigma_t\|_{L^r(\mathbb{F}_q^d, dx)} \leq A_{S_t}(p \rightarrow r) \|f\|_{L^p(\mathbb{F}_q^d, dx)}$$

holds for all functions f on \mathbb{F}_q^d .

Problem 2.1 (Spherical averaging problem) *Determine all exponents $1 \leq p, r \leq \infty$ such that*

$$A_{S_t}(p \rightarrow r) \lesssim 1.$$

Notation 2.2 *From now on, we simply write A for the spherical averaging operator A_{S_t} .*

By testing (2.3) with $f = \delta_0$ and by using the duality of the averaging operator, it is not hard to notice that the necessary conditions for the boundedness of $A(p \rightarrow r)$ are as follows: $(1/p, 1/r)$ is contained in the convex hull of points $(0, 0)$, $(0, 1)$, $(1, 1)$, and $(\frac{d}{d+1}, \frac{1}{d+1})$.

Using the Fourier decay estimate on S_t and its cardinality, it can be shown that these necessary conditions are sufficient. For the reader's convenience, we give a detail proof although the argument is standard, as is well known in the literature such as [4, 12].

Theorem 2.3 *Let $1 \leq p, r \leq \infty$ be numbers such that $(1/p, 1/r)$ lies on the convex hull of points $(0, 0)$, $(0, 1)$, $(1, 1)$, and $(\frac{d}{d+1}, \frac{1}{d+1})$. Then we have $A(p \rightarrow r) \lesssim 1$.*

Proof Since both $d\sigma_t$ and dx have total mass 1, it follows from Young's inequality for convolution functions that if $1 \leq r \leq p \leq \infty$, then

$$(2.4) \quad \|f * d\sigma_t\|_{L^r(\mathbb{F}_q^d, dx)} \leq \|f\|_{L^p(\mathbb{F}_q^d, dx)}.$$

We notice that these results do not hold for the Euclidean Averaging problem.

By the interpolation and the duality, we only need to establish the following critical estimate:

$$A\left(\frac{d+1}{d} \rightarrow d+1\right) \lesssim 1.$$

It is well known that for nonzero t ,

$$|(d\sigma_t)^\vee(m)| := \left| \frac{1}{|S_t|} \sum_{x \in S_t} \chi(m \cdot x) \right| \lesssim q^{-\frac{(d-1)}{2}} \quad \text{for all } m \neq (0, \dots, 0),$$

where χ denotes a nontrivial additive character of \mathbb{F}_q (see the proof of Lemma 2.2 in [10]).

Since $|S_t| \sim q^{d-1}$, we complete the proof by combining this Fourier decay estimate with the following well-known lemma (see Lemma 6.1 in [12]).

Lemma 2.4 *Let $d\sigma$ be the normalized surface measure on an variety S in \mathbb{F}_q^d with $|S| \sim q^{d-1}$. If $|(d\sigma)^\vee(m)| \lesssim q^{-\frac{k}{2}}$ for all $m \in \mathbb{F}_q^d \setminus (0, \dots, 0)$ and for some $k > 0$, then we have*

$$A\left(\frac{k+2}{k+1} \rightarrow k+2\right) \lesssim 1. \quad \blacksquare$$

The boundary points of the convex hull play an important role in the application of Theorem 2.3. More precisely, we will apply the following result, which is a direct consequence of Theorem 2.3.

Lemma 2.5 *Let $1 \leq p, r \leq \infty$, and let A denote the averaging operator over the sphere S_t , $t \neq 0$, in \mathbb{F}_q^d , $d \geq 2$.*

- (i) *If $1 \leq \frac{1}{p} \leq \frac{d}{d+1}$ and $\frac{1}{r} = \frac{1}{dp}$, then $A(p \rightarrow r) \lesssim 1$.*
- (ii) *If $\frac{d}{d+1} \leq \frac{1}{p} \leq 1$ and $\frac{1}{r} = \frac{d}{p} - d + 1$, then $A(p \rightarrow r) \lesssim 1$.*

3 Sharp mapping properties for the K_2 form

In this section, we provide the sharp mapping properties of the operator associated with the graph K_2 . To this end, as described below, we relate the problem to the spherical averaging problem.

As usual, the inner product of the nonnegative real-valued functions f, g on \mathbb{F}_q^d is defined as

$$\langle f, g \rangle := \|fg\|_1 = \frac{1}{q^d} \sum_{x \in \mathbb{F}_q^d} f(x)g(x).$$

Let $t \in \mathbb{F}_q^*$, and let f_1, f_2 be nonnegative real-valued functions on \mathbb{F}_q^d , $d \geq 2$. Then the K_2 form Λ_{K_2} on \mathbb{F}_q^d is defined by

$$(3.1) \quad \Lambda_{K_2}(f_1, f_2) = \frac{1}{q^d |S_t|} \sum_{x^1, x^2 \in \mathbb{F}_q^d} S_t(x^1 - x^2) f_1(x^1) f_2(x^2).$$

Here, the quantity $q^d |S_t|$ represents the normalizing factor $\mathcal{N}(G)$ in (1.4) when the graph G is K_2 . In other words, $\mathcal{N}(G) = q^d |S_t|$, which is the number of the pair $(x^1, x^2) \in \mathbb{F}_q^d \times \mathbb{F}_q^d$ such that $S_t(x^1 - x^2) = 1$.

By a change of variables, we can write

$$(3.2) \quad \Lambda_{K_2}(f_1, f_2) = \frac{1}{q^d} \sum_{x^1 \in \mathbb{F}_q^d} f_1(x^1) \left(\frac{1}{|S_t|} \sum_{x^2 \in \mathbb{F}_q^d} f_2(x^1 - x^2) S_t(x^2) \right) = \langle f_1, A f_2 \rangle,$$

where A denotes the averaging operator related to the sphere S_t . Likewise, we also obtain that $\Lambda_{K_2}(f_1, f_2) = \langle A f_1, f_2 \rangle$.

The main goal of this section is to address all numbers $1 \leq p_1, p_2 \leq \infty$ satisfying $\Lambda_{K_2}(p_1, p_2) \lesssim 1$.

We begin with the necessary conditions for the boundedness of the K_2 form Λ_{K_2} on \mathbb{F}_q^d .

Proposition 3.1 *Let $1 \leq p_1, p_2 \leq \infty$. Suppose that $\Lambda_{K_2}(p_1, p_2) \lesssim 1$. Then we have*

$$\frac{1}{p_1} + \frac{d}{p_2} \leq d \quad \text{and} \quad \frac{d}{p_1} + \frac{1}{p_2} \leq d.$$

Proof By symmetry, it is clear that $\Lambda_{K_2}(p_1, p_2) \lesssim 1 \iff \Lambda_{K_2}(p_2, p_1) \lesssim 1$. Hence, it suffices to prove the first listed conclusion that $\frac{1}{p_1} + \frac{d}{p_2} \leq d$.

From (3.2) and our assumption that $\Lambda_{K_2}(p_1, p_2) \lesssim 1$, we must have

$$\Lambda_{K_2}(f_1, f_2) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2}.$$

We test this inequality with $f_1 = 1_{S_t}$ and $f_2 = \delta_0$. Then

$$\Lambda_{K_2}(f_1, f_2) = \frac{1}{q^d} \sum_{x^1 \in S_t} \frac{1}{|S_t|} = q^{-d},$$

and

$$\|f_1\|_{p_1} \|f_2\|_{p_2} \sim (q^{-d} |S_t|)^{1/p_1} (q^{-d})^{1/p_2} \sim q^{-\frac{1}{p_1} - \frac{d}{p_2}}.$$

By a direct comparison, we get the desired result. \blacksquare

Remark 3.2 For $1 \leq p_1, p_2 \leq \infty$, one can note that $\frac{1}{p_1} + \frac{d}{p_2} \leq d$ and $\frac{d}{p_1} + \frac{1}{p_2} \leq d$ if and only if $(1/p_1, 1/p_2) \in [0, 1] \times [0, 1]$ lies on the convex hull of points $(0, 0)$, $(0, 1)$, $(\frac{d}{d+1}, \frac{d}{d+1})$, $(1, 0)$.

Let us move to the sufficient conditions on the exponents $1 \leq p_1, p_2 \leq \infty$ such that $\Lambda_{K_2}(p_1, p_2) \lesssim 1$. We now show that the necessary conditions are in fact sufficient conditions for $\Lambda_{K_2}(p_1, p_2) \lesssim 1$.

Theorem 3.3 (Sharp boundedness result for the K_2 form on \mathbb{F}_q^d) *Let $1 \leq p_1, p_2 \leq \infty$. Then we have*

$$\Lambda_{K_2}(p_1, p_2) \lesssim 1 \quad \text{if and only if} \quad \frac{1}{p_1} + \frac{d}{p_2} \leq d, \quad \frac{d}{p_1} + \frac{1}{p_2} \leq d.$$

Proof By Proposition 3.1, it will be enough to prove that $\Lambda_{K_2}(p_1, p_2) \lesssim 1$ for all $1 \leq p_1, p_2 \leq \infty$ satisfying

$$\frac{1}{p_1} + \frac{d}{p_2} \leq d, \quad \frac{d}{p_1} + \frac{1}{p_2} \leq d.$$

By the interpolation theorem and the nesting property of the norm, it suffices to establish the estimates on the critical endpoints $(1/p_1, 1/p_2) \in [0, 1] \times [0, 1]$, which are $(0, 1)$, $(1, 0)$, and $(d/(d+1), d/(d+1))$. In other words, it remains to prove the following estimates:

$$\Lambda_{K_2}(\infty, 1) \lesssim 1, \quad \Lambda_{K_2}(1, \infty) \lesssim 1, \quad \Lambda_{K_2}\left(\frac{d+1}{d}, \frac{d+1}{d}\right) \lesssim 1.$$

Since $\Lambda_{K_2}(f_1, f_2) = \langle f_1, Af_2 \rangle$, it follows by Hölder's inequality that if $A(p_2 \rightarrow p'_1) \lesssim 1$ with $1 \leq p_1, p_2 \leq \infty$, then $\Lambda_{K_2}(p_1, p_2) \lesssim 1$. Thus, matters are reduced to establishing the following averaging estimates:

$$A(1 \rightarrow 1) \lesssim 1, \quad A(\infty \rightarrow \infty) \lesssim 1, \quad A\left(\frac{d+1}{d} \rightarrow d+1\right) \lesssim 1.$$

However, these averaging estimates are clearly valid by Theorem 2.3, and thus the proof is complete. \blacksquare

The following result is a special case of Theorem 3.3, but it is very useful in practice.

Corollary 3.4 *For any dimensions $d \geq 2$, we have $\Lambda_{K_2}\left(\frac{d+1}{d}, \frac{d+1}{d}\right) \lesssim 1$.*

Proof Notice that if $p_1 = p_2 = \frac{d+1}{d}$, then it satisfies that $\frac{1}{p_1} + \frac{d}{p_2} \leq d$ and $\frac{d}{p_1} + \frac{1}{p_2} \leq d$. Hence, the statement follows immediately from Theorem 3.3. \blacksquare

4 Boundedness problem for the K_3 form

Let $t \in \mathbb{F}_q^*$. The K_3 form Λ_{K_3} on \mathbb{F}_q^d can be defined as

(4.1)

$$\Lambda_{K_3}(f_1, f_2, f_3) = \frac{1}{q^d} \frac{1}{|S_t| |S_t^{d-2}|} \sum_{x^1, x^2, x^3 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) S_t(x^3 - x^1) \prod_{i=1}^3 f_i(x^i),$$

where each f_i , $i = 1, 2, 3$, is a nonnegative real-valued function on \mathbb{F}_q^d , and the quantity $q^d |S_t| |S_t^{d-2}|$ stands for the normalizing factor $\mathcal{N}(G)$ in (1.4) when $G = K_3$. More precisely, $\mathcal{N}(G)$ is the number of $(x^1, x^2, x^3) \in (\mathbb{F}_q^d)^3$ such that $S_t(x^1 - x^2) S_t(x^2 - x^3) S_t(x^3 - x^1) = 1$.

The purpose of this section is to find the numbers $1 \leq p_1, p_2, p_3 \leq \infty$ such that $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$.

When the dimension d is 2, we will settle this problem up to the endpoint estimate. To this end, we relate our problem to the estimate of the Bilinear Averaging Operator (see (4.3)) for which we establish the sharp bound.

On the other hand, as we shall see, in three and higher dimensions $d \geq 3$, it is not easy to deduce the sharp results. However, when one of the exponents p_1, p_2, p_3 is ∞ , we will be able to obtain the optimal results. This will be done by applying Theorem 3.3, the boundedness result for the K_2 form Λ_{K_2} on \mathbb{F}_q^d .

We begin by deducing necessary conditions for our problem in \mathbb{F}_q^d , $d \geq 2$. Recall that for $d = 2$, we pose an additional restriction that $3 \in \mathbb{F}_q$ is a square number.

Proposition 4.1 (Necessary conditions for the boundedness of Λ_{K_3}) *Let $1 \leq p_1, p_2, p_3 \leq \infty$. Suppose that $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$. Then we have*

$$\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} \leq d, \quad \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d, \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq d.$$

In particular, when $d = 2$, it can be shown by Polymake¹ [1, 6] that $(1/p_1, 1/p_2, 1/p_3)$ is contained in the convex hull of the points: $(0, 0, 1), (0, 1, 0), (2/3, 2/3, 0), (1/2, 1/2, 1/2), (2/3, 0, 2/3), (1, 0, 0), (0, 0, 0), (0, 2/3, 2/3)$.

Proof We only prove the first inequality in the conclusion since we can establish other inequalities by symmetric property of $\Lambda_{K_3} f_1, f_2, f_3$. We will use the simple fact that $x \in S_t$ if and only if $-x \in S_t$. In the definition (4.1), taking $f_1 = \delta_0, f_2 = 1_{S_t}$, and $f_3 = 1_{S_t}$, we see that

$$\|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \sim q^{-\frac{d}{p_1}} q^{-\frac{1}{p_2}} q^{-\frac{1}{p_3}},$$

$$\Lambda_{K_3}(f_1, f_2, f_3) = \frac{1}{q^d} \frac{1}{|S_t| |S_t^{d-2}|} \left(\sum_{x^2, x^3 \in S_t: \|x^2 - x^3\| = t} 1 \right) \sim q^{-d},$$

where the last similarity above follows from Corollary A.4 in the Appendix with our assumption that $3 \in \mathbb{F}_q$ is a square number for $d = 2$.

By the direct comparison of these estimates, we obtain the required necessary condition. ■

Remark 4.2 In order to prove that the necessary conditions in Proposition 4.1 are sufficient conditions for $d = 2$, we only need to establish the following critical endpoint estimates: $\Lambda_{K_3}(2, 2, 2) \lesssim 1, \Lambda_{K_3}(\infty, \infty, \infty) \lesssim 1, \Lambda_{K_3}(1, \infty, \infty) \lesssim 1, \Lambda_{K_3}(\infty, 1, \infty) \lesssim 1, \Lambda_{K_3}(\infty, \infty, 1) \lesssim 1, \Lambda_{K_3}(\frac{3}{2}, \frac{3}{2}, \infty) \lesssim 1, \Lambda_{K_3}(\frac{3}{2}, \infty, \frac{3}{2}) \lesssim 1, \Lambda_{K_3}(\infty, \frac{3}{2}, \frac{3}{2}) \lesssim 1$. In fact, this claim follows by interpolating the critical points given in the second part of Proposition 4.1.

4.1 Boundedness results for Λ_{K_3} on \mathbb{F}_q^d

The graph K_2 can be obtained by removing any one of three vertices in the graph K_3 . Therefore, the boundedness of $\Lambda_{K_2}(p_1, p_2)$ can determine the boundedness of

¹Polymake is software for the algorithmic treatment of convex polyhedra.

$\Lambda_{K_3}(p_1, p_2, \infty)$. Using this observation, in the case when one of p_1, p_2, p_3 is ∞ , we are able to obtain sharp boundedness results for $\Lambda_{K_3}(p_1, p_2, p_3)$.

Theorem 4.3 *Let $1 \leq a, b \leq \infty$ satisfy that $\frac{1}{a} + \frac{d}{b} \leq d$ and $\frac{d}{a} + \frac{1}{b} \leq d$. Then we have $\Lambda_{K_3}(a, b, \infty) \lesssim 1$, $\Lambda_{K_3}(a, \infty, b) \lesssim 1$, and $\Lambda_{K_3}(\infty, a, b) \lesssim 1$.*

Proof The statement of the theorem follows immediately by combining Theorem 3.3 and the following claim: If $\Lambda_{K_2}(a, b) \lesssim 1$, then

$$\Lambda_{K_3}(a, b, \infty) \lesssim 1, \quad \Lambda_{K_3}(a, \infty, b) \lesssim 1, \quad \text{and} \quad \Lambda_{K_3}(\infty, a, b) \lesssim 1.$$

It suffices by symmetry to prove that if $\Lambda_{K_2}(a, b) \lesssim 1$, then $\Lambda_{K_3}(a, b, \infty) \lesssim 1$.

Since $f_i, i = 1, 2, 3$, are nonnegative real-number functions on \mathbb{F}_q^d , it follows that

$$\Lambda_{K_3}(f_1, f_2, f_3) \leq \frac{1}{|S_t^{d-2}|} \left(\max_{\|x^1 - x^2\| = t} \sum_{x^3 \in \mathbb{F}_q^d: \|x^2 - x^3\| = t = \|x^3 - x^1\|} f_3(x^3) \right) \Lambda_{K_2}(f_1, f_2).$$

Since $|S_t^{d-2}| \sim q^{d-2}$ and $\Lambda_{K_2}(f_1, f_2) \lesssim \|f_1\|_a \|f_2\|_b$, it suffices to prove that the maximum value in the above parenthesis is $\lesssim q^{d-2} \|f_3\|_\infty$. Let us denote by I the maximum above.

By a change of variables, $x = x^1, y = x^1 - x^2$, we see that

$$I \leq \left(\max_{x \in \mathbb{F}_q^d, y \in S_t} \sum_{x^3 \in \mathbb{F}_q^d: \|x - y - x^3\| = t = \|x^3 - x\|} 1 \right) \|f_3\|_\infty.$$

By another change of variables by putting $z = x - x^3$, we get

$$I \leq \left(\max_{x \in \mathbb{F}_q^d, y \in S_t} \sum_{z \in S_t: \|z - y\| = t} 1 \right) \|f_3\|_\infty = \left(\max_{y \in S_t} \sum_{z \in S_t: \|z - y\| = t} 1 \right) \|f_3\|_\infty.$$

Now, applying Corollary A.4 in the Appendix, we conclude that $I \lesssim q^{d-2} \|f_3\|_\infty$ as required. \blacksquare

It is not hard to see from Proposition 4.1 that Theorem 4.3 cannot be improved in the case when one of the exponents p_1, p_2, p_3 is ∞ . In particular, the following critical endpoint estimates follows immediately from Theorem 4.3:

$$\begin{aligned} \Lambda_{K_3}\left(\frac{d+1}{d}, \frac{d+1}{d}, \infty\right) &\lesssim 1, \quad \Lambda_{K_3}\left(\frac{d+1}{d}, \infty, \frac{d+1}{d}\right) \lesssim 1, \quad \Lambda_{K_3}\left(\infty, \frac{d+1}{d}, \frac{d+1}{d}\right) \lesssim 1, \\ \Lambda_{K_3}(\infty, \infty, \infty) &\lesssim 1, \quad \Lambda_{K_3}(1, \infty, \infty) \lesssim 1, \quad \Lambda_{K_3}(\infty, 1, \infty) \lesssim 1, \quad \Lambda_{K_3}(\infty, \infty, 1) \lesssim 1. \end{aligned} \tag{4.2}$$

Remark 4.4 From (4.2) and Remark 4.2, we see that to completely solve the problem on the boundedness of $\Lambda_{K_3}(p_1, p_2, p_3)$ for $d = 2$, we only need to establish the following critical endpoint estimate

$$\Lambda_{K_3}\left(\frac{d+2}{d}, \frac{d+2}{d}, \frac{d+2}{d}\right) = \Lambda_{K_3}(2, 2, 2) \lesssim 1.$$

4.2 Sharp restricted strong-type estimates in two dimensions

Although Theorem 4.3 is valid for all dimensions $d \geq 2$, it is not sharp, compared to the necessary conditions given in Proposition 4.1. In this subsection, we will deduce the sharp boundedness results up to the endpoints for $\Lambda_{K_3}(p_1, p_2, p_3)$ in two dimensions. To this end, we need the following theorem, which can be proven by modifying the Euclidean argument introduced in Section 7 of [7].

Theorem 4.5 *Let Λ_{K_3} be the K_3 form on \mathbb{F}_q^2 . Then, for all subsets E, F, H of \mathbb{F}_q^2 , the following estimate holds: $\Lambda_{K_3}(E, F, H) \lesssim \|E\|_2 \|F\|_2 \|H\|_2$.*

For $1 \leq p_1, p_2, p_3 \leq \infty$, we say that the restricted strong-type $\Lambda_{K_3}(p_1, p_2, p_3)$ estimate holds if the estimate

$$\Lambda_{K_3}(E, F, H) \lesssim \|E\|_{p_1} \|F\|_{p_2} \|H\|_{p_3}$$

is valid for all subsets E, F, H of \mathbb{F}_q^2 . In this case, we write $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$.

Proof The proof proceeds with some reduction. When $d = 2$, by a change of variables by letting $x = x^3, y = x^3 - x^1, z = x^3 - x^2$, (4.1) becomes

$$\Lambda_{K_3}(f_1, f_2, f_3) = \frac{1}{q^2} \sum_{x \in \mathbb{F}_q^2} f_3(x) \left[\frac{1}{|S_t|} \sum_{y, z \in \mathbb{F}_q^2} S_t(z - y) S_t(z) S_t(y) f_1(x - y) f_2(x - z) \right].$$

We define $B(f_1, f_2)(x)$ as the value in the bracket above, namely,

$$(4.3) \quad B(f_1, f_2)(x) := \frac{1}{|S_t|} \sum_{y, z \in S_t: \|z - y\| = t} f_1(x - y) f_2(x - z).$$

We refer to this operator B as “the bilinear averaging operator.” It is clear that

$$\Lambda_{K_3}(f_1, f_2, f_3) = \frac{1}{q^2} \sum_{x \in \mathbb{F}_q^2} B(f_1, f_2)(x) f_3(x) = \langle B(f_1, f_2), f_3 \rangle.$$

By Hölder’s inequality, we have

$$\Lambda_{K_3}(f_1, f_2, f_3) \leq \|B(f_1, f_2)\|_2 \|f_3\|_2.$$

Thus, Theorem 4.5 follows immediately from the reduction lemma below ■

Lemma 4.6 *Let $B(f_1, f_2)$ be the bilinear averaging operator defined as in (4.3). Then, for all subsets E, F of \mathbb{F}_q^2 , we have*

$$\|B(E, F)\|_2 \lesssim \|E\|_2 \|F\|_2.$$

Proof We begin by representing the bilinear averaging operator $B(f_1, f_2)$. From (4.3), note that

$$B(f_1, f_2)(x) = \frac{1}{|S_t|} \sum_{y \in S_t} f_1(x - y) \left(\sum_{z \in S_t: \|z - y\| = t} f_2(x - z) \right).$$

For each $y \in S_t$, let $\Theta(y) := \{z \in S_t : \|z - y\| = t\}$. With this notation, the bilinear averaging operator is written as

$$B(f_1, f_2)(x) = \frac{1}{|S_t|} \sum_{y \in S_t} f_1(x - y) \left(\sum_{z \in \Theta(y)} f_2(x - z) \right).$$

Let η denote the quadratic character of \mathbb{F}_q^* . Recall that $\eta(s) = 1$ for a square number s in \mathbb{F}_q^* , and $\eta(s) = -1$ otherwise. Notice from Corollary A.4 in the Appendix that $\Theta(y)$ is the empty set for all $y \in S_t$ if $d = 2$ and $\eta(3) = -1$. In this case, the problem is trivial since $B(f_1, f_2)(x) = 0$ for all $x \in \mathbb{F}_q^2$. Therefore, when $d = 2$, we always assume that $\eta(3) = 1$.

Notice that $|\Theta(y)| = 2$ for all $y \in S_t$, which follows from the last statement of Corollary A.4 in the Appendix. More precisely, for each $y \in S_t$, we can write

$$\Theta(y) = \{\theta y, \theta^{-1}y\},$$

where θy denotes the rotation of y by “60 degrees,” and $\|y - \theta y\| = t = \|y - \theta^{-1}y\|$.

From these observations, the bilinear averaging operator $B(f_1, f_2)$ can be represented as follows:

$$(4.4) \quad B(f_1, f_2)(x) = B_\theta(f_1, f_2)(x) + B_{\theta^{-1}}(f_1, f_2)(x).$$

Here, we define

$$(4.5) \quad B_\theta(f_1, f_2)(x) := \frac{1}{|S_t|} \sum_{y \in S_t} f_1(x - y) f_2(x - \theta y),$$

and

$$B_{\theta^{-1}}(f_1, f_2) := \frac{1}{|S_t|} \sum_{y \in S_t} f_1(x - y) f_2(x - \theta^{-1}y).$$

In order to complete the proof of the lemma, it suffices to establish the following two estimates: for all subsets E, F of \mathbb{F}_q^d ,

$$(4.6) \quad \|B_\theta(E, F)\|_2 \lesssim \|E\|_2 \|F\|_2,$$

and

$$(4.7) \quad \|B_{\theta^{-1}}(E, F)\|_2 \lesssim \|E\|_2 \|F\|_2.$$

We will only provide the proof of the estimate (4.6) since the proof of (4.7) is the same.

Now we start proving the estimate (4.6). Since $\|E\|_2^2 = q^{-2}|E|$ and $\|F\|_2^2 = q^{-2}|F|$, it is enough to prove that

$$(4.8) \quad \|B_\theta(E, F)\|_2^2 \lesssim q^{-4}|E||F|.$$

Without loss of generality, we may assume that $|E| \leq |F|$. By the definition, it follows that

$$\|B_\theta(E, F)\|_2^2 = q^{-2}|S_t|^{-2} \sum_{x \in \mathbb{F}_q^2} \sum_{y, y' \in S_t} E(x - y) E(x - y') F(x - \theta y) F(x - \theta y') = I + II,$$

where the first term I is the value corresponding to the case where $y = y'$, whereas the second term II is corresponding to the case where $y \neq y'$. We have

$$I = q^{-2}|S_t|^{-2} \sum_{y \in S_t} \sum_{x \in \mathbb{F}_q^2} E(x - y)F(x - \theta y).$$

Applying a change of variables by replacing x with $x + y$, we see that

$$I = q^{-2}|S_t|^{-2} \sum_{y \in S_t} \sum_{x \in \mathbb{F}_q^2} E(x)F(x + y - \theta y) = q^{-2}|S_t|^{-2} \sum_{x \in E} \left(\sum_{y \in S_t} F(x + y - \theta y) \right).$$

Observe that $y - \theta y \neq y' - \theta y'$ for all y, y' in S_t with $y \neq y'$. Then we see that the value in the parentheses above is bounded above by $|S_t \cap F| \leq |F|$. Therefore, we obtain the desired estimate:

$$I \leq q^{-2}|S_t|^{-2}|E||F| \sim q^{-4}|E||F|.$$

Next, it remains to show that $II \lesssim q^{-4}|E||F|$. Since we have assumed that $|E| \leq |F|$, it suffices to show that $II \lesssim q^{-4}|E|^2$.

By the definition of II , it follows that

$$II = q^{-2}|S_t|^{-2} \sum_{x \in \mathbb{F}_q^2} \sum_{y, y' \in S_t; y \neq y'} E(x - y)E(x - y')F(x - \theta y)F(x - \theta y').$$

It is obvious that

$$II \leq q^{-2}|S_t|^{-2} \sum_{y, y' \in S_t; y \neq y'} \sum_{x \in \mathbb{F}_q^2} E(x - y)E(x - y').$$

We use a change of variables by replacing x with $x + y$. Then we have

$$\begin{aligned} II &\leq q^{-2}|S_t|^{-2} \sum_{x \in \mathbb{F}_q^2} E(x) \left(\sum_{y, y' \in S_t; y \neq y'} E(x + y - y') \right) \\ &= q^{-2}|S_t|^{-2} \sum_{x \in \mathbb{F}_q^2} E(x) \left(\sum_{0 \neq u \in \mathbb{F}_q^2} E(x + u)W(u) \right), \end{aligned}$$

where $W(u)$ denotes the number of pairs $(y, y') \in S_t \times S_t$ such that $u = y - y'$ and $y \neq y'$. It is not hard to see that for any nonzero vector $u \in \mathbb{F}_q^2$, we have $W(u) \leq 2$. So we obtain that

$$II \lesssim q^{-2}|S_t|^{-2} \sum_{x \in \mathbb{F}_q^2} E(x) \left(\sum_{0 \neq u \in \mathbb{F}_q^2} E(x + u) \right) \lesssim q^{-4}|E|^2,$$

as required. ■

In two dimensions, we are able to obtain the optimal boundedness of $\Lambda_{K_3}(p_1, p_2, p_3)$ except for one endpoint. Indeed, we have the following result.

Theorem 4.7 Let $1 \leq p_1, p_2, p_3 \leq \infty$, and let Λ_{K_3} be the K_3 form on \mathbb{F}_q^2 .

(i) If $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$, then

$$(4.9) \quad \frac{2}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} \leq 2, \quad \frac{1}{p_1} + \frac{2}{p_2} + \frac{1}{p_3} \leq 2, \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{2}{p_3} \leq 2.$$

(ii) Conversely, if (p_1, p_2, p_3) satisfies all three inequalities (4.9), then $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$ for $(p_1, p_2, p_3) \neq (2, 2, 2)$, and we have $\Lambda_{K_3}(2, 2, 2) \lesssim 1$.

Proof The first part of the theorem is the special case of Proposition 4.1 with $d = 2$. Now we prove the second part. As stated in Proposition 4.1, one can notice by using Polymake [1, 6] that all the points $\left(\frac{1}{p_1}, \frac{1}{p_2}, \frac{1}{p_3}\right) \in [0, 1]^3$ satisfying all three inequalities (4.9) are contained in the convex hull of the critical points

$$(1/2, 1/2, 1/2), (0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1), \left(\frac{2}{3}, \frac{2}{3}, 0\right), \left(\frac{2}{3}, 0, \frac{2}{3}\right), \left(0, \frac{2}{3}, \frac{2}{3}\right).$$

Notice from Theorem 4.5 with $d = 2$ that the restricted strong-type estimate for the operator Λ_{K_3} holds for the point $(1/p_1, 1/p_2, 1/p_3) = (1/2, 1/2, 1/2)$. In addition, notice from the estimates (4.2) with $d = 2$ that $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$ for the above critical points $(1/p_1, 1/p_2, 1/p_3)$ except for $(1/2, 1/2, 1/2)$. Hence, the statement of the second part follows immediately by invoking the interpolation theorem. ■

5 Boundedness results for the P_2 form

For $t \in \mathbb{F}_q^*$ and functions f_i , $i = 1, 2, 3$, on \mathbb{F}_q^d , the P_2 form Λ_{P_2} on \mathbb{F}_q^d is defined by

$$(5.1) \quad \Lambda_{P_2}(f_1, f_2, f_3) = \frac{1}{q^d |S_t|^2} \sum_{x^1, x^2, x^3 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) f_1(x^1) f_2(x^2) f_3(x^3),$$

where the quantity $q^d |S_t|^2$ stands for the normalizing factor $\mathcal{N}(G)$ in (1.4) when $G = P_2$. Note that this can be written as

$$(5.2) \quad \Lambda_{P_2}(f_1, f_2, f_3) = \langle f_2, A f_1 \cdot A f_3 \rangle.$$

In this section, we study the problem determining all numbers $1 \leq p_1, p_2, p_3 \leq \infty$ satisfying $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$. Compared to the K_3 form Λ_{K_3} , this problem is much hard to find the optimal answers. Based on the formula (5.2) with the averaging estimates in Lemma 2.5, we are able to address partial results on this problem (see Theorem 5.3).

Proposition 5.1 (Necessary conditions for the boundedness of Λ_{P_2}) Let $1 \leq p_1, p_2, p_3 \leq \infty$. Suppose that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$. Then we have

$$\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d, \quad \frac{d}{p_1} + \frac{1}{p_2} \leq d, \quad \frac{1}{p_2} + \frac{d}{p_3} \leq d, \quad \frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq 2d - 1.$$

Also, under this assumption when $d = 2$, it can be shown by Polymake [1, 6] that $(1/p_1, 1/p_2, 1/p_3)$ is contained in the convex hull of points: $(0, 1, 0), (1/2, 0, 1), (1, 0, 1/2), (1, 0, 0), (5/6, 1/3, 1/2), (1/2, 1/3, 5/6), (2/3, 2/3, 0), (0, 2/3, 2/3), (0, 0, 0), (0, 0, 1)$.

Proof Suppose that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$. Then, for all functions $f_i, i = 1, 2, 3$, on \mathbb{F}_q^d , we have

$$\begin{aligned}\Lambda_{P_2}(f_1, f_2, f_3) &= \frac{1}{q^d |S_t|^2} \sum_{x^1, x^2, x^3 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) f_1(x^1) f_2(x^2) f_3(x^3) \\ &\lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3}.\end{aligned}$$

We test the above inequality with $f_2 = \delta_0, f_1 = f_3 = 1_{S_t}$. It is plain to note that

$$\|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} = (q^{-d} |S_t|)^{\frac{1}{p_1}} q^{-\frac{d}{p_2}} (q^{-d} |S_t|)^{\frac{1}{p_3}} \sim q^{-\frac{1}{p_1} - \frac{d}{p_2} - \frac{1}{p_3}},$$

and

$$\Lambda_{P_2}(f_1, f_2, f_3) = q^{-d}.$$

Therefore, we obtain that $q^{-d} \lesssim q^{-\frac{1}{p_1} - \frac{d}{p_2} - \frac{1}{p_3}}$. This implies the first inequality in the conclusion that $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d$.

To obtain the second inequality in the conclusion, we choose $f_1 = \delta_0, f_2 = 1_{S_t}$, and $f_3 = 1_{\mathbb{F}_q^d}$. Then it is easy to check that

$$\|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \sim q^{-\frac{d}{p_1} - \frac{1}{p_2}},$$

and

$$\Lambda_{P_2}(f_1, f_2, f_3) = \frac{1}{q^d |S_t|^2} \sum_{x^2 \in S_t} \sum_{x^3 \in \mathbb{F}_q^d} S_t(x^2 - x^3) = q^{-d}.$$

Comparing these estimates gives the second inequality in the conclusion.

The third inequality in the conclusion can be easily obtained by switching the roles of f_1 and f_3 in the proof of the second one.

To deduce the last inequality in the conclusion, we take $f_1 = f_3 = \delta_0$ and $f_2 = 1_{S_t}$. Then

$$\|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \sim q^{-\frac{d}{p_1} - \frac{1}{p_2} - \frac{d}{p_3}},$$

and

$$\Lambda_{P_2}(f_1, f_2, f_3) = \frac{1}{q^d |S_t|} \sim q^{-2d+1}.$$

From these, we have the required result that $\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq 2d - 1$. ■

By symmetry, it is not hard to note that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1 \iff \Lambda_{P_2}(p_3, p_2, p_1) \lesssim 1$. In the following lemma, we prove that the boundedness question for the P_2 form Λ is closely related to the spherical averaging problem over finite fields.

Lemma 5.2 Suppose that $\frac{1}{r_1} + \frac{1}{p_2} + \frac{1}{r_3} = 1, A(p_1 \rightarrow r_1) \lesssim 1$, and $A(p_3 \rightarrow r_3) \lesssim 1$ for some $1 \leq p_1, p_2, p_3, r_1, r_3 \leq \infty$. Then we have $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$.

Proof Since $\Lambda_{P_2}(f_1, f_2, f_3) = \langle f_2, A f_1 \cdot A f_3 \rangle$, we obtain by Hölder's inequality with the first assumption that

$$\Lambda_{P_2}(f_1, f_2, f_3) \leq \|A f_1\|_{r_1} \|f_2\|_{p_2} \|A f_3\|_{r_3} \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3},$$

where the averaging assumption was used for the last inequality. Hence, $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$, as required. \blacksquare

Now we state and prove our boundedness results of $\Lambda_{P_2}(p_1, p_2, p_3)$ on \mathbb{F}_q^d .

Theorem 5.3 *Let $1 \leq p_1, p_2, p_3 \leq \infty$. Then, for the P_2 form Λ_{P_2} on \mathbb{F}_q^d , the following four statements hold:*

- (i) *If $0 \leq \frac{1}{p_1}, \frac{1}{p_3} \leq \frac{d}{d+1}$ and $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d$, then $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$.*
- (ii) *If $0 \leq \frac{1}{p_1} \leq \frac{d}{d+1} \leq \frac{1}{p_3} \leq 1$ and $\frac{1}{dp_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq d$, then $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$.*
- (iii) *If $0 \leq \frac{1}{p_3} \leq \frac{d}{d+1} \leq \frac{1}{p_1} \leq 1$ and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{dp_3} \leq d$, then $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$.*
- (iv) *If $\frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_3} \leq 1$ and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq 2d - 1$, then $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$.*

Proof We proceed as follows.

- (i) By the nesting property of the norm, it suffices to prove it in the case when $0 \leq \frac{1}{p_1}, \frac{1}{p_3} \leq \frac{d}{d+1}$ and $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} = d$. This equation can be rewritten as $\frac{1}{dp_1} + \frac{1}{p_2} + \frac{1}{dp_3} = 1$. Since $0 \leq \frac{1}{p_1}, \frac{1}{p_3} \leq \frac{d}{d+1}$, we see from Lemma 2.5 (i) that letting $\frac{1}{r_1} = \frac{1}{dp_1}$, $\frac{1}{r_3} = \frac{1}{dp_3}$, we have $A(p_1 \rightarrow r_1) \lesssim 1$ and $A(p_3 \rightarrow r_3) \lesssim 1$. Since $\frac{1}{r_1} + \frac{1}{p_2} + \frac{1}{r_3} = 1$, applying Lemma 5.2 gives the required result.
- (ii) As in the proof of the first part of the theorem, it will be enough to prove $\Lambda(p_1, p_2, p_3) \lesssim 1$ in the case when $0 \leq \frac{1}{p_1} \leq \frac{d}{d+1} \leq \frac{1}{p_3} \leq 1$ and $\frac{1}{dp_1} + \frac{1}{p_2} + \frac{d}{p_3} = d$. Let $\frac{1}{r_1} = \frac{1}{dp_1}$ and $\frac{1}{r_3} = \frac{d}{p_3} - d + 1$. Then, by Lemma 2.5, it follows that $A(p_1 \rightarrow r_1) \lesssim 1$ and $A(p_3 \rightarrow r_3) \lesssim 1$. Also, notice that $\frac{1}{r_1} + \frac{1}{p_2} + \frac{1}{r_3} = 1$. Hence, Theorem 5.3(ii) follows from Lemma 5.2.
- (iii) Switching the roles of p_1, p_2 , the proof is exactly the same as that of the second part of this theorem.
- (iv) As before, it suffices to prove the case when $\frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_3} \leq 1$ and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} = 2d - 1$. Put $\frac{1}{r_k} = \frac{d}{p_k} - d + 1$ for $k = 1, 3$. Then we see from Lemma 2.5(ii) that $A(p_k \rightarrow r_k) \lesssim 1$ for $k = 1, 3$. Notice that $\frac{1}{r_1} + \frac{1}{p_2} + \frac{1}{r_3} = 1$. Therefore, using Lemma 5.2, we finish the proof. \blacksquare

As a special case of Theorem 5.3, we obtain the following.

Corollary 5.4 *For any dimensions $d \geq 2$, we have $\Lambda_{P_2}\left(\frac{d+1}{d}, \frac{d+1}{d-1}, \frac{d+1}{d}\right) \lesssim 1$.*

Proof This clearly follows from Theorem 5.3 by taking $p_1 = p_3 = \frac{d+1}{d}$, and $p_2 = \frac{d+1}{d-1}$. \blacksquare

When $d = 2$, Theorem 5.3 does not cover some points such as $(1, 0, 1/2)$ from the convex hull in the necessary conditions by Proposition 5.1. However, it cannot be concluded that Theorem 5.3 is not sharp because the necessary conditions can be improved. While we do not know whether Theorem 5.3 is optimal or not, the result will play a crucial role in proving the following theorem which implies that Conjecture 1.5 is true for the graph K_3 and its subgraph P_2 in all dimensions $d \geq 2$ (see Corollary 5.7 below).

Theorem 5.5 *Let Λ_{K_3} and Λ_{P_2} be the operators associated with K_3 and P_2 , respectively, on \mathbb{F}_q^d . Then, if $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$ for $1 \leq p_1, p_2, p_3 \leq \infty$, we have $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$.*

Proof Suppose that $\Lambda_{K_3}(p_1, p_2, p_3) \lesssim 1$ for $1 \leq p_1, p_2, p_3 \leq \infty$. Then, by Proposition 4.1, the exponents p_1, p_2, p_3 satisfy the following three inequalities:

$$(5.3) \quad \frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} \leq d, \quad \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d, \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq d.$$

To complete the proof, it remains to show that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$. We will prove this by considering the four cases depending on the sizes of p_1 and p_3 .

Case 1: Suppose that $0 \leq \frac{1}{p_1}, \frac{1}{p_3} \leq \frac{d}{d+1}$. The condition (5.3) clearly implies that $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d$. Thus, by Theorem 5.3(i), we obtain the required conclusion that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$.

Case 2: Suppose that $0 \leq \frac{1}{p_1} \leq \frac{d}{d+1} \leq \frac{1}{p_3} \leq 1$. By Theorem 5.3(ii), to prove that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$, it will be enough to show that

$$\frac{1}{dp_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq d.$$

However, this inequality clearly follows from the third inequality in (5.3) since $d \geq 2$.

Case 3: Suppose that $0 \leq \frac{1}{p_3} \leq \frac{d}{d+1} \leq \frac{1}{p_1} \leq 1$. By Theorem 5.3(iii), it suffices to show that $\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{dp_3} \leq d$. However, this inequality can be easily obtained from the first inequality in (5.3).

Case 4: Suppose that $\frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_3} \leq 1$. By Theorem 5.3(iv), to show that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$, we only need to prove that

$$\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq 2d - 1.$$

However, this inequality can be easily proven as follows:

$$\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} = \left(\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} \right) + \frac{d-1}{p_3} \leq d + \frac{d-1}{p_3} \leq 2d - 1,$$

where the first inequality follows from the first inequality in (5.3), and the last inequality follows from a simple fact that $1 \leq p_3 \leq \infty$. \blacksquare

Remark 5.6 The reverse statement of Theorem 5.5 cannot be true. Indeed, we know by Corollary 5.4 that $\Lambda_{P_2}\left(\frac{d+1}{d}, \frac{d+1}{d-1}, \frac{d+1}{d}\right) \lesssim 1$. However, $\Lambda_{K_3}\left(\frac{d+1}{d}, \frac{d+1}{d-1}, \frac{d+1}{d}\right)$ cannot be bounded, which can be easily shown by considering Proposition 4.1, namely, the necessary conditions for the boundedness of $\Lambda_{K_3}(p_1, p_2, p_3)$.

We invoke Theorem 5.5 to deduce the following result.

Corollary 5.7 *Conjecture 1.5 is true for the graph K_3 and its subgraph P_2 in \mathbb{F}_q^d , $d \geq 2$.*

Proof It is clear that P_2 is a subgraph of K_3 in \mathbb{F}_q^d . Since $\delta(K_3) = 2$, $d \geq 2$, and $\delta(P_2) = 1$, we have $\min\{\delta(K_3), d\} = 2 > \delta(P_2) = 1$. Hence, all assumptions of Conjecture 1.5 are satisfied for K_3 and P_2 . Then the statement of the corollary follows immediately from Theorem 5.5. \blacksquare

6 Mapping properties for the $(C_4 + t)$ form

We investigate the mapping properties of the operator associated with the graph $C_4 +$ diagonal. Throughout the remaining sections, we assume that t is a nonzero element in \mathbb{F}_q^* . Let $f_i, 1 \leq i \leq 4$, be nonnegative real-valued functions on \mathbb{F}_q^d .

The operator Λ_{\diamond_t} is associated with the graph $C_4 +$ diagonal t (Figure 1d), and we define $\Lambda_{\diamond_t}(f_1, f_2, f_3, f_4)$ as the quantity

$$(6.1) \quad \frac{1}{q^d |S_t| |S_t^{d-2}|^2} \sum_{x^1, x^2, x^3, x^4 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) S_t(x^3 - x^4) S_t(x^4 - x^1) S_t(x^1 - x^3) \prod_{i=1}^4 f_i(x^i).$$

The operator Λ_{\diamond_t} is referred to as the $(C_4 + t)$ form on \mathbb{F}_q^d . Here, notice that we take the quantity $q^d |S_t| |S_t^{d-2}|^2$ as the normalizing factor $\mathcal{N}(G)$ in (1.4).

Applying a change of variables by letting $x = x^1, u = x^1 - x^2, v = x^1 - x^3, w = x^1 - x^4$, we see that

$$(6.2) \quad \Lambda_{\diamond_t}(f_1, f_2, f_3, f_4) = \frac{1}{q^d} \sum_{x \in \mathbb{F}_q^d} f_1(x) T(f_2, f_3, f_4)(x) = \langle f_1, T(f_2, f_3, f_4) \rangle,$$

where the operator $T(f_2, f_3, f_4)$ is defined by

$$(6.3) \quad T(f_2, f_3, f_4)(x) := \frac{1}{|S_t| |S_t^{d-2}|^2} \sum_{u, v, w \in S_t} S_t(v - u) S_t(w - v) f_2(x - u) f_3(x - v) f_4(x - w).$$

We are asked to find $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that

$$(6.4) \quad \Lambda_{\diamond_t}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_{p_4}$$

holds for all nonnegative real-valued functions $f_i, 1 \leq i \leq 4$, on \mathbb{F}_q^d . In other words, our main problem is to determine all numbers $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$.

Lemma 6.1 (Necessary conditions for the boundedness of $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4)$) *Let Λ_{\diamond_t} be the $(C_4 + t)$ form on \mathbb{F}_q^d . If $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$, then we have*

$$\frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d, \quad \frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} + \frac{1}{p_4} \leq d, \quad \text{and} \quad \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 2.$$

Also, under this assumption when $d = 2$, it can be shown by Polymake [1, 6] that $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the points $(0, 0, 1, 0), (0, 1, 0, 0), (0, 0, 0, 1), (1/2, 0, 1/2, 1/2), (2/3, 2/3, 0, 0), (1, 0, 0, 0), (2/3, 0, 2/3, 0), (1/2, 1/2, 1/2, 0), (2/3, 0, 0, 2/3), (0, 2/3, 2/3, 0), (0, 0, 0, 0), (0, 0, 2/3, 2/3)$.

Proof Taking $f_1 = f_2 = f_4 = 1_{S_t}$, and $f_3 = \delta_0$ in (6.4), we obtain the first conclusion

$$(6.5) \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d.$$

The second conclusion follows by symmetry from the first conclusion. Finally, one can easily prove the third conclusion, that is, $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 2$, by testing the inequality (6.4) with $f_1 = f_3 = 1_{S_t}$ and $f_2 = f_4 = \delta_0$. ■

6.1 Boundedness results for Λ_{\diamond_t} on \mathbb{F}_q^d

Given a rhombus with a fixed diagonal (the graph $C_4 + \text{diagonal}$), we will show that by removing the vertex x^2 or the vertex x^4 ,

$$\Lambda_{\diamond_t}(f_1, f_2, f_3, f_4) \lesssim \|f_2\|_{\infty} \Lambda_{K_3}(f_1, f_3, f_4) \quad \text{and} \quad \Lambda_{\diamond_t}(f_1, f_2, f_3, f_4) \lesssim \|f_4\|_{\infty} \Lambda_{K_3}(f_1, f_2, f_3).$$

Hence, upper bounds of $\Lambda_{\diamond_t}(p_1, \infty, p_3, p_4)$ and $\Lambda_{\diamond_t}(p_1, p_2, p_3, \infty)$ can be controlled by upper bounds of the $\Lambda_{K_3}(p_1, p_3, p_4)$ and $\Lambda_{K_3}(p_1, p_2, p_3)$, respectively. More precisely, we have the following relation.

Proposition 6.2 *Suppose that $\Lambda_{K_3}(p, s, r) \lesssim 1$ for $1 \leq p, s, r \leq \infty$. Then we have*

$$\Lambda_{\diamond_t}(p, \infty, s, r) \lesssim 1 \quad \text{and} \quad \Lambda_{\diamond_t}(p, s, r, \infty) \lesssim 1.$$

Proof Since $\Lambda_{K_3}(p, s, r) \lesssim 1$ for $1 \leq p, s, r \leq \infty$, we see that for all nonnegative functions f, g, h on \mathbb{F}_q^d , $\Lambda_{K_3}(f, g, h) \lesssim \|f\|_p \|g\|_s \|h\|_r$. Thus, to complete the proof, it will be enough to establish the following estimates: For all nonnegative functions f_i , $i = 1, 2, 3, 4$,

$$(6.6) \quad \Lambda_{\diamond_t}(f_1, f_2, f_3, f_4) \lesssim \|f_2\|_{\infty} \Lambda_{K_3}(f_1, f_3, f_4)$$

and

$$(6.7) \quad \Lambda_{\diamond_t}(f_1, f_2, f_3, f_4) \lesssim \|f_4\|_{\infty} \Lambda_{K_3}(f_1, f_2, f_3).$$

Since the proofs of both (6.6) and (6.7) are the same, we only provide the proof of the estimate (6.7). Notice by the definition of $\Lambda_{\diamond_t}(f_1, f_2, f_3, f_4)$ in (6.1) that $\Lambda_{\diamond_t}(f_1, f_2, f_3, f_4)$ can be written as the form

$$\frac{1}{q^d |S_t| |S_t^{d-2}|} \sum_{\substack{x^1, x^2, x^3 \in \mathbb{F}_q^d \\ : \|x^1 - x^2\| = \|x^2 - x^3\| = \|x^1 - x^3\| = t}} \left(\prod_{i=1}^3 f_i(x^i) \right) \left[\frac{1}{|S_t^{d-2}|} \sum_{x^4 \in \mathbb{F}_q^d} S_t(x^3 - x^4) S_t(x^4 - x^1) f_4(x^4) \right].$$

For each $x^1, x^3 \in \mathbb{F}_q^d$ with $\|x^1 - x^3\| = t$, we define $M(x^1, x^3)$ as the value in the above bracket. Then, recalling the definition of $\Lambda_{K_3}(f_1, f_2, f_3)$ given in (4.1), we see that

$$\Lambda_{\diamond_t}(f_1, f_2, f_3, f_4) \leq \left(\max_{\substack{x^1, x^3 \in \mathbb{F}_q^d \\ : \|x^1 - x^3\| = t}} M(x^1, x^3) \right) \Lambda_{K_3}(f_1, f_2, f_3).$$

Hence, the estimate (6.7) follows immediately by proving the following claim:

$$(6.8) \quad M := \max_{\substack{x^1, x^3 \in \mathbb{F}_q^d \\ : \|x^1 - x^3\| = t}} \frac{1}{|S_t^{d-2}|} \sum_{x^4 \in \mathbb{F}_q^d} S_t(x^3 - x^4) S_t(x^4 - x^1) \lesssim 1.$$

To prove this claim, we first apply a change of variables by letting $x = x^1$, $y = x^1 - x^3$. Then it follows that

$$M = \max_{x \in \mathbb{F}_q^d, y \in S_t} \frac{1}{|S_t^{d-2}|} \sum_{x^4 \in \mathbb{F}_q^d} S_t(x - y - x^4) S_t(x^4 - x).$$

Letting $z = x - x^4$, we have

$$M = \max_{x \in \mathbb{F}_q^d, y \in S_t} \frac{1}{|S_t^{d-2}|} \sum_{z \in S_t: \|z - y\| = t} 1 \sim \frac{1}{q^{d-2}} \max_{y \in S_t} \sum_{z \in S_t: \|z - y\| = t} 1.$$

By Corollary A.4 in the Appendix, we conclude that $M \lesssim 1$, as required. \blacksquare

In arbitrary dimensions $d \geq 2$, we have the following consequences.

Theorem 6.3 Suppose that $1 \leq a, b \leq \infty$ satisfy that

$$(6.9) \quad \frac{1}{a} + \frac{d}{b} \leq d \quad \text{and} \quad \frac{d}{a} + \frac{1}{b} \leq d.$$

Namely, let $(1/a, 1/b)$ be contained in the convex hull of points $(0, 0), (0, 1), (d/(d+1), d/(d+1)), (1, 0)$. Then we have $\Lambda_{\diamond_t}(a, \infty, b, \infty) \lesssim 1$, $\Lambda_{\diamond_t}(a, b, \infty, \infty) \lesssim 1$, $\Lambda_{\diamond_t}(a, \infty, \infty, b) \lesssim 1$, $\Lambda_{\diamond_t}(\infty, \infty, a, b) \lesssim 1$, $\Lambda_{\diamond_t}(\infty, a, b, \infty) \lesssim 1$.

Proof From Theorem 4.3, we know that the assumption (6.9) implies that $\Lambda_{K_3}(a, b, \infty) \lesssim 1$, $\Lambda_{K_3}(a, \infty, b) \lesssim 1$, and $\Lambda_{K_3}(\infty, a, b) \lesssim 1$. Hence, the statement of the theorem follows immediately by combining these and Proposition 6.2. \blacksquare

6.2 Sharp boundedness results up to endpoints for Λ_{\diamond_t} on \mathbb{F}_q^2

In this subsection, we collect our boundedness results for the operator Λ_{\diamond_t} in two dimensions.

Theorem 6.4 Let Λ_{\diamond_t} be the $(C_4 + t)$ form on \mathbb{F}_q^2 . Let $1 \leq p_1, p_2, p_3 \leq \infty$.

(i) Suppose that $(p_1, p_2, p_3) \neq (2, 2, 2)$ satisfies the following equations:

$$\frac{2}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} \leq 2, \quad \frac{1}{p_1} + \frac{2}{p_2} + \frac{1}{p_3} \leq 2, \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{2}{p_3} \leq 2.$$

Then we have $\Lambda_{\diamond_t}(p_1, \infty, p_2, p_3) \lesssim 1$ and $\Lambda_{\diamond_t}(p_1, p_2, p_3, \infty) \lesssim 1$.

(ii) In addition, we have $\Lambda_{\diamond_t}(2, \infty, 2, 2) \lesssim 1$ and $\Lambda_{\diamond_t}(2, 2, 2, \infty) \lesssim 1$, where \lesssim is used to denote that the boundedness of \diamond_t holds for all indicator test functions.

Proof Notice that Proposition 6.2 still holds after replacing \lesssim by \lesssim . Hence, the statement of the theorem is directly obtained by combining Proposition 6.2 and Theorem 4.7(ii). \blacksquare

Theorem 6.4 guarantees the sharp boundedness for the operator \diamond_t up to endpoints. Indeed, we have the following result.

Theorem 6.5 Let \diamond_t be the $(C_4 + t)$ form on \mathbb{F}_q^2 . The necessary conditions for $\diamond_t(p_1, p_2, p_3, p_4) \lesssim 1$ given in Lemma 6.1 are sufficient except for the two points $(p_1, p_2, p_3, p_4) = (2, 2, 2, \infty), (2, \infty, 2, 2)$.

In addition, we have

$$(6.10) \quad \diamond_t(2, \infty, 2, 2) \lesssim 1 \quad \text{and} \quad \diamond_t(2, 2, 2, \infty) \lesssim 1.$$

Proof The statement (6.10) was already proven in Theorem 6.4(ii). Hence, using the interpolation theorem and the second part of Lemma 6.1, the matter is reducing to proving $\diamond_t(p_1, p_2, p_3, p_4) \lesssim 1$ for the critical endpoints $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ including all the following points: $(0, 0, 0, 0), (1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1), (2/3, 2/3, 0, 0), (2/3, 0, 2/3, 0), (2/3, 0, 0, 2/3), (0, 2/3, 2/3, 0), (0, 0, 2/3, 2/3)$.

In other words, the proof will be complete by proving the following estimates: $\diamond_t(\infty, \infty, \infty, \infty) \lesssim 1, \diamond_t(1, \infty, \infty, \infty) \lesssim 1, \diamond_t(\infty, 1, \infty, \infty) \lesssim 1, \diamond_t(\infty, \infty, 1, \infty) \lesssim 1, \diamond_t(\infty, \infty, \infty, 1) \lesssim 1, \diamond_t(3/2, 3/2, \infty, \infty) \lesssim 1, \diamond_t(3/2, \infty, 3/2, \infty) \lesssim 1, \diamond_t(3/2, \infty, \infty, 3/2) \lesssim 1, \diamond_t(\infty, 3/2, 3/2, \infty) \lesssim 1, \diamond_t(\infty, \infty, 3/2, 3/2) \lesssim 1$.

However, by a direct computation, these estimates follow immediately from Theorem 6.4(i). ■

7 Boundedness problem for the C_4 form

Let $t \in \mathbb{F}_q^*$. Given nonnegative real-valued functions $f_i, 1 \leq i \leq 4$, on \mathbb{F}_q^d , we define $\Lambda_{C_4}(f_1, f_2, f_3, f_4)$ to be the following value:

(7.1)

$$\frac{1}{q^d |S_t|^2 |S_t^{d-2}|} \sum_{x^1, x^2, x^3, x^4 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) S_t(x^3 - x^4) S_t(x^4 - x^1) \prod_{i=1}^4 f_i(x^i),$$

where the quantity $q^d |S_t|^2 |S_t^{d-2}|$ stands for the normalizing factor $\mathcal{N}(G)$ in (1.4) when $G = C_4$.

Main problem is to find all exponents $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that the inequality

$$(7.2) \quad \Lambda_{C_4}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_{p_4}$$

holds for all nonnegative real-valued functions $f_i, 1 \leq i \leq 4$, on \mathbb{F}_q^d . In other words, our main problem is to determine all numbers $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$.

Lemma 7.1 (Necessary conditions for the boundedness of $\Lambda_{C_4}(p_1, p_2, p_3, p_4)$) Suppose that (7.2) holds, namely $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$. Then we have

$$\begin{aligned} \frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} &\leq d+1, \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d+1, \quad \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{1}{p_4} \leq d+1, \\ \frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} + \frac{1}{p_4} &\leq d+1, \quad \frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq 2d-2, \quad \text{and} \quad \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d-2. \end{aligned}$$

In particular, when $d = 2$, it can be shown by Polymake [1, 6] that $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the points $(0, 0, 1, 0), (0, 0, 0, 1), (0, 1, 0, 0), (2/3, 0, 0, 2/3), (2/3, 2/3, 0, 0), (1, 0, 0, 0), (0, 0, 0, 0), (0, 2/3, 2/3, 0), (0, 0, 2/3, 2/3)$.

Remark 7.2 When $d = 2, 3$, the first four inequalities in the conclusion are not necessary. We only need the last two.

Proof The six inequalities in the conclusion can be easily deduced by testing the inequality (7.2) with the following specific functions, respectively: We leave the proofs to the readers.

- 1) $f_1 = f_2 = f_3 = 1_{S_t}$, and $f_4 = \delta_0$.
- 2) $f_1 = f_2 = f_4 = 1_{S_t}$, and $f_3 = \delta_0$.
- 3) $f_1 = f_3 = f_4 = 1_{S_t}$, and $f_2 = \delta_0$.
- 4) $f_2 = f_3 = f_4 = 1_{S_t}$, and $f_1 = \delta_0$.
- 5) $f_2 = f_4 = 1_{S_t}$, and $f_1 = f_3 = \delta_0$.
- 6) $f_1 = f_3 = 1_{S_t}$, and $f_2 = f_4 = \delta_0$.

■

7.1 Boundedness results for Λ_{C_4} on \mathbb{F}_q^d

In this subsection, we provide some exponents $1 \leq p_i \leq \infty, 1 \leq i \leq 4$, such that $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$ in the specific case when one of p_i is ∞ , but it is valid for all dimensions $d \geq 2$. In general, it is very hard to deduce nontrivial boundedness results for the C_4 form on \mathbb{F}_q^d .

We begin by observing that an upper bound of $\Lambda_{C_4}(f_1, f_2, f_3, f_4)$ can be controlled by estimating for both the K_2 form and the P_2 form.

Lemma 7.3 *For all nonnegative functions $f_i, i = 1, 2, 3, 4$, on $\mathbb{F}_q^d, d \geq 2$, we have*

$$\Lambda_{C_4}(f_1, f_2, f_3, f_4) \lesssim \begin{cases} \left(\frac{1}{|S_t^{d-2}|} L(f_1 f_3, f_2) + \Lambda(f_1, f_2, f_3) \right) \|f_4\|_\infty, \\ \left(\frac{1}{|S_t^{d-2}|} L(f_2 f_4, f_1) + \Lambda(f_4, f_1, f_2) \right) \|f_3\|_\infty, \\ \left(\frac{1}{|S_t^{d-2}|} L(f_1 f_3, f_4) + \Lambda(f_3, f_4, f_1) \right) \|f_2\|_\infty, \\ \left(\frac{1}{|S_t^{d-2}|} L(f_2 f_4, f_3) + \Lambda(f_2, f_3, f_4) \right) \|f_1\|_\infty. \end{cases}$$

Proof We only provide the proof of the first inequality,

$$(7.3) \quad \Lambda_{C_4}(f_1, f_2, f_3, f_4) \lesssim \frac{1}{|S_t^{d-2}|} \Lambda_{K_2}(f_1 f_3, f_2) \|f_4\|_\infty + \Lambda_{P_2}(f_1, f_2, f_3) \|f_4\|_\infty,$$

since other inequalities can be easily proven in the same way by replacing the role of f_4 with f_3, f_2, f_1 , respectively. By definition, the value of $\Lambda_{C_4}(f_1, f_2, f_3, f_4)$ is equal to

$$\begin{aligned} & \frac{1}{q^d |S_t|^2 |S_t^{d-2}|} \sum_{x^1, x^2, x^3 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) \left(\prod_{i=1}^3 f_i(x^i) \right) \\ & \times \left(\sum_{x^4 \in \mathbb{F}_q^d} S_t(x^3 - x^4) S_t(x^4 - x^1) f_4(x^4) \right). \end{aligned}$$

For fixed $x^1, x^3 \in \mathbb{F}_q^d$, the sum in the above bracket can be estimated as follows:

$$\sum_{x^4 \in \mathbb{F}_q^d} S_t(x^3 - x^4) S_t(x^4 - x^1) f_4(x^4) \lesssim \begin{cases} |S_t| \|f_4\|_\infty, & \text{if } x^1 = x^3, \\ q^{d-2} \|f_4\|_\infty, & \text{if } x^1 \neq x^3. \end{cases}$$

Notice that this estimates are easily obtained by invoking Corollary A.4 in the Appendix after using a change of variables.

Let $\Lambda_{C_4}(f_1, f_2, f_3, f_4) =: \diamondsuit_1 + \diamondsuit_2$, where \diamondsuit_1 denotes the contribution to $\Lambda_{C_4}(f_1, f_2, f_3, f_4)$ when $x^1 = x^3$, and \diamondsuit_2 does it when $x^1 \neq x^3$. Then it follows that

$$\begin{aligned}\diamondsuit_1 &\lesssim \frac{\|f_4\|_\infty}{q^d |S_t| |S_t^{d-2}|} \sum_{x^1, x^2 \in \mathbb{F}_q^d} S_t(x^1 - x^2)(f_1 f_3)(x^1) f_2(x^2) = \frac{1}{|S_t^{d-2}|} \Lambda_{K_2}(f_1 f_3, f_2) \|f_4\|_\infty, \\ \diamondsuit_2 &\lesssim \frac{\|f_4\|_\infty}{q^d |S_t|^2} \sum_{x^1, x^2, x^3 \in \mathbb{F}_q^d: x^1 \neq x^3} S_t(x^1 - x^2) S_t(x^2 - x^3) f_1(x^1) f_2(x^2) f_3(x^3) \lesssim \Lambda_{P_2}(f_1, f_2, f_3) \|f_4\|_\infty.\end{aligned}$$

Hence, we obtain the required estimate (7.3). \blacksquare

In Lemma 7.3, we obtained four different kinds of the upper bounds of the $\Lambda_{C_4}(f_1, f_2, f_3, f_4)$. Using each of them, we are able to deduce exponents p_1, p_2, p_3, p_4 with $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$, where at least one of $p_j, j = 1, 2, 3, 4$, takes ∞ .

The following result can be proven by applying the first upper bound of $\Lambda_{C_4}(f_1, f_2, f_3, f_4)$ in Lemma 7.3 together with Theorems 3.3 and 5.3.

Proposition 7.4 *Let $1 \leq p_1, p_2, p_3 \leq \infty$. For the C_4 form Λ_{C_4} on \mathbb{F}_q^d , $d \geq 2$, the following statements are true.*

- (i) *If $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d$ and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq d$, then $\Lambda_{C_4}(p_1, p_2, p_3, \infty) \lesssim 1$.*
- (ii) *If $\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_4} \leq d$ and $\frac{1}{p_1} + \frac{d}{p_2} + \frac{d}{p_4} \leq d$, then $\Lambda_{C_4}(p_1, p_2, \infty, p_4) \lesssim 1$.*
- (iii) *If $\frac{1}{p_1} + \frac{1}{p_3} + \frac{d}{p_4} \leq d$ and $\frac{d}{p_1} + \frac{d}{p_3} + \frac{1}{p_4} \leq d$, then $\Lambda_{C_4}(p_1, \infty, p_3, p_4) \lesssim 1$.*
- (iv) *If $\frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d$ and $\frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq d$, then $\Lambda_{C_4}(\infty, p_2, p_3, p_4) \lesssim 1$.*

Proof We will only provide the proof of the first part of the theorem since the proofs of other parts are the same in the sense that the proof of the first part uses the first upper bound of Lemma 7.3 and the proofs of other parts can also use their corresponding upper bounds of Lemma 7.3 to complete the proofs.

Let us start proving the first part of the theorem. To complete the proof, we aim to show that for all nonnegative functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^d ,

$$\Lambda_{C_4}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_\infty,$$

whenever the exponents $1 \leq p_1, p_2, p_3 \leq \infty$ satisfy the following conditions:

$$(7.4) \quad \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d \quad \text{and} \quad \frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq d.$$

By the first part of Lemma 7.3, it follows that

$$\Lambda_{C_4}(f_1, f_2, f_3, f_4) \lesssim \left(\frac{1}{|S_t^{d-2}|} L(f_1 f_3, f_2) + \Lambda(f_1, f_2, f_3) \right) \|f_4\|_\infty.$$

Therefore, under the assumptions (7.4), our problem is reducing to establishing the following two estimates:

$$(7.5) \quad \Lambda_{K_2}(f_1 f_3, f_2) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3},$$

$$(7.6) \quad \Lambda_{P_2}(f_1, f_2, f_3) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3}.$$

For $1 \leq p_1, p_3 \leq \infty$, let $1/r = 1/p_1 + 1/p_3$. Then the conditions (7.4) are the same as

$$\frac{1}{r} + \frac{d}{p_2} \leq d \quad \text{and} \quad \frac{d}{r} + \frac{1}{p_2} \leq d.$$

So these conditions enable us to invoke Theorem 3.3 so that we obtain the estimate (7.5) as follows:

$$\Lambda_{K_2}(f_1 f_3, f_2) \lesssim \|f_1 f_3\|_r \|f_2\|_{p_2} \leq \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3},$$

where we used Hölder's inequality in the last inequality.

It remains to prove the estimate (7.6) under the assumptions (7.4). To do this, we shall use Theorem 5.3, which gives sufficient conditions for $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$. We directly compare the conditions (7.4) with the assumptions of Theorem 5.3. Then it is not hard to observe the following statements.

- (Case 1) In the case when $0 \leq \frac{1}{p_1}, \frac{1}{p_3} \leq \frac{d}{d+1}$, the conditions (7.4) imply the hypothesis of the first part of Theorem 5.3.
- (Case 2) In the case when $0 \leq \frac{1}{p_1} \leq \frac{d}{d+1} \leq \frac{1}{p_3} \leq 1$, the conditions (7.4) imply the hypothesis of the second part of Theorem 5.3. To see this, notice that if $d/p_1 + 1/p_2 + d/p_3 \leq d$, then $1/(dp_1) + 1/p_2 + d/p_3 \leq d$.
- (Case 3) In the case when $0 \leq \frac{1}{p_3} \leq \frac{d}{d+1} \leq \frac{1}{p_1} \leq 1$, the conditions (7.4) imply the hypothesis of the third part of Theorem 5.3.
- (Case 4) In the case when $\frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_3} \leq 1$, the conditions (7.4) imply the hypothesis of the fourth part of Theorem 5.3.

Hence, we conclude from Theorem 5.3 that $\Lambda_{P_2}(p_1, p_2, p_3) \lesssim 1$ under the assumptions (7.4), as desired. \blacksquare

7.2 Sharp boundedness results for Λ_{C_4} on \mathbb{F}_q^2

Recall that Proposition 7.4 provides sufficient conditions for $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$ in any dimensions $d \geq 2$. In this section, we show that Proposition 7.4 is sharp in two dimensions. More precisely, using Proposition 7.4, we will prove the following optimal result.

Theorem 7.5 *Let Λ_{C_4} be the C_4 form on \mathbb{F}_q^2 . For $1 \leq p_i \leq \infty, 1 \leq i \leq 4$, we have*

$$\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1 \text{ if and only if } \frac{2}{p_1} + \frac{1}{p_2} + \frac{2}{p_3} + \frac{1}{p_4} \leq 2, \text{ and } \frac{1}{p_1} + \frac{2}{p_2} + \frac{1}{p_3} + \frac{2}{p_4} \leq 2.$$

Proof The necessary conditions for $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$ follow immediately from Lemma 7.1 for $d = 2$ (see Remark 7.2).

Conversely, suppose that $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ satisfy the following two inequalities:

$$(7.7) \quad \frac{2}{p_1} + \frac{1}{p_2} + \frac{2}{p_3} + \frac{1}{p_4} \leq 2, \text{ and } \frac{1}{p_1} + \frac{2}{p_2} + \frac{1}{p_3} + \frac{2}{p_4} \leq 2.$$

Then, as mentioned in Lemma 7.1, it can be shown by Polymake [1, 6] that $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the points $(0, 0, 1, 0)$,

$(0, 0, 0, 1), (0, 1, 0, 0), (2/3, 0, 0, 2/3), (2/3, 2/3, 0, 0), (1, 0, 0, 0), (0, 0, 0, 0), (0, 2/3, 2/3, 0), (0, 0, 2/3, 2/3)$.

By interpolating the above nine critical points, to prove $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$ for all $p_i, 1 \leq i \leq 4$ satisfying the inequalities in (7.7), it will be enough to prove it for the nine critical points $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$. This can be easily proven by using Proposition 7.4. For example, for the point $(1/p_1, 1/p_2, 1/p_3, 1/p_4) = (2/3, 0, 0, 2/3)$, a direct computation shows that the assumptions in Proposition 7.4(ii) are satisfied and thus $\Lambda_{C_4}(p_1, p_2, p_3, p_4) = \Lambda_{C_4}(3/2, \infty, \infty, 3/2) \lesssim 1$. For other critical points, we can easily prove them in the same way so that we omit the detail proofs. ■

Notice that the graph C_4 is a subgraph of the graph $C_4 + \text{diagonal}$, and they are associated with the operators Λ_{C_4} and Λ_{\diamond_t} , respectively. Hence, the following proposition shows that the answer to Question 1.2 is negative when G is the $C_4 + \text{diagonal}$, and G' is the C_4 . However, this does not mean that Conjecture 1.5 is not true since the C_4 and the $C_4 + \text{diagonal}$ do not satisfy the main hypothesis (1.6) of Conjecture 1.5.

Proposition 7.6 *Let $\Lambda_{\diamond_t}, \Lambda_{C_4}$ be the $(C_4 + t)$ form and the C_4 form on \mathbb{F}_q^2 , respectively. Let $1 \leq p_1, p_2, p_3, p_4 \leq \infty$. Then the following statements hold.*

- (i) *If $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$, then $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$.*
- (ii) *Moreover, there exist exponents $1 \leq a, b, c, d \leq \infty$ such that $\Lambda_{\diamond_t}(a, b, c, d) \lesssim 1$ but $\Lambda_{C_4}(a, b, c, d)$ is not bounded.*

Proof First, let us prove the statement (ii) in the conclusion. To prove this, we choose $(a, b, c, d) = (3/2, \infty, 3/2, \infty)$. From Theorem 6.4(i), we can easily note that $\Lambda_{\diamond_t}(3/2, \infty, 3/2, \infty) \lesssim 1$. However, it is impossible that $\Lambda_{C_4}(3/2, \infty, 3/2, \infty) \lesssim 1$, which can be shown from Theorem 7.5.

Next, let us prove the first conclusion of the theorem. Suppose that $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$ for $1 \leq p_i \leq \infty, 1 \leq i \leq 4$. Then, as mentioned in the second conclusion of Lemma 7.1, the point $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ lies on the convex body with the critical endpoints: $(0, 0, 1, 0), (0, 0, 0, 1), (0, 1, 0, 0), (2/3, 0, 0, 2/3), (2/3, 2/3, 0, 0), (1, 0, 0, 0), (0, 0, 0, 0), (0, 2/3, 2/3, 0), (0, 0, 2/3, 2/3)$.

Invoking the interpolation theorem, to prove the conclusion that $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$, it will be enough to establish the boundedness only for those nine critical points $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$. More precisely, it remains to establish the following estimates:

$\Lambda_{\diamond_t}(\infty, \infty, \infty, \infty) \lesssim 1, \Lambda_{\diamond_t}(1, \infty, \infty, \infty) \lesssim 1, \Lambda_{\diamond_t}(\infty, 1, \infty, \infty) \lesssim 1, \Lambda_{\diamond_t}(\infty, \infty, 1, \infty) \lesssim 1, \Lambda_{\diamond_t}(\infty, \infty, \infty, 1) \lesssim 1, \Lambda_{\diamond_t}(3/2, 3/2, \infty, \infty) \lesssim 1, \Lambda_{\diamond_t}(3/2, \infty, \infty, 3/2) \lesssim 1, \Lambda_{\diamond_t}(\infty, 3/2, 3/2, \infty) \lesssim 1, \Lambda_{\diamond_t}(\infty, \infty, 3/2, 3/2) \lesssim 1$.

However, these estimates follow by applying Theorem 6.4(i). ■

8 Boundedness problem for the P_3 form

For $t \in \mathbb{F}_q^*$ and nonnegative real-valued functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^d , we define $\Lambda_{P_3}(f_1, f_2, f_3, f_4)$ as the following value:

$$(8.1) \quad \frac{1}{q^d |S_t|^3} \sum_{x^1, x^2, x^3, x^4 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) S_t(x^3 - x^4) \prod_{i=1}^4 f_i(x^i).$$

This operator Λ_{P_3} will be named the P_3 form on \mathbb{F}_q^d since it is related to the graph P_3 with vertices in \mathbb{F}_q^d , $d \geq 2$. Note that in the definition of $\Lambda_{P_3}(f_1, f_2, f_3, f_4)$, we take the normalizing fact $q^d |S_t|^3$, which is corresponding to $\mathcal{N}(G)$ in (1.4) when G is the P_3 .

We want to determine $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that

$$(8.2) \quad \Lambda_{P_3}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_{p_4}$$

holds for all nonnegative real-valued functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^d . In other words, our main problem is to find all numbers $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$.

Lemma 8.1 (Necessary conditions for $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$) Suppose that $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$. Then we have $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d$, $\frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d$, $\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_4} \leq d + 2$, $\frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} \leq d + 2$, $\frac{1}{p_1} + \frac{d}{p_2} + \frac{d}{p_4} \leq 2d - 1$, $\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq 2d - 1$, and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d$.

In particular, when $d = 2$, by using Polymake [1, 6], it can be shown that $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the points: $(0, 1, 0, 1/2)$, $(0, 1, 0, 0)$, $(1/2, 0, 1/2, 1)$, $(1/2, 0, 1, 0)$, $(1, 1/2, 0, 0)$, $(1, 0, 0, 0)$, $(1, 1/3, 1/3, 0)$, $(1, 0, 1/2, 0)$, $(1/2, 1/3, 5/6, 0)$, $(1, 0, 0, 1)$, $(1/2, 1/2, 1/2, 1/2)$, $(1, 1/2, 0, 1/2)$, $(0, 5/6, 1/3, 1/2)$, $(0, 0, 0, 1)$, $(0, 1/2, 0, 1)$, $(0, 2/3, 2/3, 0)$, $(0, 1/3, 1/3, 1)$, $(0, 0, 1/2, 1)$, $(0, 0, 1, 0)$, $(0, 0, 0, 0)$.

Remark 8.2 When $d = 2$, the third, fourth, and seventh inequalities above are not necessary. When $d = 3$, the third and fourth inequalities above are not necessary.

Proof As in the proofs of Propositions 3.1, 4.1, and 5.1, the conclusions of the statement follow by testing the inequality (8.2) with the following specific functions, respectively:

- 1) $f_1 = f_3 = 1_{S_t}, f_2 = \delta_0$, and $f_4 = 1_{\mathbb{F}_q^d}$.
- 2) $f_1 = 1_{\mathbb{F}_q^d}, f_2 = f_4 = 1_{S_t}$, and $f_3 = \delta_0$.
- 3) $f_2 = f_3 = f_4 = 1_{S_t}$, and $f_1 = \delta_0$.
- 4) $f_1 = f_2 = f_3 = 1_{S_t}$, and $f_4 = \delta_0$.
- 5) $f_1 = f_3 = 1_{S_t}$, and $f_2 = f_4 = \delta_0$.
- 6) $f_2 = f_4 = 1_{S_t}$, and $f_1 = f_3 = \delta_0$.
- 7) $f_2 = f_3 = 1_{S_t}$, and $f_1 = f_4 = \delta_0$.

■

8.1 Boundedness results for Λ_{P_3} on \mathbb{F}_q^d

We begin by observing that an upper bound of $\Lambda_{P_3}(f_1, f_2, f_3, f_4)$ can be controlled by the value $\Lambda_{P_2}(f_1, f_2, f_3)$.

Proposition 8.3 Let $1 \leq a, b, c \leq \infty$. If $\Lambda_{P_2}(a, b, c) \lesssim 1$, then $\Lambda_{P_3}(a, b, c, \infty)$, $\Lambda_{P_3}(\infty, a, b, c) \lesssim 1$.

Proof For all nonnegative functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^d , our task is to prove the following inequalities:

$$(8.3) \quad \Lambda_{P_3}(f_1, f_2, f_3, f_4) \lesssim \begin{cases} \Lambda_{P_2}(f_1, f_2, f_3) \|f_4\|_\infty, \\ \|f_1\|_\infty \Lambda(f_2, f_3, f_4). \end{cases}$$

We will only prove the first inequality, that is,

$$(8.4) \quad \Lambda_{P_3}(f_1, f_2, f_3, f_4) \lesssim \Lambda_{P_2}(f_1, f_2, f_3) \|f_4\|.$$

By symmetry, the second inequality can be easily proven in the same way. By definition in (8.1), we can write $\Lambda_{P_3}(f_1, f_2, f_3, f_4)$ as

$$\frac{1}{q^d |S_t|^2} \sum_{x^1, x^2, x^3 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) \left(\prod_{i=1}^3 f_i(x^i) \right) \left(\frac{1}{|S_t|} \sum_{x^4 \in \mathbb{F}_q^d} f_4(x^4) S_t(x^3 - x^4) \right).$$

Since the value in the above bracket is $Af_4(x^3)$, which is clearly dominated by $\|Af_4\|_\infty$, the required estimate (8.4) follows immediately from the definition of $\Lambda_{P_2}(f_1, f_2, f_3)$ in (5.1). \blacksquare

The following lemma can be deduced from Proposition 8.3 and Theorem 5.3.

Lemma 8.4 Consider the P_3 form Λ_{P_3} on \mathbb{F}_q^d . Suppose that the exponents $1 \leq a, b, c \leq \infty$ satisfy one of the following conditions:

- (i) $0 \leq \frac{1}{a}, \frac{1}{c} \leq \frac{d}{d+1}$ and $\frac{1}{a} + \frac{d}{b} + \frac{1}{c} \leq d$,
- (ii) $0 \leq \frac{1}{a} \leq \frac{d}{d+1} \leq \frac{1}{c} \leq 1$, and $\frac{1}{da} + \frac{1}{b} + \frac{d}{c} \leq d$,
- (iii) $0 \leq \frac{1}{c} \leq \frac{d}{d+1} \leq \frac{1}{a} \leq 1$, and $\frac{d}{a} + \frac{1}{b} + \frac{1}{dc} \leq d$,
- (iv) $\frac{d}{d+1} \leq \frac{1}{a}, \frac{1}{c} \leq 1$ and $\frac{d}{a} + \frac{1}{b} + \frac{d}{c} \leq 2d - 1$.

Then we have $\Lambda_{P_3}(a, b, c, \infty) \lesssim 1$ and $\Lambda_{P_3}(\infty, a, b, c) \lesssim 1$.

Proof Using Theorem 5.3 with $p_1 = a$, $p_2 = b$, $p_3 = c$, it is clear that $\Lambda_{P_2}(a, b, c) \lesssim 1$ for all exponents a, b, c in our assumption. Hence, the statement follows immediately from Proposition 8.3. \blacksquare

Now we prove that the value $\Lambda_{P_3}(f_1, f_2, f_3, f_4)$ can be expressed in terms of the averaging operator over spheres. For functions f, g, h on \mathbb{F}_q^d , let us denote

$$\langle f, g, h \rangle := \|fgh\|_1 = \frac{1}{q^d} \sum_{x \in \mathbb{F}_q^d} f(x)g(x)h(x).$$

Proposition 8.5 Let $f_i, i = 1, 2, 3, 4$, be nonnegative real-valued functions on \mathbb{F}_q^d . Then we have

$$\Lambda_{P_3}(f_1, f_2, f_3, f_4) = \langle Af_1, f_2, A(f_3 \cdot Af_4) \rangle = \langle A(f_2 \cdot Af_1), f_3, Af_4 \rangle.$$

Proof By symmetry, to complete the proof, it suffices to prove the first equality, that is,

$$\Lambda_{P_3}(f_1, f_2, f_3, f_4) = \langle Af_1, f_2, A(f_3 \cdot Af_4) \rangle.$$

Combining the definition in (8.1) and the definition of the spherical averaging operator A , it follows that

$$\begin{aligned}\Lambda_{P_3}(f_1, f_2, f_3, f_4) &= \frac{1}{q^d} \sum_{x^2 \in \mathbb{F}_q^d} f_2(x^2) A f_1(x^2) \left[\frac{1}{|S_t|} \sum_{x^3 \in \mathbb{F}_q^d} f_3(x^3) S_t(x^2 - x^3) A f_4(x^3) \right] \\ &= \frac{1}{q^d} \sum_{x^2 \in \mathbb{F}_q^d} f_2(x^2) A f_1(x^2) A(f_3 \cdot A f_4)(x^2).\end{aligned}$$

This gives the required estimate. \blacksquare

Combining Proposition 8.5 and the averaging estimate over spheres, we are able to deduce sufficient conditions for the boundedness of the P_3 form Λ_{P_3} on \mathbb{F}_q^d .

Lemma 8.6 *Let $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ be exponents satisfying one of the following conditions:*

- (i) $0 \leq \frac{1}{p_1}, \frac{1}{p_4}, \frac{1}{p_3} + \frac{1}{dp_4} \leq \frac{d}{d+1}$, and $\frac{1}{dp_1} + \frac{1}{p_2} + \frac{1}{dp_3} + \frac{1}{d^2 p_4} \leq 1$.
- (ii) $0 \leq \frac{1}{p_1}, \frac{1}{p_4} \leq \frac{d}{d+1} \leq \frac{1}{p_3} + \frac{1}{dp_4} \leq 1$, and $\frac{1}{dp_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d$.
- (iii) $0 \leq \frac{1}{p_1}, \frac{1}{p_3} + \frac{d}{p_4} - d + 1 \leq \frac{d}{d+1} \leq \frac{1}{p_4} \leq 1$, and $\frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 1$.
- (iv) $0 \leq \frac{1}{p_1} \leq \frac{d}{d+1} \leq \frac{1}{p_4}, \frac{1}{p_3} + \frac{d}{p_4} - d + 1 \leq 1$, and $\frac{1}{dp_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{d^2}{p_4} \leq d^2$.
- (v) $0 \leq \frac{1}{p_4}, \frac{1}{p_3} + \frac{1}{dp_4} \leq \frac{d}{d+1} \leq \frac{1}{p_1} \leq 1$, and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{dp_3} + \frac{1}{d^2 p_4} \leq d$.
- (vi) $0 \leq \frac{1}{p_4} \leq \frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_3} + \frac{1}{dp_4} \leq 1$, and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq 2d - 1$.
- (vii) $0 \leq \frac{1}{p_3} + \frac{d}{p_4} - d + 1 \leq \frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_4} \leq 1$, and $\frac{d^2}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq d^2 + d - 1$.
- (viii) $\frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_4}, \frac{1}{p_3} + \frac{d}{p_4} - d + 1 \leq 1$, and $\frac{d}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{d^2}{p_4} \leq d^2 + d - 1$.

Then we have $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$ and $\Lambda_{P_3}(p_4, p_3, p_2, p_1) \lesssim 1$.

Proof By symmetry, it will be enough to prove the first part of conclusions, that is, $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$. To complete the proof, we will first find the general conditions that guarantee this conclusion. Next, we will demonstrate that each of the hypotheses in the theorem satisfies the general conditions.

To derive the first general condition, we assume that $1 \leq r_1, p_2, r \leq \infty$ satisfy that

$$(8.5) \quad \frac{1}{r_1} + \frac{1}{p_2} + \frac{1}{r} \leq 1.$$

Then, by Proposition 8.5 and Hölder's inequality,

$$\Lambda_{P_3}(f_1, f_2, f_3, f_4) \leq \|A f_1\|_{r_1} \|f_2\|_{p_2} \|A(f_3 \cdot A f_4)\|_r,$$

where we also used the nesting property of norms associated with the normalizing counting measure. Assume that $1 \leq p_1, s \leq \infty$ satisfy the following averaging estimates over spheres:

$$(8.6) \quad A(p_1 \rightarrow r_1) \lesssim 1 \quad \text{and} \quad A(s \rightarrow r) \lesssim 1.$$

It follows that $\Lambda_{P_3}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|Af_4\|_s$. Now we assume that $1 \leq p_3, t \leq \infty$ satisfy that

$$(8.7) \quad \frac{1}{s} = \frac{1}{p_3} + \frac{1}{t}.$$

Then, by Hölder's inequality, we see that

$$\Lambda_{P_3}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|Af_4\|_t.$$

Finally, if we assume that $1 \leq p_4 \leq \infty$ satisfies the following averaging estimate

$$(8.8) \quad A(p_4 \rightarrow t) \lesssim 1,$$

then we obtain that $\Lambda_{P_3}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_{p_4}$.

In summary, we see that $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$ provided that the numbers $1 \leq p_i \leq \infty, i = 1, 2, 3, 4$, satisfy all the conditions (8.5)–(8.8). Thus, to finish the proof, we will show that each of the eight hypotheses in the theorem satisfies all these conditions.

Given $1 \leq p_1, p_4 \leq \infty$, by Lemma 2.5, we can chose $1 \leq r_1, t \leq \infty$ such that the first averaging estimate in (8.6) and the averaging estimate (8.8) hold, respectively. More precisely, we can select $0 \leq 1/r_1, 1/t \leq 1$ as follows:

- If $0 \leq \frac{1}{p_1} \leq \frac{d}{d+1}$, then we take $1/r_1 = 1/(dp_1)$.
- If $0 \leq \frac{1}{p_4} \leq \frac{d}{d+1}$, then we take $1/t = 1/(dp_4)$.
- If $\frac{d}{d+1} \leq \frac{1}{p_1} \leq 1$, then we choose $1/r_1 = d/p_1 - d + 1$.
- If $\frac{d}{d+1} \leq \frac{1}{p_4} \leq 1$, then we choose $1/t = d/p_4 - d + 1$.

In the next step, we determine $1 \leq r \leq \infty$ by using the condition (8.7) and the second averaging estimate in (8.6). Since two kinds of t values can be chosen as above, the condition (8.7) becomes

$$\frac{1}{s} = \frac{1}{p_3} + \frac{1}{dp_4} \quad \text{or} \quad \frac{1}{s} = \frac{1}{p_3} + \frac{d}{p_4} - d + 1.$$

Combining these s values with the second averaging estimate in (8.6), the application of Lemma 2.5 enables us to choose $1/r$ values as follows:

- If $0 \leq \frac{1}{s} = \frac{1}{p_3} + \frac{1}{dp_4} \leq \frac{d}{d+1}$, then we take $\frac{1}{r} = \frac{1}{dp_3} + \frac{1}{d^2p_4}$.
- If $\frac{d}{d+1} \leq \frac{1}{s} = \frac{1}{p_3} + \frac{1}{dp_4} \leq 1$, then we take $\frac{1}{r} = \frac{d}{p_3} + \frac{1}{p_4} - d + 1$.
- If $0 \leq \frac{1}{s} = \frac{1}{p_3} + \frac{d}{p_4} - d + 1 \leq \frac{d}{d+1}$, then we take $\frac{1}{r} = \frac{1}{dp_3} + \frac{1}{p_4} - 1 + \frac{1}{d}$.
- If $\frac{d}{d+1} \leq \frac{1}{s} = \frac{1}{p_3} + \frac{d}{p_4} - d + 1 \leq 1$, then we take $\frac{1}{r} = \frac{d}{p_3} + \frac{d^2}{p_4} - d^2 + 1$.

Finally, use the condition (8.5) together with previously selected two values for r_1 and four values for r . Then we obtain the required remaining conditions. ■

Remark 8.7 Notice that Lemma 8.4 is a special case of Lemma 8.6. However, the proof of Lemma 8.4 is much simpler than that of Lemma 8.6.

We do not know if the consequences from Lemmas 8.4 and 8.6 imply the sharp boundedness results for the P_3 form Λ_{P_3} on \mathbb{F}_q^d . However, they play an important role in proving the proposition below, which states that the exponents for

$\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$ are less restricted than those for $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$. The precise statement is as follows.

Theorem 8.8 *Let Λ_{\diamond_t} and Λ_{P_3} be the operators acting on the functions on \mathbb{F}_q^2 . If $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$ for $1 \leq p_1, p_2, p_3, p_4 \leq \infty$, then $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$.*

Proof Assume that $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$ for $1 \leq p_1, p_2, p_3, p_4 \leq \infty$. Then, by Lemma 6.1, the point $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the following points: $(0, 0, 1, 0), (0, 1, 0, 0), (0, 0, 0, 1), (1/2, 0, 1/2, 1/2), (2/3, 2/3, 0, 0), (1, 0, 0, 0), (2/3, 0, 2/3, 0), (1/2, 1/2, 1/2, 0), (2/3, 0, 0, 2/3), (0, 2/3, 2/3, 0), (0, 0, 0, 0), (0, 0, 2/3, 2/3)$.

To complete the proof, by the interpolation theorem, it suffices to show that for each of the above critical points $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$, we have

$$\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1.$$

To prove this, we will use Lemma 8.6 and Lemma 8.4. By Lemma 8.6 with the hypothesis (i), one can notice that $\Lambda_{P_3}(3/2, \infty, \infty, 3/2) \lesssim 1$, which is corresponding to the point $(1/p_1, 1/p_2, 1/p_3, 1/p_4) = (2/3, 0, 0, 2/3)$. Similarly, Lemma 8.6 with the hypothesis (ii) can be used for the point $(1/2, 0, 1/2, 1/2)$, namely, $\Lambda_{P_3}(2, \infty, 2, 2) \lesssim 1$.

For any other points, we can invoke Lemma 8.4. More precisely, we can apply Lemma 8.4 with the hypothesis (i) for the points $(0, 1, 0, 0), (2/3, 2/3, 0, 0), (2/3, 0, 2/3, 0), (1/2, 1/2, 1/2, 0), (0, 2/3, 2/3, 0), (0, 0, 0, 0), (0, 0, 2/3, 2/3)$. The points $(0, 0, 1, 0), (0, 0, 0, 1)$ can be obtained by Lemma 8.4 with the hypothesis (ii). Finally, for the point $(1, 0, 0, 0)$, we can prove that $\Lambda_{P_3}(1, \infty, \infty, \infty) \lesssim 1$ by using Lemma 8.4 with the hypothesis (iii). This completes the proof. ■

Remark 8.9 The reverse statement of Theorem 8.8 is not true in general. As a counterexample, we can take $p_1 = 3/2, p_2 = 3, p_3 = 3/2, p_4 = \infty$. Indeed, the assumption (i) of Lemma 8.4 with $d = 2$ implies that $\Lambda_{P_3}(3/2, 3, 3/2, \infty) \lesssim 1$. However, $\Lambda_{\diamond_t}(3/2, 3, 3/2, \infty)$ cannot be bounded, which follows from Lemma 6.1.

We obtain the following consequence of Theorem 8.8.

Corollary 8.10 *Conjecture 1.5 is valid for the graph $C_4 + \text{diagonal}$ and its subgraph P_3 in \mathbb{F}_q^2 .*

Proof It is obvious that the P_3 is a subgraph of $C_4 + \text{diagonal}$ in \mathbb{F}_q^2 . For $d = 2$, it is plain to notice that $\min\{\delta(C_4 + \text{diagonal}), d\} = 2 > \delta(P_3) = 1$. Thus, the graph $C_4 + \text{diagonal}$ and its subgraph P_3 satisfy all assumptions of Conjecture 1.5. Then the statement of the corollary follows immediately from Theorem 8.8 since the operators \diamond_t and Λ_{P_3} are related to the $C_4 + \text{diagonal}$ and its subgraph P_3 , respectively. ■

The following theorem provides a concrete example for a positive answer to Question 1.2 since the operators \diamond and Λ_{P_3} are related to the graph C_4 and its subgraph P_3 , respectively. Furthermore, the graphs also satisfy Conjecture 1.5 (see Corollary 8.13 below).

Theorem 8.11 *Let Λ_{C_4} and Λ_{P_3} be the operators acting on the functions on \mathbb{F}_q^2 . If $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1, 1 \leq p_1, p_2, p_3, p_4 \leq \infty$, then $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$.*

Proof By Proposition 7.6(i), if $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$, then $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$. By Theorem 8.8, if $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$, then $\Lambda_{P_3}(p_1, p_2, p_3, p_4) \lesssim 1$. Hence, the statement follows. \blacksquare

Remark 8.12 The reverse statement of Theorem 8.11 cannot hold. As in Remark 8.9, if we can take $p_1 = 3/2, p_2 = 3, p_3 = 3/2, p_4 = \infty$, then $\Lambda_{P_3}(3/2, 3, 3/2, \infty) \lesssim 1$. However, $\Lambda_{C_4}(3/2, 3, 3/2, \infty)$ cannot be bounded, which follows from Theorem 7.5.

Corollary 8.13 *Conjecture 1.5 holds true for the graph C_4 and its subgraph P_3 on \mathbb{F}_q^2 .*

Proof The main hypothesis (1.6) of Conjecture 1.5 is satisfied for the graph C_4 and its subgraph P_3 on \mathbb{F}_q^2 :

$$\min\{\delta(C_4), 2\} = 2 > 1 = \delta(P_3).$$

Since the operators Λ_{C_4} and Λ_{P_3} are associated with the graph C_4 and its subgraph P_3 , respectively, the statement of the corollary follows from Theorem 8.11. \blacksquare

9 Operators associated with the graph $K_3 + \text{tail}$ (a kite)

Given $t \in \mathbb{F}_q^*$ and functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^d , we define $\Lambda_{\triangleleft}(f_1, f_2, f_3, f_4)$ as the following value:

$$\frac{1}{q^d |S_t|^2 |S_t^{d-2}|} \sum_{x^1, x^2, x^3, x^4 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) S_t(x^3 - x^4) S_t(x^4 - x^1) \prod_{i=1}^4 f_i(x^i).$$

Note that this operator Λ_{\triangleleft} is related to the graph $K_3 + \text{tail}$ (Figure 1g), and so the normalizing factor $\mathcal{N}(G)$ in (1.4) can be taken as the quantity $q^d |S_t|^2 |S_t^{d-2}|$.

Here, our main problem is to determine all exponents $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that

$$(9.1) \quad \Lambda_{\triangleleft}(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_{p_4}$$

holds for all nonnegative real-valued functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^d . In other words, we are asked to determine all numbers $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$.

Recall that when $d = 2$, we assume that $3 \in \mathbb{F}_q$ is a square number.

Lemma 9.1 (Necessary conditions for the boundedness of $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4)$) *Suppose that (9.1) holds, namely $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$. Then we have*

$$\begin{aligned} \frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} &\leq d, \quad \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq d, \quad \frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} \leq d, \\ \frac{1}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} &\leq 2d - 1, \quad \text{and} \quad \frac{d}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 1. \end{aligned}$$

In particular, if $d = 2$, then it can be shown by Polymake [1, 6] that $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the points: $(0, 0, 1, 0), (0, 1, 0, 0), (0, 1, 0, 1/2), (2/3, 0, 2/3, 0), (1/2, 1/2, 1/2, 0), (5/6, 0, 1/3, 1/2), (1, 0, 0, 0), (1/3, 0, 1/3, 1), (5/8, 5/8, 1/8, 1/2), (1/2, 0, 0, 1), (1/4, 1/4, 1/4, 1), (1/3, 1/3, 0, 1), (1, 0, 0, 1/2)$,

$(2/3, 2/3, 0, 1/2), (2/3, 2/3, 0, 0), (0, 1/2, 0, 1), (0, 1/3, 1/3, 1), (0, 0, 0, 1), (0, 5/6, 1/3, 1/2), (0, 0, 1/2, 1), (0, 0, 0, 0), (0, 2/3, 2/3, 0)$.

Proof To deduce the first inequality, we test (9.1) with $f_1 = f_2 = f_4 = 1_{S_t}$ and $f_3 = \delta_0$. To obtain the second one, we test (9.1) with $f_1 = f_3 = 1_{S_t}$, $f_2 = \delta_0$, and $f_4 = 1_{\mathbb{F}_q^d}$. To get the third one, we test (9.1) with $f_1 = \delta_0$, $f_2 = f_3 = 1_{S_t}$, and $f_4 = 1_{\mathbb{F}_q^d}$. To prove the fourth one, we test (9.1) with $f_1 = f_3 = 1_{S_t}$ and $f_2 = f_4 = \delta_0$. Finally, to obtain the fifth inequality, we test (9.1) with $f_1 = f_4 = \delta_0$ and $f_2 = f_3 = 1_{S_t}$. \blacksquare

9.1 Sufficient conditions for the boundedness of $\Lambda_{\trianglelefteq}$ on \mathbb{F}_q^d

When one of exponents p_1, p_2, p_4 is ∞ , the boundedness problem of $\Lambda_{\trianglelefteq}(p_1, p_2, p_3, p_4)$ can be reduced to that for the K_3 form Λ_{K_3} or the P_2 form Λ_{P_2} .

Proposition 9.2 *Let $1 \leq a, b, c \leq \infty$.*

- (i) *If $\Lambda_{K_3}(a, b, c) \lesssim 1$, then $\Lambda_{\trianglelefteq}(a, b, c, \infty) \lesssim 1$.*
- (ii) *If $\Lambda_{P_2}(a, b, c) \lesssim 1$, then $\Lambda_{\trianglelefteq}(\infty, a, b, c) \lesssim 1$ and $\Lambda_{\trianglelefteq}(a, \infty, b, c) \lesssim 1$.*

Proof For all nonnegative functions f_i , $i = 1, 2, 3, 4$, on \mathbb{F}_q^d , we aim to prove the following inequalities:

$$(9.2) \quad \Lambda_{\trianglelefteq}(f_1, f_2, f_3, f_4) \lesssim \begin{cases} \Lambda_{K_3}(f_1, f_2, f_3) \|f_4\|_{\infty}, \\ \|f_1\|_{\infty} \Lambda_{P_2}(f_2, f_3, f_4), \\ \|f_2\|_{\infty} \Lambda_{P_2}(f_1, f_3, f_4). \end{cases}$$

By the definition, $\Lambda_{\trianglelefteq}(f_1, f_2, f_3, f_4)$ can be expressed as

$$\begin{aligned} & \frac{1}{q^d |S_t| |S_t^{d-2}|} \sum_{x^1, x^2, x^3 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^2 - x^3) S_t(x^3 - x^1) \left(\prod_{i=1}^3 f_i(x^i) \right) \\ & \times \left(\frac{1}{|S_t|} \sum_{x_4 \in \mathbb{F}_q^d} S_t(x^3 - x^4) f_4(x^4) \right). \end{aligned}$$

The sum in the above bracket is clearly dominated by $\|f_4\|_{\infty}$ for all $x^3 \in \mathbb{F}_q^d$. Hence, recalling the definition of $\Lambda_{K_3}(f_1, f_2, f_3)$ in (4.1), we get the first inequality in (9.2):

$$\Lambda_{\trianglelefteq}(f_1, f_2, f_3, f_4) \leq \Lambda_{K_3}(f_1, f_2, f_3) \|f_4\|_{\infty}.$$

Now we prove the second and third inequalities in (9.2). We will only provide the proof of the second inequality, that is,

$$(9.3) \quad \Lambda_{\trianglelefteq}(f_1, f_2, f_3, f_4) \leq \|f_1\|_{\infty} \Lambda_{P_2}(f_2, f_3, f_4).$$

The third inequality can be similarly proved by switching the roles of variables x^1, x^2 . We write $\Lambda_{\trianglelefteq}(f_1, f_2, f_3, f_4)$ as follows:

$$\begin{aligned} & \frac{1}{q^d |S_t|^2} \sum_{x^2, x^3, x^4 \in \mathbb{F}_q^d} S_t(x^2 - x^3) S_t(x^3 - x^4) \left(\prod_{i=2}^4 f_i(x^i) \right) \\ & \times \left(\frac{1}{|S_t^{d-2}|} \sum_{x^1 \in \mathbb{F}_q^d} S_t(x^1 - x^2) S_t(x^3 - x^1) f_1(x^1) \right). \end{aligned}$$

Recall the definition of $\Lambda_{P_2}(f_2, f_3, f_4)$ in (5.1). Then, to prove the inequality (9.3), it will be enough to show that for all $x^2, x^3 \in \mathbb{F}_q^d$ with $\|x^2 - x^3\| = t \neq 0$, the value in the above bracket is $\lesssim \|f_1\|_\infty$. Now, by a simple change of variables, the value in the above bracket is the same as

$$\frac{1}{|S_t^{d-2}|} \sum_{x^1 \in S_t} S_t((x^3 - x^2) - x^1) f_1(x^1 + x^2).$$

This is clearly dominated by

$$\frac{1}{|S_t^{d-2}|} \sum_{x^1 \in S_t} S_t((x^3 - x^2) - x^1) \|f_1\|_\infty.$$

Since $\|x^3 - x^2\| = t \neq 0$, applying Corollary A.4 in the Appendix gives us the desirable estimate. \blacksquare

We address sufficient conditions for the boundedness of Λ_{\triangleleft} on \mathbb{F}_q^d .

The following result can be obtained from Proposition 9.2(i).

Lemma 9.3 *Let Λ_{\triangleleft} be defined on the functions on \mathbb{F}_q^d , $d \geq 2$. Suppose that $1 \leq a, b \leq \infty$ satisfies the following equations:*

$$\frac{1}{a} + \frac{d}{b} \leq d \quad \text{and} \quad \frac{d}{a} + \frac{1}{b} \leq d.$$

Then we have $\Lambda_{\triangleleft}(a, b, \infty, \infty) \lesssim 1$, $\Lambda_{\triangleleft}(a, \infty, b, \infty) \lesssim 1$, $\Lambda_{\triangleleft}(\infty, a, b, \infty) \lesssim 1$.

Proof The statement follows immediately by combining Proposition 9.2(i) with Theorem 4.3. \blacksquare

Proposition 9.2(ii) can be used to deduce the following result.

Lemma 9.4 *Let Λ_{\triangleleft} be defined on the functions on \mathbb{F}_q^d , $d \geq 2$. Suppose that $1 \leq a, b, c \leq \infty$ satisfies one of the following conditions:*

- (i) $0 \leq \frac{1}{a}, \frac{1}{c} \leq \frac{d}{d+1}$ and $\frac{1}{a} + \frac{d}{b} + \frac{1}{c} \leq d$,
- (ii) $0 \leq \frac{1}{a} \leq \frac{d}{d+1} \leq \frac{1}{c} \leq 1$, and $\frac{1}{da} + \frac{1}{b} + \frac{d}{c} \leq d$,
- (iii) $0 \leq \frac{1}{c} \leq \frac{d}{d+1} \leq \frac{1}{a} \leq 1$, and $\frac{d}{a} + \frac{1}{b} + \frac{1}{dc} \leq d$,
- (iv) $\frac{d}{d+1} \leq \frac{1}{a}, \frac{1}{c} \leq 1$ and $\frac{d}{a} + \frac{1}{b} + \frac{d}{c} \leq 2d - 1$.

Then we have $\Lambda_{\triangleleft}(\infty, a, b, c) \lesssim 1$ and $\Lambda_{\triangleleft}(a, \infty, b, c) \lesssim 1$.

Proof From our assumptions on the numbers a, b, c , Theorem 5.3 implies that $\Lambda_{P_2}(a, b, c) \lesssim 1$. Hence, the statement follows by applying Proposition 9.2(ii). \blacksquare

9.2 Boundedness of Λ_{\triangleleft} in two dimensions

Lemmas 9.3 and 9.4 provide nontrivial results available in higher dimensions. In this section, we will show that further improvements can be made in two dimensions. Before we state and prove the improvements, we collect the results in two dimensions, which can be direct consequences of Lemmas 9.3 and 9.4.

To deduce the following result, we will apply Lemma 9.3 with $d = 2$.

Corollary 9.5 Let Λ_{\triangleleft} be defined on functions on \mathbb{F}_q^2 . Then we have $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ provided that (p_1, p_2, p_3, p_4) is one of the following points: $(\infty, \infty, \infty, \infty), (1, \infty, \infty, \infty), (\infty, 1, \infty, \infty), (\infty, \infty, 1, \infty), (3/2, 3/2, \infty, \infty), (3/2, \infty, 3/2, \infty), (\infty, 3/2, 3/2, \infty)$.

Proof Using the first conclusion of Lemma 9.3 with $d = 2$, we see that $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ whenever (p_1, p_2, p_3, p_4) takes the following points: $(\infty, \infty, \infty, \infty), (1, \infty, \infty, \infty), (\infty, 1, \infty, \infty), (3/2, 3/2, \infty, \infty)$.

Next, the second conclusion of Lemma 9.3 with $d = 2$ implies that $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ for the points $(p_1, p_2, p_3, p_4) = (\infty, \infty, 1, \infty), (3/2, \infty, 3/2, \infty)$. Finally, it follows from the third conclusion of Lemma 9.3 with $d = 2$ that $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ for $(p_1, p_2, p_3, p_4) = (\infty, 3/2, 3/2, \infty)$. Hence, the proof is complete. \blacksquare

The following theorem will be proven by applying Lemma 9.4 with $d = 2$.

Corollary 9.6 Let Λ_{\triangleleft} be defined on the functions on \mathbb{F}_q^2 . Suppose that (p_1, p_2, p_3, p_4) is one of the following points: $(\infty, \infty, \infty, 1), (2, \infty, 2, 2), (3/2, \infty, \infty, 3/2), (\infty, \infty, 3/2, 3/2)$. Then we have $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$.

Proof We get that $\Lambda_{\triangleleft}(\infty, \infty, \infty, 1) \lesssim 1$ by using the assumption (ii) and the first conclusion of Lemma 9.4. Invoking the assumption (i) and the second conclusion of Lemma 9.4, one can directly note that $\Lambda_{\triangleleft}(2, \infty, 2, 2) \lesssim 1$ and $\Lambda_{\triangleleft}(3/2, \infty, \infty, 3/2) \lesssim 1$. Finally, to prove that $\Lambda_{\triangleleft}(\infty, \infty, 3/2, 3/2) \lesssim 1$, one can use the assumption (i) and the first conclusion of Lemma 9.4. \blacksquare

We now introduce the connection between $\Lambda_{\triangleleft}(f_1, f_2, f_3, f_4)$ and the bilinear averaging operator.

Proposition 9.7 Let B be the bilinear operator defined as in (4.3). Then, for any nonnegative real-valued functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^2 , we have

$$\Lambda_{\triangleleft}(f_1, f_2, f_3, f_4) = \|B(f_1, f_2) \cdot f_3 \cdot Af_4\|_1,$$

where A denotes the averaging operator over the circle in \mathbb{F}_q^2 .

Proof In two dimensions, $\Lambda_{\triangleleft}(f_1, f_2, f_3, f_4)$ can be rewritten as the following form:

$$\frac{1}{q^2|S_t|^2} \sum_{x^1, x^2, x^3, x^4 \in \mathbb{F}_q^2} S_t(x^1 - x^2) S_t(x^3 - x^2) S_t(x^3 - x^4) S_t(x^3 - x^1) \prod_{i=1}^4 f_i(x^i).$$

By the change of variables by putting $y^1 = x^3 - x^1$, $y^2 = x^3 - x^2$, $y^3 = x^3$, $y^4 = x^3 - x^4$, the value $\Lambda_{\triangleleft}(f_1, f_2, f_3, f_4)$ becomes

$$\frac{1}{q^2|S_t|^2} \sum_{y^1, y^2, y^3, y^4 \in \mathbb{F}_q^2} S_t(y^2 - y^1) S_t(y^2) S_t(y^4) S_t(y^1) f_1(y^3 - y^1) f_2(y^3 - y^2) f_3(y^3) f_4(y^3 - y^4).$$

This can be expressed as follows:

$$\frac{1}{q^2} \sum_{y^3 \in \mathbb{F}_q^2} f_3(y^3) \left(\frac{1}{|S_t|} \sum_{y^4 \in S_t} f_4(y^3 - y^4) \right) \left(\frac{1}{|S_t|} \sum_{y^1, y^2 \in S_t: \|y^2 - y^1\| = t} f_1(y^3 - y^1) f_2(y^3 - y^2) \right).$$

Recalling the definitions of the averaging operator in (2.1) and the bilinear averaging operator in (4.3), it follows that

$$\Lambda_{\triangleleft}(f_1, f_2, f_3, f_4) = \frac{1}{q^2} \sum_{y^3 \in \mathbb{F}_q^2} f_3(y^3) A f_4(y^3) B(f_1, f_2)(y^3).$$

By the definition of the normalized norm $\|\cdot\|_1$, the statement follows. \blacksquare

For $1 \leq p_1, p_2, p_3, p_4 \leq \infty$, recall that the notation $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ is used if the following estimate holds for all subsets E, F, G, H of \mathbb{F}_q^2 :

$$\Lambda_{\triangleleft}(E, F, G, H) \lesssim \|E\|_{p_1} \|F\|_{p_2} \|G\|_{p_3} \|H\|_{p_4},$$

and this estimate is referred to as the restricted strong-type $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4)$ estimate.

The following theorem is our main result in two dimensions, which gives a new restricted strong-type estimate for the boundedness on the operator Λ_{\triangleleft} .

Theorem 9.8 *Let Λ_{\triangleleft} be defined on functions on \mathbb{F}_q^2 . Let $1 \leq p_3, p_4 \leq \infty$. Then the following statements are valid for all subsets E, F of \mathbb{F}_q^2 and all nonnegative functions f_3, f_4 on \mathbb{F}_q^2 .*

(i) *If $2 \leq p_3 \leq \infty$, $3/2 \leq p_4 \leq \infty$, and $\frac{1}{p_3} + \frac{1}{2p_4} \leq \frac{1}{2}$, then we have*

$$\Lambda_{\triangleleft}(E, F, f_3, f_4) \lesssim \|E\|_2 \|F\|_2 \|f_3\|_{p_3} \|f_4\|_{p_4}.$$

(ii) *If $2 \leq p_3 \leq \infty$, $4/3 \leq p_4 \leq 3/2$, and $\frac{1}{p_3} + \frac{2}{p_4} \leq \frac{3}{2}$, then we have*

$$\Lambda_{\triangleleft}(E, F, f_3, f_4) \lesssim \|E\|_2 \|F\|_2 \|f_3\|_{p_3} \|f_4\|_{p_4}.$$

Proof Let E, F be subsets of \mathbb{F}_q^2 and f, g be nonnegative real-valued functions on \mathbb{F}_q^2 . By Proposition 9.7 and Hölder's inequality, it follows that for $2 \leq p_3 \leq \infty$,

$$\Lambda_{\triangleleft}(E, F, f_3, f_4) \leq \|B(E, F)\|_2 \|f_3\|_{p_3} \|A f_4\|_{\frac{2p_3}{p_3-2}}.$$

Here, we also notice that $2 \leq \frac{2p_3}{p_3-2} \leq \infty$. Since $\|B(E, F)\|_2 \lesssim \|E\|_2 \|F\|_2$ by Lemma 4.6, we see that

$$\Lambda_{\triangleleft}(E, F, f_3, f_4) \leq \|E\|_2 \|F\|_2 \|f_3\|_{p_3} \|A f_4\|_{\frac{2p_3}{p_3-2}}.$$

Hence, to complete the proof, it suffices to show that for all exponents p_3, p_4 satisfying the assumptions of the theorem, we have

$$(9.4) \quad A\left(p_4 \rightarrow \frac{2p_3}{p_3-2}\right) \lesssim 1.$$

To prove this, we first recall from Theorem 2.3 with $d = 2$ that $A(p \rightarrow r) \lesssim 1$ for any numbers $1 \leq p, r \leq \infty$ such that $(1/p, 1/r)$ lies on the convex hull of points $(0, 0), (0, 1), (1, 1)$, and $(\frac{2}{3}, \frac{1}{3})$. Also, invoke Lemma 2.5 to find the equations indicating the endpoint estimates for $A(p \rightarrow r) \lesssim 1$. Using those averaging estimates with

$p = p_4, r = \frac{2p_3}{p_3-2}$, the inequality (9.4) can be obtained by a direct computation, where we also use the fact that $2 \leq r = \frac{2p_3}{p_3-2} \leq \infty$. \blacksquare

The following corollary is a direct consequence of Theorem 9.8.

Corollary 9.9 *Let Λ_{\triangleleft} be defined on functions on \mathbb{F}_q^2 . Then we have $\Lambda_{\triangleleft}(2, 2, 2, \infty) \lesssim 1$.*

Proof The statement follows by a direct application of Theorem 9.8(i). \blacksquare

The lemma below shows that the exponents for $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$ are more restricted than those for $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ up to the endpoints. This also provides a positive answer to Question 1.2 since the graph $K_3 + \text{tail}$ is a subgraph of the graph $C_4 + \text{diagonal}$.

Lemma 9.10 *Let Λ_{\diamond_t} and Λ_{\triangleleft} be the operators acting on the functions on \mathbb{F}_q^2 . Suppose that $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$ for $1 \leq p_1, p_2, p_3, p_4 \leq \infty$. Then we have $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ except for the point $(2, 2, 2, \infty)$. In addition, we have $\Lambda_{\triangleleft}(2, 2, 2, \infty) \lesssim 1$.*

Proof Assume that $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$ for $1 \leq p_1, p_2, p_3, p_4 \leq \infty$. Then, by Lemma 6.1, the point $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the following points: $(0, 0, 1, 0), (0, 1, 0, 0), (0, 0, 0, 1), (1/2, 0, 1/2, 1/2), (2/3, 2/3, 0, 0), (1, 0, 0, 0), (2/3, 0, 2/3, 0), (1/2, 1/2, 1/2, 0), (2/3, 0, 0, 2/3), (0, 2/3, 2/3, 0), (0, 0, 0, 0), (0, 0, 2/3, 2/3)$.

By Corollaries 9.5 and 9.6, the strong-type estimate $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$ holds for all the above points $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ except for $(1/2, 1/2, 1/2, 0)$. Moreover, we know from Corollary 9.9 that $\Lambda_{\triangleleft}(2, 2, 2, \infty) \lesssim 1$. Hence, the statement follows by interpolating those points. \blacksquare

Remark 9.11 The reverse statement of Lemma 9.10 is not true. To see this, observe from Theorem 9.8(ii) that $\Lambda_{\triangleleft}(2, 2, 6, 3/2) \lesssim 1$. In addition, by Lemma 6.1, notice that $\Lambda_{\diamond_t}(2, 2, 6, 3/2)$ cannot be bounded.

Corollary 9.12 *Conjecture 1.5 holds up to endpoints for the graph $C_4 + \text{diagonal}$ and its subgraph $K_3 + \text{tail}$ in \mathbb{F}_q^2 .*

Proof The operators Λ_{\diamond_t} and Λ_{\triangleleft} are associated with the $C_4 + \text{diagonal}$ and its subgraph $K_3 + \text{tail}$ in \mathbb{F}_q^2 , respectively. Hence, invoking Lemma 9.10, the proof is reduced to showing that the $C_4 + \text{diagonal}$ and its subgraph $K_3 + \text{tail}$ satisfy the main hypothesis (1.6) of Conjecture 1.5. However, it is clear that

$$\min\{\delta(C_4 + \text{diagonal}), 2\} = 2 > 1 = \delta(K_3 + \text{tail}).$$

Thus, the proof is complete. \blacksquare

The following result shows that there exists an inclusive relation between boundedness exponents for the operators corresponding to the graphs C_4 and $K_3 + \text{tail}$, although they are not subgraphs of each other.

Lemma 9.13 *Let Λ_{C_4} and Λ_{\triangleleft} be defined on functions on \mathbb{F}_q^2 and let $1 \leq p_1, p_2, p_3, p_4 \leq \infty$. Then if $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$, we have $\Lambda_{\triangleleft}(p_1, p_2, p_3, p_4) \lesssim 1$.*

Proof First, by Theorem 7.5, note that $\Lambda_{C_4}(2, 2, 2, \infty)$ cannot be bounded. Now suppose that $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$. Then $(p_1, p_2, p_3, p_4) \neq (2, 2, 2, \infty)$. Using Proposition 7.6, we get $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$. Then the statement follows immediately from Lemma 9.10. \blacksquare

By combining Remark 9.11 and Proposition 7.6, it is clear that the reverse of Lemma 9.13 does not hold. Notice that Lemma 9.13 provides an example to satisfy Conjecture 1.5 without the hypothesis that G' is a subgraph of the graph G .

10 Boundedness problems for the Y-shaped graph

In this section, we study the boundedness of the operator for the Y-shaped graph in Figure 1h. For $t \in \mathbb{F}_q^*$, the Y-shaped form Λ_Y is defined by

(10.1)

$$\Lambda_Y(f_1, f_2, f_3, f_4) = \frac{1}{q^d |S_t|^3} \sum_{x^1, x^2, x^3, x^4 \in \mathbb{F}_q^d} S_t(x^3 - x^1) S_t(x^3 - x^2) S_t(x^3 - x^4) \prod_{i=1}^4 f_i(x^i),$$

where functions f_i , $i = 1, 2, 3, 4$, are defined on \mathbb{F}_q^d . Note that this operator Λ_Y is related to the Y-shaped graph, and so the normalizing factor $\mathcal{N}(G)$ in (1.4) can be taken as $q^d |S_t|^3$.

We aim to find all numbers $1 \leq p_1, p_2, p_3, p_4 \leq \infty$ such that $\Lambda_Y(p_1, p_2, p_3, p_4) \lesssim 1$.

Lemma 10.1 (Necessary conditions for the boundedness of $\Lambda_Y(p_1, p_2, p_3, p_4)$) *Let $1 \leq p_i \leq \infty$, $1 \leq i \leq 4$. Suppose that $\Lambda_Y(p_1, p_2, p_3, p_4) \lesssim 1$. Then all the following inequalities are satisfied: $\frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d$, $\frac{d}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 3d - 2$, $\frac{d}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} \leq 2d - 1$, $\frac{d}{p_1} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 1$, $\frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 1$, $\frac{d}{p_1} + \frac{1}{p_3} \leq d$, $\frac{d}{p_2} + \frac{1}{p_3} \leq d$, $\frac{1}{p_3} + \frac{d}{p_4} \leq d$.*

In particular, if $d = 2$, then it can be shown by Polymake [1, 6] that $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the points: $(0, 0, 1, 0)$, $(1, 0, 0, 1/2)$, $(1, 0, 1/2, 0)$, $(1/2, 1, 0, 1/2)$, $(1, 1/2, 0, 1/2)$, $(1/2, 1/2, 0, 1)$, $(1, 0, 1/3, 1/3)$, $(1/2, 5/6, 1/3, 0)$, $(1, 1/2, 0, 0)$, $(1/2, 0, 1/3, 5/6)$, $(1/2, 1, 0, 0)$, $(1, 1/3, 1/3, 0)$, $(1/2, 0, 0, 1)$, $(1, 0, 0, 0)$, $(0, 0, 0, 0)$, $(0, 0, 0, 1)$, $(0, 1, 0, 0)$, $(0, 0, 2/3, 2/3)$, $(0, 5/6, 1/3, 1/2)$, $(0, 1, 0, 1/2)$, $(0, 1/2, 1/3, 5/6)$, $(0, 1/2, 0, 1)$, $(0, 2/3, 2/3, 0)$.

Proof By a direct computation, the conclusions of the lemma easily follow by testing the inequality

$$\Lambda_Y(f_1, f_2, f_3, f_4) \leq \Lambda_Y(p_1, p_2, p_3, p_4) \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_{p_4},$$

with the following specific functions, respectively:

1) $f_1 = f_2 = f_4 = 1_{S_t}$, $f_3 = \delta_0$.	2) $f_1 = f_2 = f_4 = \delta_0$, $f_3 = 1_{S_t}$.
3) $f_1 = f_2 = \delta_0$, $f_3 = 1_{S_t}$, $f_4 = 1_{\mathbb{F}_q^d}$.	4) $f_1 = f_2 = \delta_0$, $f_2 = 1_{\mathbb{F}_q^d}$, $f_3 = 1_{S_t}$.
5) $f_1 = 1_{\mathbb{F}_q^d}$, $f_2 = f_4 = \delta_0$, $f_3 = 1_{S_t}$.	6) $f_1 = \delta_0$, $f_2 = f_4 = 1_{\mathbb{F}_q^d}$, $f_3 = 1_{S_t}$.
7) $f_1 = f_4 = 1_{\mathbb{F}_q^d}$, $f_2 = \delta_0$, $f_3 = 1_{S_t}$.	8) $f_1 = f_2 = 1_{\mathbb{F}_q^d}$, $f_3 = 1_{S_t}$, $f_4 = \delta_0$.

\blacksquare

10.1 Sufficient conditions for the boundedness of Λ_Y on \mathbb{F}_q^d

It is not hard to observe that the boundedness problem for the Y-shaped form can be reduced to the spherical averaging estimate. Indeed, the value $\Lambda_Y(f_1, f_2, f_3, f_4)$ in (10.1) can be written by

$$\Lambda_Y(f_1, f_2, f_3, f_4) = \frac{1}{q^d} \sum_{x^3 \in \mathbb{F}_q^d} f_3(x^3) \prod_{i=1,2,4} \left(\frac{1}{|S_t|} \sum_{x^i \in \mathbb{F}_q^d} S_t(x^3 - x^i) f_i(x^i) \right).$$

Invoking the definition of the averaging operator $A = A_{S_t}$ in (2.2), we get

$$\Lambda_Y(f_1, f_2, f_3, f_4) = \frac{1}{q^d} \sum_{x^3 \in \mathbb{F}_q^d} f_3(x^3) A f_1(x^3) A f_2(x^3) A f_4(x^4) = \|A f_1 \cdot A f_2 \cdot f_3 \cdot A f_4\|_1.$$

By Hölder's inequality and the nesting property of the norm $\|\cdot\|_p$, we get

$$(10.2) \quad \Lambda_Y(f_1, f_2, f_3, f_4) \leq \|A f_1\|_{r_1} \|A f_2\|_{r_2} \|f_3\|_{p_3} \|A f_4\|_{r_4} \quad \text{if } \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{p_3} + \frac{1}{r_4} \leq 1.$$

Proposition 10.2 *Let $1 \leq p_1, p_2, p_3, p_4, r_1, r_2, r_4 \leq \infty$ be extended real numbers which satisfy the following assumptions: $\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{p_3} + \frac{1}{r_4} \leq 1$ and $A(p_i \rightarrow r_i) \lesssim 1$ for all $i = 1, 2, 4$. Then we have*

$$\Lambda_Y(p_1, p_2, p_3, p_4) \lesssim 1.$$

Proof By combining the inequality (10.2) with our assumptions on the averaging estimates, it follows that for all functions $f_i, i = 1, 2, 3, 4$, on \mathbb{F}_q^d ,

$$\Lambda_Y(f_1, f_2, f_3, f_4) \lesssim \|f_1\|_{p_1} \|f_2\|_{p_2} \|f_3\|_{p_3} \|f_4\|_{p_4}.$$

This completes the proof. ■

The following result provides lots of sufficient conditions for the boundedness of the Y-shaped form.

Proposition 10.3 *Let $1 \leq p_1, p_2, p_3, p_4 \leq \infty$, and let Λ_Y be the Y-shaped form on \mathbb{F}_q^d . Then $\Lambda_Y(p_1, p_2, p_3, p_4) \lesssim 1$ provided that one of the following conditions is satisfied:*

- (i) $0 \leq \frac{1}{p_1}, \frac{1}{p_2}, \frac{1}{p_4} \leq \frac{d}{d+1}$ and $\frac{1}{p_1} + \frac{1}{p_2} + \frac{d}{p_3} + \frac{1}{p_4} \leq d$.
- (ii) $0 \leq \frac{1}{p_1}, \frac{1}{p_2} \leq \frac{d}{d+1} \leq \frac{1}{p_4}$ and $\frac{1}{d p_1} + \frac{1}{d p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq d$.
- (iii) $0 \leq \frac{1}{p_1}, \frac{1}{p_4} \leq \frac{d}{d+1} \leq \frac{1}{p_2} \leq 1$ and $\frac{1}{d p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{1}{d p_4} \leq d$.
- (iv) $0 \leq \frac{1}{p_2}, \frac{1}{p_4} \leq \frac{d}{d+1} \leq \frac{1}{p_1} \leq 1$ and $\frac{d}{p_1} + \frac{1}{d p_2} + \frac{1}{p_3} + \frac{1}{d p_4} \leq d$.
- (v) $0 \leq \frac{1}{p_1} \leq \frac{d}{d+1} \leq \frac{1}{p_2}, \frac{1}{p_4} \leq 1$ and $\frac{1}{d p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 1$.
- (vi) $0 \leq \frac{1}{p_2} \leq \frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_4} \leq 1$ and $\frac{d}{p_1} + \frac{1}{d p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 2d - 1$.
- (vii) $0 \leq \frac{1}{p_4} \leq \frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_2} \leq 1$ and $\frac{d}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{1}{d p_4} \leq 2d - 1$.
- (viii) $\frac{d}{d+1} \leq \frac{1}{p_1}, \frac{1}{p_2}, \frac{1}{p_4} \leq 1$ and $\frac{d}{p_1} + \frac{d}{p_2} + \frac{1}{p_3} + \frac{d}{p_4} \leq 3d - 2$.

Proof The proof uses Proposition 10.2 and the sharp averaging estimates in Lemma 2.5. The proof of this theorem is similar to that of Theorem 5.3. Therefore, we leave the detail of the proof to readers. ■

Conjecture 1.5 is also supported by the following theorem.

Theorem 10.4 Let Λ_{\diamond_t} and Λ_Y be the operators acting on the functions on \mathbb{F}_q^2 . If $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$ with $1 \leq p_1, p_2, p_3, p_4 \leq \infty$, then $\Lambda_Y(p_1, p_2, p_3, p_4) \lesssim 1$.

Proof Assume that $\Lambda_{\diamond_t}(p_1, p_2, p_3, p_4) \lesssim 1$. Then, by Lemma 6.1, $(1/p_1, 1/p_2, 1/p_3, 1/p_4)$ is contained in the convex hull of the points $(0, 0, 1, 0), (0, 1, 0, 0), (0, 0, 0, 1), (1/2, 0, 1/2, 1/2), (2/3, 2/3, 0, 0), (1, 0, 0, 0), (2/3, 0, 2/3, 0), (1/2, 1/2, 1/2, 0), (2/3, 0, 0, 2/3), (0, 2/3, 2/3, 0), (0, 0, 0, 0), (0, 0, 2/3, 2/3)$. By interpolating those critical points, it suffices to check that each critical point above satisfies one of the eight hypotheses of Proposition 10.3 with $d = 2$. However, this can be easily shown by a direct computation. For example, for the critical point $(1/p_1, 1/p_2, 1/p_3, 1/p_4) = (1/2, 1/2, 1/2, 0)$, we can invoke the hypothesis (i) of Proposition 10.3 with $d = 2$ and obtain that $\Lambda_Y(2, 2, 2, \infty) \lesssim 1$. In the same way, it can be easily proven for other critical points. ■

Remark 10.5 The reverse statement of Theorem 10.4 is not true. To find a counterexample, we can take $p_1 = p_3 = \infty, p_2 = p_4 = 3/2$. Indeed, by the hypothesis (5) of Proposition 10.3 with $d = 2$, we see that $\Lambda_Y(\infty, 3/2, \infty, 3/2) \lesssim 1$. However, $\Lambda_{\diamond_t}(\infty, 3/2, \infty, 3/2)$ is not bounded, which follows from Lemma 6.1 with $d = 2$.

The following corollary proposes some possibility that the assumption of the subgraph in Conjecture 1.5 can be dropped.

Corollary 10.6 Let Λ_{C_4} and Λ_Y be the operators acting on the functions on \mathbb{F}_q^2 . If $\Lambda_{C_4}(p_1, p_2, p_3, p_4) \lesssim 1$ with $1 \leq p_1, p_2, p_3, p_4 \leq \infty$, then $\Lambda_Y(p_1, p_2, p_3, p_4) \lesssim 1$.

Proof The statement of the corollary follows immediately by combining Proposition 7.6 and Theorem 10.4. ■

Combining all the results obtained so far, we get the following theorem:

Theorem 10.7 When $d = 2$ and $n = 3, 4$, Conjecture 1.5 is true, where we accept boundedness results up to endpoints in the case when G is the $C_4 +$ diagonal and its subgraph G' is the $K_3 +$ tail.

Proof By Corollary 5.7 for $n = 3$, and by Corollaries 8.10, 8.13, 9.12, and 10.6 for $n = 4$, we have proven that for $d = 2$ and $n = 3, 4$, there is the required inclusive boundedness relationship between any two operators corresponding to arbitrary connected ordered graph G and its subgraph G' except for the following three cases:

- (I) $G = C_4 +$ diagonal and $G' = C_4$.
- (II) $K_3 +$ tail and $G' = Y$ -shape.
- (III) $G = K_3 +$ tail and $G' = P_3$.

However, since $\delta(G) = \delta(G')$ for each case of (I), (II), and (III), they do not satisfy the main hypothesis (1.6) of Conjecture 1.5. Hence, they cannot be counterexamples contradicting Conjecture 1.5 and so there is no counterexample against Conjecture 1.5, as required. ■

Acknowledgments A. Iosevich, D. Koh, and T. Pham would like to thank the VIASM for the hospitality and for the excellent working conditions.

A Appendix

In this appendix, we introduce the number of intersection points of two spheres in \mathbb{F}_q^d . Let η denote the quadratic character of \mathbb{F}_q^* , namely, $\eta(s) = 1$ for a square number s in \mathbb{F}_q^* , and $\eta(s) = -1$ otherwise.

Definition A.1 Given a nonzero vector m in \mathbb{F}_q^d , and $t, b \in \mathbb{F}_q$, we define $N(m, t, b)$ to be the number of common solutions $x \in \mathbb{F}_q^d$ of the following equations: $\|x\| = t$, $m \cdot x = b$.

Notice that the value of $N(m, t, b)$ is the number of all intersection points between the sphere S_t and the plane $\{x \in \mathbb{F}_q^d : m \cdot x = b\}$. The explicit value of it is well known as follows.

Lemma A.2 Let $b, t \in \mathbb{F}_q$, and let m be a nonzero element in \mathbb{F}_q^d , $d \geq 2$. Then the following statements hold:

(i) If $\|m\| \neq 0$ and $b^2 - t\|m\| = 0$, then

$$N(m, t, b) = \begin{cases} q^{d-2}, & \text{if } d \text{ is even,} \\ q^{d-2} + q^{\frac{d-3}{2}}(q-1)\eta\left((-1)^{\frac{d-1}{2}}\|m\|\right), & \text{if } d \text{ is odd.} \end{cases}$$

(ii) If $\|m\| \neq 0$ and $b^2 - t\|m\| \neq 0$, then

$$N(m, t, b) = \begin{cases} q^{d-2} + q^{\frac{d-2}{2}}\eta\left((-1)^{\frac{d}{2}}(b^2 - t\|m\|)\right), & \text{if } d \text{ is even,} \\ q^{d-2} - q^{\frac{d-3}{2}}\eta\left((-1)^{\frac{d-1}{2}}\|m\|\right), & \text{if } d \text{ is odd.} \end{cases}$$

(iii) If $\|m\| = 0 = b^2 - t\|m\|$, then

$$N(m, t, b) = \begin{cases} q^{d-2} + v(t)q^{\frac{d-2}{2}}\eta\left((-1)^{\frac{d}{2}}\right), & \text{if } d \text{ is even,} \\ q^{d-2} - q^{\frac{d-1}{2}}\eta\left((-1)^{\frac{d-1}{2}}t\right), & \text{if } d \text{ is odd,} \end{cases}$$

where $v(t) = -1$ if $t \in \mathbb{F}_q^*$ and $v(0) = q-1$.

(iv) If $\|m\| = 0$ and $b^2 - t\|m\| \neq 0$, then $N(m, t, b) = q^{d-2}$.

Proof See Exercises 6.31–6.34 in [14], or one can prove it by using the discrete Fourier analysis with the explicit value of the Gauss sum. ■

By a direct application of Lemma A.2, one can find the explicit number of the intersections of two spheres over finite fields. Precisely, we have the following result.

Theorem A.3 Given a nonzero vector $m \in \mathbb{F}_q^d$ and $t, j \in \mathbb{F}_q$, let

$$\Theta(m, t, j) := \{x \in S_t : \|x - m\| = j\}.$$

If $m \in S_\ell$, then $|\Theta(m, t, j)| = N\left(m, t, \frac{t+\ell-j}{2}\right)$.

Proof Since $\|x - m\| = t + \ell - 2m \cdot x$ for $x \in S_t, m \in S_\ell$, it is clear that $\Theta(m, t, j)$ is the number of common solutions x of the following equations:

$$\|x\| = t, \quad m \cdot x = \frac{t + \ell - j}{2}.$$

Hence, by the definition of N , we obtain the required conclusion. \blacksquare

Corollary A.4 Let $t \in \mathbb{F}_q^*$ and $\ell \in \mathbb{F}_q$. Then, for every nonzero vector $m \in S_\ell$, we have

$$\sum_{x \in S_t: \|x - m\| = t} 1 \sim q^{d-2}$$

excepting for the following three cases:

- 1) $d = 2, \ell \neq 0, \eta(t\ell - \ell^2/4) = -1$. 2) $d = 2, \ell = 0, \eta(-1) = 1$.
- 3) $d = 3, \ell = 0, \eta(-t) = 1$.

For each of those three cases, the value in the above sum takes zero. On the other hand, if $d = 2, \ell \neq 0$, and $\eta(t\ell - \ell^2/4) = 1$, the value in the above sum is exactly two.

Proof It follows from Theorem A.3 that for any $\|m\| = \ell$,

$$\sum_{x \in S_t: \|x - m\| = t} 1 = N\left(m, t, \frac{\ell}{2}\right),$$

and so the corollary is a direct consequence of Lemma A.2(i)–(iii). \blacksquare

References

- [1] B. Assarf, E. Gawrilow, K. Herr, M. Joswig, B. Lorenz, A. Paffenholz, and T. Rehn, *Computing convex hulls and counting integer points with polymake*. Math. Program. Comput. 9(2017), no. 1, 1–38.
- [2] M. Bennett, J. Chapman, D. Covert, D. Hart, A. Iosevich, and J. Pakianathan, *Long paths in the distance graph over large subsets of vector spaces over finite fields*. J. Korean Math. 53(2016), 115–126.
- [3] M. Bennett, D. Hart, A. Iosevich, J. Pakianathan, and M. Rudnev, *Group actions and geometric combinatorics in F_q^d* . Forum Math. 29(2017), no. 1, 91–110.
- [4] A. Carbery, B. Stones, and J. Wright, *Averages in vector spaces over finite fields*. Math. Proc. Camb. Philos. Soc. 144(2008), no. 1, 13–27.
- [5] J. Chapman, M. Erdo\u{g}an, D. Hart, A. Iosevich, and D. Koh, *Pinned distance sets, Wolff's exponent in finite fields and sum-product estimates*. Math. Z. 271(2012), no. 1–2, 63–93.
- [6] E. Gawrilow and M. Joswig, *Polymake: a framework for analyzing convex polytopes*. In: Polytopes – combinatorics and computation (Oberwolfach, 1997), Birkhäuser, Basel, pp. 43–73.
- [7] A. Greenleaf, A. Iosevich, B. Krause, and A. Liu, *L^p estimates for bilinear generalized radon transforms in the plane*. In: Combinatorial and additive number theory, Springer, Cham, 2022, pp. 179–198.
- [8] B. Hanson, B. Lund, and O. Roche-Newton, *On distinct perpendicular bisectors and pinned distances in finite fields*. Finite Fields Appl. 37(2016), 240–264.
- [9] A. Iosevich, G. Jardine, and B. McDonald, *Cycles of arbitrary length in distance graphs on F_q^d* . Proc. Steklov Inst. Math. 314(2021), no. 1, 27–43.
- [10] A. Iosevich and M. Rudnev, *Erdős distance problem in vector spaces over finite fields*. Trans. Amer. Math. Soc. 359(2007), no. 12, 6127–6142.
- [11] D. Koh and S. Lee, *Averages and maximal averages over product j -varieties in finite fields*. Finite Fields Appl. 79(2022), 101984.

- [12] D. Koh and C. Y. Shen, *Extension and averaging operators for finite fields*. Proc. Edinb. Math. Soc. 56(2013), no. 2, 599–614.
- [13] D. Koh, C. Y. Shen, and I. Shparlinski, *Averaging operators over homogeneous varieties over finite fields*. J. Geom. Anal. 26(2016), no. 2, 1415–1441.
- [14] R. Lidl and H. Niederreiter, *Finite fields*, Cambridge University Press, Cambridge, 1997.
- [15] B. Murphy, G. Petridis, T. Pham, M. Rudnev, and S. Stevenson, *On the pinned distances problem in positive characteristic*. J. Lond. Math. Soc. 105(2022), no. 1, 469–499.
- [16] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. In: With the assistance of Timothy S. Murphy, Princeton Mathematical Series, 43, Monographs in Harmonic Analysis, III, Princeton University Press, Princeton, NJ, 1993, xiv + 695 pp.

Department of Mathematics, University of Rochester, Rochester, NY, United States

e-mail: pbhowmik@ur.rochester.edu

Department of Mathematics, University of Rochester, Rochester, NY, United States

e-mail: alex.iosevich@rochester.edu

Department of Mathematics, Chungbuk National University, Cheongju, South Korea

e-mail: koh131@chungbuk.ac.kr

University of Science, Vietnam National University Hanoi, Hanoi, Vietnam

e-mail: thangham.math@vnu.edu.vn