

The weight two compactly supported Euler characteristic of moduli spaces of curves

SAM PAYNE AND THOMAS WILLWACHER

Dedicated to Enrico Arbarello with gratitude and admiration

Abstract: We derive a formula for the generating function for the weight two compactly supported \mathbb{S}_n -equivariant Euler characteristics of the moduli spaces of curves $\mathcal{M}_{g,n}$, using graph complexes and calculations inspired by operadic methods.

1. Introduction

The rational cohomology of moduli spaces of curves is of central interest not only in algebraic geometry but also in topology, geometric group theory, and mathematical physics. In the stable range, the cohomology is completely understood, thanks to the celebrated proof of Mumford’s conjecture via stable homotopy theory [17]. Outside of the stable range, even the most basic properties, such as the parameters k, g, n for which the rational cohomology $H^k(\mathcal{M}_{g,n})$ is zero or non-zero, have remained largely mysterious [18, §9].

Algebraic geometry techniques have recently yielded new nonvanishing results and lower bounds on dimensions of cohomology groups in the unstable range [4, 5, 19]. The algebraic structure of $\mathcal{M}_{g,n}$ as a Deligne-Mumford stack endows its rational cohomology with a mixed Hodge structure. Nonvanishing results and lower bounds on dimensions of cohomology groups are then obtained by studying the associated graded of the weight filtration, one weight at a time. The first two nontrivial graded pieces of the weight filtration may be expressed in terms of the cohomology of commutative graph complexes. The weight zero cohomology is naturally identified with the cohomology of the original commutative graph complex studied by Kontsevich [13]. In weight two, which is the focus of the present paper, one finds more general graph complexes closely related to those appearing in recent work on the embedding calculus [6]. The identification of weight two cohomology with

Received January 17, 2022.

the cohomology of these graph complexes relies fundamentally on the results and methods of Arbarello and Cornalba [1].

The cohomology groups of the graph complexes that arise in this context are far from fully understood. In such cases where cohomology groups cannot be computed directly, Euler characteristics often yield valuable insights. For instance, Harer and Zagier famously computed the Euler characteristics of \mathcal{M}_g and deduced that these moduli spaces have cohomology groups that grow superexponentially in both even and odd degrees [11]. This is especially striking since the stable cohomology grows subexponentially and, at the time of their writing and for nearly twenty years after, there were no known examples where $H^k(\mathcal{M}_g)$ is nonzero for odd k . The first such example was $H^5(\mathcal{M}_4)$ [22]. All other known examples are proved using the associated graded of the weight filtration and graph complex techniques; see [4, 19].

Even if one is primarily interested in the cohomology of \mathcal{M}_g , for inductive arguments using the boundary structure of the moduli space of stable curves as in [1, 9], it is essential to understand the cohomology of moduli spaces of curves with marked points $\mathcal{M}_{g,n}$, and to understand these not only as vector spaces but as representations of the symmetric group \mathbb{S}_n , with the action induced by permuting marked points. Thus, one is naturally led to investigate the \mathbb{S}_n -equivariant Euler characteristic of the associated graded of the weight filtration on the rational cohomology of $\mathcal{M}_{g,n}$. By Poincaré duality [20, Theorem 6.3], it is equivalent to study the associated graded of the weight filtration on compactly supported cohomology.

A closed formula for the weight 0 compactly supported Euler characteristic of moduli spaces of curves is given in [3], and the weight 1 compactly supported cohomology vanishes. Our main result is a formula for the generating function for the \mathbb{S}_n -equivariant Euler characteristic of the weight 2 compactly supported cohomology of $\mathcal{M}_{g,n}$.

1.1. Statement of main result

Let V_λ denote the irreducible representation of the symmetric group corresponding to a Young diagram λ . For any graded \mathbb{Q} -vector space $W = \bigoplus_k W^k$ with finite dimensional graded pieces and an action of the symmetric group \mathbb{S}_n , we consider the decomposition into isotypical components

$$W^k \cong \bigoplus_\lambda V_\lambda \otimes \mathbb{Q}^{n_{\lambda,k}}.$$

Then the equivariant Euler characteristic of W is defined to be the symmetric function

$$\chi^{\mathbb{S}}(W) = \sum_{k,\lambda} (-1)^k n_{\lambda,k} s_{\lambda},$$

with s_{λ} the Schur function corresponding to λ .

We are particularly interested in the associated graded of the weight filtration

$$\text{gr}_i H_c^{\bullet}(\mathcal{M}_{g,n}) := W_i H_c^{\bullet}(\mathcal{M}_{g,n}) / W_{i-1} H_c^{\bullet}(\mathcal{M}_{g,n}),$$

which carries a natural \mathbb{S}_n -action induced by permuting the marked points. Let

$$\chi_i^{\mathbb{S}}(\mathcal{M}_{g,n}) := \chi^{\mathbb{S}}(\text{gr}_i H_c^{\bullet}(\mathcal{M}_{g,n})).$$

We will give a formula for the generating function for the weight 2 equivariant Euler characteristics

$$(1) \quad \omega_2 := \sum_{\substack{g,n \\ 2g+n \geq 3}} \chi_2^{\mathbb{S}}(\mathcal{M}_{g,n}) \hbar^g$$

in the ring of formal symmetric power series $\Lambda = \mathbb{Q}[[p_1, p_2, \dots]]$. Here $p_{\ell} = x_1^{\ell} + x_2^{\ell} + x_3^{\ell} + \dots$ is the degree ℓ power sum. Let $P_{\ell} := 1 + p_{\ell}$ be the inhomogeneous power sum. We define

$$(2) \quad Z_{\ell} := \frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) \frac{P_d}{\hbar^d},$$

where μ is the Möbius function and \hbar is a formal variable that will count the genus, as well as

$$(3) \quad \psi_0(z) := - \sum_{j=1}^{\infty} \frac{B_j}{j} \frac{1}{(-z)^j} \quad \text{and} \quad \psi_1(z) := - \sum_{j=0}^{\infty} B_j \frac{1}{(-z)^{j+1}}.$$

Note that $\psi_0(z) + \log z$ and $\psi_1(z)$ are the asymptotic expansions as $z \rightarrow \infty$ of the digamma and trigamma function respectively. We identify the logarithm with its power series expansion

$$\log(1-z) = - \sum_{j=1}^{\infty} z^j.$$

Theorem 1.1. *The generating function for \mathbb{S} -equivariant weight 2 compactly supported Euler characteristics of moduli spaces of curves is*

$$\begin{aligned}
(4) \quad \omega_2 = & \frac{(\hbar - 1)P_1}{2} \left[\left(-\frac{\hbar}{P_1} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \left(\log(\ell \hbar^\ell Z_\ell) + \psi_0(-Z_\ell) \right) \right)^2 \right. \\
& + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \left(\log(2\ell \hbar^{2\ell} Z_{2\ell}) + \psi_0(-Z_{2\ell}) \right) \\
& + \sum_{\ell \geq 1} \left(\frac{\mu(\ell)}{\ell} \right)^2 \psi_1(-Z_\ell) - \frac{\hbar^2}{P_1^2} \left. \right] \\
& - \hbar + (\hbar^2 - 1)P_1 + \frac{1}{2} (P_1^2 + P_2).
\end{aligned}$$

Let us discuss the convergence and structure of the formula. First note that

$$\frac{1}{Z_\ell} = \frac{\ell \hbar^\ell}{P_\ell(1 + X_\ell)} = \frac{\ell \hbar^\ell}{P_\ell} \sum_{j \geq 0} (-X_\ell)^j = O(\hbar^\ell)$$

with

$$(5) \quad X_\ell := \sum_{\substack{d \mid \ell \\ d \neq \ell}} \mu(\ell/d) \frac{\hbar^{\ell-d} P_d}{P_\ell} = O(\hbar^{\frac{\ell}{2}}).$$

We can hence expand the terms within the square brackets in (4) as power series in \hbar . First,

$$\begin{aligned}
(6) \quad \log(\ell \hbar^\ell Z_\ell) + \psi_0(-Z_\ell) &= \log P_\ell - \sum_{j \geq 1} \frac{(-1)^j}{j} X_\ell^j - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{B_k}{k} \binom{-k}{j} \left(\frac{\ell \hbar^\ell}{P_\ell} \right)^k X_\ell^j \\
(7) \quad \psi_1(-Z_\ell) &= - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} B_{k-1} \binom{-k}{j} \left(\frac{\ell \hbar^\ell}{P_\ell} \right)^k X_\ell^j.
\end{aligned}$$

Note in particular that, with the exception of the term $\log P_\ell$, the expressions (6) and (7) are (infinite) linear combinations of Laurent monomials of the form

$$(8) \quad \hbar^a \frac{P_{d_1}^{b_1} P_{d_2}^{b_2} \cdots P_{d_s}^{b_s}}{P_\ell^c} \quad \text{such that} \quad \begin{cases} d_i \mid \ell, & \text{for } 1 \leq i \leq s, \\ b_1 + \cdots + b_s \leq c, \\ a + b_1 d_1 + \cdots + b_s d_s = \ell c. \end{cases}$$

In particular, if we say that P_d has degree d , then the coefficient of \hbar^a for $a > 0$ in each of the expressions (6) and (7) is a finite linear combination of

Laurent monomials of degree $-a$ in the inhomogeneous power sum symmetric functions, each of the special form prescribed by (8). Inserting (6) and (7) into (4) we find

(9)

$$\begin{aligned}
 \omega_2 = & \frac{(\hbar - 1)P_1}{2} \left[\left(-\frac{\hbar}{P_1} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_\ell + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \right. \right. \\
 & \times \left(-\sum_{j \geq 1} \frac{(-1)^j}{j} X_\ell^j - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{B_k}{k} \binom{-k}{j} \left(\frac{\ell \hbar^\ell}{P_\ell} \right)^k X_\ell^j \right) \left. \right)^2 \\
 & + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{2\ell} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \\
 & \times \left(-\sum_{j \geq 1} \frac{(-1)^j}{j} X_{2\ell}^j - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{B_k}{k} \binom{-k}{j} \left(\frac{2\ell \hbar^{2\ell}}{P_{2\ell}} \right)^k X_{2\ell}^j \right) \\
 & \left. - \sum_{\ell \geq 1} \left(\frac{\mu(\ell)}{\ell} \right)^2 \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} B_{k-1} \binom{-k}{j} \left(\frac{\ell \hbar^\ell}{P_\ell} \right)^k X_\ell^j - \frac{\hbar^2}{P_1^2} \right] \\
 & - \hbar + (\hbar^2 - 1)P_1 + \frac{1}{2} \left(P_1^2 + P_2 \right).
 \end{aligned}$$

By expanding the square and collecting terms, we arrive at the following equivalent version of Theorem 1.1.

Corollary 1.2. *Let A_g and C_g be the coefficients of \hbar^g in the following power series, with X_ℓ as in (5):*

(10)

$$\begin{aligned}
 \sum_g A_g \hbar^g &:= -P_1 \left(-\frac{\hbar}{P_1} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \right. \\
 &\quad \times \left. \left(-\sum_{j \geq 1} \frac{(-1)^j}{j} X_\ell^j - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{B_k}{k} \binom{-k}{j} \left(\frac{\ell \hbar^\ell}{P_\ell} \right)^k X_\ell^j \right) \right); \\
 \sum_g C_g \hbar^g &:= -\frac{P_1}{2} \left[\left(-\frac{\hbar}{P_1} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \right. \right. \\
 &\quad \times \left. \left. \left(-\sum_{j \geq 1} \frac{(-1)^j}{j} X_\ell^j - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{B_k}{k} \binom{-k}{j} \left(\frac{\ell \hbar^\ell}{P_\ell} \right)^k X_\ell^j \right) \right)^2
 \end{aligned}$$

$$\begin{aligned}
& + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \left(- \sum_{j \geq 1} \frac{(-1)^j}{j} X_{2\ell}^j \right. \\
& - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{B_k}{k} \binom{-k}{j} \left(\frac{2\ell \hbar^{2\ell}}{P_{2\ell}} \right)^k X_{2\ell}^j \Big) \\
& \left. - \sum_{\ell \geq 1} \left(\frac{\mu(\ell)}{\ell} \right)^2 \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} B_{k-1} \binom{-k}{j} \left(\frac{\ell \hbar^{\ell}}{P_{\ell}} \right)^k X_{\ell}^j - \frac{\hbar^2}{P_1^2} \right].
\end{aligned}$$

Then A_g is a finite linear combination of monomials

$$\frac{P_{d_1}^{b_1} P_{d_2}^{b_2} \cdots P_{d_s}^{b_s}}{P_{\ell}^c} \quad \text{such that} \quad \begin{cases} d_i | \ell, \text{ for } 1 \leq i \leq s, \\ b_1 + \cdots + b_s \leq c + 1, \\ g + b_1 d_1 + \cdots + b_s d_s = \ell c + 1. \end{cases}$$

Similarly, C_g is a finite linear combinations of monomials

$$\frac{P_{d_1}^{b_1} P_{d_2}^{b_2} \cdots P_{d_s}^{b_s}}{P_{\ell_1}^{c_1} P_{\ell_2}^{c_2}} \quad \text{such that} \quad \begin{cases} d_i | \ell_1 \text{ or } d_i | \ell_2, \text{ for } 1 \leq i \leq s, \\ b_1 + \cdots + b_s \leq c_1 + c_2 + 1, \\ g + b_1 d_1 + \cdots + b_s d_s = \ell_1 c_1 + \ell_2 c_2 + 1. \end{cases}$$

Furthermore, the equivariant Euler characteristics $\chi_2^{\mathbb{S}}(\mathcal{M}_{g,\bullet}) := \sum_n \chi_2^{\mathbb{S}}(\mathcal{M}_{g,n})$ are expressed as follows:

$$\begin{aligned}
\chi_2^{\mathbb{S}}(\mathcal{M}_{0,\bullet}) &= -\frac{P_1}{2} \left[\left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{\ell} \right)^2 + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{2\ell} \right] - P_1 + \frac{P_1^2}{2} + \frac{P_2}{2} \\
\chi_2^{\mathbb{S}}(\mathcal{M}_{1,\bullet}) &= +\frac{P_1}{2} \left[\left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{\ell} \right)^2 + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{2\ell} \right] \\
&\quad + \left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{\ell} \right) A_1 + C_1 - 1 \\
\chi_2^{\mathbb{S}}(\mathcal{M}_{2,\bullet}) &= \left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{\ell} \right) (A_2 - A_1) + C_2 - C_1 + P_1 \\
\chi_2^{\mathbb{S}}(\mathcal{M}_{g,\bullet}) &= \left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{\ell} \right) (A_g - A_{g-1}) + C_g - C_{g-1} \quad \text{for } g \geq 3
\end{aligned}$$

We remark that $X_1 = 0$, and hence the terms involving positive powers of X_1 can be dropped from the sums above. Furthermore, the individual

terms appearing in the formulas for the Euler characteristics have natural interpretations in terms of graphs. See §5, for details. The first few Laurent polynomials A_g and C_g are:

$$\begin{aligned}
A_1 &= -\frac{P_1^2 - P_2}{2P_2}, \quad C_1 = \frac{P_1^2 + P_2}{2P_2}, \quad A_2 = \frac{1}{12P_1} - \frac{P_1^3}{4P_2^2} + \frac{P_1}{2P_2} - \frac{P_1^2}{3P_3}, \\
C_2 &= \frac{1}{8P_1} + \frac{P_1^3}{8P_2^2} - \frac{P_1P_2}{4P_4}, \quad A_3 = -\frac{P_1^4}{6P_2^3} + \frac{P_1^2}{2P_2^2} + \frac{P_1}{2P_3} + \frac{P_1P_3}{6P_6}, \\
C_3 &= \frac{1}{24P_1^2} + \frac{P_1^4}{24P_2^3} + \frac{P_1^2}{8P_2^2} - \frac{5}{24P_2} + \frac{P_1}{3P_3} - \frac{P_1^3}{6P_2P_3} - \frac{P_1P_3}{6P_6}, \\
A_4 &= -\frac{1}{120P_1^3} - \frac{P_1^5}{8P_2^4} + \frac{P_1^3}{2P_2^3} - \frac{P_1}{6P_2^2} - \frac{P_1^3}{6P_3^2} - \frac{P_1^2}{5P_5} + \frac{P_1P_2}{6P_6}, \\
C_4 &= -\frac{1}{288P_1^3} + \frac{P_1^5}{96P_2^4} + \frac{5P_1^3}{24P_2^3} - \frac{7P_1}{16P_2^2} - \frac{1}{24P_1P_2} - \frac{P_1^3}{18P_3^2} - \frac{2}{9P_3} \\
&\quad + \frac{P_1^4}{12P_2^2P_3} + \frac{5P_1^2}{12P_2P_3} - \frac{P_1P_2^2}{8P_4^2} + \frac{P_1}{2P_4} - \frac{P_1P_2}{6P_6} - \frac{P_3}{12P_6} + \frac{P_1^2P_3}{12P_2P_6}, \\
A_5 &= -\frac{P_1^6}{10P_2^5} + \frac{P_1^4}{2P_2^4} - \frac{P_1^2}{6P_6} + \frac{P_1^2}{2P_3^2} - \frac{P_1^2}{3P_2^3} + \frac{P_1}{2P_5} + \frac{P_5P_1}{10P_{10}}, \\
C_5 &= -\frac{P_1^6}{240P_2^5} - \frac{P_1^5}{18P_2^3P_3} - \frac{P_1^4}{12P_2P_3^2} + \frac{13P_1^4}{48P_2^4} + \frac{7P_1^3}{24P_2^2P_3} - \frac{P_1^3}{10P_2P_5} \\
&\quad + \frac{P_3P_1^3}{24P_2^2P_6} + \frac{11P_1^2}{36P_6} + \frac{5P_1^2}{12P_3^2} - \frac{53P_1^2}{72P_2^3} - \frac{P_1}{4P_2P_3} + \frac{P_1}{5P_5} \\
&\quad - \frac{P_3P_1}{12P_2P_6} - \frac{P_5P_1}{10P_{10}} - \frac{P_2}{12P_6} + \frac{1}{24P_2^2} - \frac{1}{24P_3P_1} - \frac{P_3}{72P_6P_1} \\
&\quad - \frac{1}{240P_2P_1^2} - \frac{1}{80P_1^4}.
\end{aligned}$$

1.2. Relations to the topological and weight zero Euler characteristics

For fixed genus g , by work of Gorsky, the generating function $\sum_n \chi^{\mathbb{S}}(\mathcal{M}_{g,n})$ is a finite sum of Laurent monomials of degree $2-2g$ in the inhomogeneous power sum symmetric functions P_d , each of whose denominators is a pure power P_{ℓ}^c ,

satisfying conditions analogous to (8) [10]. The coefficients involve certain orbifold Euler characteristics $\chi^{orb}(\mathcal{M}_{h,s})$, and these have natural expressions in terms of Bernoulli numbers, by the work of Harer and Zagier [11].

Gorsky's method of proof is beautiful and elementary: one stratifies \mathcal{M}_g according to the topological type of the action of the automorphism group and considers the action of automorphisms on the configuration spaces of n points. Each topological type of a curve with an automorphism contributes to precisely one monomial of the special form described above. The final formula is then deduced using the basic properties of compactly supported topological Euler characteristic, namely that it is additive for stratifications and multiplicative for fibrations.

Weight graded Euler characteristics are more subtle than topological Euler characteristics. In particular, although weight graded Euler characteristics of algebraic varieties and Deligne-Mumford stacks are additive with respect to stratifications, they are not multiplicative with respect to algebraic fibrations (flat families), even when the fibers are smooth and proper. Nevertheless, all of the essential ideas in Gorsky's computation can be adapted to compute the \mathbb{S}_n -equivariant weight zero Euler characteristic of $\mathcal{M}_{g,n}$, using the interpretation of weight zero cohomology as the singular cohomology of a moduli space of stable tropical curves [3]. As a result, the weight zero Euler characteristic has a structure closely analogous to that of Gorsky's formula: for fixed genus, the generating function is a finite sum of monomials of degree $1-g$ in the inhomogeneous power sum symmetric functions P_d , each of whose denominators is a pure power, satisfying conditions analogous to (8). Roughly speaking, each pair of a graph with an automorphism contributes to precisely one monomial of the prescribed special form; see [3, Proposition 3.2]. In the final formula, the coefficients are given by certain orbifold Euler characteristics of spaces of graphs, which have natural expressions in terms of Bernoulli numbers; see [3, Lemma 7.4], which is proved by induction starting from the $n=0$ case, due to Kontsevich [7, 13].

We do not have such a topological interpretation for weight two cohomology of moduli spaces of curves, and the weight two Euler characteristic is of a somewhat different form. In particular, for fixed genus $g \geq 2$, the generating function $\sum_n \chi_2^{\mathbb{S}}(\mathcal{M}_{g,n})$ is not a finite sum of Laurent monomials in the inhomogeneous power sum symmetric functions. Lacking a topological interpretation for the weight two Euler characteristic, we express it as the Euler characteristic of a complex of decorated graphs, and use basic properties of symmetric sequences and an operadic identity (Proposition 3.2) to produce the formula for the all genus generating function stated as Theorem 1.1. Specializing to a fixed genus, we obtain an expression involving logarithmic terms

as well as Laurent monomials in the inhomogeneous power sum symmetric functions, of two different degrees, with denominators that are products of either one or two pure powers (Corollary 1.2).

1.3. Outline of proof

To show Theorem 1.1 we proceed in three steps. First, we show a general formula, see Proposition 3.2, computing the Euler characteristic of graph complexes spanned by graphs with one special vertex, decorated by an arbitrary symmetric sequence. Next, in Sections 3.5–3.6, we recall a recent result of the authors, identifying the cohomology $\text{gr}_2 H_c^\bullet(\mathcal{M}_{g,n})$ with the cohomology of a certain graph complex $X_{g,n}$, spanned by graphs with one suitably decorated special vertex. Finally, in Section 4 we apply Proposition 3.2 to (a close variant of) the graph complex $X_{g,n}$. After some algebraic rewriting and simplifications one then arrives at the formula of Theorem 1.1.

1.4. Numerical computations

The generating function in Theorem 1.1 can be implemented to compute the equivariant Euler characteristic of $\text{gr}_2 H_c^\bullet(\mathcal{M}_{g,n})$ for small values of g and n , as shown in Figure 1.

g,n	0	1	2	3	4	5
0 0	0	0	0	0	s_4	$-s_{32}$
1 0	0	s_1	0	0	$-s_{211}$	$s_{311} + s_{32} + s_{41}$
2 0	0	0	0	$-s_{21} - s_3$	s_4	$-s_{2111} - s_{221} - s_{311} + 2s_{31} + 2s_5$
3 0	s_1	$s_{11} + s_2$	0	$-2s_3$	$2s_{22} - s_{31} - 2s_4$	$-3s_{2111} - s_{221} - 6s_{311} + 2s_{32} + 2s_5$
4 0	0	0	0	$2s_{111}$	$3s_{1111} + s_{211} - 3s_{22} - 4s_{31} - 3s_4$	$5s_{11111} + 2s_{21111} + 2s_{221} - 6s_{311} + s_{32} - s_{41} + 3s_5$
5 0	s_1	$s_{11} + 3s_2$	0	$5s_{21} + 2s_3$	$-s_{1111} + 5s_{211} - 4s_{31} - 9s_4$	$2s_{11111} + 10s_{2111} + 11s_{221} - 14s_{31} - 8s_5$
6 0	0	$-3s_1$	$-4s_{11} - s_2$	$-2s_{111} + 8s_3$	$4s_{1111} + 8s_{211} - 6s_{31} + 10s_{31} + 3s_4$	$6s_{11111} + 21s_{2111} - 4s_{221} + 15s_{311} - 26s_{32} - 22s_{41} - 19s_5$
7 0	s_1	$2s_{11}$	0	$-7s_{1111} + 4s_{21} + 3s_3$	$-15s_{1111} + 7s_{211} + 15s_{221} + 24s_{31} + 42s_4$	$-25s_{11111} + s_{2111} - 7s_{221} + 35s_{311} - 22s_{32} - 25s_{41} - 37s_5$
8 -1	$-4s_1$	$-8s_{11} - 9s_2$	$-6s_{111} - 18s_{21} + 4s_3$	$-8s_{1111} - 28s_{211} - 10s_{221} + 24s_{31} + 42s_4$	$-19s_{11111} - 19s_{221} + 91s_{311} + 22s_{32} + 87s_{41} + 20s_5$	
9 4	$9s_1$	$8s_{11} + s_2$	$-2s_{111} + s_{21} - 17s_3$	$-44s_{1111} - 64s_{211} + 17s_{221} - 17s_{31} + 5s_4$	$-71s_{11111} - 80s_{2111} + 55s_{221} + 110s_{311} + 193s_{32} + 193s_{41} + 69s_5$	
10 -4	$-9s_1$	$-6s_{11} - 10s_2$	$12s_{111} - 46s_{21} - 29s_3$	$38s_{1111} - 68s_{211} - 35s_{221} - 79s_{31} + 28s_4$	$3s_{11111} - 227s_{21111} - 155s_{221} - 234s_{311} + 50s_{32} + 196s_{41} + 209s_5$	

Figure 1: The equivariant Euler characteristic of $\text{gr}_2 H_c^\bullet(\mathcal{M}_{g,n})$ for $g \leq 10$ and $n \leq 5$, computed using Mathematica and Sage from Theorem 1.1 and expressed in terms of Schur functions.

Specializing to $n = 0$, and writing out the terms for $g \leq 50$, we find that $\sum_g \chi_2(\mathcal{M}_g) \hbar^g$ is equal to:

$$\begin{aligned}
& -\hbar^8 + 4\hbar^9 - 4\hbar^{10} + 4\hbar^{11} - 7\hbar^{12} + 8\hbar^{13} - 16\hbar^{14} + 10\hbar^{15} - 10\hbar^{16} \\
& + 28\hbar^{17} + 17\hbar^{18} + 94\hbar^{19} - 12\hbar^{20} - 382\hbar^{21} - 196\hbar^{22} + 2181\hbar^{23} \\
& - 4304\hbar^{24} - 43135\hbar^{25} + 50281\hbar^{26} + 737650\hbar^{27} - 676300\hbar^{28} \\
& - 13670125\hbar^{29} + 13593898\hbar^{30} + 301176776\hbar^{31} - 303061590\hbar^{32}
\end{aligned}$$

$$\begin{aligned}
& - 7579395579\hbar^{33} + 7577048267\hbar^{34} + 215276816193\hbar^{35} \\
& - 215186827379\hbar^{36} - 6867916230403\hbar^{37} + 6867642573519\hbar^{38} \\
& + 244534892486924\hbar^{39} - 244525053194639\hbar^{40} \\
& - 9660791088309245\hbar^{41} + 9660393501146309\hbar^{42} + 421316847700113027\hbar^{43} \\
& - 421303860389448419\hbar^{44} - 20188618782635720913\hbar^{45} \\
& + 20188137225679260098\hbar^{46} + 1058435119750859223989\hbar^{47} \\
& - 1058415071189257113479\hbar^{48} - 60478235401984833358102\hbar^{49} \\
& + 60477318015911247931156\hbar^{50} + O(\hbar^{51}) .
\end{aligned}$$

For $n = 0$ and $g \leq 200$, we display $|\chi_2(\mathcal{M}_g)|$ on a logarithmic scale, along with a plot of the sign of $\chi_2(\mathcal{M}_g)$, which seems to be eventually periodic. See Figure 2.




Figure 2: Plots of $\log(1 + |\chi_2(\mathcal{M}_g)|)$ and the sign of $\chi_2(\mathcal{M}_g)$ for $g \leq 200$, on the left and right, respectively.

Conjecture 1.3. *For $g \geq 23$, the sign of $\chi_2(\mathcal{M}_g)$ is -1 for $g \equiv 0, 1 \pmod{4}$ and $+1$ for $g \equiv 2, 3 \pmod{4}$.*

2. Preliminaries

2.1. Notation and conventions

We work over the rational numbers \mathbb{Q} . All vector spaces are understood to be \mathbb{Q} -vector spaces, and likewise all homology and cohomology groups are taken with \mathbb{Q} -coefficients. If V is a vector space with the action of a finite group G , we identify the invariant subspace V^G with the coinvariant space V_G , by averaging over the group action. The phrase differential graded is abbreviated

dg. For a dg vector space V and an integer k , let $V[k]$ be the graded vector space obtained by shifting all degrees by k , i.e., the degree j part of V is the degree $j - k$ part of $V[k]$.

2.2. Symmetric sequences and products

A symmetric sequence is a collection $\mathcal{C} = \{\mathcal{C}(r)\}_{r \geq 0}$ of \mathbb{S}_r -modules in dg vector spaces. Similarly, a bisymmetric sequence is a collection $\mathcal{B} = \{\mathcal{B}(r, s)\}_{r, s \geq 0}$ of $\mathbb{S}_r \times \mathbb{S}_s$ -modules. We say that $\mathcal{C}(r)$ is the part of \mathcal{C} of *arity* r , and $\mathcal{B}(r, s)$ is the part of \mathcal{B} of *arity* (r, s) .

The category of symmetric sequences \mathbf{Seq} has several monoidal products. First, for two symmetric sequences \mathcal{C}, \mathcal{D} we have the symmetric sequence $\mathcal{C} \otimes \mathcal{D}$, given by

$$(\mathcal{C} \otimes \mathcal{D})(r) = \mathcal{C}(r) \otimes \mathcal{D}(r),$$

with the diagonal \mathbb{S}_r -action. Second, we have the symmetric sequence $\mathcal{C} \boxtimes \mathcal{D}$ given by

$$(\mathcal{C} \boxtimes \mathcal{D})(r) = \bigoplus_{j+k=r} \text{Ind}_{\mathbb{S}_j \times \mathbb{S}_k}^{\mathbb{S}_r} \mathcal{C}(j) \otimes \mathcal{D}(k).$$

Let us note in particular that the symmetric sequence

$$\mathcal{C}^{\boxtimes k} = \underbrace{\mathcal{C} \boxtimes \cdots \boxtimes \mathcal{C}}_{k \times}$$

carries an action of \mathbb{S}_k by permuting the factors. Thus, $\{(\mathcal{C}^{\boxtimes k})(r)\}_{k \geq 0, r \geq 0}$ is a bisymmetric sequence.

We then have a third monoidal product on symmetric sequences, the plethysm product $\mathcal{D} \circ \mathcal{C}$, given by

$$(\mathcal{D} \circ \mathcal{C})(r) = \bigoplus_{k \geq 0} \mathcal{D}(k) \otimes_{\mathbb{S}_k} \mathcal{C}^{\boxtimes k}(r).$$

The category of symmetric sequences \mathbf{Seq} is monoidal with respect to the plethysm product, with unit $\mathbf{1}$ given by

$$\mathbf{1}(r) = \begin{cases} \mathbb{Q} & \text{if } r = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Finally, given \mathcal{C} and \mathcal{D} symmetric sequences of dg vector spaces we define the dg vector space

$$\mathcal{C} \otimes_{\mathbb{S}} \mathcal{D} := \bigoplus_{r \geq 0} \mathcal{C}(r) \otimes_{\mathbb{S}_r} \mathcal{D}(r).$$

2.3. Equivariant Euler characteristics

For F an endomorphism of a finite dimensional graded vector space $V = \bigoplus_i V_i$, we denote by $\text{STr}(F)$ the super-trace of F , that is, the alternating sum

$$\text{STr}(F) = \sum_i (-1)^i \text{Tr}(F|_{V_i}).$$

For $\sigma \in \mathbb{S}_r$ a permutation we denote by $i(\sigma) = (i_1, i_2, \dots)$ its cycle type, with i_k the number of cycles of length k . For ρ a graded \mathbb{S}_r -representation on V , we denote the \mathbb{S}_r -equivariant Euler characteristic in $\Lambda = \mathbb{Q}[[p_1, p_2, \dots]]$ by

$$\chi^{\mathbb{S}}(V) = \frac{1}{r!} \sum_{\sigma \in \mathbb{S}_r} \text{STr}(\rho(\sigma)) p^{i(\sigma)},$$

using the multi-index notation $p^{i(\sigma)} = p_1^{i_1(\sigma)} p_2^{i_2(\sigma)} \dots$. This equivariant Euler characteristic is actually a polynomial in p_1, p_2, \dots ; we define it as a power series in view of the case of symmetric sequences that we consider below. If V has an additional (non-negative) grading with grading generator G (i.e., where G is equal to k times the identity on the subspace of degree k) we use the notation

$$(11) \quad \chi^u(V) = \frac{1}{r!} \sum_{\sigma \in \mathbb{S}_r} \text{STr}(u^G \rho(\sigma)) p^{i(\sigma)} \in \Lambda[[u]].$$

We freely generalize this notation to multiple gradings and other variables counting the multidegree. For example, if V has an additional \mathbb{Z}^2 -grading, with grading generators G_1, G_2 , then we write

$$\chi^{u,v}(V) = \frac{1}{r!} \sum_{\sigma \in \mathbb{S}_r} \text{STr}(u^{G_1} v^{G_2} \rho(\sigma)) p^{i(\sigma)} \in \Lambda[[u, v]],$$

and similarly for a \mathbb{Z}^3 -grading.

For \mathcal{A} a symmetric sequence such that $\mathcal{A}(r)$ is finite dimensional for each r , we write

$$\chi^{\mathbb{S}}(\mathcal{A}) := \sum_{r \geq 0} \chi^{\mathbb{S}}(\mathcal{A}(r))$$

and similarly for $\chi^u(\mathcal{A})$, when \mathcal{A} has an additional grading.

The operations on symmetric sequences discussed in §2.2 induce operations on symmetric functions. First,

$$\chi^{\mathbb{S}}(\mathcal{A} \boxtimes \mathcal{B}) = \chi^{\mathbb{S}}(\mathcal{A}) \chi^{\mathbb{S}}(\mathcal{B}).$$

Second, there is a plethysm product \circ on symmetric functions $\circ : \Lambda \otimes \Lambda \rightarrow \Lambda$ such that

$$\chi^{\mathbb{S}}(\mathcal{A} \circ \mathcal{B}) = \chi^{\mathbb{S}}(\mathcal{A}) \circ \chi^{\mathbb{S}}(\mathcal{B}).$$

Concretely, the plethysm product on symmetric functions may be computed by the following rules:

$$(12) \quad \begin{aligned} (f_1 + f_2) \circ g &= f_1 \circ g + f_2 \circ g & (f_1 f_2) \circ g &= (f_1 \circ g)(f_2 \circ g) \\ p_n \circ f &= f(p_n, p_{2n}, p_{3n}, \dots) \quad \text{for } f = f(p_1, p_2, p_3, \dots) \end{aligned}$$

In the presence of an additional grading, encoded by a variable u as in (11), the last rule is replaced by

$$(13) \quad p_n \circ f = f(u^n, p_n, p_{2n}, p_{3n}, \dots) \quad \text{for } f = f(u, p_1, p_2, p_3, \dots)$$

For more details we refer to [16] or the concise introduction to symmetric functions in [9].

On the subspace of Λ consisting of symmetric polynomials (finite series), there is an inner product given by

$$(14) \quad \begin{aligned} \langle p^i, p^j \rangle &= \delta_{ij} \prod_k k^{i_k} i_k! = \prod_k \left(\left(\frac{\partial}{\partial p_k} \right)^{i_k} k^{j_k} p_k^{j_k} \right)_{p_k=0} \\ &= \left(\prod_k \left(k \frac{\partial}{\partial p_k} \right)^{i_k} p^j \right)_{p_1=p_2=\dots=0}. \end{aligned}$$

Here, $i = (i_1, \dots)$ and $j = (j_1, \dots)$ are multi-indices, $p^i := p_1^{i_1} p_2^{i_2} \dots$, and δ_{ij} is the Kronecker symbol. This inner product has the property that for \mathcal{A}, \mathcal{B} finite dimensional symmetric sequences

$$(15) \quad \chi(\mathcal{A} \otimes_{\mathbb{S}} \mathcal{B}) = \langle \chi^{\mathbb{S}}(\mathcal{A}), \chi^{\mathbb{S}}(\mathcal{B}) \rangle.$$

2.4. Examples of symmetric sequences and their equivariant Euler characteristics

For later use we shall recall the Euler characteristics of some symmetric sequences from the literature.

The commutative operad is a simple and natural symmetric sequence, which we consider in two variations:

$$\text{Com}_1(r) = \mathbb{Q} \quad \text{for } r \geq 0, \quad \text{Com}(r) = \begin{cases} \mathbb{Q} & \text{for } r \geq 1, \\ 0 & \text{for } r = 0. \end{cases}$$

Here, \mathbb{Q} denotes the trivial \mathbb{S}_r -representation. The corresponding equivariant Euler characteristics are [8, §5.1]:

$$(16) \quad \chi^{\mathbb{S}}(\mathbf{Com}_1) = \exp \left(\sum_{\ell} \frac{p_{\ell}}{\ell} \right) \quad \chi^{\mathbb{S}}(\mathbf{Com}) = \exp \left(\sum_{\ell} \frac{p_{\ell}}{\ell} \right) - 1.$$

The Lie operad is the symmetric sequence defined such that $\mathbf{Lie}(r)$ is the subspace of the free Lie algebra $\text{FreeLie}(x_1, \dots, x_r)$ in r letters consisting of linear combinations of Lie words in which each x_j occurs exactly once. We shall also need a degree shifted version \mathbf{Lie}_0 defined in the same way, except that the generators x_j are considered of cohomological degree one, and one has an overall degree shift by -1 , i.e.,

$$\mathbf{Lie}_0(r) = \mathbf{Lie}(r) \otimes \text{sgn}_r[1 - r].$$

The corresponding equivariant Euler characteristics are

$$(17) \quad \chi^{\mathbb{S}}(\mathbf{Lie}) = - \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log(1 - p_{\ell}), \quad \chi^{\mathbb{S}}(\mathbf{Lie}_0) = \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log(1 + p_{\ell}).$$

See [8, Proposition 5.3 and Lemma 5.4].

The Poisson operad is the symmetric sequence $\mathbf{Pois} = \mathbf{Com} \circ \mathbf{Lie}$. The elements of $\mathbf{Pois}(r)$ can be understood as linear combinations of Poisson words in letters x_1, \dots, x_r such that each letter occurs exactly once. We shall also use a degree shifted version

$$\mathbf{Pois}_0 = \mathbf{Com} \circ \mathbf{Lie}_0.$$

Both \mathbf{Pois} and \mathbf{Pois}_0 carry a *complexity grading* by the number of Lie brackets. More precisely, we define $\mathbf{Lie}(r)$ and $\mathbf{Lie}_0(r)$ to be concentrated in complexity $r - 1$, and this descends to give the complexity grading on \mathbf{Pois} and \mathbf{Pois}_0 . By [8, Proposition 5.5] and [2, Proposition 6.3], the graded equivariant Euler characteristics, with the variable u counting the complexity as in (11), are:

$$(18) \quad \chi^u(\mathbf{Pois}) = \prod_{\ell \geq 1} (1 - u^{\ell} p_{\ell})^{\frac{1}{\ell} \sum_{d \mid \ell} \mu(\ell/d) \frac{1}{u^d}} - 1,$$

$$(19) \quad \chi^u(\mathbf{Pois}_0) = \prod_{\ell \geq 1} (1 + u^{\ell} p_{\ell})^{\frac{1}{\ell} \sum_{d \mid \ell} \mu(\ell/d) \frac{1}{u^d}} - 1.$$

Also note that the complexity grading on \mathbf{Pois}_0 is, by definition, equal to the cohomological grading.

Next consider the symmetric sequence concentrated in arity one

$$HS(r) = \begin{cases} H(S^1) = \mathbb{Q} \oplus \mathbb{Q}[-1] & \text{for } r = 1 \\ 0 & \text{for } r \neq 1 \end{cases}.$$

Then one defines the symmetric sequence

$$\mathbf{BV}_0 := \mathbf{Pois}_0 \circ HS.$$

This is a version of the Batalin-Vilkovisky operad (see [15, §13.7]), hence the notation. We declare the complexity grading on \mathbf{BV}_0 to be the same as the cohomological grading.

It is clear that $\chi^u(HS) = p_1 - up_1$. From (12), (13), and (19) one then finds

$$(20) \quad \chi^u(\mathbf{BV}_0) = \prod_{\ell \geq 1} (1 + u^\ell (1 - u^\ell) p_\ell)^{\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) \frac{1}{u^d}} - 1.$$

Elements of $\mathbf{BV}_0(r)$ are identified with linear combinations of Poisson words in variables $x_1, \dots, x_r, Dx_1, \dots, Dx_r$, with every index occurring exactly once, for example

$$x_1 \wedge Dx_2 \wedge [x_3, Dx_4] \in \mathbf{BV}_0(4).$$

For technical reasons we also define the reduced sub-symmetric sequence $\mathbf{BV}_0^{\text{red}} \subset \mathbf{BV}_0$ to be spanned by those Poisson words that do not have a factor x_j for any j . In other words,

$$\mathbf{BV}_0 = (\mathbf{BV}_0^{\text{red}} \boxtimes \mathbf{Com}_1) \oplus \mathbf{Com}.$$

Hence

$$(21) \quad \begin{aligned} \chi^u(\mathbf{BV}_0^{\text{red}}) &= (\chi^u(\mathbf{BV}_0) + 1) \chi(\mathbf{Com}_1)^{-1} - 1 \\ &= \prod_{\ell \geq 1} e^{-\frac{1}{\ell} p_\ell} (1 + u^\ell (1 - u^\ell) p_\ell)^{\frac{1}{\ell} \sum_{d|\ell} \mu(d/\ell) \frac{1}{u^d}} - 1. \end{aligned}$$

Finally, note that the definition of the \mathbb{S} -equivariant Euler characteristic above has a natural generalization for bisymmetric sequences. We shall only need the following example. Define the bisymmetric sequence Δ_0 such that

$$(22) \quad \Delta_0(r, s) = \begin{cases} \mathbb{Q}[\mathbb{S}_r] & \text{for } r = s \\ 0 & \text{otherwise} \end{cases}.$$

Here the group ring $\mathbb{Q}[\mathbb{S}_r]$ is considered as an $\mathbb{S}_r \times \mathbb{S}_r$ -module by left and right multiplication. We then have

$$\begin{aligned} \chi^{\mathbb{S}}(\Delta_0) := \sum_{r,s} \frac{1}{r!s!} \sum_{\substack{\sigma \in \mathbb{S}_r \\ \nu \in \mathbb{S}_s}} \text{STr}_{\Delta_0}(\rho(\sigma, \nu)) p^{i(\sigma)} q^{i(\nu)} &= \exp \left(\sum_{n \geq 1} \frac{1}{n} p_n q_n \right) \\ &\in \Lambda_{p,q} := \mathbb{Q}[[p_1, q_1, p_2, q_2, \dots]]. \end{aligned}$$

3. Graph complexes associated to moduli spaces of curves

3.1. The graph complexes $G^{(g,n)}$ and weight zero compactly supported cohomology of $\mathcal{M}_{g,n}$

Let g and n be non-negative integers. The graph complex $G^{(g,n)}$ studied in [5, 19] is generated by connected graphs of genus g with no loop edges and no vertices of valence 2. Here, g is the first Betti number of the geometric realization of the graph. The vertices of valence 1 are called *external*, and the other vertices are *internal*. Each generator has precisely n external vertices, labeled in bijection with $\{1, \dots, n\}$, and the symmetric group \mathbb{S}_n acts by permuting these labels. The edges not connecting to univalent vertices are called *structural*, and the cohomological degree of a generator is the number of structural edges. Each generator comes with a total ordering of its structural edges, and we impose the relation that reordering is multiplication by the sign of the induced permutation. The differential δ_{split} on $G^{(g,n)}$ is defined by splitting internal vertices.

$$(23) \quad \delta_{\text{split}} \Gamma = \sum_{v \text{ vertex}} \Gamma \text{ split } v \quad \begin{array}{c} \text{Diagram: } \text{A vertex } v \text{ with } k \text{ edges} \\ \text{Mapping: } v \mapsto \sum \text{Diagram: } \text{A vertex } v \text{ with } k-1 \text{ edges, plus } k \text{ separate edges} \end{array}$$

Theorem 3.1 ([5]). *For each g, n such that $2g + n \geq 3$ there is an \mathbb{S}_n -equivariant isomorphism*

$$H(G^{(g,n)}) \cong W_0 H_c(\mathcal{M}_{g,n})$$

with the weight zero part of the compactly supported cohomology of the moduli space of curves.

The differential does not affect the Euler characteristic. In this paper we will work with $G^{(g,n)}$ and its variants primarily as graded representations of symmetric groups.

3.2. Two variants of $G^{(g,n)}$

Distinguishing structural edges from nonstructural edges in the definition of $G^{(g,n)}$ will be inconvenient for the calculations that follow. For this reason, we define the variant $\check{G}^{(g,n)}$ exactly as above, except that the degree of a graph is the number of edges, and the orientation is given by a total ordering of all of the edges, not just the structural ones. For $(g,n) \neq (0,2)$ one can readily see that

$$\check{G}^{(g,n)} \cong G^{(g,n)} \otimes \text{sgn}_n[-n].$$

We also consider an enlargement of $\check{G}^{(g,n)}$ obtained by relaxing the connectedness assumption. Generators of the resulting graph complex $\check{fG}^{(g,n)}$ are possibly disconnected graphs of genus g with no loop edges and no vertices of valence 2. The external vertices are labeled in bijection with $\{1, \dots, n\}$, and the orientation is given by a total ordering of all edges. The genus g of a possibly disconnected graph with e edges and v vertices is

$$g := e - v + 1.$$

Note that some disconnected graphs have negative genus.

From $\check{fG}^{(g,n)}$, we build the symmetric sequence \check{fG} defined by

$$\check{fG}(r) := \bigoplus_{g \in \mathbb{Z}} \check{fG}^{(g,r)}.$$

On this symmetric sequence we consider two different gradings: the *genus grading*, in which $\check{fG}^{(g,r)}$ is of genus g , and the *complexity grading* in which $\check{fG}^{(g,r)}$ is of complexity $g + r - 1$. This is the genus of the connected graph obtained by fusing all external vertices into one. Note that the complexity grading is non-negative, and the number of isomorphism classes of graphs of a given complexity is finite.


3.3. Graph complexes with one special vertex

Let \mathcal{A} be a symmetric sequence. Consider

$$\mathcal{A} \otimes_{\mathbb{S}} \check{fG}.$$

This is a graded vector space whose elements are naturally interpreted as linear combinations of graphs as above, but with one special vertex that

carries a decoration in \mathcal{A} .



More precisely, the decoration of the special vertex is in $\mathcal{A}(r)$, with r the valence of the special vertex. The complexity grading on \widetilde{fG} induces a complexity grading on $\mathcal{A} \otimes_S \widetilde{fG}$, in which the complexity of a graph with one vertex decorated by \mathcal{A} is equal to its genus. The following result gives a general formula for the complexity-graded Euler characteristic of such \mathcal{A} -decorated graph complexes.

Proposition 3.2. *Let \mathcal{A} be any symmetric sequence in finite dimensional graded vector spaces. Then we have*


$$\chi^u(\mathcal{A} \otimes_S \widetilde{fG}) = \chi^u((\mathcal{A} \circ \text{Lie}_0) \otimes_S \text{BV}_0^{\text{red}}) = \langle \chi^S(\mathcal{A}) \circ \chi^S(\text{Lie}_0), \chi^u(\text{BV}_0^{\text{red}}) \rangle,$$

with u counting the complexity, and $\chi^u(-)$ being the graded Euler characteristic.

Note that the graded Euler characteristic is well-defined since the number of graphs of any given complexity is finite. Note also that the complexity grading is derived solely from the factor \widetilde{fG} , or respectively BV_0^{red} . The other factors Lie_0 or \mathcal{A} do not contribute to the complexity, i.e., they are considered to be of complexity zero. Finally, recall that $\chi^S(\text{Lie}_0)$ and $\chi^u(\text{BV}_0^{\text{red}})$ are given in (17) and (21), respectively.

3.4. Proof of Proposition 3.2

In the proof, we will use the Kontsevich graphical operad \mathbf{Graphs}_0 . We briefly sketch its construction and main properties. For details we refer to Kontsevich's original paper [12], or recollections elsewhere, for example [6, §2.9f]. Elements of $\mathbf{Graphs}_0(r)$ are series of graphs with r labeled external vertices and an arbitrary number of internal vertices such as in the following picture.



The symmetric group \mathbb{S}_r acts by permuting the labels $1, \dots, r$ of external vertices. The graph is oriented by a total ordering of its edges. The sign conventions are thus identical to those for $\check{G}^{(g,n)}$ above. The degree is the number of edges. One allows loops at both internal and external vertices. Note that the symmetric sequence \mathbf{Graphs}_0 carries a dg operad structure in which the differential is given by vertex splitting; however, the differential does not matter for the purposes of computing Euler characteristics.

We also consider the variant $\mathbf{BVGraphs}_0$, obtained as a quotient of \mathbf{Graphs}_0 by setting to zero graphs with loop edges at internal vertices. (Loops are still allowed at external vertices.) Both \mathbf{Graphs}_0 and $\mathbf{BVGraphs}_0$ have a complexity grading: the complexity of a graph with v internal vertices and e edges is $e - v$.

Furthermore, we consider the subspaces

$$\mathbf{Graphs}_0^{\text{red}} \subset \mathbf{Graphs}_0 \quad \mathbf{BVGraphs}_0^{\text{red}} \subset \mathbf{BVGraphs}_0$$

spanned by graphs all of whose external vertices have valence ≥ 1 .

Theorem 3.3 (Kontsevich [12], Lambrechts-Volic [14]¹). *There are operad quasi-isomorphisms $\mathbf{Pois}_0 \rightarrow \mathbf{Graphs}_0$ and $\mathbf{BV}_0 \rightarrow \mathbf{BVGraphs}_0$ that preserve the complexity gradings and restrict to quasi-isomorphisms on the reduced parts $(-)^{\text{red}}$. In particular,*

$$\chi^u(\mathbf{BVGraphs}_0^{\text{red}}) = \chi^u(\mathbf{BV}_0^{\text{red}}).$$

The connection between $\mathbf{BVGraphs}_0^{\text{red}}$ and our $\check{f}G$ is as follows.

Lemma 3.4. *Let \mathcal{B} be any symmetric sequence and $C \in \mathbb{Z}$. Then*

$$(24) \quad \mathcal{B} \otimes_{\mathbb{S}} \text{gr}^C \mathbf{BVGraphs}_0^{\text{red}} \cong (\mathcal{B} \circ \text{Com}) \otimes_{\mathbb{S}} \text{gr}^C \check{f}G,$$


with $\text{gr}^C(-)$ referring to the part of complexity C . In particular, if \mathcal{B} is finite-dimensional in each arity then

$$\chi^u(\mathcal{B} \otimes_{\mathbb{S}} \mathbf{BVGraphs}_0^{\text{red}}) = \chi^u((\mathcal{B} \circ \text{Com}) \otimes_{\mathbb{S}} \check{f}G).$$


Proof. As in the previous section the right-hand side of (24) may be interpreted as the space of linear combinations of graphs of genus (i.e., loop number) C , with one special vertex decorated by $\mathcal{B} \circ \text{Com}$. Equivalently such

¹Only the result for \mathbf{Graphs}_0 is stated in the given references, but the version for $\mathbf{BVGraphs}_0$ is an easy consequence.

graphs may be interpreted as 3-level graphs with one vertex decorated by \mathcal{B} , connected to another layer of vertices contributed by the factor Com .



But the left-hand side of (24) also has the interpretation as linear combinations of three-level graphs of genus C . In this case the middle layer of vertices is contributed by the external vertices of graphs in $\text{BVGraphs}_0^{\text{red}}$.



Hence we conclude that both sides of (24) are identical. \square

As a final ingredient we will use the Koszul property of the commutative operad [15, Theorem 7.6.4(i) and Proposition 13.1.7]. This means that the symmetric sequence (the Koszul complex) $\text{Lie}_0^* \circ \text{Com}$ carries a differential, such that the resulting cohomology is concentrated in arity 1. Here Lie_0^* is the linear dual of Lie_0 . In particular, this implies that on the level of Euler characteristics we have that

$$(25) \quad \begin{aligned} p_1 &= \chi^{\mathbb{S}}(\text{Lie}_0^* \circ \text{Com}) = \chi^{\mathbb{S}}(\text{Lie}_0^*) \circ \chi^{\mathbb{S}}(\text{Com}) \\ &= \chi^{\mathbb{S}}(\text{Lie}_0) \circ \chi^{\mathbb{S}}(\text{Com}) = \chi^{\mathbb{S}}(\text{Lie}_0 \circ \text{Com}). \end{aligned}$$

Here we also used that the equivariant Euler characteristic of a symmetric sequence and its dual are identical, due to the fact that all representations of the symmetric group are self-dual. We also note that one can alternatively derive (25) directly from (16) and (17), using the plethysm rules (12).

Proof of Proposition 3.2. The plethysm product with p_1 , $f \mapsto f \circ p_1$, is the identity. Hence, from (25) it follows that on the level of Euler characteristics

$$\chi^u(\mathcal{A} \otimes_{\mathbb{S}} \widetilde{f}G) = \chi^u((\mathcal{A} \circ \text{Lie}_0 \circ \text{Com}) \otimes_{\mathbb{S}} \widetilde{f}G).$$

Using Lemma 3.4 this is the same as

$$\chi^u((\mathcal{A} \circ \text{Lie}_0) \otimes_{\mathbb{S}} \text{BVGraphs}_0^{\text{red}}).$$

Using Theorem 3.3 this in turn is identified with

$$\chi^u((\mathcal{A} \circ \text{Lie}_0) \otimes_{\mathbb{S}} \text{BV}_0^{\text{red}}),$$

Applying (the graded version of) (15) we can evaluate this further to

$$\begin{aligned} \chi^u((\mathcal{A} \circ \text{Lie}_0) \otimes_{\mathbb{S}} \text{BV}_0^{\text{red}}) &= \langle \chi^{\mathbb{S}}(\mathcal{A} \circ \text{Lie}_0), \\ \chi^u(\text{BV}_0^{\text{red}}) \rangle &= \langle \chi^{\mathbb{S}}(\mathcal{A}) \circ \chi^{\mathbb{S}}(\text{Lie}_0), \chi^u(\text{BV}_0^{\text{red}}) \rangle, \end{aligned}$$

and hence Proposition 3.2 follows. \square

3.5. The graph complexes $X_{g,n}$ and weight two compactly supported cohomology of $\mathcal{M}_{g,n}$

The main result of [19] is the identification of $\text{gr}_2 H_c^\bullet(\mathcal{M}_{g,n})$ with the cohomology of a graph complex $X_{g,n}$ that is close to graph complexes arising in knot theory and the embedding calculus. We shall use this identification to show our main Theorem 1.1, by computing the Euler characteristic of $X_{g,n}$.

Recall from [19] that the generators for $X_{g,n}$ are simple graphs without loops or multiple edges, in which no vertices have valence 2. The vertices of valence at least 3 are *internal* and those of valence 1 are *external*. Each external vertex is decorated with an element from the set $\{\epsilon, \omega, 1, \dots, n\}$, such that:

- Each label $1, \dots, n$ appears exactly once and the label ω appears exactly twice;
- The graph obtained by joining all external vertices labeled ϵ or ω is connected and has genus g ;

Say that an edge with two external vertices labeled a and b is an (a, b) -edge. We further require that

- No connected component is an (ϵ, ω) or (ω, ω) edge;

An edge is *structural* if it does not contain an external vertex with label from $\{1, \dots, n\}$. The degree of a graph is the number of structural edges plus one. Each generator comes with a total ordering of the structural edges, and we impose the relation that permuting the structural edges is multiplication by the sign of the permutation.

The differential δ on $X_{g,n}$ is a sum of two parts $\delta = \delta_{\text{split}} + \delta_{\text{join}}$, defined by splitting internal vertices

$$(26) \quad \delta_{\text{split}} \Gamma = \sum_{v \text{ vertex}} \Gamma \text{ split } v \quad \text{Diagram: } \text{A vertex } v \text{ with } k \text{ edges} \mapsto \sum \text{Diagram showing } v \text{ split into } k \text{ edges.}$$

and joining external vertices

$$(27) \quad \delta_{\text{join}} \text{Diagram: } \Gamma \text{ with } k \text{ legs} = \sum_{\substack{S \subset \{\epsilon\text{- and } \omega\text{- legs}\} \\ |S| \geq 2}} \text{Diagram: } \Gamma \text{ with } k \text{ legs, one leg } \epsilon \text{ or } \omega \text{ is joined.}$$

We refer to [19] for details.

Theorem 3.5 (Theorem 1.1 of [19]). *There is an \mathbb{S}_n -equivariant isomorphism*

$$H(X_{g,n}) \cong \text{gr}_2 H_c(\mathcal{M}_{g,n})$$

for each $(g, n) \neq (1, 1)$ with $2g + n \geq 3$.

To show Theorem 1.1 we may hence compute the equivariant Euler characteristic of the graph complexes $X_{g,n}$, that is, we may use the following corollary.

Corollary 3.6.

$$\omega_2 = \hbar p_1 + \sum_{\substack{g, n \\ 2g + n \geq 3}} \hbar^n \chi^{\mathbb{S}}(X_{g,n})$$

3.6. Two variants of $X_{g,n}$

We shall also consider variant of the complexes $X_{g,n}$ above. First, we define the complexes $\check{X}_{g,n}$ in the same manner as above, except that we treat all edges in the same way, and declare the degree to be the total number of edges and the ordering of edges to be defined on all edges. In other words,

$$(28) \quad \check{X}_{g,n} = X_{g,n} \otimes \text{sgn}_n[-n - 1].$$

These graph complexes assemble naturally into symmetric sequences

$$X(r) := \bigoplus_g X_{g,r} \quad \text{and} \quad \check{X}(r) := \bigoplus_g \check{X}_{g,r}.$$

We consider also the enlargement $\widetilde{fX}_{g,n} \supset \check{X}_{g,n}$ obtained by relaxing the connectedness assumption on generating graphs. In other words, $\widetilde{fX}_{g,n}$ is defined just as $\check{X}_{g,n}$ except that the generators are allowed to be possibly disconnected graphs of genus g without vertices of valence 2, with external vertices labeled by symbols $\{1, \dots, n, \epsilon, \omega\}$, such that each label $1, \dots, n$ appears exactly once and every connected component contains at least one external vertex. The generators are oriented by a total ordering of the edges set. Note that we allow graphs with any number of ϵ - or ω -legs, with (ϵ, ω) - and (ω, ω) -edges, and we also allow disconnected graphs. The complexity of a generator for $\widetilde{fX}_{g,n}$ is the genus of the graph obtained by fusing all external vertices. We again assemble the graph complexes $\widetilde{fX}_{g,n}$ into a symmetric sequence

$$\widetilde{fX}(r) := \bigoplus_g \widetilde{fX}_{g,r}.$$

Lemma 3.7. *We have an isomorphism of symmetric sequences*

$$\widetilde{fX} \cong (\Delta_0 \boxtimes \text{Com}_1 \boxtimes \text{Com}_1) \otimes_{\mathbb{S}} \widetilde{fG}.$$

with Δ_0 the bisymmetric sequence (22), and with \boxtimes and $\otimes_{\mathbb{S}}$ operating on the first symmetric sequence structure.

Proof. Unpacking the notation, both sides are the same. \square

4. Euler characteristics

The goal of this section is to prove Theorem 1.1. The strategy of the proof is to first compute the equivariant Euler characteristic of the enlarged complexes $\widetilde{fX}_{g,n}$, and then extract from this the Euler characteristic of $\check{X}_{g,n}$. We then deduce the Euler characteristic of $X_{g,n}$, and hence that of $\text{gr}_2 H_c(\mathcal{M}_{g,n})$.

The proof uses some elementary but tedious computations with power series arising from the asymptotic expansions of polygamma functions. We perform these computations in §4.1, in order not to clutter the main argument line of §4.2. The reader is encouraged to skip §4.1 on the first reading.

4.1. Auxiliary computations of derivatives

We will use the following identity involving gamma functions.

Proposition 4.1 (after Songhafouo Tsopméné and Turchin [21]). *Let X and u be formal variables and $\ell \in \mathbb{N}$. Abbreviate*

$$(29) \quad E_\ell := \frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) \frac{1}{u^d} \quad \lambda_\ell := u^\ell (1 - u^\ell) \ell.$$

Then we have the equality of formal power series in u

$$(30) \quad (1 + \partial_a)^X e^{-a} (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \frac{(-\lambda_\ell)^X \Gamma(-E_\ell + X)}{\Gamma(-E_\ell)} =: U_\ell(X, u),$$

where on the right-hand side one inserts the asymptotic expansions as $u \rightarrow 0-$, and on the left-hand side one defines $(1 + \partial_a)^X := \exp(-X \sum_{j \geq 1} \frac{(-1)^j}{j} \partial_a^j)$. Furthermore, the coefficient of each power of u in the series on either side is a polynomial in X .

Let us note that, from the standard recurrence relation $\Gamma(z+1) = z\Gamma(z)$, one obtains the formula

$$(31) \quad \begin{aligned} U_\ell(X + p, u) \\ = (-\lambda_\ell)^p (-E_\ell + X) (-E_\ell + X + 1) \cdots (-E_\ell + X + p - 1) U_\ell(X, u), \end{aligned}$$

for any non-negative integer p .

The formula (30) is a variation of a formula found in [21, §2.2] and [23, Proposition 15.7]. The only difference is that in loc. cit. one replaces λ_ℓ by ℓu^ℓ . However, this change does not alter the proof of the formula. Hence we shall only sketch the derivation (due to Turchin) here.

Proof of Proposition 4.1. First, one checks that both sides of (30) are power series in u with coefficients that are polynomials in X . Since any polynomial is completely determined by its values on non-negative integers, it is sufficient to show (30) for $X = 0, 1, 2, \dots$. For $X = 0$ the identity (30) is trivial – both sides are 1. For $X = p$ a non-negative integer one has, using (31),

$$U_\ell(p, u) = \lambda_\ell^p E_\ell (E_\ell - 1) \cdots (E_\ell - p + 1).$$

Similarly, the left-hand side of (30) becomes

$$(1 + \partial_a)^p e^{-a} (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \partial_a^p (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \lambda_\ell^p E_\ell (E_\ell - 1) \cdots (E_\ell - p + 1),$$

and the proposition follows. \square

Recall Stirling's asymptotic expansion of the Γ function

$$(32) \quad \log \Gamma(z) \sim \left(z - \frac{1}{2} \right) \log z - z + \frac{1}{2} \log(2\pi) + B(z) \quad \text{as } z \rightarrow \infty$$

with

$$(33) \quad B(z) = \sum_{r \geq 2} \frac{B_r}{r(r-1)} \frac{1}{z^{r-1}}.$$

Using this expansion one can write $U_\ell(X, u)$ more explicitly as

$$\begin{aligned} \log U_\ell(X, u) &= \log \frac{(-\lambda_\ell)^X \Gamma(-E_\ell + X)}{\Gamma(-E_\ell)} \\ &= X(\log(\lambda_\ell E_\ell) - 1) + \left(-E_\ell + X - \frac{1}{2} \right) \log \left(1 - \frac{X}{E_\ell} \right) \\ &\quad + B(-E_\ell + X) - B(-E_\ell). \end{aligned}$$

Furthermore, note that we may compute, for non-negative integers k ,

$$(\log(1 + \partial_a))^k (1 + \partial_a)^X e^{-a} (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \partial_X^k U_\ell(X, u).$$

For later use, let us list the following special cases:

$$(34) \quad (\log(1 + \partial_a))(1 + \partial_a)^X e^{-a} (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \partial_X U_\ell(X, u) = (\log(\lambda_\ell(E_\ell - X)) + \psi_0(-E_\ell + X)) U_\ell(X, u)$$

$$(35) \quad (\log(1 + \partial_a))(1 + \partial_a)^{X+1} e^{-a} (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \partial_X U_\ell(X+1, u) \\ = \left(\log(\lambda_\ell(E_\ell - X)) + \psi_0(-E_\ell + X) + \frac{1}{-E_\ell + X} \right) \\ \cdot (-\lambda_\ell)(-E_\ell + X) U_\ell(X, u)$$

$$(36) \quad (\log(1 + \partial_a))^2 (1 + \partial_a)^X e^{-a} (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \partial_X^2 U_\ell(X, u) \\ = \left((\log(\lambda_\ell(E_\ell - X)) + \psi_0(-E_\ell + X))^2 + \psi_1(-E_\ell + X) \right) U_\ell(X, u)$$

$$(37) \quad (\log(1 + \partial_a))^2 (1 + \partial_a)^{X+1} e^{-a} (1 + \lambda_\ell a)^{E_\ell} |_{a=0} = \partial_X^2 U_\ell(X+1, u) \\ = \left(\left(\log(\lambda_\ell(E_\ell - X)) + \psi_0(-E_\ell + X) + \frac{1}{-E_\ell + X} \right)^2 + \psi_1(-E_\ell + X) - \frac{1}{(-E_\ell + X)^2} \right) \\ \cdot (-\lambda_\ell)(-E_\ell + X) U_\ell(X, u)$$

Here we use the recurrence relation (31) and the notation (3) for the digamma and trigamma series.

4.2. Euler characteristic of \widetilde{fX}

The graded vector space \widetilde{fX} has additional gradings from the complexity-grading, and from the number of ϵ and ω -legs. We can hence define the tri-graded \mathbb{S} -equivariant Euler characteristic

$$\chi^{u,v,w}(\widetilde{fX}) \in \Lambda[[u, v, w]],$$

with u, v, w being the formal variables tracking the complexity and the number of ϵ - and ω -legs, respectively. In other words, the Euler characteristic of the subcomplex with k ϵ - and l ω -legs and complexity m is the coefficient of $u^m v^k w^l$ in the formal power series. Then we have:

Proposition 4.2. *The equivariant Euler characteristic of \widetilde{fX} is*

$$\chi^{u,v,w}(\widetilde{fX}) = \prod_{\ell} U_{\ell} \left(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) (p_d + v^d + w^d), u \right)$$

with the function U_{ℓ} defined in Proposition 4.1 above.

Proof. We note that

$$\chi^{v,w}(\Delta_0 \boxtimes \text{Com}_1 \boxtimes \text{Com}_1) = \exp \left(\sum_{\ell \geq 1} \frac{1}{\ell} (q_{\ell} + v^{\ell} + w^{\ell}) p_{\ell} \right) \in \Lambda_{p,q}[[v, w]].$$

Hence

$$\begin{aligned} \chi^{v,w}((\Delta_0 \boxtimes \text{Com}_1 \boxtimes \text{Com}_1) \circ \text{Lie}_0) \\ &= \exp \left(\sum_{\ell \geq 1} \sum_{k \geq 1} \frac{\mu(k)}{\ell k} (q_{\ell} + v^{\ell} + w^{\ell}) \log(1 + p_{k\ell}) \right) \\ &= \exp \left(\sum_{\ell \geq 1} \frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) (q_d + v^d + w^d) \log(1 + p_{\ell}) \right) \\ &= \prod_{\ell \geq 1} (1 + p_{\ell})^{\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) (q_d + v^d + w^d)}. \end{aligned}$$

We hence compute, using Lemma 3.7 and Proposition 3.2:

$$\begin{aligned}
& \chi^{u,v,w}(\widetilde{fX}) \\
&= \chi^{u,w}((\Delta_0 \boxtimes \text{Com}_1 \boxtimes \text{Com}_1) \otimes_{\mathbb{S}} \widetilde{fG}) \\
&= \chi^{u,v,w}((\Delta_0 \boxtimes \text{Com}_1 \boxtimes \text{Com}_1) \circ \text{Lie}_0) \otimes_{\mathbb{S}} \text{BV}_0^{\text{red}} \\
&= \langle \chi^{v,w}(\Delta_0 \boxtimes \text{Com}_1 \boxtimes \text{Com}_1) \circ \chi(\text{Lie}_0), \chi^u(\text{BV}_0^{\text{red}}) \rangle \\
&= \prod_{\ell} (1 + \partial_a)^{\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)(q_d + v^d + w^d)} e^{-a} (1 + u^{\ell} (1 - u^{\ell}) \ell a)^{\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) \frac{1}{u^d}} |_{a=0} \\
&= \prod_{\ell} U_{\ell} \left(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)(q_d + v^d + w^d), u \right).
\end{aligned}$$

In the first three steps, we use Proposition 3.2, (15), and (14), respectively. In the last step, we use the function U_{ℓ} of Proposition 4.1. Note that here we had to work with bisymmetric sequences and hence two sets of power sums p_j, q_j , so that in the final expression the formula is in $\Lambda_q = \mathbb{Q}[[q_1, q_2, \dots]]$ instead of Λ . But the trivial replacement $q_d \rightarrow p_d$ yields the proposition. \square

We note that the part of \widetilde{fX} spanned by graphs without any ϵ - or ω -legs agrees with \widetilde{fG} . Hence we obtain, by setting $v = w = 0$ in the formula of Proposition 4.2:

Corollary 4.3.

$$\chi^u(\widetilde{fG}) = \chi^{u,v,w}(\widetilde{fX})(v = w = 0) = \prod_{\ell} U_{\ell} \left(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u \right).$$

We note that this corollary has been found earlier in [21].

4.3. Euler characteristic of the connected part with two ω -legs

Next we reduce the computation of the Euler characteristic of $\check{X}_{g,n}$ to that of $\widetilde{fX}_{g,n}$. As a first step we defined the sub-symmetric sequence $\widetilde{fX}^{\text{conn}} \subset \widetilde{fX}$ generated by graphs in which each connected component has at least one ϵ - or ω -leg. We then have

$$\widetilde{fX} = \widetilde{fX}^{\text{conn}} \boxtimes \widetilde{fG}.$$

Hence, by Corollary 4.3, the Euler characteristic of \widetilde{fX}^{conn} is

$$\begin{aligned}\chi^{u,v,w}(\widetilde{fX}^{conn}) &= \frac{\chi^{u,v,w}(\widetilde{fX})}{\chi^{u,v,w}(\widetilde{fX})(v=w=0)} \\ &= \prod_{\ell} \frac{U_{\ell}(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)(p_d + v^d + w^d), u)}{U_{\ell}(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)p_d, u)}\end{aligned}$$

Furthermore, we shall only be interested in the subcomplexes

$$\widetilde{fX}^{conn,j} \subset \widetilde{fX}^{conn}$$

spanned by graphs with $j = 0, 1, 2$ ω -legs, and we do not want to fix the number of ϵ -legs since the latter is not invariant under the differential. The relevant Euler characteristic is computed by the coefficient of w^j , evaluated at $v = 1$. For example, for $j = 0$ we set $w = 0$ to obtain

$$\begin{aligned}\chi^u(\widetilde{fX}^{conn,0}) &= \chi^{u,v,w}(\widetilde{fX}^{conn})(v=1, w=0) \\ &= \prod_{\ell \geq 1} \frac{U_{\ell}(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)(p_d + 1), u)}{U_{\ell}(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)p_d, u)} \\ &= \frac{U_1(p_1 + 1, u)}{U_1(p_1, u)} = u(1-u)Y_1^-, \end{aligned}$$

where we used that $\sum_{d|\ell} \mu(\ell/d) = \delta_{1\ell}$, equation (31), and the abbreviation

$$Y_{\ell}^{\pm} := \sum_{d|\ell} \mu(\ell/d)(u^{-d} \pm p_d).$$

Next, for $j = 1$:

$$\begin{aligned}\chi^u(\widetilde{fX}^{conn,1}) &= \partial_w|_{w=0} \chi^{u,v,w}(\widetilde{fX}^{conn})(v=1) \\ &= \frac{\partial_X U_1(p_1 + 1, u)}{U_1(p_1, u)} + \frac{U_1(p_1 + 1, u)}{U_1(p_1, u)} \left(\sum_{\ell \geq 2} \frac{\mu(\ell)}{\ell} \frac{\partial_X U_{\ell}(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)p_d, u)}{U_{\ell}(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d)p_d, u)} \right) \\ &= u(1-u)Y_1^- \left(-\frac{1}{Y_1^-} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(\ell u^{\ell} (1-u^{\ell}) Y_{\ell}^-) + \psi_0(-Y_{\ell}^-)) \right)\end{aligned}$$

To obtain the second line, we used again that $\sum_{d|\ell} \mu(\ell/d) = \delta_{1\ell}$. The first term in the second line is evaluated using (35), while the other derivatives are

computed using (34). Factoring out the common factor $u(1-u)Y_1^-$ then produces the third line. Similarly, one computes the second derivative.

$$\begin{aligned}
\chi^u(\widetilde{fX}^{conn,2}) &= \frac{1}{2}\partial_w^2|_{w=0} \chi^{u,v,w}(\widetilde{fX}^{conn})(v=1) \\
&= \frac{1}{2} \frac{\partial_X^2 U_1(p_1+1, u)}{U_1(p_1, u)} + \frac{1}{2} \frac{U_1(p_1+1, u)}{U_1(p_1, u)} \\
&\quad \times \left(\sum_{\ell \geq 2} \left(\frac{\mu(\ell)}{\ell} \right)^2 \frac{\partial_X^2 U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)}{U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)} \right) \\
&\quad + \frac{U_1(p_1+1, u)}{U_1(p_1, u)} \left(\sum_{\substack{\ell \geq 2 \\ 2|\ell}} \frac{\mu(\ell/2)}{\ell} \frac{\partial_X U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)}{U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)} \right) \\
&\quad + \frac{\partial_X U_1(p_1+1, u)}{U_1(p_1, u)} \left(\sum_{\ell \geq 2} \frac{\mu(\ell)}{\ell} \frac{\partial_X U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)}{U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)} \right) \\
&\quad + \frac{U_1(p_1+1, u)}{U_1(p_1, u)} \sum_{\ell > \ell' \geq 2} \left(\frac{\mu(\ell)}{\ell} \frac{\partial_X U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)}{U_\ell(\frac{1}{\ell} \sum_{d|\ell} \mu(\ell/d) p_d, u)} \right. \\
&\quad \times \left. \frac{\mu(\ell') \partial_X U_{\ell'}(\frac{1}{\ell'} \sum_{d|\ell'} \mu(\ell'/d) p_d, u)}{U_{\ell'}(\frac{1}{\ell'} \sum_{d|\ell'} \mu(\ell'/d) p_d, u)} \right) \\
&= \frac{1}{2} u(1-u)Y_1^- \left[\left(-\frac{1}{Y_1^-} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \left(\log(\ell u^\ell (1-u^\ell) Y_\ell^-) + \psi_0(-Y_\ell^-) \right) \right)^2 \right. \\
&\quad + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \left(\log(2\ell u^{2\ell} (1-u^{2\ell}) Y_{2\ell}^-) + \psi_0(-Y_{2\ell}^-) \right) \\
&\quad \left. - \frac{1}{(Y_1^-)^2} + \sum_{\ell \geq 1} \frac{\mu(\ell)^2}{\ell^2} \psi_1(-Y_\ell^-) \right]
\end{aligned}$$

To obtain the final equality we evaluated all terms using (31) and (34)-(37), and simplified the resulting expression. The steps of the final simplification are elementary and are omitted here, only the end result is shown.

4.4. Euler characteristic of \check{X}

Here, we show:

Proposition 4.4.

$$\chi^u(\check{X}) = \chi^u(\widetilde{fX}^{conn,2}) + u\chi^u(\widetilde{fX}^{conn,1}) + (u+u^2)\chi^u(\widetilde{fX}^{conn,0})$$

Proof. The symmetric sequence \check{X} differs from $\widetilde{fX}^{conn,2}$ only in so far that generators in \check{X} are not allowed to contain (ϵ, ω) - or (ω, ω) -edges, while we have not imposed such a condition in $\widetilde{fX}^{conn,2}$. To show the proposition we hence have to correct for those graphs.

A general graph in $\widetilde{fX}^{conn,2}$ with (ω, ω) -edge has the form

$$\omega — \omega \quad \left(\Gamma_0 \right),$$

with the right-hand part Γ_0 a general graph in $\widetilde{fX}^{conn,0}$. Hence to account for those graphs we have to subtract the expression $-u\chi^u(\widetilde{fX}^{conn,0})$ from $\chi^u(\widetilde{fX}^{conn,2})$, with the factor $-u$ accounting for the degree and genus shift introduced by the (ω, ω) -edge.

Similarly, a graph in $\widetilde{fX}^{conn,2}$ with (ϵ, ω) -edge has the form

$$\epsilon — \omega \quad \left(\Gamma_1 \right),$$

with $\Gamma_1 \in \widetilde{fX}^{conn,1}$, except for the caveat that Γ_1 must not contain an (ϵ, ω) -edge itself. But the graphs in $\widetilde{fX}^{conn,1}$ with (ϵ, ω) -edge themselves have the form

$$\epsilon — \omega \quad \left(\Gamma_0 \right),$$

with Γ_0 again a general graph in $\widetilde{fX}^{conn,0}$. Adding and subtracting the corresponding Euler characteristics one has shown the Proposition. \square

4.5. Proof of Theorem 1.1

According to (28) the complexes $\check{X}_{g,n}$ are obtained from $X_{g,n}$ by a degree shift and multiplication by the sign representation sgn_n . Hence we find that

$$\chi^u(X) = -\chi^u(\check{X})(-q_n \leftarrow q_n),$$

with the notation meaning that each occurrence of q_n should be replaced by $-q_n$ on the right-hand side. But using Proposition 4.4 and the formulas of

§4.3, we hence see that

$$\begin{aligned}
 \chi^u(X) = & \frac{u(u-1)Y_1^+}{2} \left[\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(-2\ell u^{2\ell} (1-u^{2\ell}) Y_{2\ell}^+) + \psi_0(-Y_{2\ell}^+)) \right. \\
 & + \left(-\frac{1}{Y_1^+} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(-\ell u^\ell (1-u^\ell) Y_\ell^+) + \psi_0(-Y_\ell^+)) \right)^2 \\
 & - \frac{1}{(Y_1^+)^2} + \sum_{\ell \geq 1} \frac{\mu(\ell)^2}{\ell^2} \psi_1(-Y_\ell^+) \\
 & \left. + 2u \left(-\frac{1}{Y_1^+} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(-\ell u^\ell (1-u^\ell) Y_\ell^+) + \psi_0(-Y_\ell^+)) \right) \right] \\
 & + u^2(u^2-1)Y_1^+.
 \end{aligned}$$

This expression can be simplified further. First, one has the following equality of formal power series

$$\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log(1-x^\ell) = - \sum_{\ell, n \geq 1} \frac{\mu(\ell)}{\ell} \frac{1}{n} x^{n\ell} = - \sum_{N \geq 1} \left(\sum_{d|N} \mu(d) \right) x^N = -x.$$

Using this formula three times to absorb the factors $(1-u^\ell)$ and $(1-u^{2\ell})$ inside the logarithms we obtain

$$\begin{aligned}
 \chi^u(X) = & \frac{u(u-1)Y_1^+}{2} \left[\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(-2\ell u^{2\ell} Y_{2\ell}^+) + \psi_0(-Y_{2\ell}^+)) \right. \\
 & + \left(-\frac{1}{Y_1^+} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(-\ell u^\ell Y_\ell^+) + \psi_0(-Y_\ell^+)) \right)^2 \\
 & - u^2 + u^2 - 2u^2 - \frac{1}{(Y_1^+)^2} + \sum_{\ell \geq 1} \frac{\mu(\ell)^2}{\ell^2} \psi_1(-Y_\ell^+) \left. \right] + u^2(u^2-1)Y_1^+ \\
 = & \frac{u(u-1)Y_1^+}{2} \left[\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(-2\ell u^{2\ell} Y_{2\ell}^+) + \psi_0(-Y_{2\ell}^+)) \right. \\
 & + \left(-\frac{1}{Y_1^+} + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} (\log(-\ell u^\ell Y_\ell^+) + \psi_0(-Y_\ell^+)) \right)^2 \\
 & - \frac{1}{(Y_1^+)^2} + \sum_{\ell \geq 1} \frac{\mu(\ell)^2}{\ell^2} \psi_1(-Y_\ell^+) \left. \right] + u^2(u-1)Y_1^+.
 \end{aligned}$$

Proof of Theorem 1.1. To show Theorem 1.1 we just use the above formula for $\chi^u(X)$, with the following modifications. First, to obtain the generating function in terms of genera instead of complexity one has to perform the replacements

$$(38) \quad u \rightarrow \hbar \text{ and } p_\ell \rightarrow \hbar^{-\ell} p_\ell, \quad \text{or equivalently, } u \rightarrow \hbar \text{ and } Y_\ell^+ \rightarrow Z_\ell.$$

Second, one needs to mind that the sum (1) runs only over stable indices $2g+n \geq 3$, while $X_{0,2} \cong \mathbb{Q}[-1]$ is nontrivial by our definition. Hence we need to subtract a term

$$(39) \quad -\frac{1}{2}(p_1^2 + p_2) = -\frac{1}{2}(P_1^2 + P_2) - P_1$$

from the Euler characteristic to account for this difference.

Finally, by Corollary 3.6 there is a further correction $\hbar p_1$ arising from the contribution of $g = n = 1$. Thus, we start from the formula for $\chi^u(X)$ above, then apply the substitution rule (38), add $\hbar p_1$ and subtract (39) to finally obtain the formula of Theorem 1.1. \square

5. Discussion of terms in the Euler characteristic formula

The literature contains two complementary toolsets for computing dimensions (Hilbert series) and Euler characteristics of graph complexes. The first is the calculus of symmetric functions, typically paired with operadic methods, as in [9, 21] and this paper. The second is combinatorial counting weighted by automorphisms, along the lines of the Pólya enumeration theorem, as in [3, 24]. When the first toolset is applicable, it typically yields relatively economical proofs and closed expressions for the generating functions. However, it also tends to obfuscate natural correspondences between terms in the resulting formulas and subsets of generators for the graph complex. In this section, we shall hence briefly discuss how the individual terms in the formulas of Theorem 1.1 and Corollary 1.2 relate to graph generators for $X_{g,n}$.

First recall from [19, §6.1] that $X_{g,n}$ is quasi-isomorphic to its subcomplex $X_{g,n}^*$, generated by graphs Γ that are disjoint unions of one or two connected components with ω -decorations together with one of the following:


$$(\text{empty graph}), \quad \epsilon \text{ ——— } \epsilon, \quad \epsilon \text{ ——— } j, \quad \begin{array}{c} \epsilon \text{ ——— } j \\ \epsilon \text{ ——— } \epsilon \end{array}.$$

The contribution of these four graphs to the generating function of the Euler characteristic is precisely

$$(1 - \hbar)P_1,$$

which explains the corresponding factor in (4).

Now, consider the disjoint union of the connected components that contain ω -decorations. In genus 0, the only possible graphs are pairs of two trees, as shown.



The Euler characteristic of the graph complex of rooted trees, like that of Lie_0 , is equal to $\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_\ell$. Since we have two trees in the graph, we have to take a symmetric product of two such rooted tree complexes. On the level of Euler characteristics taking the symmetric product translates to the plethysm with the symmetric function $h_2 = \frac{1}{2}(p_1^2 + p_2)$, and

$$h_2 \circ \left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_\ell \right) = \frac{1}{2} \left[\left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_\ell \right)^2 + \sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_{2\ell} \right],$$

explaining the corresponding terms in the genus 0 and 1 Euler characteristics.

In genus $g \geq 2$, there are still some graphs with one of the two ω -legs being the root of a tree. The corresponding contributions to the Euler characteristic hence have a factor $\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_\ell$ as well from the tree part, which is multiplied by the contribution from the non-tree part.

Let us next turn to the remaining terms in the Euler characteristic formulas, coming from the non-tree connected components in graphs. The characteristic feature of those terms is that they are finite linear combinations of monomials in the P_j and P_j^{-1} . This can be seen directly from using graph counting techniques to compute the Euler characteristic, using the same strategy of proof as in [3].

Consider an arbitrary graph Γ in $X_{g,n}$. We call the *core* $\gamma = [\Gamma]$ of Γ the graph obtained by the following algorithm:

- Remove all numbered external vertices and their adjacent edges.
- Recursively remove all univalent internal vertices thus created, with their adjacent edges.

- Remove the bivalent vertices thus produced and merge the two edges adjacent to them.

Let $X_{g,n}^\gamma \subset X_{g,n}$ be the subcomplex spanned by graphs with core γ , and set

$$z_\gamma := \sum_{n \geq 0} \chi^S(X_{g,n}^\gamma).$$

Then we have $\sum_{n \geq 0} \chi^S(X_{g,n}) = \sum_\gamma z_\gamma$ with the first sum being over all (isomorphism classes of) genus g cores γ . Assume for simplicity that the core γ only has connected components of genera ≥ 1 . Then one may show as in [3, Proposition 3.2] that


$$(40) \quad z_\gamma = \frac{(-1)^{e(\gamma)+1}}{|\text{Aut}(\gamma)|} \sum_{\tau \in \text{Aut}(\gamma)} \text{sgn}(\tau_E) \frac{P^{i(\tau_V)} P^{i(\tau_E)}}{P^{i(\tau_H)}}$$

with $e(\gamma)$ the number of structural edges of γ , τ_V , τ_E , and τ_H the permutations on the sets of vertices, edges and half-edges of γ , with $i(\sigma) = (i_1(\sigma), \dots)$ the cycle type of a permutation σ , and with the multi-index notation

$$P^{i(\sigma)} = P_1^{i_1(\sigma)} P_2^{i_2(\sigma)} \dots \in \Lambda.$$

Since in every genus there are only finitely many possible core graphs contributing, the Euler characteristic in this genus must hence be a finite linear combination of such Laurent monomials in the P_i , of the form given by (40).

Next suppose that the core consists of a single isolated external vertex decorated by ω , and another higher genus component. The graphs with this core are unions of trees with ω -labelled root and some other graph with only one ω -decoration.



These graphs then contribute summands of the form

$$\left(\sum_{\ell \geq 1} \frac{\mu(\ell)}{\ell} \log P_\ell \right) f(P_1, P_2, \dots),$$

with f some finite sum of Laurent monomials of the special form given by (40). This then explains the structure of the terms appearing in Corollary 1.2.

References

- [1] ENRICO ARBARELLO and MAURIZIO CORNALBA. Calculating cohomology groups of moduli spaces of curves via algebraic geometry, *Inst. Hautes Études Sci. Publ. Math.* No. **88** (1998), 97–127. [MR1733327](#)
- [2] GREGORY ARONE and VICTOR TURCHIN. Graph-complexes computing the rational homotopy of high dimensional analogues of spaces of long knots. *Ann. Inst. Fourier* **65** (2015), no. 1, 1–62. [MR3449147](#)
- [3] MELODY CHAN, CAREL FABER, SOREN GALATIUS and SAM PAYNE. The \mathbb{S}_n -equivariant top weight Euler characteristic of $M_{g,n}$, [arXiv:1904.06367](#), 2019.
- [4] MELODY CHAN, SOREN GALATIUS and SAM PAYNE. Tropical curves, graph complexes, and top weight cohomology of M_g , *J. Amer. Math. Soc.* **34** (2021), no. 2, 565–594. [MR4280867](#)
- [5] MELODY CHAN, SOREN GALATIUS and SAM PAYNE. Topology of moduli spaces of tropical curves with marked points. Preprint, [arXiv:1903.07187](#), 2019. [MR4280867](#)
- [6] BENOIT FRESSE, VICTOR TURCHIN and THOMAS WILLWACHER. On the rational homotopy type of embedding spaces of manifolds in \mathbb{R}^n . Preprint, [arXiv:2008.08146](#), 2020. [MR3870769](#)
- [7] FERENC GERLITS. The Euler characteristic of graph complexes via Feynman diagrams. Preprint, [arXiv:0412094v2](#), 2004.
- [8] EZRA GETZLER. Operads and moduli spaces of genus 0 Riemann surfaces, *The moduli space of curves* (Texel Island, 1994), 199–230, *Progr. Math.*, 129, Birkhäuser Boston, Boston, MA, 1995. [MR1363058](#)
- [9] EZRA GETZLER and MIKHAIL KAPRANOV. Modular operads. *Compositio Math.* **110** (1998), no. 1, 65–126. [MR1601666](#)
- [10] EUGENE GORSKY. The equivariant Euler characteristic of moduli spaces of curves. *Adv. Math.* **250** (2014), 588–595. [MR3122177](#)
- [11] JOHN HARER and Don Zagier. The Euler characteristic of the moduli space of curves. *Invent. Math.* **85** (1986), no. 3, 457–485. [MR0848681](#)
- [12] MAXIM KONTSEVICH. Operads and motives in deformation quantization. *Lett. Math. Phys.* **48** (1999), 35–72. [MR1718044](#)
- [13] MAXIM KONTSEVICH. Formal (non)commutative symplectic geometry. In: *The Gelfand Mathematical Seminars, 1990–1992*, 173–187, Birkhäuser Boston, Boston, MA, 1993. [MR1247289](#)

- [14] PASCAL LAMBRECHTS and ISMAR VOLIĆ. Formality of the little N -disks operad. *Mem. Amer. Math. Soc.* 230 (2014), no. 1079, viii+116 pp. [MR3220287](#)
- [15] JEAN-LOUIS LODAY and BRUNO VALLETTE. Algebraic operads. *Grundlehren Math. Wiss.*, 346, Springer, Heidelberg, 2012. [MR2954392](#)
- [16] IAN MACDONALD. Symmetric functions and Hall polynomials. Second edition. *Oxford Classic Texts in the Physical Sciences*. The Clarendon Press, Oxford University Press, New York, 2015. xii+475 pp. [MR3443860](#)
- [17] IB MADSEN and MICHAEL WEISS. The stable moduli space of Riemann surfaces: Mumford’s conjecture. *Ann. of Math.* (2) **165** (2007), no. 3, 843–941. [MR2335797](#)
- [18] DAN MARGALIT. Problems, questions, and conjectures about mapping class groups. *Breadth in contemporary topology*, 157–186, *Proc. Sympos. Pure Math.*, 102, Amer. Math. Soc., Providence, RI, 2019. [MR3967367](#)
- [19] SAM PAYNE and THOMAS WILLWACHER. Weight two compactly supported cohomology of moduli spaces of curves. Preprint, [arXiv:2110.05711](#), 2021 [MR4735654](#)
- [20] CHRIS PETERS and JOZEF STEENBRINK. Mixed Hodge structures. *Ergeb. Math. Grenzgeb.* (3), 52, Springer-Verlag, Berlin, 2008. [MR2393625](#)
- [21] PAUL ARNAUD SONGHAFOUO TSOPMÉNÉ and VICTOR TURCHIN. Euler characteristics for spaces of string links and the modular envelope of L_∞ . *Homology Homotopy Appl.* **20** (2018), no. 2, 115–144. [MR3806571](#)
- [22] ORSOLA TOMMASI. Rational cohomology of the moduli space of genus 4 curves. *Compos. Math.* **141** (2005), no. 2, 359–384. [MR2134272](#)
- [23] VICTOR TURCHIN. Hodge-type decomposition in the homology of long knots. *J. Topol.*, **3**(3):487–534, 2010. [MR2684511](#)
- [24] THOMAS WILLWACHER and MARKO ŽIVKOVIĆ. Multiple edges in M. Kontsevich’s graph complexes and computations of the dimensions and Euler characteristics. *Adv. Math.* **272** (2015), 553–578. [MR3303240](#)

Sam Payne
Department of Mathematics
2515 Speedway, PMA 8.100
Austin, TX 78722
USA
E-mail: sampayne@utexas.edu

Thomas Willwacher
Department of Mathematics
ETH Zurich
Rämistrasse 101
8092 Zurich
Switzerland
E-mail: thomas.willwacher@math.ethz.ch