

# GUBIC: The global urban biological invasions compendium for plants

Daijiang Li<sup>1</sup>  | Luke J. Potgieter<sup>2,3</sup>  | Myla F. J. Aronson<sup>4</sup> | Irena Axmanová<sup>5</sup> | Benjamin Baiser<sup>6</sup>  | Marta Carboni<sup>7</sup>  | Laura Celesti-Grapow<sup>8</sup> | Sonja Knapp<sup>9,10</sup>  | Ingolf Kühn<sup>9,10,11</sup> | Ana Carolina Lacerda de Matos<sup>12</sup> | Zdeňka Lososová<sup>5</sup>  | Flavia A. Montaño-Centellas<sup>13</sup> | Petr Pyšek<sup>14,15</sup>  | David M. Richardson<sup>3</sup> | Lauren B. Trotta<sup>6</sup> | Rafael D. Zenni<sup>12</sup>  | Sarel S. Cilliers<sup>16</sup> | Bruce D. Clarkson<sup>17</sup> | Amy J. S. Davis<sup>18</sup> | Rebecca W. Dolan<sup>19</sup> | Marcin K. Dyderski<sup>20</sup> | Franz Essl<sup>21</sup> | Orou G. Gaoue<sup>22</sup>  | Joanne Gui<sup>2</sup> | Charly Géron<sup>23,24</sup>  | Gustavo Heringer<sup>12,25,26</sup> | Cang Hui<sup>27,28</sup> | Anzar A. Khuroo<sup>29</sup>  | Stefan Klotz<sup>9</sup> | Peter M. Kotanen<sup>30</sup> | Holger Kreft<sup>31,32</sup> | Frank A. La Sorte<sup>33,34</sup>  | Jonas J. Lembrechts<sup>35</sup>  | Bernd Lenzner<sup>21</sup>  | Christopher A. Lepczyk<sup>36</sup> | Scott MacIvor<sup>2</sup>  | Cristina Martínez-Garza<sup>37</sup>  | Akira S. Mori<sup>38</sup>  | Charles Nilon<sup>39</sup> | Jan Pergl<sup>14</sup> | Stefan J. Siebert<sup>16</sup> | Alyona S. Tretyakova<sup>40</sup> | Toby P. N. Tsang<sup>2</sup>  | Kei Uchida<sup>41</sup>  | Mark van Kleunen<sup>18,42</sup> | Montserrat Vilà<sup>43,44</sup> | Hua-Feng Wang<sup>45</sup> | Patrick Weigelt<sup>46</sup>  | Peter Werner<sup>47</sup> | Nicholas S. G. Williams<sup>48</sup> | Marten Winter<sup>10</sup>  | Marc W. Cadotte<sup>2</sup> 

## Correspondence

Daijiang Li  
Email: [daijianglee@gmail.com](mailto:daijianglee@gmail.com)

Luke J. Potgieter  
Email: [lukepotgieter2@gmail.com](mailto:lukepotgieter2@gmail.com)

Marc W. Cadotte  
Email: [marc.cadotte@utoronto.ca](mailto:marc.cadotte@utoronto.ca)

## Funding information

Natural Sciences and Engineering Research Council of Canada, Grant/Award Number: #RGPIN-2022-03579 and 386151; Deutsche Forschungsgemeinschaft, Grant/Award Number: 264740629, FZT 118 and 202548816; National Science

## Abstract

1. Urban areas are foci for the introduction of non-native plant species, and they often act as launching sites for invasions into the wider environment. Although interest in biological invasions in urban areas is growing rapidly, and the extent and complexity of problems associated with invasions in these systems have increased, data on the composition and numbers of non-native plants in urbanized areas remain scattered and idiosyncratic.
2. We assembled data from multiple sources to create the Global Urban Biological Invasions Compendium (GUBIC) for vascular plants representing 553 urban centres from 61 countries across every continent except Antarctica.
3. The GUBIC repository includes 8140 non-native plant species from 253 families. The number of urban centres in which these non-native species occurred had a

Daijiang Li and Luke J. Potgieter contributed equally to this work.

For affiliations refer to page 9.

This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Ecological Solutions and Evidence* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

Foundation, Grant/Award Number: DEB-2213567; The Italian Ministry of University and Research, Grant/Award Number: CN00000033; CNPq-Brazil, Grant/Award Number: 302643/2022-2; Czech Science Foundation, Grant/Award Number: 19-28491X and 25-15190S; Czech Academy of Sciences, Grant/Award Number: RVO 67985939

Handling Editor: Holly Jones

log-normal distribution, with 65.2% of non-native species occurring in fewer than 10 urban centres.

**4. Practical implications:** The dataset has wider applications for urban ecology, invasion biology, macroecology, conservation, urban planning and sustainability. We hope this dataset will stimulate future research in invasion ecology related to the diversity and distributional patterns of non-native flora across urban centres worldwide. Further, this information should aid the early detection and risk assessment of potential invasive species, inform policy development and assist in setting management priorities.

#### KEY WORDS

Alien species, biodiversity change, biological invasions, cities, naturalized species, non-native plants, urbanization

## 1 | INTRODUCTION

Urban areas, characterized by their high human population density and extensive landscape modification, present unique opportunities for the establishment and spread of non-native species. The convergence of global trade, transportation networks, modified microclimates and human-mediated disturbances in urban areas facilitates the introduction and proliferation of non-native species (Gallardo et al., 2016; Potgieter et al., 2024). Urban plant invasions can have profound ecological, economic and social impacts due to altered ecosystem services, impacts on human health and costs incurred from management efforts (Heringer et al., 2024; Potgieter et al., 2017). However, there is a lack of foundational data on which species occur in urban centres globally. This data gap limits our ability to assess the potential threats non-native plants pose to urban ecosystems and the services they might provide (Milanović et al., 2020), with current knowledge remaining geographically heterogeneous and focused on only a few well-studied taxa (Vaz et al., 2018).

Frameworks for understanding and managing urban plant invasions are less frequently studied than in other habitats (but see Gaertner et al., 2016; Potgieter & Cadotte, 2020). While existing frameworks integrate concepts from landscape ecology, population biology and socioecological systems, they are limited in number and scope, highlighting the need for further development to facilitate a better understanding of the mechanisms that drive invasions in urban areas as well as options for managing them. Managers in urban areas face unique challenges due to the interplay between the built environment and complex socioeconomic factors, which can significantly alter ecosystem conditions. However, these challenges have only recently been incorporated into models to predict urban invasion dynamics and impacts and identify appropriate management strategies (Gaertner et al., 2016; Potgieter et al., 2022).

Despite these advances, empirical studies on urban biological invasions remain limited, particularly in terms of taxonomic

coverage and spatial scale (Cadotte et al., 2017). Most empirical studies have focused on the ecology of particular non-native species within small urban areas. This narrow focus limits the generalizability of findings across different organisms and urban contexts. Although numerous regional and city-specific inventories of non-native species exist, these are often from uncoordinated efforts carried out independently by research groups focusing on particular research questions. Therefore, these diverse resources lack harmonization of collection methods, taxonomy and sampling effort, making them challenging to be easily used. Moreover, because some of this work is developed in collaboration with city practitioners and managers, many studies are published in the grey literature and only available in non-English languages, limiting their accessibility. While these biological inventories are crucial to advancing our understanding of urban biological invasions at the city and regional levels, a comprehensive global dataset documenting the non-native flora in urban areas around the world is required to understand the role of urban areas in shaping the patterns of plant invasions and the underlying processes. Here, we unify this diverse body of knowledge and present a global repository of non-native flora in urban centres around the globe. This repository serves as a valuable resource for improving our understanding of urban non-native floras by providing essential data, fostering collaboration, informing management and policy and facilitating coordinated global responses to the challenges they present.

## 2 | METHODS AND MATERIALS

To compile a list of non-native plant species in urban areas (see Section 2.1.2 for the methods used to delineate urban boundaries) globally, we combined multiple data sources. This approach allows for the application of standardized selection and inclusion criteria over multiple individual datasets, resulting in a harmonized and consistent dataset across urban areas and regions.

We included only established non-native plant species, which are those with self-sustaining populations, also commonly referred to as naturalized (Blackburn et al., 2011; Richardson et al., 2000; see Section 2.1.5).

## 2.1 | Data acquisition and compilation

### 2.1.1 | Data source 1: Global Urban Biological Invasions Consortium

An international workshop to address biological invasions in urban ecosystems was hosted by the Centre for Invasion Biology in Stellenbosch, South Africa, in November 2016 (Gaertner et al., 2017). This workshop led to the creation of the [Global Urban Biological Invasions Consortium](#), which hosted a coordinating meeting in June 2019 that brought together more than 70 researchers from 14 countries from all continents except Antarctica. One of the prioritized activities was to compile lists of non-native plant species for urban areas. A working group "Synthesizing Global Urban Biological Invasion Knowledge" (sGUBIK, funded by sDiv, the synthesis centre of iDiv, the German Centre for Integrative Biodiversity Research) was later established in September 2023 to synthesize these global data and examine the patterns and mechanisms driving non-native plant species' invasions in urban areas.

We compiled data using the following approaches. First, we sent a request to over 150 members of the Global Urban Biological Invasions Consortium in 2019 to upload datasets for any urban taxa to a SharePoint repository at the University of Toronto. The cut-off for the data request was December 2021. Second, during August to November 2019, we searched the published literature in English, Portuguese and Spanish as well as the Dryad data repository ([www.datadryad.org](http://www.datadryad.org)) for studies and datasets containing species lists for urban areas around the world, using keywords such as 'alien', 'animal', 'built-up', 'city', 'urban\*', 'non-native', 'exotic' and 'plant'. These approaches yielded urban datasets that encompassed various taxa and spatial scales, incorporating demographic, environmental and taxon-specific information. Additionally, we included the Urban Biodiversity Research Coordination Network (UrBioNet) dataset, a large multi-city compilation (Aronson et al., 2014), featuring 14,240 spontaneous plant species (i.e. not cultivated or planted), of which 4241 are identified as non-native, derived from published surveys across 110 urban areas in five biogeographic regions.

To ensure consistency across the datasets, we standardized city and country names by resolving variations in spelling and correcting potential typographical errors. In instances where multiple urban centres within the same country shared the same name (e.g. Madison, Wisconsin vs. Madison, Indiana in the United States), we excluded these entries from the database if it was not possible to unambiguously determine the specific city to which the data pertained. Given that most data lacked spatially explicit coordinates, precise delineations of city boundaries were unavailable.

As a result, datasets collected from data contributors, repositories or the literature were generally treated as representing areas surrounding the urban centres rather than being confined to specific urban boundaries.

### 2.1.2 | Data source 2: Global Biodiversity Information Facility

Before extracting occurrence data for each urban area from the Global Biodiversity Information Facility (GBIF), we delineated the boundaries of urban areas. We used the global urban centres data provided by the Global Human Settlement Layer (GHSL, Pesaresi et al., 2019, <https://ghsl.jrc.ec.europa.eu/ucdb2018Overview.php>), which defines urban centres as contiguous  $1\text{km}^2$  grid cells with a population density of at least 1500 inhabitants per  $\text{km}^2$  of permanent land (areas that are consistently above water and exclude bodies of water, such as oceans, seas, large rivers and lakes) or with more than 50% built-up surface shared on permanent land and with at least 50,000 inhabitants in the cluster with smoothed boundaries and small gaps ( $<15\text{km}^2$ ) filled. Overall, there are 13,189 unique urban centres worldwide. Subsequently, smaller, nearby urban centres located within a 5 km radius of the larger urban centres were integrated into the larger one, as these proximally situated centres are close enough to be considered a single urban entity and often are considered part of the metropolitan area. We refrained from further merging smaller centres that, although within a 5 km radius of the previously merged smaller centres, were situated beyond the 5 km boundary from the larger urban centre. This process resulted in 11,621 unique urban centres globally.

In August 2023, we queried GBIF and downloaded plant occurrence records from each urban centre to compile the flora of these urban areas (see Table S1 for the DOIs of downloaded datasets). The initial download comprised over 500 million records. We cleaned the GBIF data of each of the urban centres by removing records with common issues such as erroneous coordinates using R package 'CoordinateCleaner' (Zizka et al., 2019). We also removed all records with identification above species level, fossil specimens, preserved specimens, living specimens and those with locality uncertainty greater than 30 km or within a 500 m vicinity of biodiversity institutions, botanic gardens, zoos, museums, GBIF headquarters, etc.

### 2.1.3 | Quality control and merging of data

Before merging data from sources 1 and 2, we conducted preliminary filtering of these datasets. For each urban centre with GBIF data, we used the number of observations of each species as a proxy for the abundance of that species. We calculated observed species richness and estimated species richness using the Chao1 equation, which incorporates singletons and doubletons (i.e. species observed only once or twice):

$$\text{Estimated richness} = \text{Observed richness} + S^2 / (2D) \quad (1)$$

where  $S$  represents the number of singletons and  $D$  is the number of doubletons (Hsieh & Chao, 2016). We also determined the sample coverage percentage, a measure of sample completeness, based on the rarefied estimate of the total number of individuals in each urban centre using the R package 'iNEXT' for rarefaction (Chao et al., 2014; Hsieh et al., 2024).


We considered an urban centre to have robust GBIF data if: (1) it had over 1000 observed plant species; (2) the community sample coverage was >90%; and (3) the observed species richness was greater than 75% of the estimated species richness. We used these criteria to balance the number of retained urban centres and data quality. For data source 1, if an urban centre had more than 300 plant species, we retained it and further integrated it with data source 2 (GBIF data) of that urban centre regardless of the GBIF data quality. If an urban centre had fewer than 300 species from data source 1 and did not have adequate GBIF data coverage, we removed that urban centre from our database. If an urban centre had fewer than 300 species from data source 1 but had adequate GBIF data coverage (i.e. met the above three criteria), we retained both data sources for that urban centre. We removed those urban centres with only GBIF data that did not meet the three criteria above (see Figure 1 for a schematic workflow). Like the criteria we used for the GBIF data, we selected 300 species here to balance the number of urban centres and their data quality after carefully explored our datasets. The final database included 553 urban centres (Figure 2). For each of these urban centres, we derived a list of established non-native plant species using the merged data sources.


## 2.1.4 | Standardize species names

We standardized species and family names against the World Checklist of Vascular Plants (WCVP, Govaerts, 2024) for the merged database using the R package rWCVP (version 1.0.3, Brown et al., 2023). We selected WCVP as it represents one of the most comprehensive and up-to-date taxonomic resources available (Grenié et al., 2022). WCVP also serves as the taxonomic backbone for the most recent version of the Global Naturalized Alien Flora (GloNAF), which was updated following van Kleunen et al. (2019). GloNAF was used to determine whether a species is non-native in a particular region where an urban centre was located (see Section 2.1.5 below). Note that species with "unplaced names" ( $n=65$  across all species) or has not match from WCVP were excluded from the final dataset (<https://powo.science.kew.org/about-wcvp#unplacednames>), reflecting the challenges in our current taxonomic knowledge of plants worldwide. We also merged subspecies or varieties to the main species and only kept binomial species names in the final database.


## 2.1.5 | Cross-validation to determine the status of species

To distinguish between established (naturalized) and native or casual species (i.e. those that might flourish and even reproduce occasionally in an area but which do not form self-replacing populations;





**FIGURE 2** Geographic distribution of urban centres across the world (panel a;  $n=553$ ) and Europe (panel b) and the number of established non-native plant species they contain (coloured points).



**FIGURE 3** The distribution of family sizes for the 253 established non-native plant families in the dataset. The main plot contains the top 20 families which together account for 61.6% of all established non-native plant species in our dataset. The numbers after the family names represent the approximate number of total accepted species of each family. The embedded plot presents the distribution of the number of non-native plant species across all families.

TABLE 1 The most widespread (top 30) established non-native plant species in urban centres ( $n=553$ ) across the world. Note that this list was derived from different sampling efforts and has a bias in favour of non-native species in European and North American urban centres.

| Scientific name                | Family           | Number of urban centres | Number of countries | Number of GBIF records |
|--------------------------------|------------------|-------------------------|---------------------|------------------------|
| <i>Erigeron canadensis</i>     | Asteraceae       | 469                     | 47                  | 64,760                 |
| <i>Veronica persica</i>        | Plantaginaceae   | 451                     | 41                  | 41,176                 |
| <i>Oxalis corniculata</i>      | Oxalidaceae      | 434                     | 48                  | 23,721                 |
| <i>Datura stramonium</i>       | Solanaceae       | 410                     | 46                  | 12,531                 |
| <i>Robinia pseudoacacia</i>    | Fabaceae         | 404                     | 41                  | 44,657                 |
| <i>Syringa vulgaris</i>        | Oleaceae         | 393                     | 29                  | 21,767                 |
| <i>Amaranthus retroflexus</i>  | Amaranthaceae    | 381                     | 41                  | 8602                   |
| <i>Galinsoga quadriradiata</i> | Asteraceae       | 376                     | 40                  | 18,509                 |
| <i>Ailanthus altissima</i>     | Simaroubaceae    | 369                     | 35                  | 50,732                 |
| <i>Medicago sativa</i>         | Fabaceae         | 369                     | 39                  | 24,969                 |
| <i>Prunus cerasifera</i>       | Rosaceae         | 369                     | 26                  | 26,924                 |
| <i>Aesculus hippocastanum</i>  | Sapindaceae      | 367                     | 24                  | 35,349                 |
| <i>Reynoutria japonica</i>     | Polygonaceae     | 367                     | 29                  | 88,542                 |
| <i>Cymbalaria muralis</i>      | Plantaginaceae   | 366                     | 33                  | 31,151                 |
| <i>Matricaria discoidea</i>    | Asteraceae       | 363                     | 33                  | 26,254                 |
| <i>Melissa officinalis</i>     | Lamiaceae        | 359                     | 25                  | 20,768                 |
| <i>Buddleja davidii</i>        | Scrophulariaceae | 356                     | 33                  | 45,122                 |
| <i>Lunaria annua</i>           | Brassicaceae     | 340                     | 19                  | 12,135                 |
| <i>Galinsoga parviflora</i>    | Asteraceae       | 337                     | 43                  | 7306                   |
| <i>Rosa rugosa</i>             | Rosaceae         | 334                     | 22                  | 18,499                 |
| <i>Vinca major</i>             | Apocynaceae      | 326                     | 23                  | 11,597                 |
| <i>Helianthus tuberosus</i>    | Asteraceae       | 320                     | 34                  | 4244                   |
| <i>Tanacetum parthenium</i>    | Asteraceae       | 320                     | 27                  | 9527                   |
| <i>Lepidium draba</i>          | Brassicaceae     | 315                     | 29                  | 14,831                 |
| <i>Acer negundo</i>            | Sapindaceae      | 313                     | 36                  | 17,459                 |
| <i>Lysimachia punctata</i>     | Primulaceae      | 312                     | 15                  | 10,279                 |
| <i>Brassica napus</i>          | Brassicaceae     | 307                     | 25                  | 5376                   |
| <i>Impatiens glandulifera</i>  | Balsaminaceae    | 301                     | 25                  | 35,332                 |
| <i>Lepidium didymum</i>        | Brassicaceae     | 301                     | 31                  | 13,459                 |
| <i>Prunus laurocerasus</i>     | Rosaceae         | 298                     | 22                  | 48,188                 |

Richardson et al., 2000) located in a specific urban centre, we used the GloNAF database as it provides the most updated information of naturalized plant species across the world. For each urban centre, we used the delineated boundaries provided by the GHSL. For each species listed within an urban centre, we classified the species as non-native to that urban centre if its polygon intersected with the species' naturalized or invasive range. We also cross-referenced all species with local checklists of non-native plant species validated by experts (Kalusová et al., 2024). Therefore, for those urban centres (mostly in Europe), the lists of naturalized species were slightly different from those based on GloNAF alone.

### 3 | GENERAL PATTERNS


We present a global urban non-native flora for 553 cities from 61 countries across every continent with permanent human

settlements (Figure 2). These data are, however, biased towards European and North American urban centres, which together account for 80.8% of all non-native species, and 82.2% of all records within our database across the world, respectively. Our global repository includes 8140 established non-native plant species from 253 families (Figure 3). Most families contain few species, with 73 families each containing 20 or more non-native species (Figure 3). Asteraceae, Poaceae, Fabaceae and Rosaceae contain about one-third of all species ( $n=2641$ ; Table 1). The most widespread non-native plant species can be found in Table 1; the top 20 urban centres and countries with the greatest number of non-native plant species in our database can be found in Table 2. A rarefaction of species occurrences across urban centres (Figure 4) shows that we are approaching an asymptote with our sample of 553 urban centres. However, the sampling curve also suggests that more urban floral sampling is needed, especially from regions with sparse data (e.g. South Asia, northern South America).

**TABLE 2** The top 20 urban centres (left) and the top 20 countries (right) with the greatest number of non-native plant species. Note that these lists are skewed towards European and North American urban centres (see Figure 2). The numbers presented for some countries (e.g. France) also included non-native plant species from their overseas urban centres.

| Urban centre       | Number of established non-native species | Country                  | Number of established non-native species |
|--------------------|--|--------------------------|--|
| New York           | 1663                                     | United States of America | 4409                                     |
| Los Angeles        | 1534                                     | Australia                | 2596                                     |
| Sydney             | 1486                                     | France                   | 2187                                     |
| Philadelphia       | 1455                                     | New Zealand              | 1561                                     |
| Melbourne          | 1450                                     | Canada                   | 1476                                     |
| Washington D.C.    | 1414                                     | Russia                   | 1251                                     |
| Auckland           | 1310                                     | Japan                    | 1154                                     |
| Boston             | 1300                                     | Mexico                   | 1142                                     |
| San Jose (USA)     | 1231                                     | Germany                  | 1123                                     |
| Tijuana            | 1066                                     | United Kingdom           | 986                                      |
| St. Louis          | 1058                                     | Switzerland              | 966                                      |
| Tokyo              | 1038                                     | South Africa             | 947                                      |
| London (UK)        | 1014                                     | Spain                    | 916                                      |
| Christchurch       | 1009                                     | Belgium                  | 906                                      |
| Adelaide           | 995                                      | Netherlands              | 860                                      |
| Brisbane           | 994                                      | Sweden                   | 832                                      |
| Portland (OR, USA) | 990                                      | Denmark                  | 800                                      |
| Moscow             | 924                                      | Norway                   | 792                                      |
| Chicago            | 895                                      | Portugal                 | 596                                      |
| Perth              | 858                                      | Brazil                   | 588                                      |


**FIGURE 4** Rarefaction curve of the number of non-native plant species in 553 urban centres.



#### 4 | USAGE NOTES

In forming a dataset of this magnitude, we made several simplifying decisions and recognize that limitations are inevitable. Some issues to be cognizant of for analysis and interpretation include:

1. Our definition of urbanized areas delineated contiguous areas. Because of this definition, some urbanized areas span multiple regions or municipalities and form contiguous land areas. In these cases, the urbanized region is referred to as the largest administrative centre; for example, Guangzhou, China includes



**FIGURE 5** (a) Distribution of the number of non-native plant species in urban centres ( $n=553$ ) across 61 countries and (b) the relationship between the number of non-native plant species and the number of non-native species occurrence records in that dataset.

Foshan. In some cases, contiguous urbanized areas span larger administrative areas and even countries. For example, Detroit, Michigan, USA, not only includes neighbouring cities in Michigan, like Dearborn, but also the Canadian city of Windsor.

2. While most recorded species in our dataset can be confirmed as established, the status of some species could not be definitively verified with our methodology. Additionally, the dataset might include non-established non-native plant species or intentionally cultivated individuals that were not fully distinguishable from naturally occurring records. As a result, the data should be interpreted cautiously, particularly when comparing non-native species richness at broader spatial scales, such as across countries, rather than at the city level. Species in our dataset with widespread occurrences across multiple urban centres are likely to be established, whereas species recorded in only one urban centre might require further scrutiny. We recommend that users consider including these singleton records in sensitivity analyses to assess the robustness of their results. Therefore, the numbers of non-native species reported here (e.g. Table 2) are in some cases higher than those reported for individual countries in recent studies (Kalusová et al., 2024; Pyšek et al., 2017).
3. The combination of these many individual datasets means that our list is subject to numerous methodological differences, from lists being built from herbarium specimens to those observed during direct sampling. Because our goal is to compile a non-native flora of urban centres, these limitations do not significantly affect our dataset.
4. The data extracted from GBIF include geographically biased and incomplete sampling, and species counts derived from these data should not be considered exhaustive despite our strict criteria listed above. For example, many urban centres in China included fewer than 100 non-native species in our database

(Figures 2 and 5a), which are likely underestimates. Analyses of richness and diversity should include rarefaction or some other way of accounting for unequal sampling as the number of non-native species increased with the number of observations (Figure 5b). Notably, many urban centres from the Global South (e.g. India; Figure 2) were absent from our database due to the paucity of available data.

## 5 | CONCLUSION

The database presented here represents a unique and valuable resource for addressing a wide range of basic and applied ecological questions, particularly those related to biological invasions. This resource can support hypothesis testing at the macro- and global scale (e.g. biotic resistance or invasion debt). It can also be used to model non-native plant species invasions, underscoring its utility not only in scientific research but also in conservation planning and practice. Lastly, it has the potential to guide more informed decision-making in biodiversity conservation, ecosystem restoration, environmental sustainability and invasive species management across diverse ecological, biogeographical and urban contexts.

## AUTHOR CONTRIBUTIONS

The Global Urban Biological Invasions Consortium, led by Marc Cadotte and including all co-authors, conceived the initial idea, which was further developed by the sGUBIK working group (Daijiang Li, Luke J. Potgieter, Myla F. J. Aronson, Irena Axmanová, Benjamin Baiser, Marta Carboni, Laura Celesti-Grapow, Sonja Knapp, Ingolf Kühn, Ana Carolina Lacerda de Matos, Zdeňka Lososová, Flavia A. Montaño-Centellas, Petr Pyšek, David M. Richardson, Lauren B. Trotta, Rafael D. Zenni and Marc W. Cadotte). Luke J. Potgieter led the data collection

effort, gathering information from the literature and contributors, who were invited to join as co-authors if they provided additional contributions to the manuscript. Daijiang Li led the data compilation from GBIF. The sGUBIK team supported data integration and standardization. Luke J. Potgieter and Daijiang Li drafted the manuscript, with all co-authors contributing to its editing and revision.

## AFFILIATIONS

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, Arizona, USA; <sup>2</sup>Department of Biological Sciences, University of Toronto-Scarborough, Toronto, Ontario, Canada; <sup>3</sup>Centre for Invasion Biology, Department of Botany & Zoology, Stellenbosch University, Stellenbosch, South Africa; <sup>4</sup>Department of Ecology, Evolution and Natural Resources, Rutgers, The State University of New Jersey, New Brunswick, New Jersey, USA; <sup>5</sup>Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic; <sup>6</sup>Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, Florida, USA; <sup>7</sup>Department of Sciences, Roma Tre University, Rome, Italy; <sup>8</sup>Department of Environmental Biology, Sapienza University of Rome, Rome, Italy; <sup>9</sup>Department of Community Ecology, Helmholtz Centre for Environmental Research-UFZ, Halle (Saale), Germany; <sup>10</sup>German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Leipzig, Germany; <sup>11</sup>Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany; <sup>12</sup>Departamento de Ecología e Conservação, Instituto de Ciências Naturais, Federal University of Lavras, Lavras, Minas Gerais, Brazil; <sup>13</sup>Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana, USA; <sup>14</sup>Department of Invasion Ecology, Institute of Botany, Czech Academy of Sciences, Průhonice, Czech Republic; <sup>15</sup>Department of Ecology, Faculty of Science, Charles University, Prague, Czech Republic; <sup>16</sup>Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa; <sup>17</sup>Environmental Research Institute, University of Waikato, Hamilton, New Zealand; <sup>18</sup>Ecology Group, Department of Biology, University of Konstanz, Konstanz, Germany; <sup>19</sup>Friesner Herbarium, Butler University, Indianapolis, Indiana, USA; <sup>20</sup>Institute of Dendrology, Polish Academy of Sciences, Kórnik, Poland; <sup>21</sup>Division of BiolInvasions, Global Change & Macroecology, Department of Botany and Biodiversity Research, University of Vienna, Vienna, Austria; <sup>22</sup>Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, Tennessee, USA; <sup>23</sup>Biodiversity and Landscape, TERRA Research Centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, Belgium; <sup>24</sup>Plants and Ecosystems, University of Antwerp, Wilrijk, Belgium; <sup>25</sup>Nürtingen-Geislingen University (HfWU), Nürtingen, Germany; <sup>26</sup>Thünen Institute, Institute of Biodiversity, Braunschweig, Germany; <sup>27</sup>Centre for Invasion Biology, Department of Mathematical Sciences, Stellenbosch University, Stellenbosch, South Africa; <sup>28</sup>African Institute for Mathematical Sciences, National Institute for National Institute for Theoretical and Computational Sciences, Cape Town, South Africa; <sup>29</sup>Centre for Biodiversity & Taxonomy, Department of Botany, University of Kashmir, Srinagar, Jammu and Kashmir, India; <sup>30</sup>Department of Ecology and Evolutionary Biology, University of Toronto Mississauga, Mississauga, Ontario, Canada; <sup>31</sup>Biodiversity, Macroecology & Biogeography, University of Göttingen, Göttingen, Germany; <sup>32</sup>Centre of Biodiversity and Sustainable Land Use (CBL), University of Göttingen, Göttingen, Germany; <sup>33</sup>Department of Ecology and Evolutionary Biology, Yale University, New Haven, Connecticut, USA; <sup>34</sup>Center for Biodiversity and Global Change, Yale University, New Haven, Connecticut, USA; <sup>35</sup>Ecology & Biodiversity, Department of Biology, Utrecht University, Utrecht, The Netherlands; <sup>36</sup>College of Forestry, Wildlife and Environment, Auburn University, Auburn, Alabama, USA; <sup>37</sup>Centro de Investigación en Biodiversidad y Conservación, Universidad Autónoma del Estado de Morelos, Cuernavaca, Mexico; <sup>38</sup>Research Center for Advanced Science and Technology, The University of Tokyo, Tokyo, Japan; <sup>39</sup>School of Natural Resources, University of Missouri, Columbia, Missouri, USA; <sup>40</sup>Ekaterinburg, Russia; <sup>41</sup>Faculty of Environmental Studies, Tokyo City University, Tokyo, Japan; <sup>42</sup>Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation, Taizhou University, Taizhou, China; <sup>43</sup>Departamento de Biología Vegetal y Ecología,

Universidad de Sevilla, Sevilla, Spain; <sup>44</sup>Estación Biológica de Doñana (EBD-CSIC), Sevilla, Spain; <sup>45</sup>Key Laboratory of Tropical Biological Resources of Ministry of Education, School of Life and Pharmaceutical Sciences, Hainan University, Haikou, China; <sup>46</sup>Department of Environmental Science, Radboud Institute for Biological and Environmental Sciences, Radboud University, Nijmegen, The Netherlands; <sup>47</sup>Institute for Housing and Environment, Darmstadt, Germany and <sup>48</sup>School of Agriculture Food and Ecosystem Science, University of Melbourne, Parkville, Victoria, Australia

## FUNDING INFORMATION

This paper is a joint effort of the working group sGUBIK kindly supported by sDiv, the Synthesis Centre of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118, 202548816). The Global Urban Biological Invasion Consortium was initially funded by the Connaught Global Challenges Award, the Office of the Vice-President International, the School of Graduate Studies, University of Toronto, and the Office of the Vice-Principal Research at the University of Toronto Scarborough. L.J.P. and D.M.R. acknowledge support from the Centre for Invasion Biology and Stellenbosch University. D.L. was supported by US NSF DEB-2213567. M.W.C. was supported by the Natural Sciences and Engineering Research Council of Canada (#386151). Z.L. and P.P. were supported by grant no. 25-15190S (Czech Science Foundation) and long-term research development project RVO 67985939 (Czech Academy of Sciences). Z.L. and I.A. were supported by EXPRO grant no. 19-28491X (Czech Science Foundation). M.C. acknowledges the support of NBFC, funded by the Italian Ministry of University and Research, PNRR, Missione 4 Componente 2, "Dalla ricerca all'impresa", Investimento 1.4, Project CN00000033. R.D.Z. acknowledges the support of CNPq-Brazil (302643/2022-2). M.v.K., A.D., and M.W. acknowledge funding of the German Research Foundation (M.v.K., A.D.: 264740629, M.W. via iDiv). P.M.K. was supported by the Natural Sciences and Engineering Research Council of Canada (Discovery Grant #RGPIN-2022-03579). M.V. by (PID2021-122690OB-I00) funded by MCIN/AEI/10.13039/501100011033/FEDER, UE. G.H. acknowledges the support of Alexander von Humboldt Foundation and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior-Brasil (Capes)-Finance code 001. M.K.D. acknowledges the support of the Institute of Dendrology, Polish Academy of Sciences. A.S.M was supported by the Mitsui & Co., Ltd. Environment Fund (funding no. R17-0062). F.E. and B.L. acknowledge funding by the Austrian Science Fundation FWF (project I5825).b.

## CONFLICT OF INTEREST STATEMENT

Marc Cadotte is the Senior Editor of Ecological Solutions and Evidence, but took no part in the peer review and decision-making processes for this paper. The authors declare no further conflicts of interest.

## PEER REVIEW

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1002/2688-8319.70020>.

## DATA AVAILABILITY STATEMENT

Data can be found on Zenodo: <https://doi.org/10.5281/zenodo.14559926> (Li, 2024).

## ORCID

Daijiang Li  <https://orcid.org/0000-0002-0925-3421>  
 Luke J. Potgieter  <https://orcid.org/0000-0001-7790-2721>  
 Benjamin Baiser  <https://orcid.org/0000-0002-3573-1183>  
 Marta Carboni  <https://orcid.org/0000-0002-9348-4758>  
 Sonja Knapp  <https://orcid.org/0000-0001-5792-4691>  
 Zdeňka Lososová  <https://orcid.org/0000-0001-9152-7462>  
 Petr Pyšek  <https://orcid.org/0000-0001-8500-442X>  
 Rafael D. Zenni  <https://orcid.org/0000-0002-4315-7986>  
 Orou G. Gaoue  <https://orcid.org/0000-0002-0946-2741>  
 Charly Géron  <https://orcid.org/0000-0001-7912-4708>  
 Anzar A. Khuroo  <https://orcid.org/0000-0002-0251-2793>  
 Frank A. La Sorte  <https://orcid.org/0000-0001-8521-2501>  
 Jonas J. Lembrechts  <https://orcid.org/0000-0002-1933-0750>  
 Bernd Lenzner  <https://orcid.org/0000-0002-2616-3479>  
 Scott MacIvor  <https://orcid.org/0000-0002-2443-8192>  
 Cristina Martínez-Garza  <https://orcid.org/0000-0002-9310-564X>  
 Akira S. Mori  <https://orcid.org/0000-0002-8422-1198>  
 Toby P. N. Tsang  <https://orcid.org/0000-0002-8176-7777>  
 Kei Uchida  <https://orcid.org/0000-0002-5955-6017>  
 Patrick Weigelt  <https://orcid.org/0000-0002-2485-3708>  
 Marten Winter  <https://orcid.org/0000-0002-9593-7300>  
 Marc W. Cadotte  <https://orcid.org/0000-0002-5816-7693>

## REFERENCES

Aronson, M. F., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A., Warren, P. S., Williams, N. S., Cilliers, S., Clarkson, B., Dobbs, C., Dolan, R., Hedblom, M., Klotz, S., Kooijmans, J. L., Kühn, I., Macgregor-Fors, I., McDonnell, M., Mörtingberg, U., ... Winter, M. (2014). A global analysis of the impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers. *Proceedings of the Royal Society B: Biological Sciences*, 281(1780), 20133330. <https://doi.org/10.1098/rspb.2013.3330>

Blackburn, T. M., Pyšek, P., Bacher, S., Carlton, J. T., Duncan, R. P., Jarošík, V., Wilson, J. R. U., & Richardson, D. M. (2011). A proposed unified framework for biological invasions. *Trends in Ecology & Evolution*, 26, 333–339. <https://doi.org/10.1016/j.tree.2011.03.023>

Brown, M. J. M., Walker, B. E., Black, N., Govaerts, R., Ondo, I., Turner, R., & Nic Lughadha, E. (2023). rWVCVP: A companion R package to the World Checklist of Vascular Plants. *New Phytologist*, 240, 1355–1365. <https://doi.org/10.1111/nph.18919>

Cadotte, M. W., Yasui, S. L. E., Livingstone, S., & MacIvor, J. S. (2017). Are urban systems beneficial, detrimental, or indifferent for biological invasion? *Biological Invasions*, 19, 3489–3503. <https://doi.org/10.1007/s10530-017-1586-y>

Chao, A., Gotelli, N., Hsieh, T., Sander, E., Ma, K., Colwell, R., & Ellison, A. (2014). Rarefaction and extrapolation with Hill numbers: A framework for sampling and estimation in species diversity studies. *Ecological Monographs*, 84, 45–67. <https://doi.org/10.1890/13-0133.1>

Gaertner, M., Larson, B. M., Irlich, U. M., Holmes, P. M., Stafford, L., van Wilgen, B. W., & Richardson, D. M. (2016). Managing invasive species in cities: A framework from Cape Town, South Africa. *Landscape and Urban Planning*, 151, 1–9. <https://doi.org/10.1016/j.landurbplan.2016.03.010>

Gaertner, M., Wilson, J. R. U., Cadotte, M. W., MacIvor, J. S., Zenni, R. D., & Richardson, D. M. (2017). Non-native species in urban environments: Patterns, processes, impacts and challenges. *Biological Invasions*, 19, 3461–3469. <https://doi.org/10.1007/s10530-017-1598-7>

Gallardo, B., Clavero, M., Sánchez, M. I., & Vilà, M. (2016). Global ecological impacts of invasive species in aquatic ecosystems. *Global Change Biology*, 22, 151–163. <https://doi.org/10.1111/gcb.13004>

Govaerts, R. (2024). The World Checklist of Vascular Plants (WCVP). Royal Botanic Gardens, Kew. Checklist dataset. <https://doi.org/10.15468/6h8ucr>

Grenié, M., Berti, E., Carvajal-Quintero, J., Dädlow, G. M. L., Sagouis, A., & Winter, M. (2022). Harmonizing taxon names in biodiversity data: A review of tools, databases and best practices. *Methods in Ecology and Evolution*, 14, 1–14. <https://doi.org/10.1111/2041-210X.13802>

Heringer, G., Fernandez, R. D., Bang, A., Cordonnier, M., Novoa, A., Lenzner, B., Capinha, C., Renault, D., Roiz, D., Moodley, D., Tricarico, E., Holenstein, K., Kourantidou, M., Kirichenko, N. I., Adelino, J. R. P., Dimarco, R. D., Bodey, T. W., Watari, Y., & Courchamp, F. (2024). Economic costs of invasive non-native species in urban areas: An underexplored financial drain. *Science of the Total Environment*, 917, 170336. <https://doi.org/10.1016/j.scitotenv.2024.170336>

Hsieh, T., Ma, K., & Chao, A. (2024). iNEXT: Interpolation and Extrapolation for Species Diversity. R package version 3.0.1. [http://chao.stat.nthu.edu.tw/wordpress/software\\_download/](http://chao.stat.nthu.edu.tw/wordpress/software_download/)

Hsieh, T. C., Ma, K., & Chao, A. (2016). iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution*, 7(12), 1451–1456.

Kalusová, V., Čeplová, N., Danihelka, J., Večeřa, M., Pyšek, P., Albert, A., Anastasiu, P., Biurrun, I., Boch, S., Cottaz, C., Essl, F., Kuzemko, A., Maslo, S., Mifsud, S., Protopopova, V. V., Shevera, M., Širbu, C., Svenning, J.-C., Welk, E., & Axmanová, I. (2024). Alien plants of Europe: An overview of national and regional inventories. *Preslia*, 96, 149–182. <https://doi.org/10.23855/preslia.2024.149>

Li, D. (2024). daijiang/GUBIC\_data: Release for Zenodo (v1.0.1). Zenodo. <https://doi.org/10.5281/zenodo.14559926>

Milanović, M., Knapp, S., Pyšek, P., & Kühn, I. (2020). Linking traits of invasive plants with ecosystem services and disservices. *Ecosystem Services*, 42, 101072.

Pesaresi, M., Florczyk, A., Schiavina, M., Melchiorri, M., & Maffenini, L. (2019). GHS settlement grid, updated and refined REGIO model 2014 in application to GHS-BUILT R2018A and GHS-POP R2019A, multi-temporal (1975–1990–2000–2015), R2019A. European Commission, Joint Research Centre (JRC), 10. [Dataset] PID <http://data.europa.eu/89h/42e8be89-54ff-4646-be7b-bf9e64da5218>

Potgieter, L. J., & Cadotte, M. W. (2020). The application of selected invasion frameworks to urban ecosystems. *NeoBiota*, 62, 365–386. <https://doi.org/10.3897/neobiota.62.50661>

Potgieter, L. J., Gaertner, M., Kueffer, C., Larson, B. M. H., Livingston, S., O'Farrell, P., & Richardson, D. M. (2017). Alien plants as mediators of ecosystem services and disservices in urban systems: A global review. *Biological Invasions*, 19, 3571–3588. <https://doi.org/10.1007/s10530-017-1589-8>

Potgieter, L. J., Li, D., Aronson, M. F. J., Baiser, B., Kühn, I., Aronson, M. F. J., Carboni, M., Celesti-Grapow, L., de Matos, A. C. L., Lososová, Z., Montaño-Centellas, F. A., Pyšek, P., Richardson, D. M., Tsang, T. P. N., Zenni, R. D., & Cadotte, M. W. (2024). Cities shape the diversity and spread of nonnative species. *Annual Review of Ecology, Evolution, and Systematics*, 55, 157–180. <https://doi.org/10.1146/annurev-ecolsys-102722-012749>

Potgieter, L. J., Shrestha, N., & Cadotte, M. W. (2022). Prioritizing sites for terrestrial invasive alien plant management in urban ecosystems.

*Ecological Solutions and Evidence*, 3, e12160. <https://doi.org/10.1002/2688-8319.12160>

Pyšek, P., Pergl, J., Essl, F., Lenzner, B., Dawson, W., Kreft, H., Weigelt, P., Winter, M., Kartesz, J., Nishino, M., Antonova, L. A., Barcelona, J. F., Cabezas, F. J., Cárdenas, D., Cárdenas-Toro, J., Castaño, N., Chacón, E., Chatelain, C., Dullinger, S., ... van Kleunen, M. (2017). Naturalized alien flora of the world. *Preslia*. Czech Botanical Society. <https://doi.org/10.23855/preslia.2017.203>

Richardson, D. M., Pyšek, P., Rejmanek, M., Barbour, M. G., Panetta, F. D., & West, C. J. (2000). Naturalization and invasion of alien plants: Concepts and definitions. *Diversity and Distributions*, 6, 93–107. <https://doi.org/10.1046/j.1472-4642.2000.00083.x>

van Kleunen, M., Pyšek, P., Dawson, W., Essl, F., Kreft, H., Pergl, J., Weigelt, P., Stein, A., Dullinger, S., König, C., Lenzner, B., Maurel, N., Moser, D., Seebens, H., Kartesz, J., Nishino, M., Alek-Sanyan, A., Ansong, M., Antonova, L. A., ... Winter, M. (2019). The Global Naturalized Alien Flora (GloNAF) database. *Ecology*, 100, 1–2. <https://doi.org/10.1002/ecy.2542>

Vaz, A. S., Castro-Díez, P., Godoy, O., Alonso, Á., Vilà, M., Saldaña, A., Marchante, H., Bayón, Á., Silva, J. S., Vicente, J. R., & Honrado, J. P. (2018). An indicator-based approach to analyse the effects of non-native tree species on multiple cultural ecosystem services. *Ecological Indicators*, 85, 48–56. <https://doi.org/10.1016/j.ecolind.2017.10.009>

Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq, H., Herdean, A., Ariza, M., Scharn, R., Svantesson, S., Wengström, N., Zizka, V., & Antonelli, A. (2019). CoordinateCleaner:

Standardized cleaning of occurrence records from biological collection databases. *Methods in Ecology and Evolution*, 10, 744–751. R package version 3.0.1. <https://doi.org/10.1111/2041-210X.13152>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**Table S1.** Global Biodiversity Information Facility (GBIF) data used to derive the Global Urban Biological Invasions Compendium database.

**How to cite this article:** Li, D., Potgieter, L. J., Aronson, M. F. J., Axmanová, I., Baiser, B., Carboni, M., Celesti-Grapow, L., Knapp, S., Kühn, I., Lacerda de Matos, A. C., Lososová, Z., Montaño-Centellas, F. A., Pyšek, P., Richardson, D. M., Trotta, L. B., Zenni, R. D., Cilliers, S. S., Clarkson, B. D., Davis, A. J. S., ... Cadotte, M. W. (2025). GUBIC: The global urban biological invasions compendium for plants. *Ecological Solutions and Evidence*, 6, e70020. <https://doi.org/10.1002/2688-8319.70020>