

Absence of bound states for quantum walks and CMV matrices via reflections

Christopher Cedzich and Jake Fillman

Abstract. We give a criterion based on reflection symmetries in the spirit of Jitomirskaya–Simon to show absence of point spectrum for (split-step) quantum walks and Cantero–Moral–Velázquez (CMV) matrices. To accomplish this, we use some ideas from a recent paper by the authors and collaborators to implement suitable reflection symmetries for such operators. We give several applications. For instance, we deduce arithmetic delocalization in the phase for the unitary almost-Mathieu operator and singular continuous spectrum for generic CMV matrices generated by the Thue–Morse subshift.

1. Introduction

One of the fundamental issues in the theory of ergodic operators is to establish the presence of Anderson localization (i.e., pure point spectrum with exponentially decaying eigenfunctions) whenever one suspects it to be present, and furthermore, one often suspects Anderson localization occurs for operator families in the regime of positive Lyapunov exponents. Indeed, positivity of the Lyapunov exponent is often the first step on a journey towards localization and provides a key input: namely, (via Ruelle’s theorem [43]) positivity of the Lyapunov exponent ensures that for any energy for which the pointwise Lyapunov exponent exists, one has an exponential dichotomy of solutions and therefore generalized eigenvalues must decay exponentially. However, the phenomenon in question is in general very delicate, since one must be careful of the exceptional set on which the pointwise Lyapunov exponent fails to exist. This can be overcome, and hence the desired localization can be proved, using a variety of techniques, such as multi-scale analysis, fractional moment analysis, large deviations, repulsion of singular clusters, elimination of double resonances, and others; for references and further reading, we direct the reader to [4, 16, 17, 35].

Mathematics Subject Classification 2020: 47B93 (primary); 47A10, 47B36, 58J51, 81Q10 (secondary).

Keywords: unitary almost Mathieu operator, singular continuous spectrum, resonances, quantum walks, CMV matrices, delocalization, quasi-periodicity.

These approaches have been pursued and implemented in a number of works concerning quasi-periodic [12, 27, 46, 47], almost-periodic [11], skew-shift [14, 37, 39] and random [2, 8, 28, 34, 48] CMV matrices.

One is also naturally interested in understanding structures that preclude localization in the regime of positive exponents (either globally, or locally in energy). The present paper accomplishes precisely this task for CMV operators in the presence of suitable reflection symmetries by synthesizing ideas of [11] to exploit gauge-invariance in order to implement the approach of Jitomirskaya and Simon [31]. The main results of this work in particular show that several localization results recently obtained for the unitary almost-Mathieu operator [12, 47] and mosaic model [11] are sharp in the sense that localization for almost every phase cannot be strengthened to localization for all phases. Furthermore, the method gives generic absence of eigenvalues for CMV matrices and quantum walks generated by subshifts containing many palindromes. In particular, we obtain the first results on singular continuous spectrum for CMV matrices generated by the Thue–Morse subshift, to our knowledge.

Let us briefly recall some of the background literature. In the setting of self-adjoint operators Jitomirskaya–Simon gave a criterion for Schrödinger operators to have purely continuous spectrum [31]. This was adapted to palindromic subshift operators by Hof, Knill, and Simon [29] and developed further in works such as Koslover [36] and Jitomirskaya and Liu [30]. As was noted in [11], arguments based on reflections to be more delicate (even outright impossible) for standard CMV operators, due to a lack of appropriate reflection symmetries. However, the authors and collaborators observed in [11] that one can overcome this by passing to the setting of so-called *generalized* CMV matrices via complexification and relating the complexified CMV matrix to the initial operator via a diagonal gauge transformation that *preserves* the Verblunsky coefficients. Let us also mention a different obstruction to localization, known as the Gordon criterion [25], is based on the presence of suitable repetitions, rather than reflections. This is significantly more straightforward and has already been worked out for CMV matrices in [21]; see also [38] for an arithmetic strengthening and [40, 41] for earlier results. Recently, Avila and Damanik have announced another approach to delocalization that applies to ergodic self-adjoint operators in arbitrary dimensions [6].

CMV matrices have physical significance, as they are (equivalent to) one-dimensional split-step quantum walks, which was first observed in [9] and elaborated upon in [11]. Split-step quantum walks are the simplest type of the rich class of quantum walks which have gained popularity in recent years as models of discrete time quantum dynamics. In this sense, quantum walks play a role similar to that of Jacobi matrices for continuous time evolutions generated by general self-adjoint operators, and one is therefore naturally interested in their dynamical [1–3, 5, 26, 33, 34] and spectral [7, 10, 13, 32, 42] properties.

We formulate our main results precisely in Section 2 and prove them in Section 3.

2. Results

2.1. General Results

We begin by recalling the notion of generalized extended CMV (GECMV) matrices from [11]. Let $\mathbb{S}^3 = \{(\alpha, \rho) \in \mathbb{C}^2 : |\alpha|^2 + |\rho|^2 = 1\}$ and consider a sequence $\{(\alpha_n, \rho_n)\} \in (\mathbb{S}^3)^\mathbb{Z}$. In the standard basis of $\ell^2(\mathbb{Z})$, the corresponding GECMV matrix takes the form

$$\mathcal{E} = \mathcal{E}_{\alpha, \rho} = \begin{bmatrix} \ddots & & & & & & & \\ & \ddots & & & & & & \\ & & \boxed{\begin{matrix} \overline{\alpha_0 \rho_{-1}} & -\overline{\alpha_0} \alpha_{-1} \\ \overline{\rho_0 \rho_{-1}} & -\overline{\rho_0} \alpha_{-1} \end{matrix}} & \overline{\alpha_1} \rho_0 & \rho_1 \rho_0 & & & \\ & & & -\overline{\alpha_1} \alpha_0 & -\rho_1 \alpha_0 & & & \\ & & & \overline{\alpha_2 \rho_1} & -\overline{\alpha_2} \alpha_1 & \overline{\alpha_3} \rho_2 & \rho_3 \rho_2 & \\ & & & \overline{\rho_2 \rho_1} & -\overline{\rho_2} \alpha_1 & -\overline{\alpha_3} \alpha_2 & -\rho_3 \alpha_2 & \\ & & & & \ddots & \ddots & \ddots & \ddots \end{bmatrix}, \quad (2.1)$$

where we boxed the $(0, 0)$ matrix element. A *standard* CMV matrix has the form (2.1) with $\rho_n \geq 0$ for all n ; equivalently, $\rho_n = \sqrt{1 - |\alpha_n|^2}$. The standard CMV matrix has an important role in mathematical physics, spectral theory, and orthogonal polynomials; for a textbook discussion that includes many important cases (e.g. periodic, random, and almost-periodic), see [44, 45]. For any sequence $\alpha = \{\alpha_n\} \in \mathbb{D}^\mathbb{Z}$, the associated standard CMV matrix and its cousin with complexified ρ terms will be denoted by

$$\mathcal{E}_\alpha := \mathcal{E}_{\alpha, \sqrt{1-|\alpha|^2}}, \quad \mathcal{E}_\alpha^\mathbb{C} := \mathcal{E}_{\alpha, i\sqrt{1-|\alpha|^2}}.$$

By [11], $\mathcal{E}_\alpha$ and $\mathcal{E}_\alpha^\mathbb{C}$ are equivalent via a gauge transform.

Given $u \in \mathbb{C}^\mathbb{Z}$, and $\zeta \in \frac{1}{2}\mathbb{Z}$, the *reflection* of u through the center ζ is given by

$$(\mathcal{R}_\zeta u)_n = u_{2\zeta-n}.$$

In order to get useful estimates, we will need Verblunsky pairs with exponentially good approximate reflection symmetries locally on intervals that are exponentially long (with respect to the center of reflection). To generate such symmetries for a suitably rich set of phases in the ergodic setting below, it is useful to begin with a sequence that obeys exact symmetries globally. The following notions make this precise.

We say that $\alpha = \{\alpha_n\} \in \mathbb{D}^\mathbb{Z}$ is *reflection symmetric* about the center ζ if

$$(\mathcal{R}_\zeta \alpha)_n \equiv \overline{\alpha_n}. \quad (2.2)$$

It is no loss of generality to assume that the center of reflection is 0, which we will do throughout the paper: we simply say that α is *reflection symmetric* if it is reflection symmetric about $\zeta = 0$.

The following definition captures precisely the approximate local reflection symmetries that we will need.

Definition 2.1. Given $B > 0$ and $\zeta \in \frac{1}{2}\mathbb{Z}$, we say that $\alpha \in \mathbb{D}^{\mathbb{Z}}$ is (B, ζ) -reflective if

$$|(\mathcal{R}_{\zeta}\alpha)_n - \overline{\alpha_n}| < e^{-B|\zeta|} \quad \text{for all } n \in \mathbb{Z} \text{ such that } |\zeta - n| \leq e^{B|\zeta|}.$$

We say that α is B -reflective if it is (B, ζ) -reflective for infinitely many $\zeta \in \frac{1}{2}\mathbb{Z}$. Clearly, α is (B, ζ) -reflective for every $B > 0$ whenever it is reflection symmetric about ζ , but not conversely.

Throughout, we assume that the Verblunsky coefficients are bounded away from 1 in modulus:

$$\|\alpha\|_{\infty} = r_0 < 1, \tag{2.3}$$

which in particular implies

$$|\rho_n| \geq (1 - r_0^2)^{1/2} =: c_0 > 0$$

uniformly in n .

The main result is the following.

Theorem 2.2. Assume α satisfies (2.3). There is a constant B_0 such that the following is true: if α is B -reflective with $B > B_0$, then $\mathcal{E} = \mathcal{E}_{\alpha, \rho}$ has empty point spectrum for any admissible¹ choice of ρ ; in particular, the standard CMV matrix $\mathcal{E}_{\alpha}$ has empty point spectrum.

Remark 2.3. A few comments are in order.

(a) A straightforward attempt to imitate the proof of [31] for standard CMV matrices leads to roadblocks arising from the lack of relevant symmetries. For instance, it is clear that an estimate of the form (3.7) is needed to obtain useful estimates on the quantities in, for example, (3.11) and (3.19). Thus, in addition to the identity (2.2) for the α 's, one also must have a symmetry of the type (3.7) for the ρ 's, and such a symmetry cannot hold for standard CMV matrices, so one absolutely must complexify to reveal the necessary symmetry.

(b) In fact, as the reader will see, even with the new ideas using gauge transforms to relate to the complexified ρ case, the computations are somewhat more laborious in the present setting. For instance, one may compare the proof of Lemma 3.4 to the corresponding computation in [31, Step 1]. Similarly, the absence of symplectic symmetry (even after complexification) makes the comparison of “reflected” and inverted transfer matrices quite delicate.

¹That is, any choice of ρ satisfying $|\alpha_n|^2 + |\rho_n|^2 \equiv 1$.

(c) If α satisfies (2.3), then the hypothesis of Theorem 2.2 is fulfilled whenever α is even and real-valued.

(d) The largeness condition on B is concrete. To ensure that $z \in \partial\mathbb{D}$ is not an eigenvalue of $\mathcal{E}_\alpha$, one simply needs B to be large enough that

$$\lim_{n \rightarrow \infty} e^{-Bn} \|N_{\ell+n-1,z} \cdots N_{\ell+1,z} N_{\ell,z}\| = 0$$

uniformly in $\ell \in \mathbb{Z}$, where $N_{n,z}$ denote the transfer matrices (see Lemma 3.1). Thus, for ergodic models over strictly ergodic base dynamics (for which one may apply Furman's uniform subadditive ergodic theorem [24]), one can phrase the relevant bound in terms of the Lyapunov exponent *locally* in the spectral parameter and deduce *arithmetic* delocalization; compare Theorem 2.5 and Corollary 2.6.

(e) When applying the results to general quantum walks that are not given by GEVM matrices, some care is needed. For instance, the electric walks in [13, 14] have symmetric coins, but application of the gauge transform leads to a CMV matrix that does not satisfy the necessary symmetries, so our results do not apply in that setting.

(f) It follows readily from the result of Damanik and Killip [19] that for any almost-periodic sequence α that is not limit-periodic, the set of reflective $\omega \in \text{hull}(\alpha)$ has zero Haar measure.

Theorem 2.2 can be applied to almost-periodic CMV matrices. Recall that $\alpha \in \mathbb{C}^{\mathbb{Z}}$ is called *almost-periodic* if its orbit

$$\text{orb}(\alpha) := \{\alpha(\cdot - n) : n \in \mathbb{Z}\}$$

is relatively compact in $\ell^\infty(\mathbb{Z})$. In that case, the closure (which is compact) is called the *hull*:

$$\text{hull}(\alpha) := \overline{\text{orb}(\alpha)}^{\|\cdot\|_\infty}.$$

Corollary 2.4. *Assume $\alpha: \mathbb{Z} \rightarrow \mathbb{D}$ satisfies (2.3). If α is reflection-symmetric and almost-periodic, then there is a dense G_δ subset $\Omega_0 \subseteq \text{hull}(\alpha)$ such that the extended CMV matrix $\mathcal{E}_\omega$ has empty point spectrum for all $\omega \in \Omega_0$.*

Note that in the setting of Corollary 2.4, $\omega \in \text{hull}(\alpha)$, so the Verblunsky coefficient sequence associated with $\mathcal{E}_\omega$ is simply the sequence ω itself, that is, $\alpha_\omega(n) = \omega(n)$. In addition, as alluded to in Remark 2.3, one can make Corollary 2.4 quantitative and can exclude eigenvalues locally in the spectral parameter. In order to do that, we need the Lyapunov exponent $L(z)$ (which will be defined in (3.22)).

Theorem 2.5. *Assume $\alpha: \mathbb{Z} \rightarrow \mathbb{D}$ satisfies (2.3). If α is almost-periodic and B -reflective, then, for any admissible ρ , $\mathcal{E}_{\alpha,\rho}$ has no eigenvalues in the region $\{z \in \partial\mathbb{D} : 2L(z) < B\}$.*

This allows us to establish a corollary giving arithmetic delocalization in the spirit of [30, Theorem 4.2]. To formulate the result, recall that a topological group is called *monothetic* if it contains a dense cyclic subgroup (e.g., this class of groups includes $\mathbb{Z}$, $\mathbb{Z}_p$, $\mathbb{T}^d$, $\mathbb{Z}_p \times \mathbb{T}^d$, and certain procyclic groups such as the p -adic integers). Given a metrizable compact monothetic Ω , a generator β of a dense cyclic subgroup, and $f \in C(\Omega, \mathbb{C})$, the sequence $(f(n\beta + \omega))_{n \in \mathbb{Z}}$ is almost-periodic² for any choice of $\omega \in \Omega$. This is a standard fact whose proof can be found in various places, e.g., [15, Section 3].

For a metrizable compact monothetic group Ω with translation-invariant metric dist , we denote

$$\|\omega\| = \|\omega\|_{\Omega} := \text{dist}(\omega, 0), \quad \omega \in \Omega.$$

For $\beta, \omega \in \Omega$, define

$$\delta(\beta, \omega) = \limsup_{|n| \rightarrow \infty} \left[-\frac{\log \|\omega + n\beta\|}{|n|} \right]. \quad (2.4)$$

Corollary 2.6. *Suppose Ω is a compact metrizable monothetic group, that $\beta, \omega \in \Omega$, and that the subgroup generated by β is dense in Ω . If $f: \Omega \rightarrow \mathbb{D}$ is Lipschitz continuous and satisfies $f(-\omega) \equiv \overline{f(\omega)}$, then for each $\omega \in \Omega$, the CMV matrix $\mathcal{E}_{\omega}$ with coefficients*

$$\alpha_{\omega}(n) = f(n\beta + \omega)$$

has no eigenvalues in the region $\{z \in \partial\mathbb{D} : L(z) < \delta(\beta, \omega)\}$.

Let us briefly point out that [30, Theorem 4.2] is formulated for $\Omega = \mathbb{T}$, but the proof they give carries through with cosmetic changes to the more general setting described in Corollary 2.6. Moreover, we need this slight generalization since the UAMO (viewed as a dynamically defined CMV matrix) is most naturally viewed as having base dynamics on $\mathbb{T} \times \mathbb{Z}_2$, as discussed below in (3.23).

2.2. Examples and applications

The criterion in the main results can be applied to models of interest. For instance, localization questions for the unitary almost-Mathieu operator (UAMO) introduced in [12, 23] have been studied in [12, 47]. In light of [11, 12], the UAMO family is gauge-equivalent to the family of GECMV matrices given by

$$\alpha_{2n-1} = \lambda_2 \cos(2\pi(n\Phi + \theta)), \quad \alpha_{2n} = \lambda_1',$$

²That is, its translates are precompact in $\ell^{\infty}(\mathbb{Z})$.

where we abbreviate $\lambda'_1 := (1 - \lambda_1^2)^{1/2}$. Let us denote the corresponding operators by $\mathfrak{E}_{\lambda_1, \lambda_2, \Phi, \theta}^{\text{UAMO}}$. Notice that we use the cosine rather than the sine (which is the definition given in [12]) since the cosine function is even and we want to emphasize reflection symmetries.

Corollary 2.7. *For any $\lambda_1, \lambda_2 \in (0, 1)$ and irrational Φ , $\mathfrak{E}_{\lambda_1, \lambda_2, \Phi, \theta}^{\text{UAMO}}$ has empty point spectrum whenever*

$$\frac{1}{2}\delta(\Phi, \theta) > \log \left[\frac{\lambda_2(1 + \lambda'_1)}{\lambda_1(1 + \lambda'_2)} \right]. \quad (2.5)$$

In particular, $\mathfrak{E}_{\lambda_1, \lambda_2, \Phi, \theta}^{\text{UAMO}}$ has empty point spectrum for generic $\theta \in \mathbb{T}$ and has purely singular continuous spectrum whenever (2.5) and $\lambda_1 < \lambda_2$ hold.

The same result can be applied to other almost-periodic CMV matrices, such as the mosaic model considered in [11], which can be given by an additional choice of parameter $s \in \mathbb{Z}_+$ and defining

$$\alpha_{2n} = \lambda'_1, \quad \alpha_{2kn-1} = \begin{cases} \lambda_2 \cos(2\pi(n\Phi + \theta)) & k \in s\mathbb{Z} \\ 0 & \text{otherwise.} \end{cases}$$

We write the resulting operator as $\mathfrak{E}_{s, \lambda_1, \lambda_2, \Phi, \theta}^{\text{mosaic}}$.

Corollary 2.8. *For any $s \in \mathbb{Z}$, $\lambda_1, \lambda_2 \in (0, 1)$, and irrational Φ , $\mathfrak{E}_{s, \lambda_1, \lambda_2, \Phi, \theta}^{\text{mosaic}}$ has empty point spectrum for generic $\theta \in \mathbb{T}$.*

Finally, let us mention that the criterion in Theorem 2.2 may be applied to ergodic CMV matrices taking finitely many values, which occur naturally in the context of subshifts. Given a finite set $\mathcal{A} \subseteq \mathbb{D}$ (the *alphabet*), a *subshift* is any compact³ set $\Omega \subseteq \mathcal{A}^{\mathbb{Z}}$ that is invariant under the action of the shift map $\text{sh}: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$ given by

$$[\text{sh } \omega](n) = \omega(n + 1).$$

It is well known that a subshift (Ω, sh) is minimal (in the sense that the shift-orbit of every $\omega \in \Omega$ is dense in Ω) if and only if there is a set

$$\mathcal{L} = \mathcal{L}(\Omega) \subseteq \bigcup_{n \in \mathbb{N}} \mathcal{A}^n$$

with the property that $\mathcal{L}$ is precisely the set of finite strings occurring in every $\omega \in \Omega$, that is,

$$\{\omega_j \omega_{j+1} \cdots \omega_{j+m-1} : j \in \mathbb{Z}, m \geq 1\} = \mathcal{L}$$

for every $\omega \in \Omega$. The set $\mathcal{L}$ is called the *language* of the subshift.

³Naturally, $\mathcal{A}$ is given the discrete topology and $\mathcal{A}^{\mathbb{Z}}$ is in turn given the product topology.

Corollary 2.9. *If (Ω, sh) is a minimal subshift whose language contains arbitrarily long palindromes, then $\mathcal{E}_\omega$ has empty point spectrum for generic $\omega \in \Omega$.*

In particular, let us mention that Corollary 2.9 can be applied to the subshift generated by the Thue–Morse substitution $S_{\text{TM}}: a \mapsto ab, b \mapsto ba$ over the alphabet $\mathcal{A} = \{a, b\}$. The corresponding CMV matrices and quantum walks were studied from a dynamical perspective [18, 22] and were known to have purely singular spectrum by Kotani theory [45] and in fact Cantor spectrum of zero Lebesgue measure [20], but singular continuous spectrum could not be established until now. Since this is an important example in the theory of aperiodic order, we formulate this explicitly as a corollary.

Corollary 2.10. *If $\Omega \subseteq \{a, b\}^{\mathbb{Z}}$ is the subshift generated by the Thue–Morse substitution, then for generic $\omega \in \Omega$, $\mathcal{E}_\omega$ has purely singular continuous spectrum.*

To the best of our knowledge, this gives the first result about singular continuity for CMV matrices or quantum walks generated by the Thue–Morse substitution. For Schrödinger operator versions of these results, see [29].

3. Proofs of the results

One studies $\mathcal{E}$ and its spectral properties via the generalized eigenvalue equation

$$\mathcal{E}u = zu, \quad z \in \mathbb{C}^* \equiv \mathbb{C} \setminus \{0\}, u \in \mathbb{C}^{\mathbb{Z}}, \quad (3.1)$$

where we emphasize that no assumption is made about u beyond being a complex-valued sequence.

For later, use note that $\mathcal{E}$ acts on coordinates via

$$\begin{aligned} [\mathcal{E}_{\alpha, \rho} u](2n) &= \overline{\alpha_{2n}} [\overline{\rho_{2n-1}} u(2n-1) - \alpha_{2n-1} u(2n)] \\ &\quad + \rho_{2n} [\overline{\alpha_{2n+1}} u(2n+1) + \rho_{2n+1} u(2n+2)] \end{aligned} \quad (3.2)$$

$$\begin{aligned} [\mathcal{E}_{\alpha, \rho} u](2n+1) &= \overline{\rho_{2n}} [\overline{\rho_{2n-1}} u(2n-1) - \alpha_{2n-1} u(2n)] \\ &\quad - \alpha_{2n} [\overline{\alpha_{2n+1}} u(2n+1) + \rho_{2n+1} u(2n+2)]. \end{aligned} \quad (3.3)$$

Solutions to $\mathcal{E}u = zu$ with $z \in \mathbb{C}^*$ and $u \in \mathbb{C}^{\mathbb{Z}}$ satisfy the iterative relation

$$\begin{bmatrix} u(2n+1) \\ u(2n) \end{bmatrix} = M_{n,z} \begin{bmatrix} u(2n-1) \\ u(2n-2) \end{bmatrix}, \quad n \in \mathbb{Z},$$

where the *transfer matrices* $M_{n,z}$ are given by

$$M_{n,z} = \mathfrak{X}(\alpha_{2n}, \alpha_{2n-1}, \alpha_{2n-2}, \rho_{2n}, \rho_{2n-1}, \rho_{2n-2}, z) \\ := \frac{1}{\rho_{2n}\rho_{2n-1}} \begin{bmatrix} z^{-1} + \alpha_{2n}\overline{\alpha_{2n-1}} + \alpha_{2n-1}\overline{\alpha_{2n-2}} + \alpha_{2n}\overline{\alpha_{2n-2}}z & \\ & -\rho_{2n}\overline{\alpha_{2n-1}} - \rho_{2n}\overline{\alpha_{2n-2}}z \\ & & -\overline{\rho_{2n-2}}\alpha_{2n-1} - \overline{\rho_{2n-2}}\alpha_{2n}z \\ & & & \rho_{2n}\overline{\rho_{2n-2}}z \end{bmatrix},$$

for $n \in \mathbb{Z}$ and $z \in \mathbb{C}^*$. This follows from direct calculations using (3.2) and (3.3), which are carried out in detail in [12, Section 4]. Note that the determinant of $M_{n,z}$ is given by

$$\det M_{n,z} = \frac{\overline{\rho_{2n-1}}\overline{\rho_{2n-2}}}{\rho_{2n}\rho_{2n-1}}.$$

Similar calculations give us the following iterative relations for starting points with reversed parity. For the reader's convenience, we sketch the argument below.

Lemma 3.1. *Suppose that $\mathcal{E}u = zu$ for $z \in \mathbb{C} \setminus \{0\}$. Then*

$$\begin{bmatrix} u(2n+2) \\ u(2n+1) \end{bmatrix} = N_{n,z} \begin{bmatrix} u(2n) \\ u(2n-1) \end{bmatrix}, \quad n \in \mathbb{Z}, \quad (3.4)$$

where

$$N_{n,z} = \mathfrak{X}(\overline{\alpha_{2n+1}}, \overline{\alpha_{2n}}, \overline{\alpha_{2n-1}}, \rho_{2n+1}, \rho_{2n}, \rho_{2n-1}, z^{-1}) \\ = \frac{1}{\rho_{2n+1}\rho_{2n}} \begin{bmatrix} z + \overline{\alpha_{2n}}\alpha_{2n-1} + \overline{\alpha_{2n+1}}\alpha_{2n-1}z^{-1} + \overline{\alpha_{2n+1}}\alpha_{2n} & \\ & -\alpha_{2n}\rho_{2n+1} - \alpha_{2n-1}\rho_{2n+1}z^{-1} \\ & & -\overline{\alpha_{2n}}\overline{\rho_{2n-1}} - \overline{\alpha_{2n+1}}\overline{\rho_{2n-1}}z^{-1} \\ & & & \overline{\rho_{2n-1}}\rho_{2n+1}z^{-1} \end{bmatrix}. \quad (3.5)$$

Moreover,

$$\det N_{n,z} = \frac{\overline{\rho_{2n}}\overline{\rho_{2n-1}}}{\rho_{2n+1}\rho_{2n}}.$$

Proof. From α_{2n} (3.2) + ρ_{2n} (3.3) and $\overline{\rho_{2n}}$ (3.2) - $\overline{\alpha_{2n}}$ (3.3) we get

$$\alpha_{2n}zu(2n) + \rho_{2n}zu(2n+1) = \overline{\rho_{2n-1}}u(2n-1) - \alpha_{2n-1}u(2n) \quad (3.6)$$

and

$$\overline{\rho_{2n}}zu(2n) - \overline{\alpha_{2n}}zu(2n+1) = \overline{\alpha_{2n+1}}u(2n+1) + \rho_{2n+1}u(2n+2),$$

respectively, where we used that $|\alpha|^2 + |\rho|^2 = 1$. Solving the first equation for $u(2n+1)$ and inserting it into the second one yields the relations in (3.4). The determinant is determined by a straightforward calculation. ■

Lemma 3.1 immediately implies that for two solutions u, v of the generalized eigenvalue equation (3.1), we have

$$\begin{bmatrix} u(2n+2) & v(2n+2) \\ u(2n+1) & v(2n+1) \end{bmatrix} = N_{n,z} N_{n-1,z} \cdots N_{0,z} \begin{bmatrix} u(0) & v(0) \\ u(-1) & v(-1) \end{bmatrix}.$$

Taking determinants on both sides and rearranging terms gives

$$\begin{aligned} W(u, v)(n) &:= \rho_{2n+1} \frac{\rho_{2n} \rho_{2n-1} \cdots \rho_1 \rho_0}{\overline{\rho_{2n}} \overline{\rho_{2n-1}} \cdots \overline{\rho_1} \overline{\rho_0}} (u(2n+2)v(2n+1) \\ &\quad - u(2n+1)v(2n+2)) \\ &= \overline{\rho_{-1}} (u(0)v(-1) - u(-1)v(0)). \end{aligned}$$

We shall call $W(u, v)$ the *Wronskian* of u and v .

Remark 3.2. The following observation will be helpful in the proof: for

$$\rho_n^{\mathbb{C}} := i \sqrt{1 - |\alpha_n|^2},$$

then

$$\begin{aligned} |\rho_n^{\mathbb{C}} - (-\overline{\rho_m^{\mathbb{C}}})| &= |(1 - |\alpha_n|^2)^{1/2} - (1 - |\alpha_m|^2)^{1/2}| \\ &\leq \frac{2r}{\sqrt{1 - r^2}} ||\alpha_n| - |\alpha_m||, \end{aligned}$$

where $r = \max\{|\alpha_n|, |\alpha_m|\}$. Thus, if α is (B, ζ) -reflective and satisfies (2.3), then $\rho^{\mathbb{C}}$ obeys

$$|(\mathcal{R}_{\zeta} \rho^{\mathbb{C}})_n - (-\overline{\rho_n^{\mathbb{C}}})| \leq \frac{2r_0}{\sqrt{1 - r_0^2}} e^{-B|\zeta|} \quad (3.7)$$

for all $n \in \mathbb{Z}$ such that $|\zeta - n| \leq \exp(B|\zeta|)$.

In view of the foregoing remark, we introduce the following.

Notation. Throughout the argument, we let $C > 0$ stand for a constant that depends only on r_0 from (2.3), and write $f \lesssim g$ to mean $f \leq Cg$ for such a constant.

Proof of Theorem 2.2. Assume α is B -reflective and B is sufficiently large. Choose $|\zeta_i| \rightarrow \infty$ such that α is (B, ζ_i) -reflective. By passing to a subsequence, we may assume that every ζ_i has the same sign and $2\zeta_i$ has the same residue modulo 4 for every i . Without loss of generality, we consider the case $\zeta_i > 0$ and $2\zeta_i \cong 2 \pmod{4}$ for every i , so we write $\zeta_i = 2m_i - 1$ with $m_i \in \mathbb{N}$. The other cases are similar. The main difference between the cases is what transfer matrix cocycle one must use. To be concrete, the current case compares N with its “reflected” variant, while the case $2\zeta_i \cong 0 \pmod{4}$ compares M with its reflected variant. The case of half-integer centers

($2\zeta_i$ odd) requires one to use transfer matrices associated with the *transpose* of a CMV matrix, but aside from that, all steps are analogous.

Due to [11, Theorem 2.1], it suffices to prove the statement of the theorem for a single admissible ρ . As discussed in Remark 3.2, the symmetry of the α_n can be passed to ρ_n , and furthermore, we need ρ to satisfy bounds such as (3.7) (up to shifting the center of reflection). Thus, we choose $\rho_n = i\sqrt{1 - |\alpha_n|^2}$. Consequently, we may assume henceforth that

$$|\rho_{4m_i-\ell-2} + \bar{\rho}_\ell| \lesssim e^{-B\zeta_i}, \quad |\alpha_{4m_i-\ell-2} - \bar{\alpha}_\ell| < e^{-B\zeta_i}, \quad (3.8)$$

for every $\ell \in \mathbb{Z}$ satisfying $|\zeta_i - \ell| \leq e^{B\zeta_i}$, and recall that our assumptions give $|\rho_\ell| \geq c_0 > 0$ uniformly in $\ell \in \mathbb{Z}$.

Remark 3.3. By a standard telescoping estimate, (3.8) implies that (since $|\rho|, |\alpha| \leq 1$)

$$|\alpha_{4m_i-\ell-2}\rho_{4m_i-\ell'-2} + \bar{\alpha}_\ell\bar{\rho}_{\ell'}| \lesssim e^{-B\zeta_i}$$

for all $\ell, \ell' \in \mathbb{Z}$ with $|\zeta_i - \ell|, |\zeta_i - \ell'| \leq e^{B\zeta_i}$, with similar bounds for other suitable combinations of ρ 's and α 's.

Let us define the reflection of u through $2m_i - \frac{1}{2}$ by

$$u_i(k) = u(4m_i - k - 1) \quad (3.9)$$

and note that

$$u_i(2n - 1) = u(4m_i - 2n), \quad u_i(2n) = u(4m_i - 2n - 1).$$

Lemma 3.4. *Under the assumptions (3.8) and with u_i as in (3.9),*

$$|W(u, u_i)(n) - W(u, u_i)(n')| \lesssim e^{-B\zeta_i}$$

for all n, n' with $|m_i - n|, |m_i - n'| \leq \frac{1}{2}e^{B\zeta_i} - 1$.

Proof. To begin, notice that the assumption on $|m_i - n|$ implies that for $\eta = -1, 0, 1$, one has

$$|\zeta_i - (2n + \eta)| = |2m_i - 1 - 2n - \eta| \leq 2|m_i - n| + 2 \leq e^{B\zeta_i},$$

and thus we can (and will) apply (3.8) with $\ell = 2n - 1, 2n, 2n + 1$ in the present argument.

By definition of the Wronskian, we have

$$\begin{aligned} & |W(u, v)(n) - W(u, v)(n - 1)| \\ &= \frac{1}{|\rho_{2n}|} \left| \rho_{2n+1}\rho_{2n}[u(2n + 2)v(2n + 1) - u(2n + 1)v(2n + 2)] \right. \\ & \quad \left. - \bar{\rho}_{2n}\bar{\rho}_{2n-1}[u(2n)v(2n - 1) - u(2n - 1)v(2n)] \right|. \end{aligned}$$

Applying the generalized eigenequation (3.2) and (3.3) for u yields

$$\begin{aligned}
 & |W(u, v)(n) - W(u, v)(n-1)| \\
 &= \frac{1}{|\overline{\rho_{2n}}|} \left| \left[zu(2n) - \overline{\alpha_{2n}}[\overline{\rho_{2n-1}}u(2n-1) - \alpha_{2n-1}u(2n)] \right. \right. \\
 &\quad \left. \left. - \rho_{2n}\overline{\alpha_{2n+1}}u(2n+1) \right] v(2n+1) \right. \\
 &\quad \left. - \rho_{2n+1}\rho_{2n}u(2n+1)v(2n+2) - \overline{\rho_{2n}}\overline{\rho_{2n-1}}u(2n)v(2n-1) \right. \\
 &\quad \left. + [zu(2n+1) + \overline{\rho_{2n}}\alpha_{2n-1}u(2n) \right. \\
 &\quad \left. + \alpha_{2n}[\overline{\alpha_{2n+1}}u(2n+1) + \rho_{2n+1}u(2n+2)] \right] v(2n) \Big| \\
 &= \frac{1}{|\overline{\rho_{2n}}|} \left| [\overline{\rho_{2n}}\alpha_{2n-1}v(2n) - \overline{\rho_{2n}}\overline{\rho_{2n-1}}v(2n-1)]u(2n) \right. \\
 &\quad \left. - [\rho_{2n}\overline{\alpha_{2n+1}}v(2n+1) + \rho_{2n+1}\rho_{2n}v(2n+2)]u(2n+1) \right. \\
 &\quad \left. - \overline{\alpha_{2n}}[\overline{\rho_{2n-1}}u(2n-1) - \alpha_{2n-1}u(2n)]v(2n+1) \right. \\
 &\quad \left. + \alpha_{2n}[\overline{\alpha_{2n+1}}u(2n+1) + \rho_{2n+1}u(2n+2)]v(2n) \right. \\
 &\quad \left. + z[u(2n)v(2n+1) + u(2n+1)v(2n)] \right|. \tag{3.10}
 \end{aligned}$$

Now, setting $v = u_i$ with u_i as in (3.9), we claim

$$\begin{aligned}
 & |W(u, u_i)(n) - W(u, u_i)(n-1)| \\
 &= \frac{1}{|\overline{\rho_{2n}}|} \left| [\overline{\rho_{2n}}\alpha_{2n-1} + \rho_{4m_i-2n-2}\overline{\alpha_{4m_i-2n-1}}]u(4m_i-2n-1)u(2n) \right. \\
 &\quad \left. - [\overline{\rho_{2n}}\overline{\rho_{2n-1}} - \rho_{4m_i-2n-2}\rho_{4m_i-2n-1}]u(4m_i-2n)u(2n) \right. \\
 &\quad \left. - [\rho_{2n}\overline{\alpha_{2n+1}} + \overline{\rho_{4m_i-2n-2}}\alpha_{4m_i-2n-3}]u(4m_i-2n-2)u(2n+1) \right. \\
 &\quad \left. + [\rho_{2n+1}\rho_{2n} - \overline{\rho_{4m_i-2n-2}}\overline{\rho_{4m_i-2n-3}}]u(4m_i-2n-3)u(2n+1) \right. \\
 &\quad \left. + z[\overline{\alpha_{4m_i-2n-2}}\alpha_{4m_i-2n-2} - \overline{\alpha_{2n}}\alpha_{2n}]u(4m_i-2n-2)u(2n) \right. \\
 &\quad \left. + z[\overline{\alpha_{4m_i-2n-2}}\rho_{4m_i-2n-2} + \overline{\rho_{2n}}\alpha_{2n}]u(4m_i-2n-1)u(2n) \right. \\
 &\quad \left. - z[\alpha_{4m_i-2n-2}\overline{\rho_{4m_i-2n-2}} + \overline{\alpha_{2n}}\rho_{2n}]u(4m_i-2n-2)u(2n+1) \right. \\
 &\quad \left. + z[\alpha_{4m_i-2n-2}\overline{\alpha_{4m_i-2n-2}} - \overline{\alpha_{2n}}\alpha_{2n}]u(4m_i-2n-1)u(2n+1) \right|. \tag{3.11}
 \end{aligned}$$

The proof of (3.11) is not straightforward and also somewhat lengthy, so we put it into Appendix A. Now, using the assumptions (3.8), (3.11), the Cauchy–Schwarz inequality, and $z \in \partial\mathbb{D}$, we get

$$|W(u, u_i)(n) - W(u, u_i)(n')| \lesssim e^{-B\xi_i}$$

for all n, n' as in the statement of the lemma. ■

Next, use the Cauchy–Schwarz inequality and $|\rho_k| \leq 1$ to see that

$$\sum_n |W(u, u_i)(n)| \leq 2. \tag{3.12}$$

This implies that there exists $n \in \mathbb{Z}$ with $|m_i - n| \leq \frac{1}{2}e^{B\xi_i} - 1$ such that

$$|W(u, u_i)(n)| \leq 3e^{-B\xi_i}$$

(for if not, summing $W(u, u_i)(n)$ over the relevant range of n 's produces a contradiction to (3.12)). Lemma 3.4 then implies that

$$|W(u, u_i)(n)| \lesssim e^{-B\xi_i} \quad \text{for all } n \in \mathbb{Z} \text{ with } |m_i - n| \leq \frac{1}{2}e^{B\xi_i} - 1.$$

Let us define $u_i^\pm = u \pm u_i$, and (using (3.9)) note that

$$u_i^\pm(2m_i - 1) = \pm u_i^\pm(2m_i). \quad (3.13)$$

Further, we define

$$\Phi_i^\pm(n) = \begin{bmatrix} u_i^\pm(2n) \\ u_i^\pm(2n - 1) \end{bmatrix}. \quad (3.14)$$

Lemma 3.5. *For each i , there exists $s = s_i \in \{+, -\}$ such that*

$$\|\Phi_i^s(m_i)\| = [|u_i^s(2m_i)|^2 + |u_i^s(2m_i - 1)|^2]^{1/2} \lesssim e^{-B\xi_i/2}. \quad (3.15)$$

Proof. First note that $W(u_i^-, u_i^+) = 2W(u, u_i)$. Together with (3.13) this implies that

$$\begin{aligned} & 2|\rho_{2m_i-1}||u_i^-(2m_i - 1)u_i^+(2m_i)| \\ &= |\rho_{2m_i-1}||u_i^-(2m_i)u_i^+(2m_i - 1) - u_i^-(2m_i - 1)u_i^+(2m_i)| \\ &= |W(u_i^-, u_i^+)(m_i - 1)| = 2|W(u, u_i)(m_i - 1)| \lesssim e^{-B\xi_i}. \end{aligned}$$

Thus,

$$|u_i^-(2m_i - 1)u_i^+(2m_i)| \lesssim e^{-B\xi_i},$$

so either

$$|u_i^-(2m_i - 1)| \lesssim e^{-B\xi_i/2} \quad \text{or} \quad |u_i^+(2m_i)| \lesssim e^{-B\xi_i/2}.$$

In either case, (3.15) follows by symmetry of the $u_i^\pm$, that is, by (3.13). ■

Next, we want to show that a similar bound holds for $\Phi_i^\pm(0)$. To this end, define

$$\Phi(n) = \begin{bmatrix} u(2n) \\ u(2n - 1) \end{bmatrix}, \quad \Phi_i(n) = \begin{bmatrix} u_i(2n) \\ u_i(2n - 1) \end{bmatrix} = \begin{bmatrix} u(4m_i - 2n - 1) \\ u(4m_i - 2n) \end{bmatrix}$$

and note that $\Phi_i^\pm = \Phi \pm \Phi_i$ on account of (3.14) and the definition of $u_i^\pm$. Let us write

$$N_z^n = N_z^{[n-1, 0]} = \prod_{k=n-1}^0 N_{k,z}, \quad (3.16)$$

where N is given by (3.5). By Lemma 3.1, we have $\Phi(n) = N_z^n \Phi(0)$ and thus $\Phi(0) = [N_z^n]^{-1} \Phi(n)$ with

$$[N_z^n]^{-1} = \prod_{k=0}^{n-1} N_{k,z}^{-1}.$$

The reader can check that $N_{n,z}^{-1}$ is given by

$$\begin{aligned} N_{n,z}^{-1} &= \frac{1}{\overline{\rho_{2n}} \overline{\rho_{2n-1}}} \begin{bmatrix} \overline{\rho_{2n-1}} \rho_{2n+1} z^{-1} \\ \alpha_{2n} \rho_{2n+1} + \alpha_{2n-1} \rho_{2n+1} z^{-1} \\ \overline{\alpha_{2n}} \overline{\rho_{2n-1}} + \overline{\alpha_{2n+1}} \overline{\rho_{2n-1}} z^{-1} \\ z + \overline{\alpha_{2n}} \alpha_{2n-1} + \overline{\alpha_{2n+1}} \alpha_{2n-1} z^{-1} + \overline{\alpha_{2n+1}} \alpha_{2n} \end{bmatrix} \\ &=: \frac{1}{\overline{\rho_{2n}} \overline{\rho_{2n-1}}} \tilde{N}_{n,z}^{-1}. \end{aligned} \quad (3.17)$$

Also, with the “mirrored transfer” matrices defined by

$$\begin{aligned} H_{n,z} &:= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} N_{n,z} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= \frac{1}{\rho_{2n+1} \rho_{2n}} \begin{bmatrix} \overline{\rho_{2n-1}} \rho_{2n+1} z^{-1} \\ -\overline{\alpha_{2n}} \overline{\rho_{2n-1}} - \overline{\alpha_{2n+1}} \overline{\rho_{2n-1}} z^{-1} \\ -\alpha_{2n} \rho_{2n+1} - \alpha_{2n-1} \rho_{2n+1} z^{-1} \\ z + \overline{\alpha_{2n}} \alpha_{2n-1} + \overline{\alpha_{2n+1}} \alpha_{2n-1} z^{-1} + \overline{\alpha_{2n+1}} \alpha_{2n} \end{bmatrix} \\ &=: \frac{1}{\rho_{2n+1} \rho_{2n}} \tilde{H}_{n,z}, \end{aligned} \quad (3.18)$$

we deduce from Lemma 3.1 that

$$\begin{aligned} \Phi_i(0) &= \begin{bmatrix} u_i(0) \\ u_i(-1) \end{bmatrix} = \begin{bmatrix} u(4m_i - 1) \\ u(4m_i) \end{bmatrix} \\ &= \underbrace{\prod_{k=2m_i-1}^{m_i} H_{k,z}}_{=: H_z^{[2m_i-1, m_i]}} \begin{bmatrix} u(2m_i - 1) \\ u(2m_i) \end{bmatrix} = H_z^{[2m_i-1, m_i]} \Phi_i(m_i). \\ &=: H_z^{[2m_i-1, m_i]} \end{aligned}$$

Lemma 3.6. *With s_i as in Lemma 3.5, $\|\Phi_i^{s_i}(0)\| \rightarrow 0$ as $m_i \rightarrow \infty$.*

Proof. We have

$$\begin{aligned} \Phi_i^\pm(0) &= \Phi(0) \pm \Phi_i(0) = [N_z^{m_i}]^{-1} \Phi(m_i) \pm H_z^{[2m_i-1, m_i]} \Phi_i(m_i) \\ &= [N_z^{m_i}]^{-1} \Phi_i^\pm(m_i) \mp [[N_z^{m_i}]^{-1} - H_z^{[2m_i-1, m_i]}] \Phi_i(m_i). \end{aligned} \quad (3.19)$$

Thus, to control $\|\Phi_i^\pm(0)\|$ we need to bound $\|[N_z^{m_i}]^{-1}\|$ and $\|[N_z^{m_i}]^{-1} - H_z^{[2m_i-1, m_i]}\|$. To do this, we use the standard telescoping estimate: for matrices $A_1, \dots, A_n$ and $B_1, \dots, B_n$ we have that

$$\prod_{k=n}^1 A_k - \prod_{k=n}^1 B_k = \sum_{i=n}^1 \left[\prod_{k=n}^{i+1} A_k \right] [A_i - B_i] \left[\prod_{k=i-1}^1 B_k \right]$$

and thus

$$\left\| \prod_{k=n}^1 A_k - \prod_{k=n}^1 B_k \right\| \leq \sum_{i=n}^1 \left\| \prod_{k=n}^{i+1} A_k \right\| \|A_i - B_i\| \left\| \prod_{k=i-1}^1 B_k \right\|.$$

Applying this to the second term of (3.19), two of the factors are bounded by the submultiplicativity of the operator norm and the observation that

$$\|N_{k,z}^{-1}\|, \|H_{2m_i-k-1,z}\| \leq C.$$

Comparing the expressions for N^{-1} and H in (3.17) and (3.18) and using the symmetries (3.8) gives

$$\|N_{k,z}^{-1} - H_{2m_i-k-1,z}\| \lesssim e^{-B\xi_i}.$$

Putting it all together, we have

$$\|[N_z^{m_i}]^{-1} - H_z^{[2m_i-1, m_i]}\| \leq m_i C^{m_i} e^{-B\xi_i} \quad (3.20)$$

Similarly,

$$\|[N_z^{m_i}]^{-1} \Phi_i^{s_i}(m_i)\| \leq C^{m_i} e^{-B\xi_i/2}. \quad (3.21)$$

Since the right-hand sides of (3.20) and (3.21) go to zero as $i \rightarrow \infty$ when B is large enough, combining these estimates with (3.19) proves the lemma. ■

With this result we can conclude the proof of Theorem 2.2 as follows. Lemma 3.6 implies that $\|\Phi(0)\| - \|\Phi(m_i)\| \rightarrow 0$. By assumption, $u \in \ell^2(\mathbb{Z})$, so we conclude that $\Phi(0) = 0$, which implies that $u = 0$ (by (3.6)). This contradicts normalization of u . ■

Let us point out that Corollary 2.4 now follows in short order by a soft argument with the Baire category theorem.

Proof of Corollary 2.4. Assume α satisfies (2.3) and is reflection-symmetric, write $\Omega = \text{hull}(\alpha)$, and fix B large.

For each $m \in \mathbb{Z}$, let $\text{Refl}(B, m) \subseteq \Omega$ denote the set of (B, m) -reflective elements of Ω . For each k , the set

$$\bigcup_{m \geq k} \text{Refl}(B, m)$$

is open by definition and dense in Ω (as it contains a translation semiorbit of α). Therefore,

$$\Omega_0 = \bigcap_{k \geq 1} \bigcup_{m \geq k} \text{Refl}(B, m)$$

is a dense G_δ by the Baire category theorem. Since $\mathcal{E}_\omega$ has purely continuous spectrum for every $\omega \in \Omega_0$ by Theorem 2.2, we are done. ■

We now move on to a discussion of the quantitative strengthening in Theorem 2.5. If α is almost-periodic, its hull $\Omega = \text{hull}(\alpha)$ is a compact abelian topological group and hence is equipped with a unique translation-invariant Borel probability measure, which we denote by μ . For each $\omega \in \Omega$, we may consider the associated transfer matrices $N_{k,z} = N_{k,z}(\omega)$ and their iterates $N_z^n(\omega)$ as in (3.5) and (3.16). The corresponding Lyapunov exponent is given by⁴

$$L(z) = \lim_{n \rightarrow \infty} \frac{1}{2n} \int_{\Omega} \log \|N_z^n(\omega)\| d\mu(\omega). \quad (3.22)$$

The argument is an adaptation of the previous argument along the contours of [30]; since it is so similar to the previous argument, we only sketch the main steps.

Proof sketch of Theorem 2.5. Fix α almost-periodic and B -reflective satisfying (2.3), assume $B > 2L(z)$, and fix $\varepsilon > 0$ small enough that $2(L(z) + \varepsilon) < B$. The initial parts of the proof proceed in precisely the same manner until one reaches (3.19). At this point, one estimates the terms using the semiuniform ergodic theorem of Furman (instead of a crude *a priori* estimate) to get improvements to (3.20) and (3.21):

$$\begin{aligned} \|[N_z^{m_i}]^{-1} \Phi_i^{s_i}(m_i)\| &\lesssim e^{2m_i(L(z)+\varepsilon)} e^{-B\zeta_i/2}, \\ \|[N_z^{m_i}]^{-1} - H_z^{[2m_i-1, m_i]}\| &\lesssim m_i e^{2m_i(L(z)+\varepsilon)} e^{-B\zeta_i}. \end{aligned}$$

Recalling that $\zeta_i = 2m_i - 1$, the condition $B > 2(L(z) + \varepsilon)$ suffices to see that the expression in (3.19) goes to zero and hence the rest of the argument can be run as before. ■

⁴The extra factor of one-half is so that the Lyapunov exponent of this cocycle matches that of the usual Szegő and Gesteş–Zinchenko cocycles, on account of the connections between cocycles discussed in the literature, e.g., [11, 18, 47].

Proof of Corollary 2.6. Notice that the assumptions imply that α_ω satisfies (2.3) for all ω . Let $\delta = \delta(\beta, \omega)$. For any $\varepsilon > 0$, we may choose $|k_i| \rightarrow \infty$ with $\|2\omega + k_i\beta\| < e^{(\delta-\varepsilon)|k_i|}$, and observe by symmetry of f that

$$\begin{aligned} |\alpha_\omega(k_i - n) - \overline{\alpha_\omega(n)}| &= |f(\omega + (k_i - n)\beta) - \overline{f(\omega + n\beta)}| \\ &= |f(\omega + (k_i - n)\beta) - f(-\omega - n\beta)| \\ &\lesssim \|2\omega + k_i\beta\| \lesssim e^{-(\delta-\varepsilon)|k_i|}. \end{aligned}$$

Thus, if $L(z) < \delta$, then we can choose $\varepsilon > 0$ with $L(z) < \delta - \varepsilon$ and it follows from the above that α_ω is $B = 2(\delta - \varepsilon)$ -reflective,⁵ so z is not an eigenvalue of $\mathcal{E}_\omega$ by Theorem 2.5. ■

With the main results proved, we may now reap our harvest of corollaries.

Proof of Corollary 2.7. Notice that $\mathcal{E}_{\lambda_1, \lambda_2, \Phi, \theta}^{\text{UAMO}}$ falls into the setting of Corollary 2.6 with

$$\Omega = \mathbb{T} \times \mathbb{Z}_2, \quad \beta = \left(\frac{\Phi}{2}, 1\right), \quad f(\theta, j) = \begin{cases} \lambda_2 \cos(2\pi\theta) & \text{if } j = 0, \\ (1 - \lambda_1^2)^{1/2} & \text{if } j = 1. \end{cases} \quad (3.23)$$

Clearly, f is Lipschitz continuous and even. Thus, according to Corollary 2.6 together with [12, Corollary 2.10], absence of eigenvalues holds throughout the spectrum for any θ such that

$$\delta((\Phi/2, 1), (\theta, 0)) > \log \left[\frac{\lambda_2(1 + \lambda_1')}{\lambda_1(1 + \lambda_2')} \right]. \quad (3.24)$$

Recalling the definition of δ from (2.4), we see that $\delta((\frac{1}{2}\Phi, 1), (\theta, 0)) = \frac{1}{2}\delta(\Phi, \theta)$, so (3.24) is equivalent to $\frac{1}{2}\delta(\Phi, \theta) > L(z)$. Since this holds for generic $\theta \in \mathbb{T}$, this shows absence of eigenvalues when (2.5) holds. Since the expression on the right-hand side of (2.5) is equal to the Lyapunov exponent on the spectrum [12] and is positive when $\lambda_1 < \lambda_2$, it follows that the absolutely continuous and point parts of the spectrum are both absent when (2.5) and $\lambda_1 < \lambda_2$ hold. ■

Proof of Corollary 2.8. This follows from similar considerations as in the previous corollary after changing $\mathbb{Z}_2$ to $\mathbb{Z}_{2s}$ and suitably modifying f . ■

Proof of Corollary 2.9. Again, observe that the assumptions imply that α_ω satisfies (2.3) for all ω . This follows immediately by combining Theorem 2.2 with work of Hof, Knill, and Simon, specifically [29, Proposition 2.1]. ■

⁵Notice that $\frac{k_i}{2}$, not k_i , is the center of reflection, which accounts for the factor of 2 here.

A. Proof of (3.11)

We need one preliminary: the $\mathcal{LM}$ factorization. Concretely, we define

$$\mathcal{L} = \bigoplus_{n \in \mathbb{Z}} \Theta(\alpha_{2n}, \rho_{2n}), \quad \mathcal{M} = \bigoplus_{n \in \mathbb{Z}} \Theta(\alpha_{2n+1}, \rho_{2n+1}), \quad \Theta(\alpha, \rho) = \begin{bmatrix} \bar{\alpha} & \rho \\ \bar{\rho} & -\alpha \end{bmatrix},$$

where $\Theta(\alpha_j, \rho_j)$ acts on the subspace $\ell^2(\{j, j+1\})$. With these definitions, one has $\mathcal{E} = \mathcal{LM}$. Since $\mathcal{L}$ and $\mathcal{M}$ are themselves unitary, this means that $\mathcal{E}u = zu$ is equivalent to $\mathcal{M}u = z\mathcal{L}^*u$. Evaluating this relation at coordinates $2n$ and $2n+1$ gives

$$\overline{\rho_{2n-1}}u(2n-1) - \alpha_{2n-1}u(2n) = z(\alpha_{2n}u(2n) + \rho_{2n}u(2n+1)), \quad (\text{A.1})$$

$$\overline{\alpha_{2n+1}}u(2n+1) + \rho_{2n+1}u(2n+2) = z(\overline{\rho_{2n}}u(2n) - \bar{\alpha}_{2n}u(2n+1)). \quad (\text{A.2})$$

We now proceed with the proof of equation (3.11). Starting from (3.10), using the eigenequation of u , and adding 0 in an inspired manner gives

$$\begin{aligned} & |W(u, u_i)(n) - W(u, u_i)(n-1)| \\ &= \frac{1}{|\rho_{2n}|} \left| [\overline{\rho_{2n}}\alpha_{2n-1}u(4m_i-2n-1) - \overline{\rho_{2n}}\overline{\rho_{2n-1}}u(4m_i-2n)]u(2n) \right. \\ &\quad + \rho_{4m_i-2n-2}[\overline{\alpha_{4m_i-2n-1}}u(4m_i-2n-1) \\ &\quad \quad + \rho_{4m_i-2n-1}u(4m_i-2n)]u(2n) \\ &\quad - \rho_{4m_i-2n-2}[\overline{\alpha_{4m_i-2n-1}}u(4m_i-2n-1) \\ &\quad \quad + \rho_{4m_i-2n-1}u(4m_i-2n)]u(2n) \\ &\quad - [\rho_{2n}\overline{\alpha_{2n+1}}u(4m_i-2n-2) + \rho_{2n+1}\rho_{2n}u(4m_i-2n-3)]u(2n+1) \\ &\quad + [\rho_{4m_i-2n-2}(\overline{\rho_{4m_i-2n-3}}u(4m_i-2n-3) \\ &\quad \quad - \alpha_{4m_i-2n-3}u(4m_i-2n-2))]u(2n+1) \\ &\quad - [\rho_{4m_i-2n-2}(\overline{\rho_{4m_i-2n-3}}u(4m_i-2n-3) \\ &\quad \quad - \alpha_{4m_i-2n-3}u(4m_i-2n-2))]u(2n+1) \\ &\quad - \overline{\alpha_{2n}}[\overline{\rho_{2n-1}}u(2n-1) - \alpha_{2n-1}u(2n)]u(4m_i-2n-2) \\ &\quad + [\alpha_{2n}(\overline{\alpha_{2n+1}}u(2n+1) + \rho_{2n+1}u(2n+2))]u(4m_i-2n-1) \\ &\quad + z(u(2n)u(4m_i-2n-2) + u(2n+1)u(4m_i-2n-1))] \Big| \\ &= \frac{1}{|\rho_{2n}|} \left| [(\overline{\rho_{2n}}\alpha_{2n-1} + \rho_{4m_i-2n-2}\overline{\alpha_{4m_i-2n-1}})u(4m_i-2n-1) \right. \\ &\quad - (\overline{\rho_{2n}}\overline{\rho_{2n-1}} - \rho_{4m_i-2n-2}\rho_{4m_i-2n-1})u(4m_i-2n)]u(2n) \\ &\quad - \rho_{4m_i-2n-2}[\overline{\alpha_{4m_i-2n-1}}u(4m_i-2n-1) \\ &\quad \quad + \rho_{4m_i-2n-1}u(4m_i-2n)]u(2n) \end{aligned}$$

$$\begin{aligned}
& - \left[(\rho_{2n} \overline{\alpha_{2n+1}} + \overline{\rho_{4m_i-2n-2} \alpha_{4m_i-2n-3}}) u(4m_i - 2n - 2) \right. \\
& \quad \left. + (\rho_{2n+1} \rho_{2n} - \overline{\rho_{4m_i-2n-2} \rho_{4m_i-2n-3}}) u(4m_i - 2n - 3) \right] u(2n + 1) \\
& - \left[\overline{\rho_{4m_i-2n-2}} (\overline{\rho_{4m_i-2n-3}} u(4m_i - 2n - 3) \right. \\
& \quad \left. - \alpha_{4m_i-2n-3} u(4m_i - 2n - 2)) \right] u(2n + 1) \\
& - \overline{\alpha_{2n}} [\overline{\rho_{2n-1}} u(2n - 1) - \alpha_{2n-1} u(2n)] u(4m_i - 2n - 2) \\
& + [\alpha_{2n} (\overline{\alpha_{2n+1}} u(2n + 1) + \rho_{2n+1} u(2n + 2))] u(4m_i - 2n - 1) \\
& + z(u(2n) u(4m_i - 2n - 2) + u(2n + 1) u(4m_i - 2n - 1)) \Big| \\
= & \frac{1}{|\rho_{2n}|} \Big| \left[(\overline{\rho_{2n}} \alpha_{2n-1} + \rho_{4m_i-2n-2} \overline{\alpha_{4m_i-2n-1}}) u(4m_i - 2n - 1) \right. \\
& \quad \left. - (\overline{\rho_{2n}} \overline{\rho_{2n-1}} - \rho_{4m_i-2n-2} \rho_{4m_i-2n-1}) u(4m_i - 2n) \right] u(2n) \\
& - \left[z u(4m_i - 2n - 2) - \overline{\alpha_{4m_i-2n-2}} (\overline{\rho_{4m_i-2n-3}} u(4m_i - 2n - 3) \right. \\
& \quad \left. - \alpha_{4m_i-2n-3} u(4m_i - 2n - 2)) \right] u(2n) \\
& - \left[(\rho_{2n} \overline{\alpha_{2n+1}} + \overline{\rho_{4m_i-2n-2} \alpha_{4m_i-2n-3}}) u(4m_i - 2n - 2) \right. \\
& \quad \left. + (\rho_{2n+1} \rho_{2n} - \overline{\rho_{4m_i-2n-2} \rho_{4m_i-2n-3}}) u(4m_i - 2n - 3) \right] u(2n + 1) \\
& - \left[z u(4m_i - 2n - 1) + \alpha_{4m_i-2n-2} (\overline{\alpha_{4m_i-2n-1}} u(4m_i - 2n - 1) \right. \\
& \quad \left. + \rho_{4m_i-2n-1} u(4m_i - 2n)) \right] u(2n + 1) \\
& - \overline{\alpha_{2n}} [\overline{\rho_{2n-1}} u(2n - 1) - \alpha_{2n-1} u(2n)] u(4m_i - 2n - 2) \\
& + [\alpha_{2n} (\overline{\alpha_{2n+1}} u(2n + 1) + \rho_{2n+1} u(2n + 2))] u(4m_i - 2n - 1) \\
& + z(u(2n) u(4m_i - 2n - 2) + u(2n + 1) u(4m_i - 2n - 1)) \Big| \\
= & \frac{1}{|\rho_{2n}|} \Big| \left[(\overline{\rho_{2n}} \alpha_{2n-1} + \rho_{4m_i-2n-2} \overline{\alpha_{4m_i-2n-1}}) u(4m_i - 2n - 1) \right. \\
& \quad \left. - (\overline{\rho_{2n}} \overline{\rho_{2n-1}} - \rho_{4m_i-2n-2} \rho_{4m_i-2n-1}) u(4m_i - 2n) \right] u(2n) \\
& - \left[(\rho_{2n} \overline{\alpha_{2n+1}} + \overline{\rho_{4m_i-2n-2} \alpha_{4m_i-2n-3}}) u(4m_i - 2n - 2) \right. \\
& \quad \left. + (\rho_{2n+1} \rho_{2n} - \overline{\rho_{4m_i-2n-2} \rho_{4m_i-2n-3}}) u(4m_i - 2n - 3) \right] u(2n + 1) \\
& + \left[\overline{\alpha_{4m_i-2n-2}} (\overline{\rho_{4m_i-2n-3}} u(4m_i - 2n - 3) \right. \\
& \quad \left. - \alpha_{4m_i-2n-3} u(4m_i - 2n - 2)) \right] u(2n) \\
& - \left[\alpha_{4m_i-2n-2} (\overline{\alpha_{4m_i-2n-1}} u(4m_i - 2n - 1) \right. \\
& \quad \left. + \rho_{4m_i-2n-1} u(4m_i - 2n)) \right] u(2n + 1) \\
& - \overline{\alpha_{2n}} [\overline{\rho_{2n-1}} u(2n - 1) - \alpha_{2n-1} u(2n)] u(4m_i - 2n - 2) \\
& + [\alpha_{2n} (\overline{\alpha_{2n+1}} u(2n + 1) + \rho_{2n+1} u(2n + 2))] u(4m_i - 2n - 1) \Big|.
\end{aligned}$$

Now, applying (A.1) and (A.2) transforms the final expression into

$$\begin{aligned}
= & \frac{1}{|\rho_{2n}|} \Big| \left[(\overline{\rho_{2n}} \alpha_{2n-1} + \rho_{4m_i-2n-2} \overline{\alpha_{4m_i-2n-1}}) u(4m_i - 2n - 1) \right. \\
& \quad \left. - (\overline{\rho_{2n}} \overline{\rho_{2n-1}} - \rho_{4m_i-2n-2} \rho_{4m_i-2n-1}) u(4m_i - 2n) \right] u(2n)
\end{aligned}$$

$$\begin{aligned}
& - \left[(\rho_{2n} \overline{\alpha_{2n+1}} + \overline{\rho_{4m_i-2n-2} \alpha_{4m_i-2n-3}}) u(4m_i - 2n - 2) \right. \\
& \quad \left. + (\rho_{2n+1} \rho_{2n} - \overline{\rho_{4m_i-2n-2} \rho_{4m_i-2n-3}}) u(4m_i - 2n - 3) \right] u(2n + 1) \\
& + \left[\overline{\alpha_{4m_i-2n-2} z} (\alpha_{4m_i-2n-2} u(4m_i - 2n - 2) \right. \\
& \quad \left. + \rho_{4m_i-2n-2} u(4m_i - 2n - 1)) \right] u(2n) \\
& - \left[\alpha_{4m_i-2n-2} z (\overline{\rho_{4m_i-2n-2}} u(4m_i - 2n - 2) \right. \\
& \quad \left. - \overline{\alpha_{4m_i-2n-2}} u(4m_i - 2n - 1)) \right] u(2n + 1) \\
& - \overline{\alpha_{2n}} z [\alpha_{2n} u(2n) + \rho_{2n} u(2n + 1)] u(4m_i - 2n - 2) \\
& + [\alpha_{2n} z (\overline{\rho_{2n}} u(2n) - \overline{\alpha_{2n}} u(2n + 1))] u(4m_i - 2n - 1) \Big|.
\end{aligned}$$

This can in turn be rearranged into

$$\begin{aligned}
& = \frac{1}{|\overline{\rho_{2n}}|} \Big| (\overline{\rho_{2n}} \alpha_{2n-1} + \rho_{4m_i-2n-2} \overline{\alpha_{4m_i-2n-1}}) u(4m_i - 2n - 1) u(2n) \\
& \quad - (\overline{\rho_{2n}} \overline{\rho_{2n-1}} - \rho_{4m_i-2n-2} \rho_{4m_i-2n-1}) u(4m_i - 2n) u(2n) \\
& \quad - (\rho_{2n} \overline{\alpha_{2n+1}} + \overline{\rho_{4m_i-2n-2} \alpha_{4m_i-2n-3}}) u(4m_i - 2n - 2) u(2n + 1) \\
& \quad + (\rho_{2n+1} \rho_{2n} - \overline{\rho_{4m_i-2n-2} \rho_{4m_i-2n-3}}) u(4m_i - 2n - 3) u(2n + 1) \\
& \quad + z [\overline{\alpha_{4m_i-2n-2}} \alpha_{4m_i-2n-2} - \overline{\alpha_{2n}} \alpha_{2n}] u(4m_i - 2n - 2) u(2n) \\
& \quad + z [\overline{\alpha_{4m_i-2n-2}} \rho_{4m_i-2n-2} + \overline{\rho_{2n}} \alpha_{2n}] u(4m_i - 2n - 1) u(2n) \\
& \quad - z [\alpha_{4m_i-2n-2} \overline{\rho_{4m_i-2n-2}} + \overline{\alpha_{2n}} \rho_{2n}] u(4m_i - 2n - 2) u(2n + 1) \\
& \quad + z [\alpha_{4m_i-2n-2} \overline{\alpha_{4m_i-2n-2}} - \overline{\alpha_{2n}} \alpha_{2n}] u(4m_i - 2n - 1) u(2n + 1) \Big|,
\end{aligned}$$

which is (3.11).

Acknowledgments. J. Fillman thanks the American Institute of Mathematics for hospitality during a recent SQuaRE program, which facilitated the work. The authors are grateful to the reviewers for carefully reading the manuscript and for helpful comments.

Funding. C. Cedzich was supported in part by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under the grant number 441423094. J. Fillman was supported in part by National Science Foundation (NSF) grant DMS-2213196.

References

- [1] A. Ahlbrecht, C. Cedzich, R. Matjeschk, V. B. Scholz, A. H. Werner, and R. F. Werner, *Asymptotic behavior of quantum walks with spatio-temporal coin fluctuations*. *Quantum Inf. Process.* **11** (2012), no. 5, 1219–1249 Zbl [1252.82080](#) MR [2979893](#)

- [2] A. Ahlbrecht, V. B. Scholz, and A. H. Werner, [Disordered quantum walks in one lattice dimension](#). *J. Math. Phys.* **52** (2011), no. 10, article no. 102201 Zbl [1272.81036](#) MR [2894584](#)
- [3] A. Ahlbrecht, H. Vogts, A. H. Werner, and R. F. Werner, [Asymptotic evolution of quantum walks with random coin](#). *J. Math. Phys.* **52** (2011), no. 4, article no. 042201 Zbl [1316.81066](#) MR [2964162](#)
- [4] M. Aizenman and S. Warzel, *Random operators. Disorder effects on quantum spectra and dynamics*. Grad. Stud. Math. 168, American Mathematical Society, Providence, RI, 2015 Zbl [1333.82001](#) MR [3364516](#)
- [5] J. Asch and A. Joye, [Lower bounds on the localisation length of balanced random quantum walks](#). *Lett. Math. Phys.* **109** (2019), no. 9, 2133–2155 Zbl [1428.82050](#) MR [3996005](#)
- [6] A. Avila and D. Damanik, Deterministic delocalization. 2024, <https://bicmr.pku.edu.cn/content/show/18-3287.html> visited on 24 September 2024
- [7] O. Bourget, J. S. Howland, and A. Joye, [Spectral analysis of unitary band matrices](#). *Comm. Math. Phys.* **234** (2003), no. 2, 191–227 Zbl [1029.47016](#) MR [1962460](#)
- [8] V. Bucaj, D. Damanik, J. Fillman, V. Gerbuz, T. VandenBoom, F. Wang, and Z. Zhang, [Localization for the one-dimensional Anderson model via positivity and large deviations for the Lyapunov exponent](#). *Trans. Amer. Math. Soc.* **372** (2019), no. 5, 3619–3667 Zbl [1422.35022](#) MR [3988621](#)
- [9] M.-J. Cantero, L. Moral, F. A. Grünbaum, and L. Velázquez, [Matrix-valued Szegő polynomials and quantum random walks](#). *Comm. Pure Appl. Math.* **63** (2010), no. 4, 464–507 Zbl [1186.81036](#) MR [2604869](#)
- [10] C. Cedzich, J. Fillman, T. Geib, and A. H. Werner, [Singular continuous Cantor spectrum for magnetic quantum walks](#). *Lett. Math. Phys.* **110** (2020), no. 6, 1141–1158 Zbl [1445.81025](#) MR [4109482](#)
- [11] C. Cedzich, J. Fillman, L. Li, D. C. Ong, and Q. Zhou, [Exact mobility edges for almost-periodic CMV matrices via gauge symmetries](#). *Int. Math. Res. Not. IMRN* (2024), no. 8, 6906–6941 MR [4735649](#)
- [12] C. Cedzich, J. Fillman, and D. C. Ong, [Almost everything about the unitary almost Mathieu operator](#). *Comm. Math. Phys.* **403** (2023), no. 2, 745–794 Zbl [1539.47080](#) MR [4645728](#)
- [13] C. Cedzich, T. Rybár, A. H. Werner, A. Alberti, M. Genske, and R. F. Werner, [Propagation and spectral properties of quantum walks in electric fields](#). *Phys. Rev. Lett.* **111** (2013), no. 16, article no. 160601
- [14] C. Cedzich and A. H. Werner, [Anderson localization for electric quantum walks and skew-shift CMV matrices](#). *Comm. Math. Phys.* **387** (2021), no. 3, 1257–1279 Zbl [1484.82043](#) MR [4324378](#)
- [15] D. Damanik and J. Fillman, [Spectral properties of limit-periodic operators](#). In *Analysis and geometry on graphs and manifolds*, pp. 382–444, London Math. Soc. Lecture Note Ser. 461, Cambridge University Press, Cambridge, 2020 Zbl [1515.47047](#) MR [4412981](#)
- [16] D. Damanik and J. Fillman, [One-dimensional ergodic Schrödinger operators – I. General theory](#). Grad. Stud. Math. 221, American Mathematical Society, Providence, RI, 2022 Zbl [1504.58001](#) MR [4567742](#)

- [17] D. Damanik and J. Fillman, [One-dimensional ergodic Schrödinger operators – II. Specific classes](#). Grad. Stud. Math., American Mathematical Society, Providence, RI, to appear
- [18] D. Damanik, J. Fillman, and D. C. Ong, [Spreading estimates for quantum walks on the integer lattice via power-law bounds on transfer matrices](#). *J. Math. Pures Appl.* (9) **105** (2016), no. 3, 293–341 Zbl 1332.81066 MR 3465806
- [19] D. Damanik and R. Killip, [Reflection symmetries of almost periodic functions](#). *J. Funct. Anal.* **178** (2000), no. 2, 251–257 Zbl 0967.43002 MR 1802894
- [20] D. Damanik and D. Lenz, [Uniform Szegő cocycles over strictly ergodic subshifts](#). *J. Approx. Theory* **144** (2007), no. 1, 133–138 Zbl 1108.37002 MR 2287381
- [21] J. Fillman, [Purely singular continuous spectrum for Sturmian CMV matrices via strengthened Gordon lemmas](#). *Proc. Amer. Math. Soc.* **145** (2017), no. 1, 225–239 Zbl 1354.47023 MR 3565375
- [22] J. Fillman, [Resolvent methods for quantum walks with an application to a Thue–Morse quantum walk](#). *Interdiscip. Inform. Sci.* **23** (2017), no. 1, 27–32 Zbl 1470.81041 MR 3637881
- [23] J. Fillman, D. C. Ong, and Z. Zhang, [Spectral characteristics of the unitary critical almost-Mathieu operator](#). *Comm. Math. Phys.* **351** (2017), no. 2, 525–561 Zbl 06702035 MR 3613513
- [24] A. Furman, [On the multiplicative ergodic theorem for uniquely ergodic systems](#). *Ann. Inst. H. Poincaré Probab. Statist.* **33** (1997), no. 6, 797–815 Zbl 0892.60011 MR 1484541
- [25] A. J. Gordon, The point spectrum of the one-dimensional Schrödinger operator. *Uspehi Mat. Nauk* **31** (1976), no. 4(190), 257–258 MR 0458247
- [26] G. Grimmett, S. Janson, and P. F. Scudo, [Weak limits for quantum random walks](#). *Phys. Rev. E* **69**, no. 2 (2004), article no. 026119
- [27] S. Guo and D. Piao, [Lyapunov behavior and dynamical localization for quasi-periodic CMV matrices](#). *Linear Algebra Appl.* **606** (2020), 68–89 Zbl 1448.37028 MR 4130514
- [28] E. Hamza, A. Joye, and G. Stolz, [Localization for random unitary operators](#). *Lett. Math. Phys.* **75** (2006), no. 3, 255–272 Zbl 1101.82014 MR 2211031
- [29] A. Hof, O. Knill, and B. Simon, [Singular continuous spectrum for palindromic Schrödinger operators](#). *Comm. Math. Phys.* **174** (1995), no. 1, 149–159 Zbl 0839.11009 MR 1372804
- [30] S. Jitomirskaya and W. Liu, [Universal reflective-hierarchical structure of quasiperiodic eigenfunctions and sharp spectral transition in phase](#). *J. Eur. Math. Soc. (JEMS)* **26** (2024), no. 8, 2797–2836 Zbl 07875499 MR 4756946
- [31] S. Jitomirskaya and B. Simon, [Operators with singular continuous spectrum. III. Almost periodic Schrödinger operators](#). *Comm. Math. Phys.* **165** (1994), no. 1, 201–205 Zbl 0830.34074 MR 1298948
- [32] A. Joye, [Density of states and Thouless formula for random unitary band matrices](#). *Ann. Henri Poincaré* **5** (2004), no. 2, 347–379 Zbl 1062.81030 MR 2057678
- [33] A. Joye, [Dynamical localization for \$d\$ -dimensional random quantum walks](#). *Quantum Inf. Process.* **11** (2012), no. 5, 1251–1269 Zbl 1252.82087 MR 2979894
- [34] A. Joye and M. Merkli, [Dynamical localization of quantum walks in random environments](#). *J. Stat. Phys.* **140** (2010), no. 6, 1025–1053 Zbl 1296.82052 MR 2684498

- [35] W. Kirsch, An invitation to random Schrödinger operators. In *Random Schrödinger operators*, pp. 1–119, Panor. Synthèses 25, Société Mathématique de France, Paris, 2008
Zbl [1162.82004](#) MR [2509110](#)
- [36] D. A. Koslover, [Jacobi operators with singular continuous spectrum](#). *Lett. Math. Phys.* **71** (2005), no. 2, 123–134 Zbl [1077.39019](#) MR [2134692](#)
- [37] H. Krüger, [Orthogonal polynomials on the unit circle with Verblunsky coefficients defined by the skew-shift](#). *Int. Math. Res. Not. IMRN* (2013), no. 18, 4135–4169 Zbl [1329.33011](#) MR [3106885](#)
- [38] L. Li, D. Damanik, and Q. Zhou, [Absolutely continuous spectrum for CMV matrices with small quasi-periodic Verblunsky coefficients](#). *Trans. Amer. Math. Soc.* **375** (2022), no. 9, 6093–6125 Zbl [07578588](#) MR [4474886](#)
- [39] Y. Lin, D. Piao, and S. Guo, [Anderson localization for the quasi-periodic CMV matrices with Verblunsky coefficients defined by the skew-shift](#). *J. Funct. Anal.* **285** (2023), no. 4, article no. 109975 Zbl [07687703](#) MR [4584984](#)
- [40] D. C. Ong, [Limit-periodic Verblunsky coefficients for orthogonal polynomials on the unit circle](#). *J. Math. Anal. Appl.* **394** (2012), no. 2, 633–644 Zbl [1263.47035](#) MR [2927484](#)
- [41] D. C. Ong, [Purely singular continuous spectrum for CMV operators generated by sub-shifts](#). *J. Stat. Phys.* **155** (2014), no. 4, 763–776 Zbl [1325.47068](#) MR [3192183](#)
- [42] S. Richard, A. Suzuki, and R. Tiedra de Aldecoa, [Quantum walks with an anisotropic coin I: spectral theory](#). *Lett. Math. Phys.* **108** (2018), no. 2, 331–357 Zbl [1384.81033](#) MR [3748367](#)
- [43] D. Ruelle, [Ergodic theory of differentiable dynamical systems](#). *Inst. Hautes Études Sci. Publ. Math.* **50** (1979), no. 50, 27–58 Zbl [0426.58014](#) MR [0556581](#)
- [44] B. Simon, [Orthogonal polynomials on the unit circle. Part 1: Classical theory](#). Amer. Math. Soc. Colloq. Publ. 54, Part 1, American Mathematical Society, Providence, RI, 2005 Zbl [1082.42020](#) MR [2105088](#)
- [45] B. Simon, [Orthogonal polynomials on the unit circle. Part 2: Spectral theory](#). Amer. Math. Soc. Colloq. Publ. 54, Part 2, American Mathematical Society, Providence, RI, 2005 Zbl [1082.42021](#) MR [2105089](#)
- [46] F. Wang and D. Damanik, [Anderson localization for quasi-periodic CMV matrices and quantum walks](#). *J. Funct. Anal.* **276** (2019), no. 6, 1978–2006 Zbl [1462.35316](#) MR [3912798](#)
- [47] F. Yang, [Anderson localization for the unitary almost Mathieu operator](#). *Nonlinearity* **37** (2024), no. 8, article no. 085010 Zbl [MR 4774416](#)
- [48] X. Zhu, [Localization for random CMV matrices](#). *J. Approx. Theory* **298** (2024), article no. 106008 Zbl [07885182](#) MR [4688052](#)

Received 26 February 2024; revised 31 August 2024.

Christopher Cedzich

Faculty of Mathematics and Natural Sciences, Heinrich Heine Universität Düsseldorf,
Universitätsstrasse 1, 40225 Düsseldorf, Germany; cedzich@hhu.de

Jake Fillman

Department of Mathematics, Texas A&M University, 3368 TAMU, College Station,
TX 77843-3368, USA; fillman@tamu.edu