

Building and Integrating Brain-Wide Maps of Nervous System Function in Invertebrates

Talya S. Kramer^{1,2}, Steven W. Flavell^{1,*}

¹Picower Institute for Learning and Memory, Department of Brain & Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA

²MIT Biology Graduate Program, Massachusetts Institute of Technology, Cambridge, MA, USA

*Correspondence: flavell@mit.edu

Abstract

The selection and execution of context-appropriate behaviors is controlled by the integrated action of neural circuits throughout the brain. However, how activity is coordinated across brain regions, and how nervous system structure enables these functional interactions, remain open questions. Recent technical advances have made it feasible to build brain-wide maps of nervous system structure and function, such as brain activity maps, connectomes, and cell atlases. Here, we review recent progress in this area, focusing on *C. elegans* and *D. melanogaster*, as recent work has produced global maps of these nervous systems. We also describe neural circuit motifs elucidated in studies of specific networks, which highlight the complexities that must be captured to build accurate models of whole-brain function.

Highlights

- New atlases define brain-wide activity and organization in *C. elegans* and *Drosophila*
- Brain-wide imaging shows consistent principles of how neural activity encodes behavior
- Maps of synaptic and neurotransmitter connectivity reveal structural organization
- Neural circuits use varied motifs to control context-appropriate behaviors

Introduction

Even in small animals, ethologically relevant behaviors can be remarkably complex. Nervous systems with a limited number of neurons can direct behaviors like foraging, courtship, and navigation, and allow animals to respond to threats, injuries, and infection. Understanding how neurons act together to direct context-appropriate behaviors is an essential question in modern neuroscience. To date, most research has focused on individual circuits or neurons controlling specific behaviors. However, recent technical advances have dramatically expanded the scope of what is possible, allowing researchers unprecedented access into the brains of animals.

In this review, we discuss recent advances in building, connecting, and interpreting brain-wide maps of nervous system function in *C. elegans* and *Drosophila*. We first discuss whole-brain neural recordings from freely-behaving animals – studies that are mapping the relationship between neural activity and behavior. We then cover new, comprehensive maps of neuronal connectivity, genetic identity, and neuromodulation that have provided insights into nervous system structure. Finally, we discuss examples of individual circuit motifs with established links between structure and function that may aid our ability to interpret these new brain-wide maps.

As this is a fast-moving field, we have largely limited our focus to developments over the past few years.

Brain-wide activity maps: how the brain encodes behavior

Recent advances have made it possible to perform whole-brain calcium imaging in behaving animals, yielding new insights into how brain-wide activity generates motor outputs. In *C. elegans*, pioneering studies examined whole-brain activity in immobilized [1]–[3] and freely moving animals [4]–[6]. These studies showed that information about behavior is distributed across the brain, with neurons representing different aspects of locomotion such as velocity and turning.

Recent work combined brain-wide imaging in moving animals with reliable, brain-wide cell identification. This development allows comparisons of neuron activity to ongoing behavior (sample data shown in Figure 1a); importantly, these relationships can then be compared across animals. Imaging the sex-specific neurons of the *C. elegans* male tail during mating behavior showed that stereotyped sets of neurons are active during different phases of mating, like sliding, turning, and copulation [7*]. While some neurons have specialized functions, others are engaged in several aspects of mating. Functional correlations between neurons changed as animals switched behavioral outputs. Another recent study used encoder models to describe how each neuron class in the head of the hermaphrodite worm encodes specific behavioral features [8**]. Many neurons encode single behavioral features, like velocity or feeding, but a surprising number of neurons conjunctively encode multiple behaviors, revealing widespread multiplexing. While many neuron classes represent behavior reliably, a stereotyped subset changed encoding upon changes in the animal's internal state, suggestive of flexible remapping. Neuronal identification allowed comparison to the *C. elegans* connectome: neurons that are connected, especially by gap junctions, are more likely to show similar activity [7]–[9]. However, anatomical predictions of activity are not perfect, suggesting additional information is needed.

Whole-brain recordings in freely-behaving *Drosophila* similarly found a vast distribution of locomotor information. Behaviors such as walking elicit changes in activity across most brain regions, while less intensive grooming behavior only recruits specific brain regions (for example, compare heatmap of brain-wide activity to simultaneous behavior in Figure 1b) [10], [11**], [12**], [13**]. Careful analysis of different locomotor features showed that specific brain regions are active during distinct behavioral components such as movement initiation, forward velocity, and turning [10]–[13]. Brain-wide activity is similar during freely-initiated and forced walking behavior, suggesting that many of these signals may be proprioceptive rather than motor commands [11]. These behavioral signals are accompanied by widespread sensory signals, which are also starting to be mapped at brain-wide scale [14].

Comparing studies from these two species reveals some consistent principles. In both animals, locomotor information is distributed across a surprisingly large area of the brain. This organization may enable coordinated behavioral outputs, allowing circuits throughout the brain to receive information about current behavior. Sleep or quiescence states consistently evoke broad downregulation in brain-wide activity [15], [16]. In addition, anatomy may dictate activity: neurons that are connected tend to have similar activity patterns in *C. elegans*; in *Drosophila*, individual brain regions often contain small functional units of neurons with related dynamics [8], [11]–[13]. Finally, activity in bilaterally symmetric neurons or brain regions is

mostly similar [8], [9], [12], [17], with a notable exception seen in turning neurons that activate on a specific side for directed turns [12], [18].

An interesting feature of brain-wide dynamics in both organisms is the timescale of behavior representation. Neuronal activity can represent current behavior or behavior in the recent past [8], [12], [13]. Neurons with related behavioral information can show varied timescales in their tuning to that behavior, allowing for an ordered recruitment of different neurons as behavior progresses [8], [12], [13]. In flies, neural activity could even predict upcoming turning behavior, revealing signals related to motor planning [18]. These studies reveal consistent principles in the organization of global brain dynamics that are conserved across evolutionarily distant species.

Maps of neuronal architecture and genetic identity

Many advances in *C. elegans* neuroscience were enabled by early mapping of the connectome [19]. Recent studies expanded that understanding, identifying connections that vary across individuals, throughout development, and based on sex [20], [21]. Notably, this work found that the greatest variability in connectivity was observed in modulatory neurons [21]. Careful analysis also revealed a previously unrecognized degree of organization in the neuropil of the *C. elegans* nerve ring, which could influence functional interactions between neurons (Figure 2a) [22], [23]. In *Drosophila*, a sophisticated electron microscopy (EM) platform [24] was instrumental in the generation of the first complete map of synaptic connectivity [25**], [26], adding to an earlier connectivity map of the hemibrain [27]. This valuable atlas revealed key principles in brain organization; for example, densely connected groups of “rich club” neurons represent about 30% of neurons [28]. The *Drosophila* connectome is also defined for larvae [29] and the Ventral Nerve Cord [30]–[33], and more targeted work has mapped the mushroom body [34] and central complex [35], providing a wealth of maps to aid studies of neural circuits. For example, analysis of the mushroom body connectome found many more visual inputs than previously known, perhaps allowing for integration with other convergent sensory cues. The *Drosophila* connectomes are limited to chemical synapses thus far, owing to technical constraints. Future annotations of electrical synapses will provide additional insights into the synaptic organization of this nervous system.

Identifying connections is only part of the battle. To understand how these synapses function, we must determine the identities of the underlying neurotransmitters and receptors. The *Drosophila* connectome benefits from artificial neural network predictions of neurotransmitter identity based on EM data [36*]. In both species, transcriptomic atlases revealed the distribution of neurotransmitters and receptors [37]–[39**]. Fluorescent reporters have also been valuable in mapping the expression of neurotransmitters [40]–[42] and receptors [43], [44*]. This work has suggested that there may be widespread extrasynaptic signaling: many receptors are expressed in cells that do not receive synapses from a cell expressing that neurotransmitter. These genetic maps are essentially complete for the main *C. elegans* neurotransmitter systems at the level of cell types (for example, neurotransmitter identity is shown in Figure 2b). Completion of single synapse resolution maps in worms and flies – a major challenge for the future – will yield additional advances.

Another factor not accounted for in connectivity maps is neuropeptide signaling. Understanding of *C. elegans* neuropeptide systems expanded enormously with the identification of 461 novel ligand-receptor pairs out of over 55,000 possible pairs tested [45**]. These findings were

combined with single-cell sequencing data to construct a brain-wide neuropeptide signaling map. Compared to synaptic signaling, neuropeptide signaling is more decentralized and far denser, with >10-fold more connections [46**]. Neuropeptide networks also link the synaptically disconnected pharyngeal network to the central brain. In addition, different peptidergic systems feature dramatically different organizations: point-to-point signaling, autocrine signaling, and broadcasting architectures, which may enable different features of emergent activity (Figure 2d-f). Mapping of the neuropeptide network in *Drosophila* showed neuropeptide expression limited to clusters of neurons but receptor expression throughout the brain [43].

A central goal of future research will be to connect brain-wide activity and connectivity maps. In *C. elegans*, a recent investigation focused on comprehensively mapping the serotonin system's structure and function [47*]. The serotonergic NSM neuron is activated by food ingestion, and its non-synaptic release of serotonin induces slow locomotion and increased feeding behavior. A combination of approaches was used to identify the contributions of each of the six serotonin receptors to these behavioral changes, determine the neurons across the connectome that express these receptors, and investigate how serotonin release impacts brain-wide activity. Different receptors mediated behavioral responses to different patterns of serotonin release. In addition, knowledge of each neuron's serotonin receptor expression could partially predict how their activity changed during serotonin release, providing links between structure and function.

Another study recently attempted to bridge the gap between architecture and activity by assembling a map of functional connectivity [48**]. This work combined cell-specific optogenetic activation, whole brain imaging, and neuronal identification to quantify how perturbing each neuron's activity affects all other neurons' activities. Many relationships were identified between neurons that are not directly connected through synapses (example shown in Figure 2c). Additionally, many of these fast, functional connections were dependent on dense core vesicle release, providing evidence for functionally important extrasynaptic signaling that may be critical to understand brain-wide dynamics.

Efforts to build increasingly precise and accurate network models of fly and worm nervous systems are ongoing. In *Drosophila*, modeling constrained by connectome and neurotransmitter data generated novel hypotheses about sensory and motor pathways [49]. Many of these predictions held true in subsequent testing; for example, novel neurons predicted to be important for water responses indeed caused a change in water intake upon optogenetic perturbation. Accurate predictions of known visual system neurons were also generated by a neural network model of a smaller region, the optic lobe, which was built using constraints from the connectome and deep learning methods [50]. Still other work used behavioral data to train a deep neural network model, cleverly using behavior recordings from flies with silenced neurons to probe the visual courtship circuitry [51]. Current modeling efforts are limited by a lack of data about electrical signaling, receptor dynamics, neuromodulation, and individual neuron characteristics such as co-neurotransmitter release.

Although the work described here has produced a wealth of information about circuit organization and function, many challenges still need to be solved in order to interrogate circuit function at brain-wide scale. In the meantime, studies of specific circuits provide valuable intuition about how nervous system structure relates to function in the context of behavior.

Neural circuit motifs in invertebrate nervous systems

Our understanding of how neural circuits generate behavior has been greatly aided by case studies of individual circuits. Here, we discuss examples of studies that uncovered network motifs that contribute to defined features of animal behavior.

At first glance, the most straightforward pathways are point to point signals, where one neuron communicates directly to another. Hundreds, if not thousands, of such connections have been identified, often underlying specific sensorimotor responses. Recent studies have shown that even neuropeptides can act in this direct manner to impact behavior. For example, in worms, the neuropeptide *flp-1* promotes avoidance of pathogenic bacteria [52*]. Upon infection, this neuropeptide is released from one neuron class, called AVK, to promote avoidance behavior through a single receptor, *dmsr-7*, which functions in RIM/RIC neurons. *flp-1* is produced in other neurons and has other receptors; *dmsr-7* is similarly broadly expressed and has many ligands, yet a single connection within this complex network has a specific behavioral function. In *Drosophila* larvae, a similar motif was found when examining nociceptive responses to heat: the neuropeptide DSK acts via one receptor on a single cell type to inhibit heat avoidance [53].

Another common network motif is broadcasting or “one to many” signaling, where a single neuron signals to several downstream partners. This network logic is effective for behavioral outputs that require synchronization, such as changes in behavioral state. In *C. elegans*, stress induced sleep is regulated by the ALA neuron, which releases multiple neuropeptides to downregulate distinct behavioral features such as feeding, head movement, and locomotion [54], [55]. Broadcasting signals are also useful as teaching signals during learning. In *Drosophila* spatial learning, dopaminergic ExR2 neurons broadly innervate the head direction network and facilitate learning during rotational movements so that animals can update their internal representations of space [56].

There are several circuit architectures that can support coincidence detection. In *C. elegans*, “hub neurons” receiving convergent signals can weigh multiple sensory inputs and generate integrated behavioral responses. For example, in response to gentle touch, several mechanosensory neurons send concurrent signals via gap junctions to a single downstream neuron, RIH, which acts as a coincidence detector to direct avoidance behavior [57], [58]. Coincidence detection circuits are also central to learning. In *Drosophila*, dopaminergic DAN neurons that contain information about motor state, internal state, and even reward and punishment converge onto defined compartments of the mushroom body; coincident activation of specific DANs with olfactory-responsive Kenyon cells changes how Kenyon cells couple to mushroom body outputs and behavior [59]–[68].

Studies of defined neural circuits have also demonstrated that behavioral outputs are not always governed by a single, linear circuit. Degenerate signaling pathways can often exist, where different neural sources can lead to the same outcome. In *C. elegans*, feeding behavior can be initiated by several independent neurons [69]. This work is reminiscent of degeneracy in the stomatogastric ganglion (STG) of crustaceans, where many underlying circuit configurations can generate the same circuit outputs [70]–[72]. Work on the STG has also demonstrated that neurons can co-release several neuropeptides that have additive or antagonistic effects (reviewed in [73]). In flies and worms, release of different neurotransmitters from the same neuron can impact distinct behavioral outputs [74], [75], effectively allowing them to participate in multiple networks. In order to accurately capture brain-wide activity, we will need to consider that the underlying signaling can be flexible, degenerate, and multiplexed and may contain many interlinked network motifs that contribute to overall circuit function.

219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248

Conclusion

The recent developments reviewed here have opened an exciting new chapter in neuroscience research. Advances in hardware and software (reviewed in [76], [77]) have yielded bountiful data on neuronal activity in freely behaving animals under a variety of conditions. While similarities between worms and flies suggest that principles of brain-wide organization may span species, examining brain-wide activity in mammals is challenging. Recent advances have allowed for recording of over a million neurons in the mouse neocortex [78], but brain-wide imaging will require additional innovations. *C. elegans* now have an atlas of how most neurons encode spontaneous behavior [8], and activity maps for the larger *Drosophila* brain provide a global view of its dynamics. In addition, both species now have genetic maps of neurotransmitter and receptor expression, as well as maps of synaptic connections. Despite this wealth of data, we are still missing information crucial to our understanding of how these nervous systems function.

Going forward, future experiments will need to address the flexibility of how neural activity encodes behavior. Examining brain-wide responses across animals in different behavioral states, in defined sensory surroundings, or during motivated behaviors will show how neuronal encoding can change based on context (see [65*]). Our current understanding of flexibility and degeneracy derived from smaller circuits suggests that brain activity maps will not be fixed.

To fully integrate the activity and molecular maps, we need a more complete understanding of neurotransmitter system dynamics. Further studies about the timescales of neurotransmitter release, the spatial organization of heterogeneous receptors on a single neuron, the extent of extrasynaptic signaling, and the kinetics of different receptors will provide valuable information. Currently, these questions are typically addressed on a case-by-case basis, but large-scale approaches [48] may be well positioned to tackle some questions at scale. In addition, sensors for neuropeptides [79] and neurotransmitters [80], [81] may be useful to further address these problems. As future work expands our view of brain-wide dynamics and organization, it will be increasingly possible to create accurate models of brain activity to generate novel, testable predictions.




Figure 1: Brain wide recordings in *C. elegans* and *Drosophila* reveal how neurons and brain regions encode behavior features

a) Whole brain calcium imaging data collection in *C. elegans*. From top to bottom: Cartoon of *C. elegans*. The worm connectome, showing synaptic connections between neuronal cells (data from [19], [21]). Sample image of whole brain calcium imaging in a freely moving worm, showing pan-neuronal GCaMP and mNeptune in the head of a worm [8]. Heatmap of brain-wide activity during spontaneous behavior, with behavior quantification for velocity, feeding rate, and angular velocity in the same animal.

258 b) Whole brain calcium imaging data collection in *Drosophila*. From top to bottom: Cartoon of a
259 *Drosophila*. The flow of information via chemical synapses between different brain regions as
260 found in the *Drosophila* connectome [26]. Sample image of a fly brain, depicting representations
261 of behavior in different regions [12]. Three views show orthogonal slices through the brain of a
262 fly. Color values show correlations for each brain region with forward velocity and left or right
263 angular velocity. Heat map of brain-wide activity during spontaneous fly behavior, with
264 behavioral annotations and speed shown for the same animal [13].




Figure 2: Structural, genetic, and network maps of the *C. elegans* connectome

a-f) Six identical arrangements of neurons from the *C. elegans* connectome, where neurons are organized by sensorimotor layer (y-axis) and connectivity (x-axis). In each panel, neurons are colored according to different structural or functional features.

a) Neurons are colored based on their neuropil layer assignments as determined in [23].

“Unassigned” neurons span multiple strata.

b) Neurotransmitter expression in all *C. elegans* neurons (data summarized in [40]). Neurons with multiple colors release multiple neurotransmitters.

c) Sample result comparing functional and anatomical connectivity for a single neuron, SAADL (shown in yellow). Neurons in green had changes in activity upon SAADL stimulation [48]. Neurons in blue are synaptically connected to SAADL [19], [21].

d-f) Examples of three different signaling motifs found in the neuropeptidergic connectome of *C. elegans* (data on neuropeptide and receptor expression patterns from [39], [45], [46]).

d) Point to point signaling, with only a few neurons expressing either the neuropeptide *nlp-23* or its cognate receptor *gnrr-3* (data from [39], [45], [46]).

e) Broadcasting expression from a single neuron pair (HSN) releasing neuropeptide *flp-23* to many downstream partners expressing receptor *dmsr-7* (data from [39], [45], [46]). Interestingly, HSN expresses both the neuropeptide and receptor, representing a possible autocrine loop.

f) Convergent signals emanating from many neurons releasing *flp-5* are integrated by only a handful of neurons expressing its receptor *egl-6* (data from [39], [45], [46]).

286 **Acknowledgments**

287 We thank members of the Flavell lab for critical comments on the manuscript. We thank Thomas
288 Felt for assistance with figures. T.S.K. acknowledges funding from the MathWorks Science
289 Fellowship. S.W.F. acknowledges funding from NIH (GM135413, NS131457, and DC020484),
290 NSF (1845663), the Alfred P. Sloan Foundation, the McKnight Scholars Program, The JPB
291 Foundation, and The Picower Institute for Learning & Memory.

292

293 **Declaration of Interests**

294 The authors have no competing interests to declare.

295

- (1) Kaplan, H. S.; Salazar Thula, O.; Khoss, N.; Zimmer, M. Nested Neuronal Dynamics Orchestrate a Behavioral Hierarchy across Timescales. *Neuron* **2020**, *105* (3), 562-576.e9. <https://doi.org/10.1016/j.neuron.2019.10.037>.
 - (2) Kato, S.; Kaplan, H. S.; Schrödel, T.; Skora, S.; Lindsay, T. H.; Yemini, E.; Lockery, S.; Zimmer, M. Global Brain Dynamics Embed the Motor Command Sequence of *Caenorhabditis Elegans*. *Cell* **2015**, *163* (3), 656–669. <https://doi.org/10.1016/j.cell.2015.09.034>.
 - (3) Schrödel, T.; Prevedel, R.; Aumayr, K.; Zimmer, M.; Vaziri, A. Brain-Wide 3D Imaging of Neuronal Activity in *Caenorhabditis Elegans* with Sculpted Light. *Nat Methods* **2013**, *10* (10), 1013–1020. <https://doi.org/10.1038/nmeth.2637>.
 - (4) Hallinen, K. M.; Dempsey, R.; Scholz, M.; Yu, X.; Linder, A.; Randi, F.; Sharma, A. K.; Shaevitz, J. W.; Leifer, A. M. Decoding Locomotion from Population Neural Activity in Moving *C. Elegans*. *eLife* **2021**, *10*, e66135. <https://doi.org/10.7554/eLife.66135>.
 - (5) Nguyen, J. P.; Shipley, F. B.; Linder, A. N.; Plummer, G. S.; Liu, M.; Setru, S. U.; Shaevitz, J. W.; Leifer, A. M. Whole-Brain Calcium Imaging with Cellular Resolution in Freely Behaving *Caenorhabditis Elegans*. *Proceedings of the National Academy of Sciences* **2016**, *113* (8), E1074–E1081. <https://doi.org/10.1073/pnas.1507110112>.
 - (6) Venkatachalam, V.; Ji, N.; Wang, X.; Clark, C.; Mitchell, J. K.; Klein, M.; Tabone, C. J.; Florman, J.; Ji, H.; Greenwood, J.; Chisholm, A. D.; Srinivasan, J.; Alkema, M.; Zhen, M.; Samuel, A. D. T. Pan-Neuronal Imaging in Roaming *Caenorhabditis Elegans*. *Proceedings of the National Academy of Sciences* **2016**, *113* (8), E1082–E1088. <https://doi.org/10.1073/pnas.1507109113>.
 - *(7) Susoy, V.; Hung, W.; Witvliet, D.; Whitener, J. E.; Wu, M.; Park, C. F.; Graham, B. J.; Zhen, M.; Venkatachalam, V.; Samuel, A. D. T. Natural Sensory Context Drives Diverse Brain-Wide Activity during *C. Elegans* Mating. *Cell* **2021**, *184* (20), 5122-5137.e17. <https://doi.org/10.1016/j.cell.2021.08.024>.
- This study examined neuronal activity in the *C. elegans* male tail during mating, identifying specific neurons active during defined phases of the behavior and others that were engaged in several components.
- ** (8) Atanas, A. A.; Kim, J.; Wang, Z.; Bueno, E.; Becker, M.; Kang, D.; Park, J.; Estrem, C.; Kramer, T. S.; Baskoylu, S.; Mansinghka, V. K.; Flavell, S. W. Brain-Wide Representations of Behavior Spanning Multiple Timescales and States in *C. Elegans*. *Cell* **2023**. <https://doi.org/10.1016/j.cell.2023.07.035>.
- This study used whole brain imaging and encoder models to determine the behavioral features encoded by most neurons in the *C. elegans* brain.
- (9) Uzel, K.; Kato, S.; Zimmer, M. A Set of Hub Neurons and Non-Local Connectivity Features Support Global Brain Dynamics in *C. Elegans*. *Current Biology* **2022**, *32* (16), 3443-3459.e8. <https://doi.org/10.1016/j.cub.2022.06.039>.
 - (10) Aimon, S.; Katsuki, T.; Jia, T.; Grosenick, L.; Broxton, M.; Deisseroth, K.; Sejnowski, T. J.; Greenspan, R. J. Fast Near-Whole-Brain Imaging in Adult *Drosophila* during Responses to Stimuli and Behavior. *PLOS Biology* **2019**, *17* (2), e2006732. <https://doi.org/10.1371/journal.pbio.2006732>.
 - ** (11) Aimon, S.; Cheng, K. Y.; Gjorgjieva, J.; Grunwald Kadow, I. C. Global Change in Brain State during Spontaneous and Forced Walk in *Drosophila* Is Composed of Combined

- Activity Patterns of Different Neuron Classes. *eLife* **2023**, *12*, e85202.
<https://doi.org/10.7554/eLife.85202>.
- This study characterized whole brain activity in walking flies, comparing activity during spontaneous and forced walk. Calcium imaging was also conducted in specific populations of neurons to examine how their activity changes with different components of behavior.
- **(12) Brezovec, B. E.; Berger, A. B.; Hao, Y. A.; Chen, F.; Druckmann, S.; Clandinin, T. R. Mapping the Neural Dynamics of Locomotion across the *Drosophila* Brain. *Current Biology* **2024**. <https://doi.org/10.1016/j.cub.2023.12.063>.
- This study examined brain-wide activity in freely behaving flies, using a combination of analytical approaches to determine which brain regions are active during different behaviors spanning various timescales.
- **(13) Schaffer, E. S.; Mishra, N.; Whiteway, M. R.; Li, W.; Vancura, M. B.; Freedman, J.; Patel, K. B.; Voleti, V.; Paninski, L.; Hillman, E. M. C.; Abbott, L. F.; Axel, R. The Spatial and Temporal Structure of Neural Activity across the Fly Brain. *Nat Commun* **2023**, *14* (1), 5572. <https://doi.org/10.1038/s41467-023-41261-2>.
- This study examined brain wide activity during several *Drosophila* behaviors, examining neural representations of behavior across multiple timescales.
- (14) Pacheco, D. A.; Thiberge, S. Y.; Pnevmatikakis, E.; Murthy, M. Auditory Activity Is Diverse and Widespread throughout the Central Brain of *Drosophila*. *Nat Neurosci* **2021**, *24* (1), 93–104. <https://doi.org/10.1038/s41593-020-00743-y>.
- (15) Nichols, A. L. A.; Eichler, T.; Latham, R.; Zimmer, M. A Global Brain State Underlies *C. Elegans* Sleep Behavior. *Science* **2017**, *356* (6344), eaam6851. <https://doi.org/10.1126/science.aam6851>.
- (16) Tainton-Heap, L. A. L.; Kirszenblat, L. C.; Notaras, E. T.; Grabowska, M. J.; Jeans, R.; Feng, K.; Shaw, P. J.; van Swinderen, B. A Paradoxical Kind of Sleep in *Drosophila Melanogaster*. *Current Biology* **2021**, *31* (3), 578-590.e6. <https://doi.org/10.1016/j.cub.2020.10.081>.
- (17) Mann, K.; Gallen, C. L.; Clandinin, T. R. Whole-Brain Calcium Imaging Reveals an Intrinsic Functional Network in *Drosophila*. *Current Biology* **2017**, *27* (15), 2389-2396.e4. <https://doi.org/10.1016/j.cub.2017.06.076>.
- (18) Brezovec, L. E.; Berger, A. B.; Druckmann, S.; Clandinin, T. R. Neural Correlates of Future Volitional Action in *Drosophila*. *bioRxiv* September 9, 2023, p 2023.09.08.556917. <https://doi.org/10.1101/2023.09.08.556917>.
- (19) White, J. G.; Southgate, J. N.; Brenner, S. The Structure of the Nervous System of the Nematode *Caenorhabditis Elegans*. *Phil. Trans. R. Soc. Lond. B* **1986**, *314* (1165), 1–340. <https://doi.org/10.1098/rstb.1986.0056>.
- (20) Cook, S. J.; Jarrell, T. A.; Brittin, C. A.; Wang, Y.; Bloniarz, A. E.; Yakovlev, M. A.; Nguyen, K. C. Q.; Tang, L. T.-H.; Bayer, E. A.; Duerr, J. S.; Bülow, H. E.; Hobert, O.; Hall, D. H.; Emmons, S. W. Whole-Animal Connectomes of Both *Caenorhabditis Elegans* Sexes. *Nature* **2019**, *571* (7763), 63–71. <https://doi.org/10.1038/s41586-019-1352-7>.
- (21) Witvliet, D.; Mulcahy, B.; Mitchell, J. K.; Meirovitch, Y.; Berger, D. R.; Wu, Y.; Liu, Y.; Koh, W. X.; Parvathala, R.; Holmyard, D.; Schalek, R. L.; Shavit, N.; Chisholm, A. D.; Lichtman, J. W.; Samuel, A. D. T.; Zhen, M. Connectomes across Development Reveal

- Principles of Brain Maturation. *Nature* **2021**, 596 (7871), 257–261.
<https://doi.org/10.1038/s41586-021-03778-8>.
- (22) Brittin, C. A.; Cook, S. J.; Hall, D. H.; Emmons, S. W.; Cohen, N. A Multi-Scale Brain Map Derived from Whole-Brain Volumetric Reconstructions. *Nature* **2021**, 591 (7848), 105–110. <https://doi.org/10.1038/s41586-021-03284-x>.
- (23) Moyle, M. W.; Barnes, K. M.; Kuchroo, M.; Gonopolskiy, A.; Duncan, L. H.; Sengupta, T.; Shao, L.; Guo, M.; Santella, A.; Christensen, R.; Kumar, A.; Wu, Y.; Moon, K. R.; Wolf, G.; Krishnaswamy, S.; Bao, Z.; Shroff, H.; Mohler, W. A.; Colón-Ramos, D. A. Structural and Developmental Principles of Neuropil Assembly in *C. Elegans*. *Nature* **2021**, 591 (7848), 99–104. <https://doi.org/10.1038/s41586-020-03169-5>.
- (24) Zheng, Z.; Lauritzen, J. S.; Perlman, E.; Robinson, C. G.; Nichols, M.; Milkie, D.; Torrens, O.; Price, J.; Fisher, C. B.; Sharifi, N.; Calle-Schuler, S. A.; Kmecova, L.; Ali, I. J.; Karsh, B.; Trautman, E. T.; Bogovic, J. A.; Hanslovsky, P.; Jefferis, G. S. X. E.; Kazhdan, M.; Khairy, K.; Saalfeld, S.; Fetter, R. D.; Bock, D. D. A Complete Electron Microscopy Volume of the Brain of Adult *Drosophila Melanogaster*. *Cell* **2018**, 174 (3), 730–743.e22. <https://doi.org/10.1016/j.cell.2018.06.019>.
- *(25) Dorkenwald, S.; Matsliah, A.; Sterling, A. R.; Schlegel, P.; Yu, S.; McKellar, C. E.; Lin, A.; Costa, M.; Eichler, K.; Yin, Y.; Silversmith, W.; Schneider-Mizell, C.; Jordan, C. S.; Brittain, D.; Halageri, A.; Kuehner, K.; Ogedengbe, O.; Morey, R.; Gager, J.; Kruk, K.; Perlman, E.; Yang, R.; Deutsch, D.; Bland, D.; Sorek, M.; Lu, R.; Macrina, T.; Lee, K.; Bae, J. A.; Mu, S.; Nehoran, B.; Mitchell, E.; Popovych, S.; Wu, J.; Jia, Z.; Castro, M.; Kemnitz, N.; Ih, D.; Bates, A. S.; Eckstein, N.; Funke, J.; Collman, F.; Bock, D. D.; Jefferis, G. S. X. E.; Seung, H. S.; Murthy, M. Neuronal Wiring Diagram of an Adult Brain. bioRxiv, p. 2023.06.27.546656, July 11, 2023, doi: 10.1101/2023.06.27.546656.
- This study released the first full connectome of an adult *Drosophila*, identifying synaptic connections for over 100,000 neurons.
- (26) Schlegel, P.; Yin, Y.; Bates, A. S.; Dorkenwald, S.; Eichler, K.; Brooks, P.; Han, D. S.; Gkantia, M.; Santos, M. dos; Munnely, E. J.; Badalamente, G.; Capdevila, L. S.; Sane, V. A.; Pleijzier, M. W.; Tamimi, I. F. M.; Dunne, C. R.; Salgarella, I.; Javier, A.; Fang, S.; Perlman, E.; Kazimiers, T.; Jagannathan, S. R.; Matsliah, A.; Sterling, A. R.; Yu, S.; McKellar, C. E.; Costa, M.; Seung, H. S.; Murthy, M.; Hartenstein, V.; Bock, D. D.; Jefferis, G. S. X. E. A Consensus Cell Type Atlas from Multiple Connectomes Reveals Principles of Circuit Stereotypy and Variation. bioRxiv July 15, 2023. <https://doi.org/10.1101/2023.06.27.546055>.
- (27) Scheffer, L. K.; Xu, C. S.; Januszewski, M.; Lu, Z.; Takemura, S.; Hayworth, K. J.; Huang, G. B.; Shinomiya, K.; Maitlin-Shepard, J.; Berg, S.; Clements, J.; Hubbard, P. M.; Katz, W. T.; Umayam, L.; Zhao, T.; Ackerman, D.; Blakely, T.; Bogovic, J.; Dolafi, T.; Kainmueller, D.; Kawase, T.; Khairy, K. A.; Leavitt, L.; Li, P. H.; Lindsey, L.; Neubarth, N.; Olbris, D. J.; Otsuna, H.; Trautman, E. T.; Ito, M.; Bates, A. S.; Goldammer, J.; Wolff, T.; Svirskas, R.; Schlegel, P.; Neace, E.; Knecht, C. J.; Alvarado, C. X.; Bailey, D. A.; Ballinger, S.; Borycz, J. A.; Canino, B. S.; Cheatham, N.; Cook, M.; Dreher, M.; Duclos, O.; Eubanks, B.; Fairbanks, K.; Finley, S.; Forknall, N.; Francis, A.; Hopkins, G. P.; Joyce, E. M.; Kim, S.; Kirk, N. A.; Kovalyak, J.; Lauchie, S. A.; Lohff, A.; Maldonado, C.; Manley, E. A.; McLin, S.; Mooney, C.; Ndam, M.; Ogundeyi, O.; Okeoma, N.; Ordish, C.; Padilla, N.; Patrick, C. M.; Paterson, T.; Phillips, E. E.; Phillips, E. M.; Rampally, N.; Ribeiro, C.;

- Robertson, M. K.; Rymer, J. T.; Ryan, S. M.; Sammons, M.; Scott, A. K.; Scott, A. L.; Shinomiya, A.; Smith, C.; Smith, K.; Smith, N. L.; Sobeski, M. A.; Suleiman, A.; Swift, J.; Takemura, S.; Talebi, I.; Tarnogorska, D.; Tenshaw, E.; Tokhi, T.; Walsh, J. J.; Yang, T.; Horne, J. A.; Li, F.; Parekh, R.; Rivlin, P. K.; Jayaraman, V.; Costa, M.; Jefferis, G. S.; Ito, K.; Saalfeld, S.; George, R.; Meinertzhagen, I. A.; Rubin, G. M.; Hess, H. F.; Jain, V.; Plaza, S. M. A Connectome and Analysis of the Adult *Drosophila* Central Brain. *eLife* **2020**, *9*, e57443. <https://doi.org/10.7554/eLife.57443>.
- (28) Lin, A.; Yang, R.; Dorkenwald, S.; Matsliah, A.; Sterling, A. R.; Schlegel, P.; Yu, S.; McKellar, C. E.; Costa, M.; Eichler, K.; Bates, A. S.; Eckstein, N.; Funke, J.; Jefferis, G. S. X. E.; Murthy, M. Network Statistics of the Whole-Brain Connectome of *Drosophila*. bioRxiv July 29, 2023, p 2023.07.29.551086. <https://doi.org/10.1101/2023.07.29.551086>.
- (29) Winding, M.; Pedigo, B. D.; Barnes, C. L.; Patsolic, H. G.; Park, Y.; Kazimiers, T.; Fushiki, A.; Andrade, I. V.; Khandelwal, A.; Valdes-Aleman, J.; Li, F.; Randel, N.; Barsotti, E.; Correia, A.; Fetter, R. D.; Hartenstein, V.; Priebe, C. E.; Vogelstein, J. T.; Cardona, A.; Zlatić, M. The Connectome of an Insect Brain. *Science* **2023**, *379* (6636), eadd9330. <https://doi.org/10.1126/science.add9330>.
- (30) Azevedo, A.; Lesser, E.; Mark, B.; Phelps, J.; Elabbady, L.; Kuroda, S.; Sustar, A.; Moussa, A.; Kandelwal, A.; Dallmann, C. J.; Agrawal, S.; Lee, S.-Y. J.; Pratt, B.; Cook, A.; Skutt-Kakaria, K.; Gerhard, S.; Lu, R.; Kemnitz, N.; Lee, K.; Halageri, A.; Castro, M.; Ih, D.; Gager, J.; Tammam, M.; Dorkenwald, S.; Collman, F.; Schneider-Mizell, C.; Brittain, D.; Jordan, C. S.; Dickinson, M.; Pacureanu, A.; Seung, H. S.; Macrina, T.; Lee, W.-C. A.; Tuthill, J. C. Tools for Comprehensive Reconstruction and Analysis of *Drosophila* Motor Circuits. bioRxiv December 15, 2022, p 2022.12.15.520299. <https://doi.org/10.1101/2022.12.15.520299>.
- (31) Cheong, H. S. J.; Eichler, K.; Stüerner, T.; Asinof, S. K.; Champion, A. S.; Marin, E. C.; Oram, T. B.; Sumathipala, M.; Venkatasubramanian, L.; Namiki, S.; Siwanowicz, I.; Costa, M.; Berg, S.; Team, J. F. P.; Jefferis, G. S. X. E.; Card, G. M. Transforming Descending Input into Behavior: The Organization of Premotor Circuits in the *Drosophila* Male Adult Nerve Cord Connectome. bioRxiv January 26, 2024. <https://doi.org/10.1101/2023.06.07.543976>.
- (32) Marin, E. C.; Morris, B. J.; Stüerner, T.; Champion, A. S.; Krzeminski, D.; Badalamente, G.; Gkantia, M.; Dunne, C. R.; Eichler, K.; Takemura, S.; Tamimi, I. F. M.; Fang, S.; Moon, S. S.; Cheong, H. S. J.; Li, F.; Schlegel, P.; Berg, S.; Team, F. P.; Card, G. M.; Costa, M.; Shepherd, D.; Jefferis, G. S. X. E. Systematic Annotation of a Complete Adult Male *Drosophila* Nerve Cord Connectome Reveals Principles of Functional Organisation. bioRxiv June 6, 2023, p 2023.06.05.543407. <https://doi.org/10.1101/2023.06.05.543407>.
- (33) Takemura, S.; Hayworth, K. J.; Huang, G. B.; Januszewski, M.; Lu, Z.; Marin, E. C.; Preibisch, S.; Xu, C. S.; Bogovic, J.; Champion, A. S.; Cheong, H. S.; Costa, M.; Eichler, K.; Katz, W.; Knecht, C.; Li, F.; Morris, B. J.; Ordish, C.; Rivlin, P. K.; Schlegel, P.; Shinomiya, K.; Stüerner, T.; Zhao, T.; Badalamente, G.; Bailey, D.; Brooks, P.; Canino, B. S.; Clements, J.; Cook, M.; Duclos, O.; Dunne, C. R.; Fairbanks, K.; Fang, S.; Finley-May, S.; Francis, A.; George, R.; Gkantia, M.; Harrington, K.; Hopkins, G. P.; Hsu, J.; Hubbard, P. M.; Javier, A.; Kainmueller, D.; Korff, W.; Kovalyak, J.; Krzemiński, D.; Lauchie, S. A.; Lohff, A.; Maldonado, C.; Manley, E. A.; Mooney, C.; Neace, E.; Nichols, M.; Ogundeyi, O.; Okeoma, N.; Paterson, T.; Phillips, E.; Phillips, E. M.; Ribeiro, C.; Ryan, S. M.; Rymer, J. T.; Scott, A. K.; Scott, A. L.; Shepherd, D.; Shinomiya, A.; Smith, C.; Smith, N.;

- Suleiman, A.; Takemura, S.; Talebi, I.; Tamimi, I. F.; Trautman, E. T.; Umayam, L.; Walsh, J. J.; Yang, T.; Rubin, G. M.; Scheffer, L. K.; Funke, J.; Saalfeld, S.; Hess, H. F.; Plaza, S. M.; Card, G. M.; Jefferis, G. S.; Berg, S. A Connectome of the Male *Drosophila* Ventral Nerve Cord. *bioRxiv* June 6, 2023, p 2023.06.05.543757. <https://doi.org/10.1101/2023.06.05.543757>.
- (34) Li, F.; Lindsey, J. W.; Marin, E. C.; Otto, N.; Dreher, M.; Dempsey, G.; Stark, I.; Bates, A. S.; Pleijzier, M. W.; Schlegel, P.; Nern, A.; Takemura, S.; Eckstein, N.; Yang, T.; Francis, A.; Braun, A.; Parekh, R.; Costa, M.; Scheffer, L. K.; Aso, Y.; Jefferis, G. S.; Abbott, L. F.; Litwin-Kumar, A.; Waddell, S.; Rubin, G. M. The Connectome of the Adult *Drosophila* Mushroom Body Provides Insights into Function. *eLife* **2020**, *9*, e62576. <https://doi.org/10.7554/eLife.62576>.
- (35) Hulse, B. K.; Haberkern, H.; Franconville, R.; Turner-Evans, D.; Takemura, S.; Wolff, T.; Noorman, M.; Dreher, M.; Dan, C.; Parekh, R.; Hermundstad, A. M.; Rubin, G. M.; Jayaraman, V. A Connectome of the *Drosophila* Central Complex Reveals Network Motifs Suitable for Flexible Navigation and Context-Dependent Action Selection. *eLife* **2021**, *10*, e66039. <https://doi.org/10.7554/eLife.66039>.
- * (36) Eckstein, N.; Bates, A. S.; Champion, A.; Du, M.; Yin, Y.; Schlegel, P.; Lu, A. K.-Y.; Rymer, T.; Finley-May, S.; Paterson, T.; Parekh, R.; Dorkenwald, S.; Matsliah, A.; Yu, S.-C.; McKellar, C.; Sterling, A.; Eichler, K.; Costa, M.; Seung, S.; Murthy, M.; Hartenstein, V.; Jefferis, G. S. X. E.; Funke, J. Neurotransmitter Classification from Electron Microscopy Images at Synaptic Sites in *Drosophila Melanogaster*. *bioRxiv* May 10, 2023, p 2020.06.12.148775. <https://doi.org/10.1101/2020.06.12.148775>.
- This study used artificial neural networks to predict neurotransmitter identity for individual *Drosophila* synapses or neurons based on EM images.
- (37) Croset, V.; Treiber, C. D.; Waddell, S. Cellular Diversity in the *Drosophila* Midbrain Revealed by Single-Cell Transcriptomics. *eLife* **2018**, *7*, e34550. <https://doi.org/10.7554/eLife.34550>.
- (38) Davie, K.; Janssens, J.; Koldere, D.; De Waegeneer, M.; Pech, U.; Kreft, Ł.; Aibar, S.; Makhzami, S.; Christiaens, V.; Bravo González-Blas, C.; Poovathingal, S.; Hulselmans, G.; Spanier, K. I.; Moerman, T.; Vanspauwen, B.; Geurs, S.; Voet, T.; Lammertyn, J.; Thienpont, B.; Liu, S.; Konstantinides, N.; Fiers, M.; Verstreken, P.; Aerts, S. A Single-Cell Transcriptome Atlas of the Aging *Drosophila* Brain. *Cell* **2018**, *174* (4), 982-998.e20. <https://doi.org/10.1016/j.cell.2018.05.057>.
- **(39) Taylor, S. R.; Santpere, G.; Weinreb, A.; Barrett, A.; Reilly, M. B.; Xu, C.; Varol, E.; Oikonomou, P.; Glenwinkel, L.; McWhirter, R.; Poff, A.; Basavaraju, M.; Rafi, I.; Yemini, E.; Cook, S. J.; Abrams, A.; Vidal, B.; Cros, C.; Tavazoie, S.; Sestan, N.; Hammarlund, M.; Hobert, O.; Miller, D. M. Molecular Topography of an Entire Nervous System. *Cell* **2021**, *184* (16), 4329-4347.e23. <https://doi.org/10.1016/j.cell.2021.06.023>.
- This study quantified gene expression for all *C. elegans* neuron classes, providing valuable information on expression patterns of neurotransmitters, neuropeptides, and their receptors.
- (40) Gendrel, M.; Atlas, E. G.; Hobert, O. A Cellular and Regulatory Map of the GABAergic Nervous System of *C. Elegans*. *eLife* **2016**, *5*, e17686. <https://doi.org/10.7554/eLife.17686>.
- (41) Pereira, L.; Kratsios, P.; Serrano-Saiz, E.; Sheftel, H.; Mayo, A. E.; Hall, D. H.; White, J. G.; LeBoeuf, B.; Garcia, L. R.; Alon, U.; Hobert, O. A Cellular and Regulatory Map of the

- Cholinergic Nervous System of *C. Elegans*. *eLife* **2015**, *4*, e12432.
<https://doi.org/10.7554/eLife.12432>.
- (42) Serrano-Saiz, E.; Poole, R. J.; Felton, T.; Zhang, F.; De La Cruz, E. D.; Hobert, O. Modular Control of Glutamatergic Neuronal Identity in *C. Elegans* by Distinct Homeodomain Proteins. *Cell* **2013**, *155* (3), 659–673. <https://doi.org/10.1016/j.cell.2013.09.052>.
- (43) Deng, B.; Li, Q.; Liu, X.; Cao, Y.; Li, B.; Qian, Y.; Xu, R.; Mao, R.; Zhou, E.; Zhang, W.; Huang, J.; Rao, Y. Chemoconnectomics: Mapping Chemical Transmission in *Drosophila*. *Neuron* **2019**, *101* (5), 876–893.e4. <https://doi.org/10.1016/j.neuron.2019.01.045>.
- *(44) Yemini, E.; Lin, A.; Nejatbakhsh, A.; Varol, E.; Sun, R.; Mena, G. E.; Samuel, A. D. T.; Paninski, L.; Venkatachalam, V.; Hobert, O. NeuroPAL: A Multicolor Atlas for Whole-Brain Neuronal Identification in *C. Elegans*. *Cell* **2021**, *184* (1), 272–288.e11.
<https://doi.org/10.1016/j.cell.2020.12.012>.
- This study combined expression of several fluorophores to enable reliable neuronal identification across animals. The strain was used to identify sites of expression for metabotropic glutamate, acetylcholine, and GABA receptors. In addition, it has been widely used in other whole-brain calcium imaging studies.
- **(45) Beets, I.; Zels, S.; Vandeweyer, E.; Demeulemeester, J.; Caers, J.; Baytemur, E.; Courtney, A.; Golinelli, L.; Hasakioğulları, İ.; Schafer, W. R.; Vértés, P. E.; Mirabeau, O.; Schoofs, L. System-Wide Mapping of Peptide-GPCR Interactions in *C. Elegans*. *Cell Reports* **2023**, *42* (9), 113058. <https://doi.org/10.1016/j.celrep.2023.113058>.
- This study used a comprehensive reverse pharmacology screen to characterize 461 novel peptide-GPCR pairs in *C. elegans*, providing the most complete map of neuropeptide signaling to date for the *C. elegans* nervous system.
- **(46) Ripoll-Sánchez, L.; Watteyne, J.; Sun, H.; Fernandez, R.; Taylor, S. R.; Weinreb, A.; Bentley, B. L.; Hammarlund, M.; Miller, D. M.; Hobert, O.; Beets, I.; Vértés, P. E.; Schafer, W. R. The Neuropeptidergic Connectome of *C. Elegans*. *Neuron* **2023**, *111* (22), 3570–3589.e5. <https://doi.org/10.1016/j.neuron.2023.09.043>.
- This study generated a neuropeptidergic connectome for *C. elegans*, combining data on gene expression, ligand-receptor pairs, and connectivity to create an atlas.
- *(47) Dag, U.; Nwabudike, I.; Kang, D.; Gomes, M. A.; Kim, J.; Atanas, A. A.; Bueno, E.; Estrem, C.; Pugliese, S.; Wang, Z.; Towlson, E.; Flavell, S. W. Dissecting the Functional Organization of the *C. Elegans* Serotonergic System at Whole-Brain Scale. *Cell* **2023**, *186* (12), 2574–2592.e20. <https://doi.org/10.1016/j.cell.2023.04.023>.
- This study mapped the worm serotonin system, examining the behavioral role of each receptor, determining their sites of expression, and identifying how whole-brain activity changes with serotonin release.
- **(48) Randi, F.; Sharma, A. K.; Dvali, S.; Leifer, A. M. Neural Signal Propagation Atlas of *Caenorhabditis Elegans*. *Nature* **2023**, *623* (7986), 406–414. <https://doi.org/10.1038/s41586-023-06683-4>.
- This study assembled an atlas of functional connectivity in *C. elegans*, examining how optogenetic perturbation of individual neurons affects the activities of all other neurons.

- (49) Shiu, P. K.; Sterne, G. R.; Spiller, N.; Franconville, R.; Sandoval, A.; Zhou, J.; Simha, N.; Kang, C. H.; Yu, S.; Kim, J. S.; Dorkenwald, S.; Matsliah, A.; Schlegel, P.; Yu, S.; McKellar, C. E.; Sterling, A.; Costa, M.; Eichler, K.; Jefferis, G. S. X. E.; Murthy, M.; Bates, A. S.; Eckstein, N.; Funke, J.; Bidaye, S. S.; Hampel, S.; Seeds, A. M.; Scott, K. A. Leaky Integrate-and-Fire Computational Model Based on the Connectome of the Entire Adult *Drosophila* Brain Reveals Insights into Sensorimotor Processing. *bioRxiv* May 2, 2023, p 2023.05.02.539144. <https://doi.org/10.1101/2023.05.02.539144>.
- (50) Lappalainen, J. K.; Tschopp, F. D.; Prakhya, S.; McGill, M.; Nern, A.; Shinomiya, K.; Takemura, S.; Gruntman, E.; Macke, J. H.; Turaga, S. C. Connectome-Constrained Deep Mechanistic Networks Predict Neural Responses across the Fly Visual System at Single-Neuron Resolution. *bioRxiv* March 13, 2023, p 2023.03.11.532232. <https://doi.org/10.1101/2023.03.11.532232>.
- (51) Cowley, B. R.; Calhoun, A. J.; Rangarajan, N.; Pillow, J. W.; Murthy, M. One-to-One Mapping between Deep Network Units and Real Neurons Uncovers a Visual Population Code for Social Behavior. *bioRxiv* October 23, 2023. <https://doi.org/10.1101/2022.07.18.500505>.
- *(52) Marquina-Solis, J.; Vandeweyer, E.; Hawk, J.; Colón-Ramos, D. A.; Beets, I.; Bargmann, C. I. Peptidergic Signaling Controls the Dynamics of Sickness Behavior in *Caenorhabditis Elegans*. *bioRxiv* April 17, 2022, p 2022.04.16.488560. <https://doi.org/10.1101/2022.04.16.488560>.
- This study examined the role of a single neuropeptide in pathogen avoidance behavior, identifying its precise sites of release and action.
- (53) Oikawa, I.; Kondo, S.; Hashimoto, K.; Yoshida, A.; Hamajima, M.; Tanimoto, H.; Furukubo-Tokunaga, K.; Honjo, K. A Descending Inhibitory Mechanism of Nociception Mediated by an Evolutionarily Conserved Neuropeptide System in *Drosophila*. *eLife* **2023**, *12*, RP85760. <https://doi.org/10.7554/eLife.85760>.
- (54) Nath, R. D.; Chow, E. S.; Wang, H.; Schwarz, E. M.; Sternberg, P. W. C. *Elegans* Stress-Induced Sleep Emerges from the Collective Action of Multiple Neuropeptides. *Current Biology* **2016**, *26* (18), 2446–2455. <https://doi.org/10.1016/j.cub.2016.07.048>.
- (55) Nelson, M. D.; Lee, K. H.; Churgin, M. A.; Hill, A. J.; Van Buskirk, C.; Fang-Yen, C.; Raizen, D. M. FMRamide-like FLP-13 Neuropeptides Promote Quiescence Following Heat Stress in *Caenorhabditis Elegans*. *Current Biology* **2014**, *24* (20), 2406–2410. <https://doi.org/10.1016/j.cub.2014.08.037>.
- (56) Fisher, Y. E.; Marquis, M.; D'Alessandro, I.; Wilson, R. I. Dopamine Promotes Head Direction Plasticity during Orienting Movements. *Nature* **2022**, *612* (7939), 316–322. <https://doi.org/10.1038/s41586-022-05485-4>.
- (57) Chatzigeorgiou, M.; Schafer, W. R. Lateral Facilitation between Primary Mechanosensory Neurons Controls Nose Touch Perception in *C. Elegans*. *Neuron* **2011**, *70* (2), 299–309. <https://doi.org/10.1016/j.neuron.2011.02.046>.
- (58) Rabinowitch, I.; Chatzigeorgiou, M.; Schafer, W. R. A Gap Junction Circuit Enhances Processing of Coincident Mechanosensory Inputs. *Current Biology* **2013**, *23* (11), 963–967. <https://doi.org/10.1016/j.cub.2013.04.030>.
- (59) Aso, Y.; Hattori, D.; Yu, Y.; Johnston, R. M.; Iyer, N. A.; Ngo, T.-T.; Dionne, H.; Abbott, L.; Axel, R.; Tanimoto, H.; Rubin, G. M. The Neuronal Architecture of the Mushroom

- Body Provides a Logic for Associative Learning. *eLife* **2014**, 3, e04577. <https://doi.org/10.7554/eLife.04577>.
- (60) Berry, J. A.; Phan, A.; Davis, R. L. Dopamine Neurons Mediate Learning and Forgetting through Bidirectional Modulation of a Memory Trace. *Cell Rep* **2018**, 25 (3), 651–662.e5. <https://doi.org/10.1016/j.celrep.2018.09.051>.
- (61) Cohn, R.; Morante, I.; Ruta, V. Coordinated and Compartmentalized Neuromodulation Shapes Sensory Processing in *Drosophila*. *Cell* **2015**, 163 (7), 1742–1755. <https://doi.org/10.1016/j.cell.2015.11.019>.
- (62) Lin, S.; Oswald, D.; Chandra, V.; Talbot, C.; Huetteroth, W.; Waddell, S. Neural Correlates of Water Reward in Thirsty *Drosophila*. *Nat Neurosci* **2014**, 17 (11), 1536–1542. <https://doi.org/10.1038/nn.3827>.
- (63) Mao, Z.; Davis, R. Eight Different Types of Dopaminergic Neurons Innervate the *Drosophila* Mushroom Body Neuropil: Anatomical and Physiological Heterogeneity. *Frontiers in Neural Circuits* **2009**, 3.
- (64) Takemura, S.; Aso, Y.; Hige, T.; Wong, A.; Lu, Z.; Xu, C. S.; Rivlin, P. K.; Hess, H.; Zhao, T.; Parag, T.; Berg, S.; Huang, G.; Katz, W.; Olbris, D. J.; Plaza, S.; Umayam, L.; Aniceto, R.; Chang, L.-A.; Lauchie, S.; Ogundeyi, O.; Ordish, C.; Shinomiya, A.; Sigmund, C.; Takemura, S.; Tran, J.; Turner, G. C.; Rubin, G. M.; Scheffer, L. K. A Connectome of a Learning and Memory Center in the Adult *Drosophila* Brain. *eLife* **2017**, 6, e26975. <https://doi.org/10.7554/eLife.26975>.
- *(65) Zolin, A.; Cohn, R.; Pang, R.; Siliciano, A. F.; Fairhall, A. L.; Ruta, V. Context-Dependent Representations of Movement in *Drosophila* Dopaminergic Reinforcement Pathways. *Nat Neurosci* **2021**, 24 (11), 1555–1566. <https://doi.org/10.1038/s41593-021-00929-y>.
- This study examined neuronal activity during olfactory navigation in the *Drosophila* mushroom body, examining how representations of behavior in dopaminergic neurons change as behavioral strategy changes.
- (66) Aso, Y.; Sitaraman, D.; Ichinose, T.; Kaun, K. R.; Vogt, K.; Belliart-Guérin, G.; Plaçais, P.-Y.; Robie, A. A.; Yamagata, N.; Schnaitmann, C.; Rowell, W. J.; Johnston, R. M.; Ngo, T.-T. B.; Chen, N.; Korff, W.; Nitabach, M. N.; Heberlein, U.; Preat, T.; Branson, K. M.; Tanimoto, H.; Rubin, G. M. Mushroom Body Output Neurons Encode Valence and Guide Memory-Based Action Selection in *Drosophila*. *eLife* **2014**, 3, e04580. <https://doi.org/10.7554/eLife.04580>.
- (67) Lim, J.; Fernandez, A. I.; Hinojos, S. J.; Aranda, G. P.; James, J.; Seong, C.-S.; Han, K.-A. The Mushroom Body D1 Dopamine Receptor Controls Innate Courtship Drive. *Genes, Brain and Behavior* **2018**, 17 (2), 158–167. <https://doi.org/10.1111/gbb.12425>.
- (68) Tsao, C.-H.; Chen, C.-C.; Lin, C.-H.; Yang, H.-Y.; Lin, S. *Drosophila* Mushroom Bodies Integrate Hunger and Satiety Signals to Control Innate Food-Seeking Behavior. *eLife* **2018**, 7, e35264. <https://doi.org/10.7554/eLife.35264>.
- (69) Trojanowski, N. F.; Padovan-Merhar, O.; Raizen, D. M.; Fang-Yen, C. Neural and Genetic Degeneracy Underlies *Caenorhabditis Elegans* Feeding Behavior. *Journal of Neurophysiology* **2014**, 112 (4), 951–961. <https://doi.org/10.1152/jn.00150.2014>.
- (70) Prinz, A. A.; Bucher, D.; Marder, E. Similar Network Activity from Disparate Circuit Parameters. *Nat Neurosci* **2004**, 7 (12), 1345–1352. <https://doi.org/10.1038/nn1352>.

- (71) Rodriguez, J. C.; Blitz, D. M.; Nusbaum, M. P. Convergent Rhythm Generation from Divergent Cellular Mechanisms. *J. Neurosci.* **2013**, *33* (46), 18047–18064. <https://doi.org/10.1523/JNEUROSCI.3217-13.2013>.
- (72) Powell, D. J.; Marder, E.; Nusbaum, M. P. Perturbation-Specific Responses by Two Neural Circuits Generating Similar Activity Patterns. *Current Biology* **2021**, *31* (21), 4831–4838.e4. <https://doi.org/10.1016/j.cub.2021.08.042>.
- (73) Nusbaum, M. P.; Blitz, D. M.; Marder, E. Functional Consequences of Neuropeptide and Small-Molecule Co-Transmission. *Nat Rev Neurosci* **2017**, *18* (7), 389–403. <https://doi.org/10.1038/nrn.2017.56>.
- (74) Huang, Y.-C.; Luo, J.; Huang, W.; Baker, C. M.; Gomes, M. A.; Meng, B.; Byrne, A. B.; Flavell, S. W. A Single Neuron in *C. Elegans* Orchestrates Multiple Motor Outputs through Parallel Modes of Transmission. *Current Biology* **2023**, *33* (20), 4430–4445.e6. <https://doi.org/10.1016/j.cub.2023.08.088>.
- (75) Imambocus, B. N.; Zhou, F.; Formozov, A.; Wittich, A.; Tenedini, F. M.; Hu, C.; Sauter, K.; Macarenhas Varela, E.; Herédia, F.; Casimiro, A. P.; Macedo, A.; Schlegel, P.; Yang, C.-H.; Miguel-Aliaga, I.; Wiegert, J. S.; Pankratz, M. J.; Gontijo, A. M.; Cardona, A.; Soba, P. A Neuropeptidergic Circuit Gates Selective Escape Behavior of *Drosophila* Larvae. *Current Biology* **2022**, *32* (1), 149–163.e8. <https://doi.org/10.1016/j.cub.2021.10.069>.
- (76) Randi, F.; Leifer, A. M. Measuring and Modeling Whole-Brain Neural Dynamics in *Caenorhabditis Elegans*. *Current Opinion in Neurobiology* **2020**, *65*, 167–175. <https://doi.org/10.1016/j.conb.2020.11.001>.
- (77) Lin, A.; Witvliet, D.; Hernandez-Nunez, L.; Linderman, S. W.; Samuel, A. D. T.; Venkatachalam, V. Imaging Whole-Brain Activity to Understand Behaviour. *Nat Rev Phys* **2022**, *4* (5), 292–305. <https://doi.org/10.1038/s42254-022-00430-w>.
- (78) Demas, J.; Manley, J.; Tejera, F.; Barber, K.; Kim, H.; Traub, F. M.; Chen, B.; Vaziri, A. High-Speed, Cortex-Wide Volumetric Recording of Neuroactivity at Cellular Resolution Using Light Beads Microscopy. *Nat Methods* **2021**, *18* (9), 1103–1111. <https://doi.org/10.1038/s41592-021-01239-8>.
- (79) Ding, K.; Han, Y.; Seid, T. W.; Buser, C.; Karigo, T.; Zhang, S.; Dickman, D. K.; Anderson, D. J. Imaging Neuropeptide Release at Synapses with a Genetically Engineered Reporter. *eLife* **2019**, *8*, e46421. <https://doi.org/10.7554/eLife.46421>.
- (80) Wan, J.; Peng, W.; Li, X.; Qian, T.; Song, K.; Zeng, J.; Deng, F.; Hao, S.; Feng, J.; Zhang, P.; Zhang, Y.; Zou, J.; Pan, S.; Shin, M.; Venton, B. J.; Zhu, J. J.; Jing, M.; Xu, M.; Li, Y. A Genetically Encoded Sensor for Measuring Serotonin Dynamics. *Nat Neurosci* **2021**, *24* (5), 746–752. <https://doi.org/10.1038/s41593-021-00823-7>.
- (81) Unger, E. K.; Keller, J. P.; Altermatt, M.; Liang, R.; Matsui, A.; Dong, C.; Hon, O. J.; Yao, Z.; Sun, J.; Banala, S.; Flanigan, M. E.; Jaffe, D. A.; Hartanto, S.; Carlen, J.; Mizuno, G. O.; Borden, P. M.; Shivange, A. V.; Cameron, L. P.; Sinning, S.; Underhill, S. M.; Olson, D. E.; Amara, S. G.; Lang, D. T.; Rudnick, G.; Marvin, J. S.; Lavis, L. D.; Lester, H. A.; Alvarez, V. A.; Fisher, A. J.; Prescher, J. A.; Kash, T. L.; Yarov-Yarovoy, V.; Gradinaru, V.; Looger, L. L.; Tian, L. Directed Evolution of a Selective and Sensitive Serotonin Sensor via Machine Learning. *Cell* **2020**, *183* (7), 1986–2002.e26. <https://doi.org/10.1016/j.cell.2020.11.040>.