

Optical and near-infrared spectroscopy of quasars at $z > 6.5$: public data release and composite spectrum

Silvia Onorato^{1,2}   Joseph F. Hennawi^{1,2}   Jan-Torge Schindler³  Jinyi Yang⁴  Feige Wang⁴  Aaron J. Barth⁵   Eduardo Bañados⁶   Anna-Christina Eilers⁷   Sarah E. I. Bosman^{6,8}   Frederick B. Davies⁶  Bram P. Venemans¹  Chiara Mazzucchelli⁹  Silvia Belladitta^{6,10}  Fabio Vito¹⁰  Emanuele Paolo Farina¹¹  Irham T. Andika^{12,13}   Xiaohui Fan¹⁴  Fabian Walter⁶  Roberto Decarli¹⁰   Masafusa Onoue^{15,16,17}   and Riccardo Nanni¹ 

¹Leiden Observatory, Leiden University, P.O. Box 9513, NL-2300 RA Leiden, the Netherlands

²Department of Physics, Broida Hall, University of California, Santa Barbara, Santa Barbara, CA 93106-9530, USA

³Hamburger Sternwarte, Universität Hamburg, Gojenbergsweg 112, D-21029 Hamburg, Germany

⁴Department of Astronomy, University of Michigan, 1085 S. University Ave., Ann Arbor, MI 48109, USA

⁵Department of Physics and Astronomy, 4129 Frederick Reines Hall, University of California, Irvine, CA 92697-4575, USA

⁶Max Planck Institut für Astronomie, Königstuhl 17, D-69117, Heidelberg, Germany

⁷MIT Kavli Institute for Astrophysics and Space Research, 77 Massachusetts Avenue, Cambridge, 02139, Massachusetts, USA

⁸Institute for Theoretical Physics, Heidelberg University, Philosophenweg 12, D-69120 Heidelberg, Germany

⁹Instituto de Estudios Astrofísicos, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Avenida Ejercito Libertador 441, Santiago, Chile

¹⁰INAF–Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via Gobetti 93/3, I-40129 Bologna, Italy

¹¹Gemini Observatory, NSF's NOIRLab, 670 N A'ohoku Place, Hilo, Hawai'i 96720, USA

¹²Technical University of Munich, TUM School of Natural Sciences, Department of Physics, James-Franck-Str 1, D-85748 Garching, Germany

¹³Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str 1, D-85748, Garching, Germany

¹⁴Steward Observatory, University of Arizona, 933 N. Cherry Ave, Tucson, AZ 85721, USA

¹⁵Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU, WPI), The University of Tokyo Institutes for Advanced Study, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

¹⁶Center for Data-Driven Discovery, Kavli IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

¹⁷Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, P.R. China

Accepted 2025 May 12. Received 2025 May 9; in original form 2024 June 11

ABSTRACT

We present optical and near-infrared (NIR) spectroscopic observations for a sample of 45 quasars at $6.50 < z \leq 7.64$ with absolute magnitudes at 1450 \AA in the range $-28.82 \leq M_{1450} \leq -24.13$ and their composite spectrum. The median redshift and M_{1450} of the quasars in the sample are $z_{\text{median}} = 6.71$ and $M_{1450,\text{median}} \simeq -26.1$, respectively. The NIR spectra are taken with Echelle spectrographs, complemented with additional data from optical long slit instruments, and then reduced consistently using the open-source Python-based spectroscopic data reduction pipeline `PypeIt`. The median of the mean signal-to-noise ratios per 110 km s^{-1} pixel in the J , H , and K band [$\text{median } \langle \text{SNR}_\lambda \rangle$] is $\text{median } \langle \text{SNR}_J \rangle = 9.7$, $\text{median } \langle \text{SNR}_H \rangle = 10.3$, and $\text{median } \langle \text{SNR}_K \rangle = 11.7$; demonstrating the good data quality. This work presents the largest medium-/moderate-resolution sample of quasars at $z > 6.5$ from ground-based instruments. Despite the diversity in instrumental set-ups and spectral quality, the data set is uniformly processed and well-characterized, making it ideally suited for several scientific goals, including the study of the quasar proximity zones and damping wings, the Ly α forest, the intergalactic medium's metal content, as well as other properties such as the distribution of SMBH masses and Eddington ratios. Our composite spectrum is compared to others at both high and low z from the literature, showing differences in the strengths of many emission lines, probably due to differences in luminosity among the samples, but a consistent continuum slope, which proves that the same spectral features are preserved in quasars at different redshift ranges.

Key words: methods: data analysis – techniques: spectroscopic – galaxies: active – quasars: supermassive black holes – cosmology : early Universe.

* E-mail: onorato@strw.leidenuniv.nl

1 INTRODUCTION

Discoveries of high-redshift quasars ($z > 6$) have uncovered the presence of exceptionally supermassive black holes (SMBHs), ranging from approximately 10^8 to $10^{10} M_\odot$, in the early stages of the Universe (Wu et al. 2015; Bañados et al. 2018; Matsuoka et al. 2019a; Onoue et al. 2019; Shen et al. 2019; Yang et al. 2020; Wang et al. 2021a; see Fan, Bañados & Simcoe 2023 for a recent review). This motivates inquiry into the rapid growth mechanisms enabling these SMBHs to reach billions of solar masses within an extraordinarily brief time frame, less than 1 Byr after the big bang.

Various theoretical models proposing different seed black hole masses (10^2 – $10^6 M_\odot$) and highly efficient accretion modes present potential explanations for the formation and growth of early SMBHs (see Woods et al. 2019 and Inayoshi, Visbal & Haiman 2020 for recent reviews). Rigorous observations of an extensive sample of $z > 6.5$ quasars are imperative to test and refine these models, advancing our comprehension of SMBH formation and evolution. Recent advancements in deep imaging surveys, coupled with the enhanced capabilities of near-infrared (NIR) spectroscopy on large telescopes, have substantially expanded the sample size of quasars with $z > 6$ to more than 200 (e.g. Fan et al. 2023). This has extended the quasar frontier beyond $z > 7.5$, moving deep into the epoch of reionization (EoR; e.g. Mortlock et al. 2011; Venemans et al. 2013, 2015; Bañados et al. 2016, 2018; Jiang et al. 2016; Mazzucchelli et al. 2017; Wang et al. 2018, 2019, 2021a; Fan et al. 2019; Matsuoka et al. 2019a, b; Reed et al. 2019; Yang et al. 2019, 2020, 2021). Furthermore, recent observations from the *JWST* have already uncovered new high- z active galactic nuclei with lower luminosity and black hole mass, promising to deliver additional breakthroughs in our understanding (e.g. Goulding et al. 2023; Kokorev et al. 2023, 2024; Larson et al. 2023; Andika et al. 2024; Fujimoto et al. 2024; Furtak et al. 2024; Greene et al. 2024; Maiolino et al. 2024, b; Matthee et al. 2024; Pérez-González et al. 2024; Labbe et al. 2025).

Our work of collecting high-quality spectra marks a significant milestone as the most extensive medium-/moderate-resolution collection of quasar data at very high redshifts obtained through ground-based instruments. Despite the diversity in spectral resolution and signal-to-noise ratios (SNR), the data set is consistently processed, ensuring reproducibility and reliability for various scientific objectives. These include the exploration of the final phases of the EoR through the study of the Ly α forest (Fan et al. 2006; Becker et al. 2015; Bosman et al. 2022), investigations into the intergalactic medium's (IGM) metal content (e.g. Davies et al. 2023), the broad-line region chemical abundances (e.g. Lai et al. 2022), and analyses of properties such as the distribution of SMBH masses and Eddington ratios (Yang et al. 2021; Farina et al. 2022; Mazzucchelli et al. 2023). Also, the creation of a composite spectrum can be extremely helpful in evaluating the continuum and line properties of quasar spectra, investigating the average rest-frame UV quasar spectral properties and their possible evolution with redshift (Vanden Berk et al. 2001; Selsing et al. 2016; Mazzucchelli et al. 2017; Meyer, Bosman & Ellis 2019; Schindler et al. 2020; Yang et al. 2021). Moreover, constructing a composite spectrum enhances the detection of spectral features that may be difficult to identify in individual spectra due to their low SNR. This approach also facilitates the identification of objects that deviate from the template and aids in modelling and fitting spectral energy distributions.

Other key analyses involving high-quality spectra aim to place constraints on the average fraction of neutral hydrogen ($\langle x_{\text{HI}} \rangle$) at the EoR and on the radiative efficiency of the earliest SMBHs. They can be performed by starting with the reconstruction of the

quasar's intrinsic blue side of the spectrum from the observed red side employing, e.g. principal component analysis (PCA) continuum modelling (e.g. Davies et al. 2018a, b; Davies, Hennawi & Eilers 2019; Bosman et al. 2021) so that we can systematically study quasar proximity zones and lifetimes t_Q (see Fan et al. 2006; Eilers et al. 2017, 2020; Davies, Hennawi & Eilers 2020; Satyavolu et al. 2023), proximate damped Ly α systems (see Bañados et al. 2019; Andika et al. 2020, 2022), and Ly α damping wings (Miralda-Escudé 1998; Davies et al. 2018a; Ďurovčíková et al. 2024; Greig et al. 2024).

This paper is the first of a series aimed at exploiting the wealth of information that could be obtained from this sample (e.g. the study of the quasar proximity zone sizes in Paper II, Onorato et al. 2025). Here, we present and make publicly available the optical and NIR spectra of 45 quasars at $6.50 < z \leq 7.64$, and the composite rest-frame UV/optical spectrum. We describe the data set including the quasar sample, the instruments, and their properties, with the data reduction in Section 2. Section 3 details the spectral calibration procedure and presents the main properties of the sample. We show a comparison between this sample and other spectroscopic data releases of high- z quasars in Section 4. We discuss the mean quasar composite spectrum in Section 5, and we conclude this work with a summary of the paper, presented in Section 6. All results below refer to a Lambda cold dark matter cosmology with parameters $\Omega_\Lambda = 0.7$, $\Omega_m = 0.3$, and $h = 0.7$; all magnitudes are reported in the AB system.

2 THE DATA SET

2.1 Quasar sample

We start this analysis by compiling a list of quasars, both known from the literature and unpublished, and applying a redshift cut at $z > 6.5$. We check the individual telescope archives for medium-/moderate-resolution spectroscopy of these sources from Echelle spectrographs, and then also long slit instruments if the Ly α region is not sufficiently covered. We avoid collecting data of faint objects ($M_{1450} \gtrsim -25$), even if we do not apply a strict M_{1450} cut as long as the luminosity of the source does not strongly compromise the data reduction. Thus, our catalogue includes 45 quasars in the redshift range $6.50 < z \leq 7.64$ ($z_{\text{median}} = 6.71$) and in the magnitude range $-28.82 \leq M_{1450} \leq -24.13$ ($M_{1450,\text{median}} \simeq -26.1$, see Fig. 1 and Table 1). The full names of the quasars are in Table 1, while short names are adopted throughout the paper. Out of the 45 objects, 11 are classified in the literature as broad absorption lines (BAL) quasars, defined as those with absorption lines with full width at half-maximum (FWHM) $\gtrsim 2000 \text{ km s}^{-1}$. These BALs sources are J0313–1806, J0038–1527, J1243+0100,¹ J0839+3900, J2348–3054, J0246–5219, J0910–0414, J0923+0402, J0706+2921, J1526–2050, and J0439+1634, as flagged in Table 1. We include three new unpublished high- z quasars, J0410–0139 (Bañados et al. 2025), J1917+5003, and J0430–1445 (Belladitta et al. 2025). This last source also seems to be a possible BAL, leading to a final number of 12 BALs in this work and hence a BAL fraction of $\simeq 27$ per cent. The fraction of BAL quasars measured by Bischetti et al. (2022) in the XQR-30 sample ($5.8 \lesssim z \lesssim 6.6$) is $\simeq 40$ per cent. They point out that other works collecting $z \gtrsim 5.7$ quasar spectra found a BAL fraction of 16–24 per cent (Shen et al. 2019; Schindler et al. 2020; Yang et al.

¹Identified as a possible BAL in the discovery paper (Matsuoka et al. 2019a), but the spectroscopy quality does not lead to a clear classification.




Figure 1. Panel (a): distribution of J -band photometry used to scale the spectra as a function of z for all the 45 quasars in this sample. The legend is unique and split between the two plots, showing the sources colour-coded with decreasing z . The full symbols mark the quasars for which J -band photometry is available, while the open symbols mark those for which it is not (we report Y -band photometry for J1243+0100 and J0923+0753, and K_p -band photometry for J1058+2930). The error bars show the uncertainties on both z and the photometric measurements. Panel (b): distribution of M_{1450} as a function of z for all the 45 quasars in this sample. The error bars show the uncertainties on z . The circles mark the M_{1450} computed from the spectrum scaled with J -, Y -, or K_p -band photometry. The stars mark the quasars for which M_{1450} from spectroscopy is not reliable because of the low SNR of the spectrum or appearance of BAL features. In the case of J1243+0100, M_{1450} is from the discovery paper (Matsuoka et al. 2019a), while for J0910–0414 and J0923+0402, we follow the method described in Appendix B. Panel (c): histogram of the redshift distribution of the sample, with bins of size 0.05. The dashed red line represents the median redshift ($z_{\text{median}} = 6.71$). Panel (d): histogram of the M_{1450} distribution of the sample, with bins of size 0.2. The dashed red line represents the median M_{1450} ($M_{1450,\text{median}} \simeq -26.1$).

2021), and suggest these could be lower limits on the actual BAL fraction, as they are based on spectra with lower resolution and SNR. Similarly, our BAL fraction may also be a lower limit due to the heterogeneous data quality. Table 2 lists the remaining *published* $z > 6.50$ sources not included in the sample, stating for each quasar

their redshift and the reason for exclusion (i.e. $M_{1450} \gtrsim -25$ or no Echelle data).

2.2 Instruments

We obtain visible (VIS) and NIR spectroscopy of our sample from the data archives of three main Echelle spectrographs: Gemini/GNIRS (Gemini Near-Infrared Spectrograph; Elias et al. 2006a, b), Keck/NIRES (Near-Infrared Echelle Spectrometer; Wilson et al. 2004), and VLT/X-Shooter (Vernet et al. 2011), both VIS and NIR arms, except for J1342+0928, whose spectrum is entirely covered by the NIR arm. For some quasars, part of the spectrum redward of Ly α is in the optical, and hence not covered by the NIR arm, so we have complemented them using observations from additional long slit instruments, such as Gemini/GMOS (Gemini Multi-Object Spectrographs; Hook et al. 2004), Keck/DEIMOS (DEep Imaging Multi-Object Spectrograph; Faber et al. 2003), Keck/LRIS (Low-Resolution Imaging Spectrometer; Oke et al. 1995; Rockosi et al. 2010), and LBT/MODS (Multi-Object Double Spectrographs; Pogge et al. 2010). A complete list of the instruments used to observe each quasar is available in Table 1.

2.2.1 Dither sequences

All GNIRS and NIRES observations were executed following an ABBA dither sequence except for the last two frames of J0313–1806, which are AABB. The X-Shooter observations were acquired with different dither sequences (i.e. ABBA, ABAB, AABB), but because of the long average exposure time per frame ($\simeq 600$ –1800 s) and the consequent change of the sky conditions, they were treated as ABCD during the data reduction instead of performing usual image differencing. The only X-Shooter frames where a dither sequence of ABBA was used while performing the reduction are those for J1110–1329 because of the short exposure time of 300 s. This applies mostly for the NIR arm since VIS data are usually not reduced doing image differencing, but using a sky model.

2.2.2 Wavelength coverage, slit widths, R , FWHM, and $d\nu_{\text{pix}}$

The GNIRS data provide full wavelength coverage of the $YJHK$ bands (0.81 – 2.52 μm). Slit widths of 0.68 or 1.00 arcsec were used, with mean resolution of $R \simeq 1200$ – 1000 or $\text{FWHM} \simeq 280$ km s^{-1} and pixel size of $d\nu_{\text{pix}} \simeq 86$ km s^{-1} . The NIRES observations have a coverage of 0.94 – 2.47 μm through a fixed 0.55 arcsec slit, where the mean resolution is $R \simeq 2700$ or $\text{FWHM} \simeq 111$ km s^{-1} and pixel size of $d\nu_{\text{pix}} \simeq 38$ km s^{-1} . The X-Shooter data cover the wavelength range 0.55 – 1.02 μm in the VIS arm and 1.02 – 2.48 μm in the NIR arm. The slit width varies among programs but is typically 0.9 arcsec in the VIS and 0.6 arcsec in the NIR arm. The mean resolution in VIS is $R \simeq 8900$ or $\text{FWHM} \simeq 34$ km s^{-1} and pixel size $d\nu_{\text{pix}} \simeq 11$ km s^{-1} , while in NIR, $R \simeq 8100$ or $\text{FWHM} \simeq 37$ km s^{-1} and $d\nu_{\text{pix}} \simeq 13$ km s^{-1} . J0319–1008 is complemented with R400 grating GMOS-N observations, covering the wavelength range 0.65 – 1.15 μm . The slit width is 1.00 arcsec, with a mean resolution of $R \simeq 3200$ or $\text{FWHM} \simeq 94$ km s^{-1} , and pixel size of $d\nu_{\text{pix}} \simeq 51$ km s^{-1} . J0706+2921 is complemented with 830G grating DEIMOS observations, covering the wavelength range 0.65 – 1.00 μm . The slit width is 1.00 arcsec, with a mean resolution of $R \simeq 7900$ or $\text{FWHM} \simeq 38$ km s^{-1} , and pixel size of $d\nu_{\text{pix}} \simeq 17$ km s^{-1} . J0218+0007 is complemented using grating 600/10000 LRIS red observations, covering the wavelength range

Table 1. Information on the 45 quasars in this sample, sorted by decreasing z . The details on the columns are provided at the end of Section 3.4.

Name	Instrument (arms)	$t_{\text{exp}} (\text{s})$	z	z_{method}	z_{ref}	J_{AB}	M_{1450}	Discovery	$\langle \text{SNR}_i \rangle$	$\langle \text{SNR}_H \rangle$	$\langle \text{SNR}_K \rangle$
J031343.89–180636.40 ^a	GNIRS/NIRES	27 300/15 840	7.6423	[C II]	1	20.92 ± 0.13	–26.208	1	15.9	18.8	27.2
J134208.110+092838.610	X-Shooter (NIR)	(81 600)	7.5413	[C II]	2	20.64 ± 0.08	–26.336	3	33.5	43.7	13
J100758.264+211529.207	GNIRS/NIRES	(21 000/7920)	7.5149	[C II]	4	20.20 ± 0.18	–26.818	4	18.5	19.5	24.2
J112001.480+064124.300	X-Shooter (NIR/VIS)	(114 000/113 440)	7.0851	[C II]	2	20.17 ± 0.07	–26.565	5	44.7	53.7	308.7
J124353.930+010038.500 ^a	GNIRS	10 944	7.07	Mg II	26	23.57 ± 0.08 ^b	–24.130 ^d	26	1.4	1	1.3
J003836.097–152723.636 ^a	GNIRS/X-Shooter (NIR/VIS)	15 300/(1 5600/17 912)	7.0340	[C II]	6	19.69 ± 0.07	–27.030	7	24.8	24.6	22.6
J025216.640–050331.810	X-Shooter (NIR/VIS)	(28 800/31 200)	7.0006	[C II]	6	20.19 ± 0.07	–26.625	8	29.1	39.7	12.7
J041009–013919 ^e	NIRES	9360	6.9964	[C II]	27	20.75 ± 0.07	–25.858	27	6	5.9	8.1
J083946.880+390011.440 ^a	GNIRS	16 800	6.9046	Mg II	9	20.39 ± 0.20	–26.214	10	27.2	29.9	43.9
J234833.340–305410.000 ^a	X-Shooter (NIR/VIS)	(9200/8783)	6.9018	[C II]	11	21.10 ± 0.08	–25.224	12	7.5	10.3	4.2
J024655.902–521949.950 ^a	X-Shooter (NIR/VIS)	(24 000/24 000)	6.8876	[C II]	6	21.20 ± 0.14	–25.301	8	6.1	11.7	4.9
J002031.470–365341.800	X-Shooter (NIR/VIS)	(4800/4800)	6.8661	[C II]	29	20.42 ± 0.10	–25.999	13	6	8.3	3.6
J191729.984+500313.540 ^e	NIRES/DEMODS	12 720/26 400	6.853	Mg II	28	20.66 ± 0.05	–26.208	28	5.2	5.4	5.3
J221100.601–632055.845	X-Shooter (NIR/VIS)	(36 000/37 961)	6.8449	[C II]	6	21.27 ± 0.18	–25.470	8	9	12.9	4.8
J031941.660–100846.000	NIRES/GMOS	18 750/15 300	6.8275	[C II]	6	20.88 ± 0.30	–25.480	8	4.9	6.1	12.2
J041128.628–090749.700	NIRES/DEMODS	57 600/30 000	6.8260	[C II]	6	20.02 ± 0.14	–26.490	10	17.4	18	19.6
J112925.368+184624.330	X-Shooter (NIR/VIS)	(12 000/12 000)	6.823	Mg II	9	20.90 ± 0.11	–25.421	14	6.6	8.2	3.2
J010953.130–304726.300	X-Shooter (NIR/VIS)	(21 600/21 600)	6.7909	[C II]	11	21.28 ± 0.14	–25.200	12	6.7	9.3	3.7
J082931.979+411740.870	GNIRS	12 600	6.7773	Mg II	9	20.26 ± 0.15	–26.154	10	12.2	9.2	13.2
J021847.040+000715.200	NIRES/LRS	57 600/35 583	6.7700	[C II]	6	21.08 ± 0.30	–25.896	9.15	2.8	3.3	3.6
J110421.580+213428.850	GNIRS	7200	6.7662	[C II]	6	19.91 ± 0.11	–26.506	10	16.1	14.3	17.6
J010133.651+165630.180	GNIRS	13 200	6.7289	[C II]	6	21.06 ± 0.13	–25.346	10	8.9	7.3	10.7
J043043.660–144541.210 ^{a,e}	GNIRS	12 000	6.7142	Mg II	28	20.78 ± 0.18	–25.656	28	8.4	7.5	11.7
J083737.830+492900.600	GNIRS	15 600	6.702	Mg II	9	20.21 ± 0.17	–26.069	10	26.4	23.7	36.2
J200241.594–301321.690	GNIRS	3600	6.6876	[C II]	6	19.97 ± 0.16	–26.622	9	9.7	6.4	8
J092358.997+075349.107	GNIRS	7200	6.6817	[C II]	6	21.25 ± 0.26 ^b	–25.687	9	3.2	2	2.9
J104819.690–010940.210	X-Shooter (NIR/VIS)	(4800/4800)	6.6759	[C II]	16	20.87 ± 0.06	–25.864	17	5.1	6.7	2.9
J22255.150+293032.040	GNIRS/X-Shooter (NIR/VIS)	48 000/(14 400/14 400)	6.6666	[C II]	31	20.46 ± 0.12	–25.831	18	10.6	9.2	8.5
J2110219.230–143853.3860	GNIRS	10 200/5760	6.6645	[C II]	6	21.15 ± 0.20	–25.421	10	9.4	8.5	10.4
J121627.584+451910.675	GNIRS	3600	6.66	Mg II	9	21.02 ± 0.13	–25.518	10	4.8	3.6	5.3
J091054.555–041406.846 ^a	GNIRS/NIRES	36 000/36 000	6.6363	[C II]	6	20.23 ± 0.07	–26.397 ^f	10	9.8	11.4	15.6
J092347.117+040254.580 ^a	X-Shooter (NIR/VIS)	(43 200/43 200)	6.6330	[C II]	6	20.14 ± 0.08	–26.524 ^f	10.19	24.4	48.5	23.1
J002429.772+391318.980	GNIRS	13 800	6.621	[C II]	20	21.28 ± 0.48	–25.316	21	8.5	6.1	9.7
J030516.920–315056.000	X-Shooter (NIR/VIS)	(16 800/16 800)	6.6139	[C II]	30	20.60 ± 0.05	–25.690	12	12.3	15.6	6.3
J070626.379+292105.460 ^a	NIRES/DEMODS	14 960/17 800	6.6037	[C II]	6	19.14 ± 0.05	–27.410	10	24.3	25.6	34.2
J213233.160+121755.260	X-Shooter (NIR/VIS)	(33 600/35 809)	6.5881	[C II]	20	19.74 ± 0.03	–26.914	20	30.6	35.3	16.5
J152637.784–203000.660 ^a	GNIRS	(43 200/46 234)	6.5864	[C II]	16	19.60 ± 0.08	–27.030	20	33	45.6	18.9
J233807.032+214358.170	GNIRS	4500	6.586	[C II]	29	20.75 ± 0.30	–25.816	9	3.4	2.4	2.8
J135088.918+5011332.600	GNIRS	7200	6.5851	[C II]	6	20.41 ± 0.16	–26.075	10	12.8	11.6	16.2
J100810.720+292044.703	NIRES/DEMODS	36 000/28 800	6.5846	[C II]	22	20.56 ± 0.05 ^e	–26.039	9	5	4.9	5.7
J092220.560+000722.900	GNIRS	9600	6.5646	[C II]	6	21.11 ± 0.11	–25.319	19	9	8.1	12

Table 1 – continued

Name	Instrument (arms)	$t_{\text{exp}}(\text{s})$	z	z_{method}	z_{ref}	J_{AB}	M_{1450}	Discovery	$\langle \text{SNR}_i \rangle$	$\langle \text{SNR}_{\text{H}} \rangle$	$\langle \text{SNR}_{\text{K}} \rangle$
J022601.870+030259.280	X-Shooter (NIR/VIS)	(14 400/23 520)	6.5405	[C II]	30	19.40 \pm 0.09	-27.192	18	26.5	30.6	11.7
J022426.540+471129.400	X-Shooter (NIR/VIS)	(31 200/33 360)	6.5222	[C II]	6	19.73 \pm 0.05	-26.663	24	27.2	40.7	20
J043947.098+163415.81 ^{a,g}	X-Shooter (NIR/VIS)	(54 720/53 656)	6.5192	[C II]	6	17.47 \pm 0.02	-28.822	25	207.7	285.8	149.5
J111033.960+132945.600	X-Shooter (NIR/VIS)	(34 800/29 520)	6.5148	[C II]	16	20.59 \pm 0.18	-26.079	18	4.8	7.6	3.7

Notes: Ref: 1 – Wang et al. (2021a); 2 – Venemans et al. (2017); 3 – Bañados et al. (2018); 4 – Yang et al. (2020); 5 – Mortlock et al. (2011); 6 – Wang et al. (2018); 7 – Yang et al. (2019); 9 – Yang et al. (2021); 10 – Wang et al. (2019); 11 – Venemans et al. (2016); 12 – Venemans et al. (2013); 13 – Reed et al. (2019); 14 – Bañados et al. (2021); 15 – Matsuoka et al. (2022); 16 – Decarli et al. (2018); 17 – Wang et al. (2017); 18 – Venemans et al. (2015); 19 – Matsuoka et al. (2018); 20 – Mazzucchelli et al. (2017); 21 – Tang et al. (2017); 22 – Wang et al. (2024); 23 – Bañados et al. (2015); 24 – Reed et al. (2017); 25 – Fan et al. (2019); 26 – Matsuoka et al. (2019a); 27 – Bañados et al. (2025); 28 – Belladitta et al. (2025); 29 – Bouwens et al. in preparation; 30 – Venemans et al. (2020); 31 – Yang et al. (2023).

^aBAL quasar. ^b Y_{AB} magnitude. ^c K_{pAB} magnitude. ^d M_{1450} from Matsuoka et al. (2019a). ^eunpublished quasar. ^f M_{1450} calculated as described in Appendix B. ^glensed quasar.

Table 2. Remaining *published* quasars at $z > 6.50$ that are not included in the sample. We report their short names, redshift, and the main reason why they are excluded from the analysis.

Quasar	z	Reason for exclusion
J2356+0017	7.012	Faint
J1317+0017	6.94	Faint
J1349+0156	6.94	Faint
J1609+5328	6.92	Faint
J2210+0304	6.88	Faint
J0214+0232	6.829	Faint
J0112+0110	6.82	Faint
J1429–0104	6.796	Faint
J1310–0050	6.796	Faint
J1612+5559	6.78	Faint
J0244–5008	6.7306	No Echelle data
J0229–0808	6.7249	No Echelle data
J0213–0626	6.7214	Faint
J1344+0128	6.72	Faint
J1338+0018	6.701	Faint
J1205+0000	6.699	Faint
J0001+0000	6.69	Faint
J1231+0052	6.69	Faint
J1130+0450	6.68	Faint
J2252+0402	6.67	Faint
J0910+0056	6.65	Faint
J1035+0324	6.63	Faint
J1023+0044	6.63	Faint
J1450–0144	6.628	Faint
J0919+0512	6.62	Faint
J1732+6531	6.6	No Echelle data
J1114+0215	6.55	Faint
J1440+0019	6.549	Faint
J0525–2406	6.5397	No Echelle data
J1545+4232	6.5	Faint

0.70–1.03 μm . The slit width is 1.00 arcsec, with a mean resolution of $R \simeq 1700$ or $\text{FWHM} \simeq 176 \text{ km s}^{-1}$, and pixel size of $d_{\text{pix}} \simeq 56 \text{ km s}^{-1}$. J0411–0907, J1917+5003, and J1058+2930 are complemented by red grating MODS observations (both MODS1 and MODS2), covering the wavelength range 0.42–1.12 μm . The slit width is 1.00 and 1.2 arcsec, with a mean resolution of $R \simeq 2700 – 2300$ or $\text{FWHM} \simeq 111 \text{ km s}^{-1}$, and pixel size of $d_{\text{pix}} \simeq 33 \text{ km s}^{-1}$. All of these properties of the instruments are summarized in Table 3.

2.3 Data reduction

All spectra are reduced with the open-source Python-based Spectroscopic Data Reduction Pipeline `PypeIt`,² using versions between 1.7.1 and 1.14.1 (Prochaska et al. 2020a, b). The pipeline performs image processing, including gain correction, bias subtraction, dark subtraction, and flat fielding. It uses supplied flat-field images to automatically trace the Echelle orders and correct for the detector illumination. Construction of the wavelength solutions and the wavelength tilt models are based on either arc (for VIS instruments) or science frames (i.e. using sky OH lines, for NIR spectrographs). Cosmic rays are removed with the L. A. COSMIC algorithm (van Dokkum 2001). The sky subtraction is based on the standard A–B mode and a B-spline fitting procedure that is performed to further clean up the sky line residuals following Bochanski et al. (2009). Optimal extraction (Horne 1986) is performed to generate

²<https://github.com/pypeit/PypeIt>

Table 3. Properties of the instruments used to collect the data. The black line separates the NIR from the VIS arm.

Instrument	Dither sequence	$\Delta\lambda$ (μm)	Slit width (arcsec)	R	FWHM (km s $^{-1}$)	dv_{pix} (km s $^{-1}$) ^a
GNIRS	ABBA	0.81–2.52	0.68/1.00	$\simeq 1200/\simeq 1000$	$\simeq 280$	$\simeq 86$
NIRES	ABBA	0.94–2.47	0.55	$\simeq 2700$	$\simeq 111$	$\simeq 38$
X-Shooter NIR	varies	1.02–2.48	0.6	$\simeq 8100$	$\simeq 37$	$\simeq 13$
X-Shooter VIS	–	0.55–1.02	0.9	$\simeq 8900$	$\simeq 34$	$\simeq 11$
GMOS	–	0.65–1.15	1.00	$\simeq 3200$	$\simeq 94$	$\simeq 51$
DEIMOS	–	0.65–1.00	1.00	$\simeq 7900$	$\simeq 38$	$\simeq 17$
LRIS	–	0.70–1.03	1.00	$\simeq 1700$	$\simeq 176$	$\simeq 56$
MODS	–	0.42–1.12	1.00/1.2	$\simeq 2700/\simeq 2300$	$\simeq 111$	$\simeq 33$

Note.^aThe pixel sizes reported are those of each individual spectrograph, while the ones used to re-bin the spectra during the co-addition procedure are reported in Section 2.4, and chosen to be coarser than the coarsest dv_{pix} between the involved instruments.

1D science spectra. We clarify that `PypeIt` uses a nearest grid point interpolation algorithm for the extraction and co-addition of spectra to avoid correlations between pixels. The extracted spectra are flux calibrated with sensitivity functions derived from the observations of spectroscopic standard stars. All flux-calibrated 1D spectra of each quasar are then co-added to achieve higher SNR and corrected for telluric absorption using `PypeIt`. In particular, during the co-addition procedure, both spectra and noise vectors are iteratively rescaled to a preliminary mean spectrum, then they are optimally weighted and combined. This procedure is continued in a loop with the rescaling factors converging to unity as the procedure is iterated. To conclude, a telluric model is fit to correct the absorbed science spectrum up to a best-fitting PCA model (Davies et al. 2018b) of said spectrum. The telluric model is based on telluric model grids produced from the Line-By-Line Radiative Transfer Model (LBLRTM4; Clough et al. 2005; Gullikson, Dodson-Robinson & Kraus 2014).

All the `PypeIt` files to reproduce the reduction are publicly available in a GitHub repository.³

2.4 Co-add of spectra from different instruments (or arms)

As reported in Table 1 and in Section 2.2, some quasars are observed with more than one spectrograph and/or in more arms. Here, we describe how we treat these spectra after the flux calibration with sensitivity function since they have different resolutions, wavelength grids, and pixel sizes.

(i) X-Shooter VIS–NIR: two different approaches were used depending on the stage of development of `PypeIt`. We co-add the majority of the quasars with versions between 1.8.2 and 1.11.1, using the Echelle `combspec` `PypeIt` routine, getting a unique spectrum in a wavelength grid of [5410, 24 770] Å and pixel size of $dv_{\text{pix}} \simeq 13$ km s $^{-1}$. Few quasars were reduced using recent `PypeIt` versions (after 1.12.3), which now support the option of editing a unique file (`.coadd1d`) to combine the 1D spectra from multiple exposures of the same object, in case we work with the same type of spectrograph (i.e. Echelle), containing both arms together. The final spectrum has the same characteristics as those obtained with the first method.

(ii) GNIRS–NIRES: we co-add them using a single `coadd1d` file. J0313–1806 and J1007+2115 are co-added on to a common grid covering [9410, 24 690] Å with a pixel size of $dv_{\text{pix}} \simeq 90$ km s $^{-1}$. We co-add J0910–0414 and J2102–1458 by requiring a pixel size of $dv_{\text{pix}} \simeq 90$ km s $^{-1}$, but without constraints on the wavelength

range. We get a final spectrum that covers [8240–25 200] Å and has $dv_{\text{pix}} \simeq 90$ km s $^{-1}$.

(iii) GNIRS–X-Shooter (VIS–NIR): we co-add them using a unique `coadd1d` file for all the GNIRS, X-Shooter VIS, and NIR frames. The final spectra cover [5410, 25 200] Å with a pixel size of $dv_{\text{pix}} \simeq 90$ km s $^{-1}$.

(iv) NIRES–GMOS: we co-add all the Echelle long slit spectra using the multi `combspec` `PypeIt` routine, which works with the final 1D spectrum in both instruments, getting an ultimate spectrum with wavelength coverage of [8000, 24 700] Å and pixel size of $dv_{\text{pix}} \simeq 55$ km s $^{-1}$.

(v) NIRES–DEIMOS: using multi `combspec` as described above, we get a final spectrum covering [8000, 24 700] Å with pixel size of $dv_{\text{pix}} \simeq 40$ km s $^{-1}$.

(vi) NIRES–LRIS: using multi `combspec` as described above, we get a final spectrum covering [8000, 24 700] Å with pixel size of $dv_{\text{pix}} \simeq 60$ km s $^{-1}$.

(vii) NIRES–MODS: using multi `combspec` as described above, we get a final spectrum covering [8000, 24 700] Å with pixel size of $dv_{\text{pix}} \simeq 40$ km s $^{-1}$.

In the end, when Echelle frames are co-added together, the telluric correction described above (at the end of Section 2.3) is applied to the final stacked spectrum; when Echelle long slit are co-added, the individual spectra of both instruments are already telluric corrected, and no further correction is required. In principle, telluric correction should be applied before heliocentric correction, since telluric absorption occurs in the Earth’s atmospheric frame, not the heliocentric vacuum frame. However, for many of the faint quasars in this work, individual exposures have low SNR, making accurate telluric correction difficult using the model-based approach in `PypeIt`. Moreover, our data were often taken on different nights, and aligning exposures in the Earth’s rest frame would shift the quasar spectrum, complicating outlier rejection. We therefore apply telluric correction to the co-added, heliocentric-frame spectra as a compromise, accepting minor line broadening to ensure sufficient SNR for accurate modelling and outlier removal. Also, this broadening effect is more relevant for bright sources, and less critical for the faint quasars in our sample.

3 QUASAR SAMPLE PROPERTIES

3.1 Redshift

In this paper, we account for uncertainties in the systemic redshift of every quasar depending on the emission line used to determine its redshift. Systemic redshifts are challenging quantities to determine because of various factors, like the broad widths of emission lines,

³https://github.com/enigma-igm/onorato24_hiz_qsos

Gunn–Peterson absorption (Gunn & Peterson 1965), and offsets between different ionization lines (Gaskell 1982; Tytler & Fan 1992; Vanden Berk et al. 2001; Richards et al. 2002; Shen et al. 2016). Also, most quasars show winds and strong internal motions, which displace many of the emission lines far from the systemic redshift of the host galaxy. Thus, following Eilers et al. (2017), we decide to assign a redshift error of $\Delta v = 100 \text{ km s}^{-1}$ for the most precise determinations of the location of the quasar, such as those using emission lines from the atomic gas reservoir of the host galaxy itself ([C II] lines). However, we emphasize that obtaining this measurement is not straightforward: the typical velocity dispersion (σ_v) observed within quasar host galaxies is around 300 km s^{-1} for detections of at least $\text{SNR} > 5$. The statistical error on the mean redshift is approximately $\sigma_v/\text{SNR} \approx 60 \text{ km s}^{-1}$. Thus, our adopted uncertainty of 100 km s^{-1} is just a conservative estimate. This assumption is based on the expectation that the quasar resides within the gravitational potential traced by the cool interstellar medium. For quasars with a redshift measurement from low-ionization broad emission lines, such as Mg II, we assume a redshift error of $\Delta v = 390 \text{ km s}^{-1}$, to account for the dispersion between the redshift of the Mg II line and the redshift of the host galaxy (e.g. Mazzucchelli et al. 2017; Schindler et al. 2020). This value is supported by previous literature and further confirmed by our own analysis on a sub-sample of our sources with both Mg II and [C II] redshifts, for which we find a median velocity shift of $\Delta v \simeq -388 \text{ km s}^{-1}$, in excellent agreement with Schindler et al. (2020). Table 1 reports redshifts, methods, and references for every quasar.

These Δv will be adopted as uncertainties on the redshift (and thus on the position of the Ly α line) to achieve our next scientific goal of studying the proximity zones of the quasars in this sample (Onorato et al. 2025). From the literature (z_{Ref} in Table 1), we also get values of Δz for every quasar, but we do not report them here. They are available online at the link at the end of Section 2.3, and we use them as uncertainties on z in Fig. 1.

3.2 Absolute flux calibration

The flux calibration of the spectra obtained by the data reduction is relative: every spectrum is corrected using a spectroscopic standard star, but the flux values at the different wavelengths may differ from the true flux. This could be due to slit losses and non-photometric sky conditions at the time of observations. Tests of absolute flux calibration have been performed in this analysis, tying the spectra to the photometric data in the Y , J , H , and K bands, where all or part of them are available. However, since there is not a good match in all the photometric bands at once, a definitive method is still an object of discussion. Also, we point out that the photometry is not taken together with the spectra, and this could introduce uncertainty in the absolute flux calibration, as some of these quasars might have varied between the spectroscopic and photometric observations.

For this paper, the reduced spectrum of each quasar is scaled using its J -band magnitude, or the Y and K_p band ones if the J band is not available (as in the case of J1243+0100, J0923+0753, and J1058+2930), all in the AB system; an example for all bands is shown in Fig. A1 and the approach is described in Appendix A. We also evaluate the consistency of this scaling by comparing synthetic photometry from the flux-scaled spectra to available Y -, H -, and K -band magnitudes. We find that the typical discrepancy is around 10 per cent, with a maximum deviation of ~ 40 per cent (see Appendix A). These NIR data come from different archival surveys: the UKIRT Infrared Deep Sky Survey (UKIDSS; Lawrence et al. 2007), the UKIRT Hemisphere Survey (UHS; Dye et al. 2018),

and the VISTA Hemisphere Survey (VHS; McMahon et al. 2013). The Y -band photometry of J1243+0100 comes from the Hyper Suprime-Cam Subaru Strategic Program survey (Aihara et al. 2018), as reported in the discovery paper (Matsuoka et al. 2019a). The J -band photometry of J0410–0139, J0430–1445, and J2132+1217 is taken from SofI (Son of ISAAC; Moorwood, Cuby & Lidman 1998), at the New Technology Telescope (NTT) in La Silla, and that of J1129+1846 and J1917+5003 comes from NOTCam,⁴ at the Nordic Optical Telescope (NOT). The K_p -band photometry of J1058+2930 is gotten from the acquisition image taken with Nires, whose guider camera uses a Mauna Kea K' (K prime) filter.⁵ Since the transmission curve of this filter is not available, we use the one from Keck/NIRC2 K_p to perform the flux scaling (see the bottom panel of Fig. A1), as it is very similar to the Mauna Kea K' . Most of these magnitudes are collected from Ross & Cross (2020) and available online. All J -band (or Y and K_p) magnitudes are displayed in panel (a) of Fig. 1 as a function of redshift z for each quasar, and the final flux-scaled spectra are plotted as a function of wavelength in Fig. 2.

A detailed list of all the filters that were used to acquire every magnitude is available at the link given at the end of Section 2.3, while the final flux-calibrated spectra are available online as explained in the Data Availability Section. An example of the format of the FITS files, selecting a random row, for two spectra reduced using different `PyPsfIt` versions is shown in Table 4.

3.3 Absolute magnitudes at 1450 Å

We calculate M_{1450} using the flux-scaled spectra⁶ of the quasars themselves. From every spectrum in the observed frame and knowing the redshift of each quasar, we compute the rest-frame wavelength.

We determine the apparent AB magnitude at 1450 Å from the median flux between 1445 and 1455 Å, converted to Jansky ($f_{\nu,1450}$), using the Pogson law: $m_{1450} = -2.5 \cdot \log_{10}(f_{\nu,1450}) + 8.9$. We continue calculating the luminosity per unit frequency (L_{ν}) at 1450 Å with equation (1), where d_L is the luminosity distance to the object at a redshift z :

$$L_{\nu} = \frac{4\pi d_L^2 \cdot f_{\nu,1450}}{1+z} \quad (1)$$

We find the absolute AB magnitude at 1450 Å using equation (2) from the luminosity per unit frequency L_{ν} , where d_0 is the reference distance (10 pc), and 3631 Jy is the zero-point flux density in the AB system:

$$M_{1450} = -2.5 \cdot \log_{10} \left(\frac{L_{\nu}}{4\pi d_0^2 \cdot 3631 \text{ Jy}} \right) \quad (2)$$

The results from this method are compared with other values from the literature, showing good agreement. For this reason, we trust the estimates we obtain for all quasars, with only three exceptions. In the case of J1243+0100, the spectrum is of poor quality as the source is faint and has a low SNR (see the values at the last three columns of Table 1). As a consequence, this makes the result difficult to trust, so we adopt the M_{1450} from Matsuoka et al. (2019a), in which they measured it from the best-fitting power-law continuum. In the

⁴Both SofI and NOTCam are dedicated follow-up.

⁵<https://www2.keck.hawaii.edu/inst/nires/guider.html>

⁶As we scale almost all the spectra using the J band, which is the photometric band closest to $1450 \cdot (1+z)$ Å, any possible uncertainties in the spectral shape should not result in a significant source of error on M_{1450} .




Figure 2. Spectra of all the 45 quasars in this sample, sorted by decreasing z (as reported in Table 1). Every subplot shows the spectrum (colour-coded with z) and the associated error (grey) in the observed wavelength range $[8245, 22980]$ Å, the short name, and z of the quasar. The light-grey bands cover $[13500, 14500]$ Å, and $[18200, 19300]$ Å, indicating the regions affected by strong telluric absorption, where the spectra are masked. All the spectra shown here are smoothed for display purposes. The box containing the name and the redshift of the quasars is coloured in light beige to mark the BALs.


case of J0910-0414 and J0923+0402, the BAL features visible in their spectra create a bias in the measurement of M_{1450} . For this reason, we correct the estimate by matching these spectra with a ‘reference spectrum’ with a trustworthy continuum shape, such as the composite spectrum created from the sample in this paper (shown in Fig. 5). The match is performed by eye and shown in Fig. B1, with a detailed description of the method followed reported in Appendix B.

We show the absolute magnitude values M_{1450} as a function of redshift z for our sample in panel (b) of Fig. 1, with its histogram in panel (d).

3.4 SNR of the spectra

We compute the SNR of the spectra in this sample considering three different wavelength ranges (the J , H , and K bands). We want to properly sample the spectra and avoid possible biases introduced by telluric absorption regions. For these reasons, we define the ranges in which we can compute the SNR in a very conservative manner, avoiding regions affected by absorption: $[11000, 13400]$, $[14500, 17950]$, and $[19650, 22400]$ Å.

To make the calculation and have a comparison among the quality of the spectra that is as fair as possible, we follow the following steps:

**Figure 2.** *Continued.*

- (i) We move the spectra to the rest frame, dividing the wavelengths by $(1 + z)$;
- (ii) We define a new wavelength grid: $[1040, 3332]\text{\AA}$ in steps of $d\nu_{\text{pix}} = 110 \text{ km s}^{-1}$ (set to be coarser than the coarsest pixel scale, in velocity, among all the instruments used to create the sample);
- (iii) We re-bin the spectra into the new wavelength grid;
- (iv) We finally compute the mean SNR in the three different wavelength ranges ($\langle \text{SNR}_{\lambda} \rangle$, where $\lambda = J, H, K$), also shifted to the rest frame according to the redshift of the quasar considered.

Two examples showing the method described above, considering the highest and lowest redshift quasar in the sample, are visible in Fig. C1. The results of this analysis are shown in the last three columns of Table 1, and in the histograms in Fig. 3, where we generate 25 logarithmically spaced bins between the minimum and maximum of

all the $\langle \text{SNR}_{\lambda} \rangle$ values in the sample. The majority of the spectra have $\langle \text{SNR}_{J,H} \rangle$ in the range [5, 10], and $\langle \text{SNR}_K \rangle$ between 0–5 and 10–15. To quantify the overall data quality we compute the median $\langle \text{SNR}_{\lambda} \rangle$ in the three wavelength ranges. We have median $\langle \text{SNR}_J \rangle = 9.7$, median $\langle \text{SNR}_H \rangle = 10.3$, and median $\langle \text{SNR}_K \rangle = 11.7$ (reported as red dashed vertical lines in Fig. 3).

The different properties for every quasar are reported in Table 1, which lists the name of the quasars, the instruments (and arms), the exposure time for each quasar in every instrument, their redshifts, the method adopted to determine the redshift, the reference for this measurement, the *J*-band photometry in the AB system, the magnitude (M_{1450}), the reference for the discovery of each quasar, and the mean SNR of the spectra in the three different wavelength ranges. The references and some notes are listed at the bottom of the table.




Figure 2. Continued.

Table 4. Example of the internal structure of the FITS files, at a random row, for two of the spectra in this sample. The first column is the wavelengths in the observed frame in units of Å. The second column is a new wavelength grid evaluated at the centres of the wavelength bins, also in units of Å (this is the quantity we usually put on the x -axis when plotting a spectrum). The third column is the flux array in units of $10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$. The fourth column is the inverse variance array in units of $10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1})^{-2}$. ‘sigma’ is a column introduced only in the most recent PyTypeIt versions and represents the noise vector (if it is not present, it is always possible to determine from $\text{ivar}^{-1/2}$). The sixth column is a good pixel mask for the spectrum, and the seventh one is the telluric model. The ‘obj model’ column is not present in the spectra reduced with the most recent PyTypeIt versions and represents the object model used for the telluric fitting. The last two columns are created after the flux scaling procedure and represent the new flux and ivar corrected with the X-band photometry of the quasar, where X could be J, Y, or K_p depending on the available photometric information, in the same units as before.

	Wave	Wave grid mid	Flux	ivar	sigma	mask	Telluric	obj model	flux scaled X	ivar scaled X
J1342+0928	10289.82613	10289.83458	0.31196	18.34511	–	1	0.99	0.61569	0.36666	13.27947
J1058+2930	10240.48127	10240.45124	1.56780	2.58264	0.62226	1	1.0	–	0.54980	21.00057




Figure 3. *Top:* histogram of the $\langle \text{SNR}_\lambda \rangle$ of the quasars in this sample, reported in Table 1 in the *J* band. We generate 25 logarithmically spaced bins between the minimum and maximum of all the $\langle \text{SNR}_\lambda \rangle$ values in the sample. The median $\langle \text{SNR}_\lambda \rangle$ is shown as a red dashed vertical line. *Middle:* same, but for the *H* band. *Bottom:* same, but for the *K* band.

4 COMPARISON WITH OTHER QUASAR SAMPLES

It is important to discuss this work by comparing it to the present literature on other high- z samples, such as Shen et al. (2019), Schindler et al. (2020), Yang et al. (2021), D’Odorico et al. (2023), and Ďurovčíková et al. (2024), where Ďurovčíková et al. (2024) and D’Odorico et al. (2023) are also spectroscopic data releases. The most relevant quantities used to characterize the samples are reported in Table 5. In Fig. 4 we show the distribution of their median z and M_{1450} colour-coded with the sample size and with the errorbars representing the extension of the samples in both z and M_{1450} . The samples that also are public data releases are marked with thicker edges and brighter shades. We want to state clearly that for all the objects where the spectra are previously published, these are not the same reductions as in earlier publications, as all data are re-reduced in this work.

The sample that shows the most similarities with this work is the one in Yang et al. (2021), with 31 quasars in common in the redshift range $6.50 < z < 7.65$ out of the 37 total sources. However, there are key differences between the two data sets. The major differences lie in the instruments used to collect the spectra: Yang et al. (2021) primarily used Gemini/GNIRS and Keck/NIRES, while in this work, a larger fraction of the data comes from VLT/X-Shooter, which offers better spectral resolution (see the R values of each instrument reported in Table 3) and, consequently, higher

quality spectra. Additionally, most of the spectra in Yang et al. (2021) have wavelength coverage only in the NIR arm, whereas our data set includes broader coverage. These differences arise due to distinct scientific objectives. Yang et al. (2021) focused on central BH masses derived from the Mg II emission lines, the Eddington ratio distribution, and chemical abundances in the rest-frame UV (such as the Fe II/Mg II ratio). Consequently, their data set did not prioritize coverage in the optical band. In contrast, we assembled a sample tailored for additional scientific goals, including the study of quasar proximity zones (Onorato et al. 2025) and the Ly α forest, which require coverage in the VIS arm, a key feature provided by our data set. Another important difference is that we have refined the redshift measurements for some quasars, incorporating more recent literature values and prioritizing [C II] measurements when available. Additionally, unlike Yang et al. (2021), which derived M_{1450} from a best-fitting power-law continuum, we independently compute these values from the spectra themselves by using the methodology described in Section 3.3.

The other three samples with only a few similarities with this work are Ďurovčíková et al. (2024) with eight sources in common in the redshift range $6.53 < z < 7.09$ out of the 18 total ones, Schindler et al. (2020) with eight quasars in common at $6.57 < z < 7.65$ out of the 38 total ones, and D’Odorico et al. (2023) with only six sources in common at $6.50 < z < 6.64$ out of the 42 total ones. In the first sample mentioned, the spectra come almost entirely from Magellan/FIRE (Simcoe et al. 2013; $R = 6000$ if the slit is 0.6 arcsec wide), which has a better data quality than the GNIRS and NIRES/MODS spectra in common, but worse than the X-Shooter ones. The other two samples have all the spectra acquired with VLT/X-Shooter.

Finally, there are no sources in common with Shen et al. (2019) as the redshift range (5.71–6.42) of their 50 quasars does not overlap with the one in this work. All their spectra come from Gemini/GNIRS.

5 COMPOSITE SPECTRUM

In this section, we present a $z > 6.5$ quasar composite spectrum based on this sample and compare it with other composite spectra known from the literature (see Fig. 5 and Table 7). We aim to study the average UV quasar spectral properties and their possible evolution through the different redshifts. We decide to exclude all the quasars that show BAL features in their spectra (flagged in Table 1) and then we generate the composite with 33 out of the 45 $z > 6.5$ quasars in the sample. This choice comes from the fact that BAL features can distort the shape of the main emission lines, producing a prominent C IV and many other high-ionization features, such as Si IV and N V.

We generate the composite spectrum following Selsing et al. (2016) as a guideline:

- (i) We move the spectra to the rest frame, dividing the wavelengths by $(1 + z)$;
- (ii) We define a new wavelength grid: [1040, 3332] Å in steps of $d\nu_{\text{pix}} = 110 \text{ km s}^{-1}$ (set to be coarser than the coarsest pixel-scale, in velocity, among all the instruments used to create the sample);
- (iii) We re-bin the spectra into the new wavelength grid;
- (iv) We normalize the spectra to the continuum flux at 1450 Å rest frame, where there are no strong broad lines or iron emission;
- (v) Only for $\lambda > 1225 \text{ Å}$, we apply the following masks to improve the overall quality of the composite, without affecting its natural shape in the Ly α region:

Table 5. Main properties of the sample in this paper, compared with those of other spectroscopic samples from the literature. The columns show, respectively, the reference of the sample, the redshift range, the M_{1450} range, the total number of quasars included in the sample (with the objects in common with this work), and the instruments used to take the spectra.

Sample	z range	M_{1450} range	N_{sources} (in common)	Instruments
Onorato et al. (2025 – this work) ^a	6.50–7.65	[-28.8, -24.1]	45	GNIRS/NIRES/X-Shooter/GMOS/LRIS/DEIMOS/MODS
Ďurovčíková et al. (2024) ^a	6.03–7.08	[-28.0, -26.5]	18 (8)	FIRE/X-Shooter/MOSFIRE/ESI
D’Odorico et al. (2023) ^a	5.77–6.63	[-27.8, -25.8]	42 (6)	X-Shooter
Yang et al. (2021)	6.30–7.65	[-27.4, -25.2]	37 (31)	GNIRS/NIRES/X-Shooter/FIRE/F2
Schindler et al. (2020)	5.78–7.54	[-29.0, -24.4]	38 (8)	X-Shooter
Shen et al. (2019)	5.71–6.42	[-27.7, -24.9]	50	GNIRS

Note. ^aPublic data release.




Figure 4. Comparison between the properties of the quasars in this work and those of the other high- z samples (Shen et al. 2019; Schindler et al. 2020; Yang et al. 2021; D’Odorico et al. 2023; Ďurovčíková et al. 2024). We plot the median z and M_{1450} for all the catalogues, with markers colour-coded with the number of sources included in each sample (N_{sources}) as indicated by the colourbar on the right. The errorbars represent the extension of the samples in both z and M_{1450} . The works where the spectra are publicly released (this work; D’Odorico et al. 2023; Ďurovčíková et al. 2024) are marked with thicker edges and brighter shades.

- (a) Telluric transmission > 0.5 : we use the individual PypeIt telluric model fits to mask out the telluric regions;
- (b) SNR > 0.5 : to mask the flux where the SNR is very low (and hence the noise is high);
- (c) $\sigma [10^{-17} \text{ erg s}^{-1} \text{cm}^{-2} \text{\AA}^{-1}] < 1.5$: to mask the flux where the noise is high (i.e. in those regions close to the right edge of the wavelength coverage of the spectra, where the noise increases exponentially);
- (d) Flux $[10^{-17} \text{ erg s}^{-1} \text{cm}^{-2} \text{\AA}^{-1}] < 40$: to mask out the outliers (e.g. hot pixels or sky lines not well subtracted during the spectra reduction).
- (vi) We create the composite spectrum as a weighted mean of the individual spectra. The weight assigned to each spectrum at each wavelength is given by equation (3). The different factors in the equation are the good pixel mask (see Table 4) that removes

bad pixels from the calculation, the combination of the four masks defined above to exclude low SNR regions, high-noise areas, and spurious outliers ($\text{masks}_{\text{tot}}$), and the number of spectra contributing at each wavelength (n_{used}). Thus, the composite spectrum at every wavelength is the sum of the fluxes of the individual spectra at that wavelength, each multiplied by its weight, and divided by the sum of the weights.

$$\text{weights} = \frac{\text{gpm} \cdot \text{masks}_{\text{tot}}}{n_{\text{used}}} \quad (3)$$

The composite spectrum that we obtain is available online as explained in the Data Availability Section, and an example of the format of the FITS file is shown in Table 6. For $\lambda \in [1700, 1900]$ Å and $\lambda \in [2300, 2600]$ Å (rest frame), the quality of the composite decreases because of the presence of strong telluric absorption at $\lambda \in [13\,500, 14\,150]$ Å and $\lambda \in [18\,200, 19\,300]$ Å (observed frame). Indeed, at the wavelengths of the highest absorption, only $\simeq 5\text{--}10$ spectra contribute to the composite because of the narrow redshift range of our sample. The quality also decreases for $\lambda > 3100$ Å, where the number of contributing spectra starts to get lower.

In the bottom panel of Fig. 5, we compare our composite with those from Vanden Berk et al. (2001), Telfer et al. (2002), Selsing et al. (2016), Shen et al. (2019), Yang et al. (2021), and D’Odorico et al. (2023); for comparison purposes, all the composites are normalized at 1450 Å, showing a better match among their continuum shape for $\lambda < 1580$ Å rather than at longer wavelengths, where there is a visible offset between the various continua. For a better comparison with our work, we consider the non-BAL composite from D’Odorico et al. (2023). We also report the main emission lines in this wavelength range, and an inset panel with a zoom-in of the region [1175, 1580] Å, where we can see differences in the Ly α and C IV lines, despite the good overlap among their continuum shape. In the middle panel of Fig. 5, we display the number of spectra that are contributing to the composite at each wavelength for our work, D’Odorico et al. (2023), Yang et al. (2021), and Shen et al. (2019) (the only three for which this piece of information is available); and finally, in the top panel, we show the distribution of the mean quasar redshift that contributes to the composite at each wavelength. We summarize the properties of our composite and the comparison ones in Table 7.

The discrepancies highlighted in the inset panel of Fig. 5 are most likely due to differences in luminosity among the samples (see the M_{1450} column in Table 7), as we know that many emission-line properties are functions of quasar luminosity. This is the case of C IV, a high-ionization line visible in the composites, where a decreasing equivalent line width (EW) is expected with increasing




Figure 5. *Bottom panel:* composite spectrum of the 33 non-BAL quasars in the sample (black) with its noise vector (grey), compared with several other composites from the literature (coloured curves). All the composites are normalized to the continuum flux at 1450 Å. The grey band for $\lambda < 1225$ Å highlights the mask-free region, where none of the masks described in the main text is applied. The main emission lines are shown as dashed blue lines. The inset panel shows a zoom-in of the region [1175, 1580] Å, where the composites differ more. *Middle panel:* number of spectra that contribute to the composite realization at each wavelength for this work, D'Odorico et al. (2023), Yang et al. (2019), and Shen et al. (2019) (same colours as in the bottom panel). *Top panel:* mean redshift of the quasars that contribute to the composite at each wavelength for this work.

Table 6. Composite spectrum of the 33 non-BAL quasars in the sample. Wavelengths are in the rest frame and units of Å. Flux density units are arbitrary, normalized to the rest-frame 1450 Å continuum flux. The third column is the noise vector. The fourth column indicates the number of quasar's spectra contributing to the composite at each wavelength. The last column is the mean redshift that contributes to the composite at each wavelength. The entire table data are available online.

Wavelengths [Å]	Flux [F_λ]	Error	N spec	Mean z
1040.00000	0.07235	0.03021	18	6.73951
1040.38167	0.14979	0.02575	18	6.73951
1040.76347	0.24631	0.02212	18	6.73951
–	–	–	–	–
1600.55798	1.36016	0.04412	30	6.72212
–	–	–	–	–
2096.78621	0.97827	0.04707	33	6.76699
–	–	–	–	–
3250.71769	0.78563	0.14352	13	6.62963
–	–	–	–	–

luminosity, according to the so-called Baldwin effect (Baldwin 1977). The biggest differences are noticeable in the strength of the C IV emission line when comparing our composite with the one from the most bright samples, such as Selsing et al. (2016), Shen et al. (2019), and D'Odorico et al. (2023). This effect is also confirmed by the test described in Section 5.3, where we divide the sample of non-BAL quasars into two luminosity bins and create a separate composite from each one. In Fig. 8, we can notice the different EW of the Ly α , C IV, and Mg II emission lines due to the difference in M_{1450} of the two sub-samples. The different strength of the Ly α line between our sample (or high- z in general) and low- z ones is due

to the increasing absorption from the IGM towards higher redshift. The overall continuum slope of the composite is consistent with those from the literature, showing that the same spectral features are preserved in quasars at different z ranges.

In the next sections, we discuss an alternative version of the composite obtained including the BAL quasars, and two tests performed by dividing the 33 non-BAL quasars into two redshift bins first, and into two M_{1450} bins later.

5.1 Including BAL quasars

As already stated at the beginning of Section 5, BAL features can affect the shape of a quasar spectrum. For this reason, we decide to exclude the contribution of all the BAL quasars in the sample to the creation of the final composite spectrum. However, for completeness purposes, we also create another version of the composite, this time considering all the quasars in the sample, so including the BALs. We show the comparison between the two versions in Fig. 6, where the blue curve is the composite created from the 33 non-BAL quasars in the sample, and the orange curve is the one obtained including also the 12 BALs flagged in Table 1, and listed in Section 2.1. The rest of the plot shows the same quantities already described in Fig. 5. From this comparison, we can see that there are no strong differences between the two versions of the composite. Both D'Odorico et al. (2023) and Yang et al. (2021) make the same investigation on the BAL contribution to their composite spectrum. They agree with this study that BALs do not cause significant differences but, unlike in this work, decide to include them in their final composite anyway.

Table 7. Main properties of the non-BAL composite spectrum created from the sample in this paper, compared with those of other composites. The columns show, respectively, the reference of the composite, the redshift range of the sample, the M_{1450} range for the sample, the total number of quasars included in the sample, and the instruments used to take the spectra.

Composite	z range	M_{1450} range	N_{tot} quasars	Instruments
Onorato et al. (2025 – this work)	6.50–7.55	[−27.4, −25.2]	33	GNIRS/NIRES/X-Shooter/GMOS/LRIS/MODS
D’Odorico et al. (2023)	5.77–6.63	[−27.8, −25.8]	24 ^a	X-Shooter
Yang et al. (2021)	6.50–7.65	[−27.4, −25.2]	38 ^b	GNIRS/NIRES/X-Shooter/FIRE/F2
Shen et al. (2019)	5.71–6.42	[−27.7, −24.9]	50	GNIRS
Selsing et al. (2016)	1.12–2.10	[−28.1, −27.2] ^c	7	X-Shooter
Telfer et al. (2002)	0.33–3.60	[−27.7, −24.7] ^c	184	FOS/GHRS/STIS
Vanden Berk et al. (2001)	0.044–4.789	[−25.8, −22.8] ^d	2204	SDSS

Notes. ^aExcluding the BALs.

^bWhere 31 come from Yang et al. (2021), and 7 from Schindler et al. (2020).

^cFrom Lusso et al. (2015), $M_{1450} = M_i(z = 2) + 1.28$.

^dConverted from $17.5 < r' < 20.5$, with $z_{\text{median}} = 1.253$.




Figure 6. Bottom panel: composite spectrum of the 33 non-BAL quasars in the sample (dark blue curve, which is the same as the black curve in Fig. 5), compared with the composite spectrum obtained including the 12 BAL quasars (orange curve), which are flagged in Table 1, and also listed in Section 2.1. There are no significant differences between the two versions of the composite. The rest of the plot shows the same quantities already described in Fig. 5.

5.2 Dividing the sample into two redshift bins

To check whether we are averaging any features in the final composite spectrum because of the relatively wide redshift distribution of the quasars in the sample, we perform another test dividing the 33 non-BAL quasars into two redshift bins and creating a composite spectrum from the spectra in each bin. If any ‘ z -related’ features are present, they might arise when dividing the sample into two groups based on the redshift. To account for enough statistics, we consider the median redshift of the 33 non-BALs ($z_{\text{median, non-BAL}} = 6.70$) and create a version of the composite from the 17 spectra having $z \geq 6.70$ and another version from the 16 spectra with $z < 6.70$. The two versions are shown in Fig. 7, where the blue curve is the composite created from the quasars at $z \geq 6.70$, and the orange curve is the one

obtained from the quasars at $z < 6.70$. The rest of the plot shows the same quantities already described in Fig. 5.

From this comparison, we notice that the high- z composite has a shallower slope and thus a redder continuum. A possible interpretation of this would be a larger number of mini BAL absorption at higher redshift. Bischetti et al. (2022, 2023) found that the BAL fraction in $z \geq 6$ quasars is two to three times higher than in quasars at $z \simeq 2$ –4.5. The presence of BALs correlates with a redder continuum, probably due to dust attenuation. If this idea is correct, it would explain why the Ly α and CIV lines look weaker. Alternatively, another explanation to justify the weaker Ly α could be the presence of more neutral hydrogen in the IGM causing more absorption and suggesting the presence of statistical IGM damping wings (Durovčková et al. 2024).




Figure 7. Bottom panel: composite spectrum of the 17 spectra having $z \geq 6.70$ (dark blue curve), compared with the composite spectrum obtained from the 16 spectra with $z < 6.70$ (orange curve), where $z_{\text{median,non-BAL}} = 6.70$ is the median redshift of the 33 non-BAL quasars. The rest of the plot shows the same quantities already described in Fig. 5.




Figure 8. Bottom panel: composite spectrum of the 16 spectra having $M_{1450} < -26.0$ (dark blue curve), compared with the composite spectrum obtained from the 17 spectra with $M_{1450} \geq -26.0$ (orange curve), where $M_{1450,\text{median,non-BAL}} = -26.0$ is the median M_{1450} of the 33 non-BAL quasars. The rest of the plot shows the same quantities already described in Fig. 5.

5.3 Dividing the sample into two magnitude bins

The last test we perform on the composite consists of dividing the 33 non-BAL quasars into two M_{1450} bins and creating a composite spectrum from the spectra in each bin. As before, to account for

enough statistics, we consider the median M_{1450} of the 33 non-BALs ($M_{1450,\text{median,non-BAL}} = -26.0$) and create a version of the composite from the 16 spectra having $M_{1450} < -26.0$ and another version from the 17 spectra with $M_{1450} \geq -26.0$. The two versions are shown in

Fig. 8, where the blue curve is the composite created from the quasars having $M_{1450} < -26.0$, and the orange curve is the one obtained from the quasars having $M_{1450} \geq -26.0$. The rest of the plot shows the same quantities already described in Fig. 5.

The differences in the strength of the emission lines come from the Baldwin effect (Baldwin 1977), already discussed in Section 5, proving that quasars' luminosity anticorrelates with their emission lines strength. From this test, we show how this effect is still visible at early times.

6 SUMMARY

In this section, we discuss the conclusions of this paper, the quantitative analysis performed on the sample, and future work. We summarize the two main results below.

(i) We report and release a sample of 45 quasars at $6.50 < z \leq 7.64$ ($z_{\text{median}} = 6.71$), in the magnitude range $-28.82 \leq M_{1450} \leq -24.13$ ($M_{1450,\text{median}} \simeq -26.1$), where 12 out of the 45 quasars show BAL features in their spectrum, and three are new unpublished quasars (Bañados et al. 2025; Belladitta et al. 2025). This represents the largest medium-/moderate-resolution released sample of quasars at high redshift from ground-based instruments. The optical and NIR spectroscopy were obtained using the Gemini/GNIRS, Keck/NIRSPEC, VLT/X-Shooter, Gemini/GMOS, Keck/DEIMOS, Keck/LRIS, and LBT/MODS instruments. The data in this sample allow us to estimate the M_{1450} of the quasars directly from the spectra, and to determine their quality from the SNR:

(a) After scaling the flux and inverse variance of each spectrum considering the J -, Y -, or K_p -band magnitude of every quasar, we calculate M_{1450} from these new flux-scaled spectra (see Section 3.3 and Appendix B). These values may be relevant in the course of the analysis on quasar proximity zones (Onorato et al. 2025), in case we have to correct the measurements for the luminosity of the quasar.

(b) We compute the SNR of the spectra in the sample, considering three different wavelength ranges that exclude the telluric regions: [11 000, 13 400], [14 500, 17 950], and [19 650, 22 400] Å (see Section 3.4 and Appendix C). We want to test the quality of the spectra and give an idea of how much a certain spectrum can be trusted.

(ii) We create a $z > 6.5$ quasar composite spectrum using 33 out of the 45 quasars of this sample: we exclude the BAL quasars to avoid biases in the analysis due to absorption features in their spectra. We compare the composite with others from low- and high-redshift quasars samples from the literature. No significant redshift evolution is found for either broad UV emission lines or quasar continuum slopes (see Section 5). An alternative version of the composite, created considering also the BAL quasars, is described in Section 5.1. We notice that there are no strong differences between the two versions. Other two tests are discussed in Sections 5.2 and 5.3. In the first one, we divide the sample of 33 non-BAL quasars into two redshift bins (17 quasars at $z \geq 6.70$ and 16 at $z < 6.70$) and create a composite from the spectra in each bin. There are small differences in the continuum shape and the strength of Ly α and C IV emission lines, visible in Fig. 7, that we tentatively interpret as due to the presence of subtle BAL absorption at higher z (Bischetti et al. 2022, 2023). Alternatively, statistical damping wings (Durovčíková et al. 2024) could also justify the weaker Ly α observed at higher z , while the difference in Mg II could arise from changes in the average black hole mass or accretion rates. In the second test, we divide the sample of 33 non-BAL quasars into two M_{1450} bins (17 quasars at

$M_{1450} \geq -26.0$ and 16 at $M_{1450} < -26.0$) and create a composite from the spectra in each bin. In Fig. 8, we see how weaker emission lines are associated with brighter quasars, confirming the presence of the Baldwin effect (Baldwin 1977), which still holds at early times.

Starting from the sample presented in this paper, in subsequent works we will reconstruct the quasar's intrinsic blue side from the observed red side, with PCA continuum modelling (e.g. Davies et al. 2018a, b; Bosman et al. 2021), to study their proximity zones (Onorato et al. 2025), and Ly α damping wings (Miralda-Escudé 1998; Davies et al. 2018a; Durovčíková et al. 2024; Greig et al. 2024). In particular, Hennawi et al. (2025) introduce a new inference approach for analysing the IGM damping wings, deriving a single Bayesian likelihood for the entire spectrum, while Kist, Hennawi & Davies (2025) quantify the precision with which these IGM damping wings analysed with the new method can measure astrophysical parameters such as $\langle x_{\text{HI}} \rangle$ and t_{Q} , and the dependence of this precision on the properties of the spectra analysed. Our final goal will be to impose more stringent constraints on $\langle x_{\text{HI}} \rangle$ during the EoR and the radiative efficiency of the earliest SMBHs.

ACKNOWLEDGEMENTS

This work is based in part on observations obtained at the international Gemini Observatory, a program of NSF NOIRLab, which is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the U.S. National Science Foundation on behalf of the Gemini Observatory partnership: the U.S. National Science Foundation (United States), National Research Council (Canada), Agencia Nacional de Investigación y Desarrollo (Chile), Ministerio de Ciencia, Tecnología e Innovación (Argentina), Ministério da Ciência, Tecnologia, Inovações e Comunicações (Brazil), and Korea Astronomy and Space Science Institute (Republic of Korea).

Some of the data presented herein were obtained at Keck Observatory, which is a private 501(c)3 non-profit organization operated as a scientific partnership among the California Institute of Technology, the University of California, and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the Native Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain. This research has made use of the Keck Observatory Archive (KOA), which is operated by the W. M. Keck Observatory and the NASA Exoplanet Science Institute (NExSci), under contract with the National Aeronautics and Space Administration.

This work is based in part on observations made with ESO telescopes at the La Silla Paranal Observatory.

This paper also uses data based on observations obtained at the LBT, an international collaboration among institutions in the United States, Italy, and Germany. The LBT Corporation partners are The University of Arizona on behalf of the Arizona university system; Istituto Nazionale di Astrofisica, Italy; LBT Beteiligungs-gesellschaft, Germany, representing the Max Planck Society, the Astrophysical Institute Potsdam, and Heidelberg University; The Ohio State University; The Research Corporation, on behalf of The University of Notre Dame, University of Minnesota and University of Virginia.

We acknowledge the use of the `PypeIt` data reduction package.

We acknowledge helpful conversations with the ENIGMA group at UC Santa Barbara and Leiden University. SO is grateful to Ben

Wang and Suk Sien Tie for help with some of the data reductions, and to Elia Pizzati and Caitlin Doughty for comments on an early version of the manuscript.

JFH acknowledges support from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No 885301), from the National Science Foundation (NSF) under Grant No. 2307180, and from NASA under the Astrophysics Data Analysis Program (ADAP, Grant No. 80NSSC21K1568). JTS is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project number 518006966. FW acknowledges support from NSF award AST-2513040. CM acknowledges support from Fondecyt Iniciacion grant 11240336 and the ANID BASAL project FB210003. EPF is supported by the international Gemini Observatory, a program of NSF’s NOIRLab, which is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation, on behalf of the Gemini partnership of Argentina, Brazil, Canada, Chile, the Republic of Korea, and the United States of America.

DATA AVAILABILITY

Data archives

The raw Gemini data (both GNIRS and GMOS) can be searched and downloaded from the Gemini Observatory Archive at link <https://archive.gemini.edu/searchform>. The user needs to set the Instrument used and the coordinates of the target (RA and Dec.). Additionally, we notice that sometimes there are superimposed features like vertical striping, horizontal banding, and quadrant offsets on GNIRS data. We use the CLEANIR⁷ Python routine to remove these artefacts.

The raw Keck data (all NIRES, DEIMOS, and LRIS) can be downloaded from a Basic Search on the Keck Observatory Archive (KOA) at the link <https://koa.ipac.caltech.edu/cgi-bin/KOA/nph-KOALogin>. The user needs to select the Instrument used, set the Object Name or Location, and decide how to return the results. In case we want to download multiple targets, we can look for More Search Options, select the desired Instruments, and add a file in the Multiple Object Table File section. A code to create such a table is provided in the GitHub repository storing the relevant tables and codes used in this paper, whose link is provided in Section 2.3.

The raw ESO data (VLT/X-Shooter) can be downloaded from the ESO Observatory Archive at the link https://archive.eso.org/eso/eso_archive_main.html. The user can type the coordinates of the target (RA and Dec.), set the Instrument used, and select the category of data desired. In case we want to download multiple targets, we can add a file in the List of Targets section. A code to create such a file is provided in the GitHub repository already mentioned.

The raw LBT data (MODS) come from private projects present in the LBT Archive at the link <http://archive.lbto.org/>. The user needs to set the coordinates of the target (RA and Dec.), the Instrument used, and to authenticate for the project.

More (and fewer) details can always be inserted in the archives when searching for targets.

Data reduction and analysis

The files to reduce the spectroscopic data with PypeIt (.pypeit, .flux, .coadd1d, and .tell), the .csv tables (containing information such as the redshift of the quasars, photometric details,

or SNR), and the relevant codes to reproduce the results in this paper are all stored in the GitHub repository previously mentioned.

Final FITS files (spectra and composites)

The data underlying this article are available in the article and in its online supplementary material.

REFERENCES

Aihara H. et al., 2018, *PASJ*, 70, S4
 Andika I. T. et al., 2020, *ApJ*, 903, 34
 Andika I. T. et al., 2022, *AJ*, 163, 251
 Andika I. T. et al., 2024, *A&A*, 685, A25
 Baldwin J. A., 1977, *ApJ*, 214, 679
 Bañados E. et al., 2016, *ApJS*, 227, 11
 Bañados E. et al., 2018, *Nature*, 553, 473
 Bañados E. et al., 2019, *ApJ*, 885, 59
 Bañados E. et al., 2021, *ApJ*, 909, 80
 Bañados E. et al., 2025, *Nat. Astron.*, 9, 293
 Bañados E., Decarli R., Walter F., Venemans B. P., Farina E. P., Fan X., 2015, *ApJ*, 805, L8
 Becker G. D., Bolton J. S., Madau P., Pettini M., Ryan-Weber E. V., Venemans B. P., 2015, *MNRAS*, 447, 3402
 Belladitta S. et al., 2025, preprint ([arXiv:2505.15923](https://arxiv.org/abs/2505.15923))
 Bischetti M. et al., 2022, *Nature*, 605, 244
 Bischetti M. et al., 2023, *ApJ*, 952, 44
 Bochanski J. J. et al., 2009, *PASP*, 121, 1409
 Bosman S. E. I. et al., 2022, *MNRAS*, 514, 55
 Bosman S. E. I., Ďurovčková D., Davies F. B., Eilers A.-C., 2021, *MNRAS*, 503, 2077
 Clough S. A., Shephard M. W., Mlawer E. J., Delamere J. S., Iacono M. J., Cady-Pereira K., Boukabara S., Brown P. D., 2005, *J. Quant. Spec. Radiat. Transf.*, 91, 233
 D’Odorico V. et al., 2023, *MNRAS*, 523, 1399
 Davies F. B. et al., 2018a, *ApJ*, 864, 142
 Davies F. B. et al., 2018b, *ApJ*, 864, 143
 Davies F. B., Hennawi J. F., Eilers A.-C., 2019, *ApJ*, 884, L19
 Davies F. B., Hennawi J. F., Eilers A.-C., 2020, *MNRAS*, 493, 1330
 Davies R. L. et al., 2023, *MNRAS*, 521, 289
 Decarli R. et al., 2018, *ApJ*, 854, 97
 Ďurovčková D. et al., 2024, *ApJ*, 969, 162
 Dye S. et al., 2018, *MNRAS*, 473, 5113
 Eilers A.-C. et al., 2020, *ApJ*, 900, 37
 Eilers A.-C., Davies F. B., Hennawi J. F., Prochaska J. X., Lukić Z., Mazzucchelli C., 2017, *ApJ*, 840, 24
 Elias J. H., Joyce R. R., Liang M., Muller G. P., Hileman E. A., George J. R., 2006b, in McLean I. S., Iye M.eds, Proc. SPIE Conf. Ser. Vol. 6269, Ground-based and Airborne Instrumentation for Astronomy. SPIE, Bellingham, p. 62694C
 Elias J. H., Rodgers B., Joyce R. R., Lazo M., Doppmann G., Winge C., Rodríguez-Ardila A., 2006a, in McLean I. S., Iye M.eds, Proc. SPIE Conf. Ser. Vol. 6269, Ground-based and Airborne Instrumentation for Astronomy. SPIE, Bellingham, p. 626914
 Faber S. M. et al., 2003, in Iye M., Moorwood A. F. M.eds, Proc. SPIE Conf. Ser. Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes. SPIE, Bellingham, p. 1657
 Fan X. et al., 2006, *AJ*, 132, 117
 Fan X. et al., 2019, *ApJ*, 870, L11
 Fan X., Bañados E., Simcoe R. A., 2023, *ARA&A*, 61, 373
 Farina E. P. et al., 2022, *ApJ*, 941, 106
 Fujimoto S. et al., 2024, *ApJ*, 977, 250
 Furtak L. J. et al., 2024, *Nature*, 628, 57
 Gaskell C. M., 1982, *ApJ*, 263, 79
 Goulding A. D. et al., 2023, *ApJ*, 955, L24
 Greene J. E. et al., 2024, *ApJ*, 964, 39
 Greig B. et al., 2024, *MNRAS*, 530, 3208
 Gullikson K., Dodson-Robinson S., Kraus A., 2014, *AJ*, 148, 53

⁷<https://www.gemini.edu/instrumentation/niri/data-reduction>

Gunn J. E., Peterson B. A., 1965, *ApJ*, 142, 1633

Hennawi J. F., Kist T., Davies F. B., Tamanas J., 2025, *MNRAS*, 539, 2621

Hook I. M., Jørgensen I., Allington-Smith J. R., Davies R. L., Metcalfe N., Murowinski R. G., Crampton D., 2004, *PASP*, 116, 425

Horne K., 1986, *PASP*, 98, 609

Inayoshi K., Visbal E., Haiman Z., 2020, *ARA&A*, 58, 27

Jiang L. et al., 2016, *ApJ*, 833, 222

Kist T., Hennawi J. F., Davies F. B., 2025, *MNRAS*, 538, 2704

Kokorev V. et al., 2023, *ApJ*, 957, L7

Kokorev V. et al., 2024, *ApJ*, 968, 38

Labbe I. et al., 2025, *ApJ*, 978, 92

Lai S. et al., 2022, *MNRAS*, 513, 1801

Larson R. L. et al., 2023, *ApJ*, 953, L29

Lawrence A. et al., 2007, *MNRAS*, 379, 1599

Lusso E., Worseck G., Hennawi J. F., Prochaska J. X., Vignali C., Stern J., O'Meara J. M., 2015, *MNRAS*, 449, 4204

Maiolino R. et al., 2024a, *Nature*, 627, 59

Maiolino R. et al., 2024b, *A&A*, 687, A67

Matsuoka Y. et al., 2018, *ApJS*, 237, 5

Matsuoka Y. et al., 2019a, *ApJ*, 872, L2

Matsuoka Y. et al., 2019b, *ApJ*, 883, 183

Matsuoka Y. et al., 2022, *ApJS*, 259, 18

Matthee J. et al., 2024, *ApJ*, 963, 129

Mazzucchelli C. et al., 2017, *ApJ*, 849, 91

Mazzucchelli C. et al., 2023, *A&A*, 676, A71

McMahon R. G., Banerji M., Gonzalez E., Koposov S. E., Bejar V. J., Lodieu N., Rebolo R., *VHS Collaboration*, 2013, *The Messenger*, 154, 35

Meyer R. A., Bosman S. E. I., Ellis R. S., 2019, *MNRAS*, 487, 3305

Miralda-Escudé J., 1998, *ApJ*, 501, 15

Moorwood A., Cuby J. G., Lidman C., 1998, *The Messenger*, 91, 9

Mortlock D. J. et al., 2011, *Nature*, 474, 616

Oke J. B. et al., 1995, *PASP*, 107, 375

Onorato S., Hennawi J. F., Pizzati E., Venemans B. P., Eilers A.-C., 2025, preprint ([arXiv:2505.09676](https://arxiv.org/abs/2505.09676))

Onoue M. et al., 2019, *ApJ*, 880, 77

Pérez-González P. G. et al., 2024, *ApJ*, 968, 4

Pogge R. W. et al., 2010, in McLean I. S., Ramsay S. K., Takami H. eds, Proc. SPIE Conf. Ser. Vol. 7735, Ground-based and Airborne Instrumentation for Astronomy III. SPIE, Bellingham, p. 77350A

Prochaska J. et al., 2020b, *J. Open Source Softw.*, 5, 2308

Prochaska J. X. et al., 2020a, Zenodo

Reed S. L. et al., 2017, *MNRAS*, 468, 4702

Reed S. L. et al., 2019, *MNRAS*, 487, 1874

Richards G. T., Vanden Berk D. E., Reichard T. A., Hall P. B., Schneider D. P., SubbaRao M., Thakar A. R., York D. G., 2002, *AJ*, 124, 1

Rockosi C. et al., 2010, in McLean I. S., Ramsay S. K., Takami H. eds, Proc. SPIE Conf. Ser. Vol. 7735, Ground-based and Airborne Instrumentation for Astronomy III. SPIE, Bellingham, p. 77350R

Ross N. P., Cross N. J. G., 2020, *MNRAS*, 494, 789

Satyavolu S. et al., 2023, *MNRAS*, 522, 4918

Schindler J.-T. et al., 2020, *ApJ*, 905, 51

Selsing J., Fynbo J. P. U., Christensen L., Krogager J. K., 2016, *A&A*, 585, A87

Shen Y. et al., 2016, *ApJ*, 831, 7

Shen Y. et al., 2019, *ApJ*, 873, 35

Simcoe R. A. et al., 2013, *PASP*, 125, 270

Tang J.-J. et al., 2017, *MNRAS*, 466, 4568

Telfer R. C., Zheng W., Kriss G. A., Davidsen A. F., 2002, *ApJ*, 565, 773

Tytlér D., Fan X.-M., 1992, *ApJS*, 79, 1

van Dokkum P. G., 2001, *PASP*, 113, 1420

Vanden Berk D. E. et al., 2001, *AJ*, 122, 549

Venemans B. P. et al., 2013, *ApJ*, 779, 24

Venemans B. P. et al., 2015, *ApJ*, 801, L11

Venemans B. P. et al., 2017, *ApJ*, 851, L8

Venemans B. P. et al., 2020, *ApJ*, 904, 130

Venemans B. P., Walter F., Zschaechner L., Decarli R., De Rosa G., Findlay J. R., McMahon R. G., Sutherland W. J., 2016, *ApJ*, 816, 37

Vernet J. et al., 2011, *A&A*, 536, A105

Wang F. et al., 2017, *ApJ*, 839, 27

Wang F. et al., 2018, *ApJ*, 869, L9

Wang F. et al., 2019, *ApJ*, 884, 30

Wang F. et al., 2021a, *ApJ*, 907, L1

Wang F. et al., 2021b, *ApJ*, 908, 53

Wang F. et al., 2024, *ApJ*, 969, 9

Wilson J. C. et al., 2004, in Moorwood A. F. M., Iye M. eds, Proc. SPIE Conf. Ser. Vol. 5492, Ground-based Instrumentation for Astronomy. SPIE, Bellingham, p. 1295

Woods T. E. et al., 2019, *Publ. Astron. Soc. Aust.*, 36, e027

Wu X.-B. et al., 2015, *Nature*, 518, 512

Yang J. et al., 2019, *AJ*, 157, 236

Yang J. et al., 2020, *ApJ*, 897, L14

Yang J. et al., 2021, *ApJ*, 923, 262

Yang J. et al., 2023, *ApJ*, 951, L5

SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

suppl_data

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX A: FLUX SCALING WITH J -, Y -, OR K_p -BAND PHOTOMETRY

As already mentioned in Section 3.2, the reduced spectrum of every quasar in this sample is scaled using the J -, Y -, or K_p -band photometry. This is because the flux calibration of the spectra performed with *TypeIt* is relative and, while we are still looking for a definitive absolute flux calibration method, we need to tie the spectroscopic measurements with the photometric ones, which are more accurate in terms of fluxes.

To correct the spectra, we scale them to a given magnitude in a specified passband filter. First, we calculate the AB magnitude of the spectrum in a certain passband filter (m_{spec}) by using the Python library *Speclite*.⁸ Then, we calculate a scale factor (SF) using the difference between the calculated magnitude from the spectrum and the true magnitude from the photometry (m_{phot}), as shown in this formula that we get from the Pogson law:

$$SF = 10^{0.4(m_{\text{spec}} - m_{\text{phot}})} \quad (A1)$$

This scale factor is used to adjust the flux and inverse variance of the spectrum to match the true magnitude in the given passband (see Fig. A1). For a subset of quasars, literature photometry is available not only in the J band, but also in the Y , H , and K bands. We compute other synthetic magnitudes, but this time from the flux-scaled spectra and compare them to the additional photometry to assess any discrepancies. We find that the typical deviation between synthetic and observed magnitudes is around 10 per cent, with a maximum discrepancy of approximately 40 per cent. Two illustrative examples, one with high and one with low discrepancy, are shown in the top two panels of Fig. A1. We do not display synthetic photometry for the Y band of the quasar J1342+0928 because the available spectrum does not fully cover the wavelength range of the Y -band filter (9635–11025 Å). As a result, it is not possible to reliably compute the synthetic magnitude for this band.

⁸<https://speclite.readthedocs.io/en/latest/>




Figure A1. *Top two panels:* example cases of the spectra of J1342+0928 and J1007+2115 before (blue) and after (orange) applying the scale factor (SF) calculated from the J -band magnitudes of these quasars (as described in equation A1), and the noise vectors. The lower plot shows (in brighter shades) the transmission curve of the passband filter used to get the photometry. We also include the Y -, H -, and K -band magnitudes, along with their respective transmission curves (in lighter shades), to compare with the synthetic photometry derived from the flux-scaled spectrum (magenta stars) and visually assess any discrepancies. *Middle:* same, but for J0923+0754 and its Y -band photometry. *Bottom:* same, but for J1058+2930 and its K_p -band photometry.




Figure B1. *Top:* match between the spectrum of J0910–0414 and the composite spectrum obtained from the sample excluding the BALs already corrected to get a better estimate of the M_{1450} . The spectrum of the BAL quasar is shown in blue, while the composite is in orange, with their noise vectors reported at the bottom of the plot. The back dashed line falls at $\lambda = 1450$ Å and the green dashed line is the normalization wavelength at 2000 Å. The magenta star is the ‘new’ continuum level assumed for the BAL quasar at 1450 Å. *Bottom:* same, but for J0923+0402.

APPENDIX B: ABSOLUTE MAGNITUDE OF J0910–0414 AND J0923+0402 (BAL QUASARS)

This appendix describes the approach followed to calculate M_{1450} in the two problematic cases of J0910–0414 and J0923+0402. These quasars are affected by strong BAL features, which mostly appear around the region at rest frame 1450 Å, making the measurement of M_{1450} with the method described in Section 3.3 difficult to trust (note the black dashed line at $\lambda = 1450$ Å in Fig. B1 falling in the area affected by the BAL). To correct this problem, we want to visualize what the continuum would be in a quasar with the same shape, but without BAL features. For this aim, we make a comparison using the composite spectrum created from the sample excluding the BAL quasars (described in Section 5 and shown in Fig. 5). We start moving the observed spectrum (already flux-scaled using its J -band magnitude) to the rest frame, while the composite spectrum is already in the rest frame. We scale the composite to the observed spectrum at a certain wavelength ($\lambda_{\text{norm}} = 2000$ Å, in both cases, produces a

good result) by multiplying for a factor A .⁹ We apply a power-law correction with an exponent α , to rotate the composite and then have a better match with the orientation of the observed spectra, according to the following formula:

$$f_{\text{composite,new}} = f_{\text{composite}} \cdot A \left(\frac{\lambda_{\text{composite}}}{\lambda_{\text{norm}}} \right)^\alpha \quad (\text{B1})$$

We set $\alpha = 0.3$ for J0910–0414 and $\alpha = 0.8$ for J0923+0402.

Finally, we can calculate M_{1450} (reported in Table 1) from the scaled composite spectrum following the method described in Section 3.3, assuming that the magenta star in Fig. B1 represents the ‘new’ continuum level for the BAL quasar at 1450 Å. Fig. B1 shows the observed spectra in blue and the composite in orange, with their noise vectors reported at the bottom of each plot; the black dashed lines represent $\lambda = 1450$ Å and the green dashed lines are the normalization wavelengths, both set at 2000 Å.

⁹It is the ratio between the flux of the BAL spectrum and the flux of the composite spectrum at the wavelength λ_{norm} .




Figure C1. *Top:* example case of the distribution of the SNR along the wavelength grid [1040, 3332]Å for the highest z quasar in the sample, J0313–1806 (dark red). The bands (J , H , and K) in which we compute the SNR are indicated according to different colours. *Bottom:* same, but for the lowest z quasar of the sample: J1110–1329.

APPENDIX C: SNR

In Fig. C1, we plot the distribution of the SNR along the rest-frame wavelength grid defined in Section 3.4 for J0313–1806 and J1110–1329 (dark red curves), and the J , H , and K bands (indicated in Section 3.4 too) are in different colours. These are two good example cases of spectra with medium/high and medium/low SNR, respectively.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.