

COLENGTH ONE DEFORMATION RINGS

DANIEL LE, BAO V. LE HUNG, STEFANO MORRA, CHOL PARK, AND ZICHENG QIAN

ABSTRACT. Let $K/\mathbb{Q}_p$ be a finite unramified extension, $\bar{\rho} : \text{Gal}(\bar{\mathbb{Q}}_p/K) \rightarrow \text{GL}_n(\bar{\mathbb{F}}_p)$ a continuous representation, and τ a tame inertial type of dimension n . We explicitly determine, under mild regularity conditions on τ , the potentially crystalline deformation ring $R_{\bar{\rho}}^{\eta, \tau}$ in parallel Hodge–Tate weights $\eta = (n-1, \dots, 1, 0)$ and inertial type τ when the *shape* of $\bar{\rho}$ with respect to τ has colength at most one. This has application to the modularity of a class of shadow weights in the weight part of Serre’s conjecture. Along the way we make unconditional the local-global compatibility results of Park and Qian [Mém. Soc. Math. Fr. (N.S.) 173 (2022), pp. vi+150].

1. INTRODUCTION

In recent years, calculations of various potentially crystalline deformation spaces have seen a number of applications to questions of local-global compatibility in the mod p and p -adic Langlands program. This includes the weight part of Serre’s conjecture, the determination of mod p multiplicities, conjectures of Breuil on integral structures in K -types, and generalizations of Colmez’s functor (see e.g. [EGS15, LLHLM18, LLHLM20, DL21, LLHLMb, LLHM⁺, LLHLMc, LLHLMa, HW22], [BHH⁺b, BHH⁺a]). Under a somewhat exotic genericity condition, [LLHLMb] shows that tamely potentially crystalline deformation spaces are equisingular to certain closed subvarieties of Pappas–Zhu local models. Not much is known about the geometry of these local models for Galois deformation spaces in general. Moreover, it is difficult in practice to make the genericity condition explicit or to work with their natural presentations.

1.1. The main result. The local model has a stratification indexed by *admissible* elements of the extended affine Weyl group called *shapes* and the complexity of the geometry increases as the length of the shape decreases. [LLHLMa] shows that when the shape is *extremal* i.e. has maximal length $\binom{n+1}{3}$ for n -dimensional representations of $\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$, then the corresponding tamely potentially crystalline deformation ring is formally smooth. The main result of this paper, which we state only for representations of $\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ in the introduction, is the following:

Received by the editors June 9, 2023, and, in revised form, January 25, 2024.

2020 *Mathematics Subject Classification*. Primary 11F33, 11F85, 11F80.

The first author was supported by the National Science Foundation under agreements Nos. DMS-1128155 and DMS-1703182 and an AMS-Simons travel grant. The second author acknowledges support from the National Science Foundation under grant Nos. DMS-1128155, DMS-1802037 and the Alfred P. Sloan Foundation. The third author was supported by the ANR-18-CE40-0026 (CLap CLap) and the Institut Universitaire de France. The fourth author was supported by Samsung Science and Technology Foundation under Project Number SSTF-BA2001-02.

Theorem 1.1.1 (Theorem 5.2.3). *Let E be a finite extension of $\mathbb{Q}_p$, with ring of integers $\mathcal{O}$ and residue field $\mathbb{F}$. Let $\bar{\rho} : \text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p) \rightarrow \text{GL}_n(\mathbb{F})$ a continuous Galois representation, τ a $3n - 7$ -generic tame inertial type, $\eta = (n - 1, \dots, 1, 0) \in \mathbb{Z}^n$, and $R_{\bar{\rho}}^{\eta, \tau}$ the lifting ring for potentially crystalline representations of type (η, τ) . If $R_{\bar{\rho}}^{\eta, \tau}$ is nonzero and the length of the shape $\tilde{w}(\bar{\rho}, \tau)$ is at least $\binom{n+1}{3} - 1$ (i.e. the colength of the shape is at most one), then $R_{\bar{\rho}}^{\eta, \tau}$ is formally smooth over $\mathcal{O}$ or $\mathcal{O}[[X, Y]]/(XY - p)$.*

Remark 1.1.2.

- (1) Replacing $\mathbb{Q}_p$ by a finite unramified extension K and requiring that the shape have colength one at each embedding $K \hookrightarrow \bar{\mathbb{Q}}_p$, we show that the deformation ring is formally smooth over a completed tensor product of rings of the form $\mathcal{O}[[X, Y]]/(XY - p)$. The number of such factors in the tensor product can furthermore be explicitly computed.
- (2) The colength one deformation spaces that have a parabolic structure were computed in [LLHLMa]. In general, colength one deformation spaces do not have a parabolic structure, making their computation far more difficult.
- (3) When $n = 3$, the tame inertial types with colength one shape are sufficient to prove the Serre weight conjecture for GL_3 [LLHLM18, LLHLM20, LLHLMc]. We generalize these ideas to prove the modularity of Serre weights of *defect at most one* under the assumptions of [LLHLMa], in particular under an explicit combinatorial genericity condition.
- (4) Using standard Taylor–Wiles techniques, Theorem 1.1.1 gives modularity lifting results similar to [LLHLMb, Theorem 9.2.1] (improving the polynomial genericity and the tameness condition at p in *loc. cit.*, but imposing specific conditions on the shape with respect to the tame inertial types).

While Theorem 1.1.1 generalizes some previous results, its proof is perhaps surprisingly subtle despite the shape having close to maximal length. We do not expect our methods to extend to shapes of smaller length. This suggests that local models for Galois deformation spaces are genuinely complicated geometric objects and that simple explicit descriptions are hard to come by.

Our principal motivation in writing this paper was to apply Theorem 1.1.1 to prove the weight elimination and mod p multiplicity one results necessary to make unconditional the local-global compatibility result of [PQ22] which states roughly that the local mod p Galois representation at p can be recovered from the $\text{GL}_n(\mathbb{Q}_p)$ -action on the Hecke isotypic part of the mod p completed cohomology of a definite unitary group. While the results of [PQ22] were superseded by those of [LLHM⁺], the method of [LLHM⁺] using only extremal shapes does not work for $\text{GSp}_4(\mathbb{Q}_p)$ while it should be possible to adapt the methods of [PQ22] (which builds on [BD14, HLM17, LMP18, MP17] in small rank) to many p -adic reductive groups over $\mathbb{Q}_p$. Indeed, this has been carried out for $\text{GSp}_4(\mathbb{Q}_p)$ [EL]. For generalizations of [LLHM⁺], an analogue of Theorem 1.1.1 should prove useful. We hope to return to this in future work.

1.2. Global applications. As mentioned in Remark 1.1.2(3), as a more immediate global application of Theorem 1.1.1 we obtain the modularity of weights of defect at most one. The notion of *defect* of a Serre weight σ for a *tame* Galois representation $\bar{\rho}$ was first introduced in [LLHLMb, §8.6]. This notion is purely combinatorial, and

encodes the maximal length for the shapes $\tilde{w}(\bar{\rho}, \tau)$ such that $\sigma \in W^?(\bar{\rho}) \cap \text{JH}(\overline{\sigma(\tau)})$. In this paper we generalize the notion of defect for any $\bar{\rho}$ in terms of specializations (as done in [LLHLMa] for extremal weights), and prove their modularity when the defect is at most one, conditional to the existence of a modular extremal weight. The result is the following, and we refer to the bulk of the paper for any undefined notion:

Theorem 1.2.1 (Theorem 6.3.1). *Let F/F^+ be a CM field. Assume that $F^+ \neq \mathbb{Q}$, that all places of F^+ above p are unramified over $\mathbb{Q}_p$ and totally split in F . Let $\bar{\tau} : G_{F^+} \rightarrow \mathcal{G}(\mathbb{F})$ be a continuous representation which is automorphic in the sense of [LLHLMa, Definition 5.5.1], with set of modular weights $W(\bar{\tau})$. Let $\bar{\tau}_p$ be the L -homomorphism attached to the collection $\{\bar{\tau}|_{G_{F_v^+}}\}_{v|p}$ and write $W_{\leq 0}^g(\bar{\tau}_p)$ and $W_{\leq 1}^g(\bar{\tau}_p)$ for the set of extremal weights and for the set of weights of defect at most one, respectively, for $\bar{\tau}_p$. Assume further that:*

- $\bar{\tau}(G_{F(\zeta_p)}) \subseteq \text{GL}_n(\mathbb{F})$ is adequate; and
- $\bar{\tau}_p$ is $6(n-1)$ -generic.

Then the following are equivalent:

- (1) $W_{\leq 0}^g(\bar{\tau}_p) \cap W(\bar{\tau}) \neq \emptyset$; and
- (2) $W_{\leq 1}^g(\bar{\tau}_p) \subset W(\bar{\tau})$.

Compared to [LLHLMa, Theorem 5.5.5], Theorem 1.2.1 assumes that F^+ is unramified above p , but it gives the modularity of weights of defect at most one. (For GL_3 , this is sufficient to prove the generic Serre weight conjecture.)

1.3. Notation. For a field K , we denote by $\bar{K}$ a fixed separable closure of K and let $G_K \stackrel{\text{def}}{=} \text{Gal}(\bar{K}/K)$. If K is defined as a subfield of an algebraically closed field, then we set $\bar{K}$ to be this field.

If K is a nonarchimedean local field, we let $I_K \subset G_K$ denote the inertial subgroup and $W_K \subset G_K$ denote the Weil group. We fix a prime $p \in \mathbb{Z}_{>0}$. Let $E \subset \bar{\mathbb{Q}}_p$ be a subfield which is finite-dimensional over $\mathbb{Q}_p$. We write $\mathcal{O}$ to denote its ring of integers, fix a uniformizer $\varpi \in \mathcal{O}$ and let $\mathbb{F}$ denote the residue field of E . We will assume throughout that E is sufficiently large.

1.3.1. Reductive groups. Let G denote a split connected reductive group (over some ring) together with a Borel B , a maximal split torus $T \subset B$, and $Z \subset T$ the center of G . Let $d = \dim G - \dim B$. When G is a product of copies of GL_n , we will take B to be upper triangular Borel and T the diagonal torus. Let $\Phi^+ \subset \Phi$ (resp. $\Phi^{\vee,+} \subset \Phi^\vee$) denote the subset of positive roots (resp. positive coroots) in the set of roots (resp. coroots) for (G, B, T) . We use the notation $\alpha > 0$ (resp. $\alpha < 0$) for a positive (resp. negative) root $\alpha \in \Phi$. Let Δ (resp. $\Delta^\vee$) be the set of simple roots (resp. coroots). Let $X^*(T)$ be the group of characters of T , and set $X^0(T)$ to be the subgroup consisting of characters $\lambda \in X^*(T)$ such that $\langle \lambda, \alpha^\vee \rangle = 0$ for all $\alpha^\vee \in \Delta^\vee$. Let $W(G)$ denote the Weyl group of (G, T) . Let w_0 denote the longest element of $W(G)$. We sometimes write W for $W(G)$ when there is no chance for confusion. Let W_a (resp. $\widetilde{W}$) denote the affine Weyl group and extended affine Weyl group

$$W_a = \Lambda_R \rtimes W(G) \quad \text{and} \quad \widetilde{W} = X^*(T) \rtimes W(G)$$

for G , respectively. We use $t_\nu \in \widetilde{W}$ to denote the image of $\nu \in X^*(T)$.

The Weyl groups $W(G)$, $\widetilde{W}$, and W_a act naturally on $X^*(T)$. If A is any ring, then the above Weyl groups act naturally on $X^*(T) \otimes_{\mathbb{Z}} A$ by extension of scalars. Let M be a free $\mathbb{Z}$ -module of finite rank (e.g. $M = X^*(T)$). The duality pairing between M and its $\mathbb{Z}$ -linear dual M^* will be denoted by $\langle \cdot, \cdot \rangle$. If A is any ring, the pairing $\langle \cdot, \cdot \rangle$ extends by A -linearity to a pairing between $M \otimes_{\mathbb{Z}} A$ and $M^* \otimes_{\mathbb{Z}} A$, and by an abuse of notation it will be denoted with the same symbol $\langle \cdot, \cdot \rangle$.

We write $G^{\vee} = G^{\vee}_{/\mathbb{Z}}$ for the split connected reductive group over $\mathbb{Z}$ determined by the root datum $(X_*(T), X^*(T), \Phi^{\vee}, \Phi)$. This defines a maximal split torus $T^{\vee} \subseteq G^{\vee}$ such that we have canonical identifications $X^*(T^{\vee}) \cong X_*(T)$ and $X_*(T^{\vee}) \cong X^*(T)$.

Let $V \stackrel{\text{def}}{=} X^*(T) \otimes_{\mathbb{Z}} \mathbb{R}$. For $(\alpha, k) \in \Phi \times \mathbb{Z}$, we have the root hyperplane $H_{\alpha, k} \stackrel{\text{def}}{=} \{x \in V \mid \langle \lambda, \alpha^{\vee} \rangle = k\}$. An *alcove* is a connected component of $V \setminus (\bigcup_{(\alpha, n)} H_{\alpha, n})$, and we denote by $\mathcal{A}$ the set of alcoves. We say that an alcove A is *restricted* if $0 < \langle \lambda, \alpha^{\vee} \rangle < 1$ for all $\alpha \in \Delta$ and $\lambda \in A$. We let A_0 denote the (dominant) base alcove, i.e. the set of $\lambda \in X^*(T) \otimes_{\mathbb{Z}} \mathbb{R}$ such that $0 < \langle \lambda, \alpha^{\vee} \rangle < 1$ for all $\alpha \in \Phi^+$. Recall that $\widetilde{W}$ acts transitively on the set of alcoves, and $\widetilde{W} \cong \widetilde{W}_a \rtimes \Omega$ where Ω is the stabilizer of A_0 . We define

$$\widetilde{W}^+ \stackrel{\text{def}}{=} \{\tilde{w} \in \widetilde{W} \mid \tilde{w}(A_0) \text{ is dominant}\}$$

$$\text{and } \widetilde{W}_1^+ \stackrel{\text{def}}{=} \{\tilde{w} \in \widetilde{W}^+ \mid \tilde{w}(A_0) \text{ is restricted}\}.$$

We fix an element $\eta \in X^*(T)$ such that $\langle \eta, \alpha^{\vee} \rangle = 1$ for all positive simple roots α and let $\tilde{w}_h$ be $w_0 t_{-\eta} \in \widetilde{W}_1^+$.

When $G = \text{GL}_n$, we fix an isomorphism $X^*(T) \cong \mathbb{Z}^n$ in the standard way, where the standard i -th basis element $\varepsilon_i \stackrel{\text{def}}{=} (0, \dots, 1, \dots, 0)$ (with the 1 in the i -th position) of the right-hand side corresponds to extracting the i -th diagonal entry of a diagonal matrix. In particular, we can write any root $\beta \in \Phi$ as $\beta = \varepsilon_i - \varepsilon_j$ for uniquely chosen $1 \leq i, j \leq n$, $i \neq j$.

Given a finite set $\mathcal{J}$ and an isomorphism $G \xrightarrow{\sim} \text{GL}_n^{\mathcal{J}}$ we use superscripts in the notations above, e.g. $\Phi^{+, \mathcal{J}} \subset \Phi^{\mathcal{J}}$, $\Delta^{\mathcal{J}}$, $X^*(T)^{\mathcal{J}}$, $W^{\mathcal{J}}$, etc., where now Φ^+ , Φ , Δ , $X^*(T)$, W , etc. are relative to GL_n . In order not to overload notations, we do not use underlined notations for the elements of $\Phi^{\mathcal{J}}$, $X^*(T)^{\mathcal{J}}$, $W^{\mathcal{J}}$, etc., so that for instance a root $\alpha \in \Phi^{\mathcal{J}}$ is in fact a collection of roots $(\alpha^{(j)})_{j \in \mathcal{J}}$ where each $\alpha^{(j)}$ is a root of GL_n . Finally, we take $\eta \in X^*(T)^{\mathcal{J}}$ to correspond to the element $(n-1, n-2, \dots, 0)_{j \in \mathcal{J}} \in (\mathbb{Z}^n)^{\mathcal{J}}$ in the identification above. When an element $j \in \mathcal{J}$ is fixed, we will abuse notation and will use the same symbol η to denote the element which corresponds to the tuple $(n-1, \dots, 1, 0)$ at j .

We let F_p^+ be a finite unramified étale $\mathbb{Q}_p$ -algebra so that F_p^+ is isomorphic to a product $\prod_{S_p} F_v^+$ over a finite set S_p where, for each $v \in S_p$, F_v^+ is a finite unramified extension of $\mathbb{Q}_p$. For each $v \in S_p$ let $\mathcal{O}_{F_v^+}$ be the ring of integers of F_v^+ , k_v the residue field and let $\mathcal{O}_p$ (resp. k_p) be the product $\prod_{v \in S_p} \mathcal{O}_{F_v^+}$ (resp. $\prod_{v \in S_p} k_v$). (This will be used in global applications, where S_p will be a finite set of places dividing p of a number field F^+ .)

If G is a split connected reductive group over $\mathbb{F}_p$, with Borel B and maximal torus T , we let $G_0 \stackrel{\text{def}}{=} \text{Res}_{k_p/\mathbb{F}_p} G/k_p$, and similarly define B_0 , T_0 . We will always assume that $\mathbb{F}$ contains the image of any ring homomorphism $k_p \rightarrow \overline{\mathbb{F}}_p$ so that we can and do fix an isomorphism $(G_0 \times_{\text{Spec } \mathbb{F}_p} \text{Spec } \mathbb{F}) \xrightarrow{\sim} (G \times_{\text{Spec } \mathbb{F}_p} \text{Spec } \mathbb{F})^{\mathcal{J}_p}$ where $\mathcal{J}_p$ denotes the set of ring homomorphisms $k_p \rightarrow \mathbb{F}$. For notational convenience, we will

write $\underline{G} \stackrel{\text{def}}{=} G_0 \times_{\text{Spec } \mathbb{F}_p} \text{Spec } \mathbb{F}$, and similarly for $\underline{B}$, $\underline{T}$. The notations $\underline{W}$, $\widetilde{\underline{W}}$, $\widetilde{\underline{W}}^+$, $\widetilde{\underline{W}}_1^+$, etc. as well as the identifications $\underline{W} \xrightarrow{\sim} W^{\mathcal{I}_p}$, $\widetilde{\underline{W}} \xrightarrow{\sim} \widetilde{W}^{\mathcal{I}_p}$, $\widetilde{\underline{W}}^+ \xrightarrow{\sim} (\widetilde{W}^+)^{\mathcal{I}_p}$, $\widetilde{\underline{W}}_1^+ \xrightarrow{\sim} (\widetilde{W}_1^+)^{\mathcal{I}_p}$, etc. should be clear.

1.3.2. Galois theory. Let K be a finite extension of $\mathbb{Q}_p$, with residue field k of degree f over $\mathbb{F}_p$. We assume that $K/\mathbb{Q}_p$ is unramified and write $W(k)$ for the ring of Witt vectors, which is also the ring of integers of K . The arithmetic Frobenius automorphism on $W(k)$, acting as raising to p -th power on the residue field will be denoted by φ . We fix an embedding σ_0 of K into E (equivalently an embedding k into $\mathbb{F}$) and define $\sigma_j = \sigma_0 \circ \varphi^{-j}$. This gives an identification between $\mathcal{J} = \text{Hom}(k, \mathbb{F})$ and $\mathbb{Z}/f\mathbb{Z}$.

We normalize Artin's reciprocity map $\text{Art}_K : K^\times \rightarrow W_K^{\text{ab}}$ so that uniformizers are sent to geometric Frobenius elements. We fix once and for all a sequence $(p_m)_{m \in \mathbb{N}} \in \overline{K}^{\mathbb{N}}$ satisfying $p_{m+1}^p = p_m$, $p_0 \stackrel{\text{def}}{=} -p \in K$ and let K_∞ be $\bigcup_{m \in \mathbb{N}} K(p_m)$.

Given an element $\pi_1 \stackrel{\text{def}}{=} (-p)^{\frac{1}{p^f-1}} \in \overline{K}$ we have a character $\omega_K : I_K \rightarrow W(k)^\times$ defined by the condition $g(\pi_1) = \omega_K(g)\pi_1$. Using our choice of embedding σ_0 this gives a fundamental character of niveau f

$$\omega_f := \sigma_0 \circ \omega_K : I_K \rightarrow \mathcal{O}^\times.$$

Let $\rho : G_K \rightarrow \text{GL}_n(E)$ be a p -adic, de Rham Galois representation. For $\sigma : K \hookrightarrow E$, we define $\text{HT}_\sigma(\rho)$ to be the multiset of σ -labeled Hodge-Tate weights of ρ , i.e. the set of integers i such that $\dim_E(\rho \otimes_{\sigma, K} \mathbb{C}_p(-i))^{G_K} \neq 0$ (with the usual notation for Tate twists). In particular, the cyclotomic character ε has Hodge-Tate weights 1 for all embedding $\sigma : K \hookrightarrow E$. For $\mu = (\mu^{(j)})_j \in X^*(\underline{T})$ we say that ρ has Hodge-Tate weights μ if for all $j \in \mathcal{J}$

$$\text{HT}_{\sigma_j}(\rho) = \{\mu_1^{(j)}, \mu_2^{(j)}, \dots, \mu_n^{(j)}\}.$$

The *inertial type* of ρ is the isomorphism class of $\text{WD}(\rho)|_{I_K}$, where $\text{WD}(\rho)$ is the Weil-Deligne representation attached to ρ as in [CDT99], Appendix B.1 (in particular, $\rho \mapsto \text{WD}(\rho)$ is *covariant*). An inertial type is a morphism $\tau : I_K \rightarrow \text{GL}_n(E)$ with open kernel and which extends to the Weil group W_K of G_K . We say that ρ has type (μ, τ) if ρ has Hodge-Tate weights μ and inertial type given by (the isomorphism class of) τ .

1.3.3. Miscellaneous. Finally, δ_P denotes the Kronecker delta function on the condition P . We also use δ for the defect function in §6.2. This shall cause no confusion.

2. PRELIMINARIES

2.1. Affine Weyl groups, tame inertial types, Serre weights.

2.1.1. Affine Weyl group. We collect here the necessary background to give a classification of colength one elements in the admissible set (Proposition 2.1.2).

Recall from §1.3.1 that G is a split reductive group with split maximal torus T . We write W for the Weyl group associated to (G, T) and $V \stackrel{\text{def}}{=} X^*(T) \otimes \mathbb{R} \cong X_*(T^\vee) \otimes \mathbb{R}$ for the apartment of (G, T) on which $\widetilde{W} \stackrel{\text{def}}{=} X^*(T) \rtimes W$ acts. We write $\mathcal{C}_0$ for the dominant Weyl chamber in V .

Recall that $\mathcal{A}$ denotes the set of alcoves of $X^*(T) \otimes \mathbb{R}$ and that $A_0 \in \mathcal{A}$ denotes the dominant base alcove. We let $\uparrow$ denote the upper arrow ordering on alcoves

as defined in [Jan03, §II.6.5] which induces the ordering $\uparrow$ on W_a via the bijection $W_a \xrightarrow{\sim} \mathcal{A}$ given by $\tilde{w} \mapsto \tilde{w}(A_0)$. The dominant base alcove A_0 defines a set of simple reflections in W_a and thus a Coxeter length function on W_a denoted $\ell(-)$ and a Bruhat order on W_a denoted by $\leq$. Given $\lambda \in X^*(T)$ we consider the set of λ -admissible elements of $\widetilde{W}$:

$$(1) \quad \text{Adm}(\lambda) \stackrel{\text{def}}{=} \left\{ \tilde{w} \in \widetilde{W} \mid \tilde{w} \leq t_{w(\lambda)} \text{ for some } w \in W \right\}.$$

If $\Omega \subset \widetilde{W}$ is the stabilizer of the base alcove, then $\widetilde{W} = W_a \rtimes \Omega$ and so $\widetilde{W}$ inherits a Bruhat order in the standard way: For $\tilde{w}_1, \tilde{w}_2 \in W_a$ and $\delta \in \Omega$, $\tilde{w}_1 \delta \leq \tilde{w}_2 \delta$ if and only if $\tilde{w}_1 \leq \tilde{w}_2$, and elements in different right W_a -cosets are incomparable. We extend $\ell(-)$ to $\widetilde{W}$ by letting $\ell(\tilde{w}\delta) \stackrel{\text{def}}{=} \ell(\tilde{w})$ for any $\tilde{w} \in W_a$, $\delta \in \Omega$.

Let $(\widetilde{W}^\vee, \leq)$ be the following partially ordered group: $\widetilde{W}^\vee$ is identified with $\widetilde{W}$ as a group, and $\ell(-)$ and $\leq$ are defined with respect to the *antidominant* base alcove. If $\tilde{w} = t_\nu w \in \widetilde{W}$ with $w \in W$ and $\nu \in X^*(T)$ we define $\tilde{w}^* \stackrel{\text{def}}{=} w^{-1} t_\nu \in \widetilde{W}^\vee$. (The assignment $\tilde{w} \mapsto \tilde{w}^*$ defines an involution which preserves length and Bruhat order, see [LLHL19, Lemma 2.1.3].) Given $\lambda \in X^*(T)$ we define the set $\text{Adm}^\vee(\lambda)$ by replacing $\widetilde{W}$ by $\widetilde{W}^\vee$ in the right hand side of (1).

We also recall that, given $m \in \mathbb{Z}$ and $\alpha \in \Phi$, the m -th α -strip is the subset of V defined by

$$\{x \in V \mid m < \langle x, \alpha^\vee \rangle < m + 1\}.$$

Finally, we say that $\tilde{w} \in \widetilde{W}$ is *regular* if it is in the sense of [LLHLMb, Definition 2.1.3].

Proposition 2.1.2. *Suppose $\tilde{w} \in \text{Adm}(\eta)$ such that $\ell(\tilde{w}) = \ell(t_\eta) - 1$. Then one of the following holds:*

- (1) $\tilde{w} = wt_\eta s_\alpha w^{-1}$ where $w \in W$ and $\alpha > 0$ a positive root such that $w(\alpha) > 0$, and there are no decompositions $\alpha = \beta_1 + \beta_2$ with $\beta_i > 0$ such that $w(\beta_i) > 0$;
- (2) $\tilde{w} = wt_{\eta-\alpha} s_\alpha w^{-1}$ where $w \in W$, $\alpha \in \Phi^+ \setminus \Delta$ such that $w(\alpha) < 0$, and for any decomposition $\alpha = \beta_1 + \beta_2$ with $\beta_i > 0$, we have $w(\beta_i) < 0$.

We say that a colength one element $\tilde{w} \in \text{Adm}(\eta)$ is of the *first form* (resp. of the *second form*) if it is as in item (1) (resp. as in item (2)) of Proposition 2.1.2. Recalling [LLHLMb, Definition 2.1.3] we note that a colength one element $\tilde{w} \in \text{Adm}(\eta)$ is irregular exactly when it is of the first form and moreover the root α appearing in (1) is a simple root.

Remark 2.1.3. If $G = \text{GL}_n$ and $\alpha = \varepsilon_i - \varepsilon_j \in \Phi^+ \setminus \Delta$ then

- (1) the condition in item (1) means that w preserves the order of i and j , and w maps no element $k \in (i, j)$ into an element in $(w(i), w(j))$;
- (2) the condition in item (2) means that w reverses the order of i and j , and maps any $k \in (i, j)$ to an element in $(w(j), w(i))$.

Proof. We assume $\tilde{w}(A_0)$ is in chamber $w(\mathcal{C}_0)$. Then there is a gallery from A_0 to $\tilde{w}(A_0)$ in the w -positive direction (cf. [HC02, §2, Definition 5.2]), hence [HH17, Corollary 4.4] shows that $\tilde{w} < wt_\eta w^{-1}$. Thus $\tilde{u} \stackrel{\text{def}}{=} w^{-1} \tilde{w}$ is dominant so that $w\tilde{u}$ is a reduced expression for $\tilde{w}$ by [LLHLMa, Lemma 2.2.1] (cf. Definition 2.1.2 in *loc. cit.* for the notion of reduced expression). We conclude that $\tilde{u} \leq t_\eta w^{-1}$ and $\ell(\tilde{u}) = \ell(t_\eta w^{-1}) - 1$. Since $\tilde{u}$ and $t_\eta w^{-1}$ are both dominant and their lengths differ

by 1 they must differ by an affine reflection in direction α for some positive root $\alpha > 0$ by [GHS18, Corollary A.1.2]. We claim that the α -strip containing $\tilde{u}$ is the lower neighbor of the α -strip containing $t_\eta w^{-1}$.

Suppose the contrary. Then for each vertex v of A_0 , the line segment joining $\tilde{u}(v)$ and $t_\eta w^{-1}(v)$ lies in the dominant Weyl chamber and contains $t_\alpha \tilde{u}(v)$. This shows that $t_\alpha \tilde{u}(A_0)$ is dominant and $\tilde{u} \uparrow t_\alpha \tilde{u} \uparrow t_\eta w^{-1}$. [Wan87, Theorem 4.3] implies that $\tilde{u} < t_\alpha \tilde{u} < t_\eta w^{-1}$ which contradicts $\ell(\tilde{u}) = \ell(t_\eta w^{-1}) - 1$.

There are two cases:

- $\tilde{u}(A_0)$ and $t_\eta w^{-1}(A_0)$ share the vertex η . In this case $\tilde{u} = t_\eta s_\alpha w^{-1}$. The conditions $\tilde{u} \leq t_\eta w^{-1}$ and $\ell(\tilde{u}) = \ell(t_\eta w^{-1}) - 1$ are equivalent to $\ell(s_\alpha w^{-1}) = \ell(w^{-1}) + 1$, equivalently $\ell(ws_\alpha) = \ell(w) + 1$. But this condition translates exactly to the condition in item (1).
- $\tilde{u}(A_0)$ has $\eta - \alpha$ as a vertex. In this case we must have $\tilde{u} = t_{\eta-\alpha} s_\alpha w^{-1}$. To see the condition on w , we note that for each positive root β we have

$$\langle \eta, \beta^\vee \rangle + \delta_{w(\beta) > 0} - 1 - (\langle \eta, \beta^\vee \rangle - \langle \alpha, \beta^\vee \rangle + \delta_{ws_\alpha(\beta) > 0} - 1) = \langle \alpha, \beta^\vee \rangle + \delta_{w(\beta) > 0} - \delta_{ws_\alpha(\beta) > 0}$$

and hence by the explicit formula for the length function (e.g. [He16, §1.7.1]) the colength 1 condition becomes

$$\sum_{\beta \neq \alpha, \beta > 0} -\delta_{w(\beta) > 0} + \delta_{ws_\alpha(\beta) > 0} = 2\langle \eta, \alpha^\vee \rangle - 2.$$

This gives the condition in item (2).

This completes the proof. $\square$

Lemma 2.1.4. *Let $\tilde{w} \in \text{Adm}^{\text{reg}}(\eta)$ satisfy $\ell(\tilde{w}) \geq \ell(t_\eta) - 1$, and let $\tilde{w}_2^{-1} w_0 t_\nu \tilde{w}_1$ be the unique up to $X^0(T)$ factorization of $\tilde{w}$ with $\tilde{w}_1, \tilde{w}_2 \in \widetilde{W}_1^+$ and $\nu \in X^+(T)$ (see [LLHLMb, Proposition 2.1.5]). Suppose that $\tilde{w} = \tilde{x}_2^{-1} s \tilde{x}_1$ with $\tilde{x}_1, \tilde{x}_2 \in \widetilde{W}^+$, $s \in W$, and that $\tilde{x}_1 \uparrow \tilde{x} \uparrow \tilde{w}_h^{-1} \tilde{x}_2$ with $\tilde{x} \in \widetilde{W}_1^+$. Then $s = w_0$ and either $\tilde{x} = \tilde{x}_1 \in \tilde{w}_1 X^0(T)$ or $\tilde{x} = \tilde{w}_h^{-1} \tilde{x}_2 \in \tilde{w}_h^{-1} \tilde{w}_2 X^0(T)$. If moreover $\ell(\tilde{w}) = \ell(t_\eta)$ we further have $\tilde{x} = \tilde{x}_1 = \tilde{w}_h^{-1} \tilde{x}_2$ and $\nu \in X^0(T)$.*

Proof. Let $\tilde{w}, \tilde{x}_1, \tilde{x}_2, s$, and $\tilde{x}$ be as above. Since $\tilde{x}_1 \uparrow \tilde{x} \uparrow \tilde{w}_h^{-1} \tilde{x}_2$, we have that $\tilde{x}_1 \leq \tilde{x}$ and $\tilde{x}_2 \leq \tilde{w}_h \tilde{x}$ by [LLHL19, Theorem 4.1.1, Proposition 4.1.2] as $\tilde{w}_h \tilde{x} \in \widetilde{W}^+$. Then we have

$$\begin{aligned} 1 &\geq \ell(t_\eta) - \ell(\tilde{w}) \\ &\geq \ell((\tilde{w}_h \tilde{x})^{-1} w_0 \tilde{x}) - [\ell(\tilde{x}_2^{-1}) + \ell(s) + \ell(\tilde{x}_1)] \\ &= [\ell(\tilde{w}_h \tilde{x}) - \ell(\tilde{x}_2)] + [\ell(w_0) - \ell(s)] + [\ell(\tilde{x}) - \ell(\tilde{x}_1)], \end{aligned}$$

where the last equality follows from [LLHL19, Lemma 4.1.9]. Since each term in the last expression is nonnegative, we immediately see that $\tilde{x} = \tilde{x}_1$ or $\tilde{w}_h^{-1} \tilde{x}_2$. (If moreover $\ell(\tilde{w}) = \ell(t_\eta)$ we further have $s = w_0$ and $\tilde{x}_1 = \tilde{x} = \tilde{w}_h^{-1} \tilde{x}_2$.) It suffices to show that $\tilde{x}_1 \in \tilde{w}_1 X^0(T)$ in the former case and $\tilde{x}_2 \in \tilde{w}_2 X^0(T)$ in the latter case.

We next show that $s = w_0$. Supposing otherwise, $\tilde{x}_1 = \tilde{x} = \tilde{w}_h^{-1}\tilde{x}_2$ and $s = w_0s_\alpha$ with $\alpha \in \Delta^+$. By Proposition 2.1.2 and the paragraph following it, we conclude that $\tilde{w}$ is not regular which is a contradiction.

Now suppose that $\tilde{x} = \tilde{x}_1$ so that we have $\tilde{x}_2^{-1}w_0\tilde{x}_1 = \tilde{w}_2^{-1}w_0t_\nu\tilde{w}_1$. The uniqueness in [LLHLMb, Proposition 2.1.5] guarantees that $\tilde{x}_1 \in \tilde{w}_1X^0(T)$ (and, if moreover $\ell(t_\eta) = \ell(\tilde{w})$, that $\nu \in X^0(T)$). Similarly, if $\tilde{x} = \tilde{w}_h^{-1}\tilde{x}_2$, then $\tilde{x}_2 \in \tilde{W}_1^+$ and uniqueness again guarantees that $\tilde{x}_2 \in \tilde{w}_2X^0(T)$. $\square$

For Lemma 2.1.5, which explicitly describes the unique up to $X^0(T)$ -decomposition of colength one regular admissible elements mentioned in Lemma 2.1.4, we specialize to the case $G = \mathrm{GL}_n$. Recall that in this case we have the elements $\varepsilon_i \in X^*(T)$ for $i = 1, \dots, n$ and the fundamental weights $\omega'_{\varepsilon_i - \varepsilon_{i+1}} \stackrel{\text{def}}{=} \sum_{k=1}^i \varepsilon_k$ for $1 \leq i \leq n-1$. We have the inclusion

$$(2) \quad W \hookrightarrow \widetilde{W}_1^+, \quad w \longmapsto t_{\eta_w}w,$$

where t_{η_w} is the dominant weight defined by

$$(3) \quad t_{\eta_w} \stackrel{\text{def}}{=} \sum_{\beta \in \Delta, w^{-1}(\beta) < 0} \omega'_\beta$$

(see [Her09, equation (5.1)]). Given $w \in W$ we write $\tilde{w}$ for the image of w via the inclusion (2).

Lemma 2.1.5. *Let $\tilde{w} \in \mathrm{Adm}^{\mathrm{reg}}(\eta)$ satisfy $\ell(\tilde{w}) = \ell(t_\eta) - 1$. Then in the decomposition $\tilde{w} = \tilde{w}_2^{-1}w_0t_\nu\tilde{w}_1$ of Lemma 2.1.4, we can take $\tilde{w}_1 = s_\alpha w^{-1}$, $\tilde{w}_2 = \widetilde{w_h w^{-1}}$ and $\nu = \eta_{w^{-1}} - \eta_{s_\alpha w^{-1}} - \varepsilon\alpha$, where ε equals 1 (resp. 0) if we are in case (2) (resp. case (1)) of Proposition 2.1.2.*

Proof. Letting $\tilde{w} = wt_{\eta - \varepsilon\alpha}s_\alpha w^{-1}$, where $\varepsilon \in \{0, 1\}$ equals 1 (resp. 0) if we are in case (2) (resp. (1)) of Proposition 2.1.2, we have

$$\tilde{w} = (\widetilde{w_h w^{-1}})^{-1}w_0t_\nu s_\alpha w^{-1}$$

by (2) and the definition of $\tilde{w}_h$. It thus suffices to prove that ν is dominant, i.e. using equation (3), that

$$\delta_{w(\beta) > 0} - \delta_{ws_\alpha(\beta) > 0} - \varepsilon\langle \alpha, \beta^\vee \rangle \geq 0$$

for all $\beta \in \Delta$. This is an elementary casewise check, based on the value of $\langle \alpha, \beta^\vee \rangle \in \{-1, 0, 1\}$ and Remark 2.1.3, and which we leave to the reader. $\square$

2.2. Local models, affine charts and monodromy conditions. For any Noetherian $\mathcal{O}$ -algebra R define

$$\begin{aligned} L\mathcal{G}(R) &\stackrel{\text{def}}{=} \{A \in \mathrm{GL}_n(R((v+p))) \mid A \text{ is upper triangular modulo } v\}; \\ L^+\mathcal{M}(R) &\stackrel{\text{def}}{=} \{A \in \mathrm{Mat}_n(R[[v+p]]) \mid A \text{ is upper triangular modulo } v\}. \end{aligned}$$

2.2.1. Affine charts. Let $\tilde{z} = zt_\nu \in \widetilde{W}^\vee$. Given an integer $h \geq 0$ define the subfunctor $\mathcal{U}(\tilde{z})^{\det, \leq h} \subseteq L\mathcal{G}$ defined on Noetherian $\mathcal{O}$ -algebras R as follows: $\mathcal{U}(\tilde{z})^{\det, \leq h}(R)$

is the set of matrices $A \in L\mathcal{G}(R)$ such that for all $1 \leq i, k \leq n$ the following holds:

- $(v+p)^h A_{ik} \in v^{\delta_{i>k}} R[v]$;
- $\deg((v+p)^h A_{ik}) \leq h + \nu_k - \delta_{i<z(k)}$;
- $(v+p)^h A_{z(k)k}$ is a monic polynomial; and
- $\det(A) = \det(z)(v+p)^{\|\nu\|}$,

where we have written $\nu = (\nu_\ell)_{1 \leq \ell \leq n} \in X^*(T)$ and $\|\nu\| \stackrel{\text{def}}{=} \sum_{\ell=1}^n \nu_\ell$.

We now introduce the subfunctors and $T^\vee$ -torsors which are relevant for our analysis of deformation rings. In the above setting, define the following subfunctors of $\mathcal{U}(\tilde{z})^{\det, \leq 0}$:

- $U^{[0, n-1]}(\tilde{z}) \subseteq \mathcal{U}(\tilde{z})^{\det, \leq 0}$ as the subfunctor whose R -valued points are those $A \in \mathcal{U}(\tilde{z})^{\det, \leq 0}$ such that both A and $(v+p)^{n-1} A^{-1}$ are in $L^+ \mathcal{M}(R)$; note that $U^{[0, n-1]}(\tilde{z})$ is representable by an affine scheme over $\mathcal{O}$;
- $U(\tilde{z}, \leq \eta) \subseteq U^{[0, n-1]}(\tilde{z})$ as the closed, $\mathcal{O}$ -flat and reduced subscheme of $U^{[0, n-1]}(\tilde{z})$ whose points on a reduced $\mathcal{O}$ -flat algebra R consist of $A \in U^{[0, n-1]}(\tilde{z})(R)$ with elementary divisors bounded by $(v+p)^\eta$ (i.e. each k by k minor of A is divisible by $(v+p)^{\frac{(k-1)k}{2}}$); and
- $\tilde{U}(\tilde{z}, \leq \eta) \stackrel{\text{def}}{=} T^\vee U(\tilde{z}, \leq \eta)$ as the subscheme of $L\mathcal{G}$ whose R -valued points are of the form DA where $D \in T^\vee(R)$ and $A \in U(\tilde{z}, \leq \eta)(R)$; note that $\tilde{U}(\tilde{z}, \leq \eta)$ is endowed with a $T^\vee_{\mathcal{O}}$ -action induced by left multiplication of matrices, and we have $[T^{\vee, \mathcal{J}} \backslash \tilde{U}(\tilde{z}, \leq \eta)] \cong U(\tilde{z}, \leq \eta)$.

The entries of $A \in U^{[0, n-1]}(\tilde{z})(R)$ will typically be written in the form

$$A_{ik} = v^{\delta_{i>k}} \left(\sum_{\ell=0}^{\nu_k - \delta_{i>k} - \delta_{i<z(k)}} a_{ik, \ell} (v+p)^\ell \right)$$

so that $U^{[0, n-1]}(\tilde{z})$ is an affine $\mathcal{O}$ -scheme whose global functions is the quotient of the polynomial $\mathcal{O}$ -algebra in the variables $a_{ik, \ell}$ (for the appropriate range of ℓ determined by $\tilde{z}$ and (i, k)) by the ideal determined by the condition $\det(A) = \det(z)(v+p)^{\|\nu\|}$. Then $U(\tilde{z}, \leq \eta)$ is a closed subscheme of $U^{[0, n-1]}(\tilde{z})$, given by the p -saturation of the condition that each k by k minor is divisible by $(v+p)^{\frac{(k-1)k}{2}}$. In general, these schemes can be somewhat hard to describe; however in Proposition 3.1.2 and Proposition 3.2.2, we will give an explicit presentations for them when $\tilde{z}$ has colength one.

2.2.2. Algebraic monodromy condition. Let $\mathbf{a} \in \mathcal{O}^n$. We define an operator $\nabla_{\mathbf{a}}$ on $L\mathcal{G}$ by $\nabla_{\mathbf{a}}(A) \stackrel{\text{def}}{=} (v \frac{d}{dv}(A) - [\text{Diag}(\mathbf{a}), A])$, and a closed subfunctor $L\mathcal{G}^{\nabla_{\mathbf{a}}} \subseteq L\mathcal{G}$ by

$$L\mathcal{G}^{\nabla_{\mathbf{a}}}(R) \stackrel{\text{def}}{=} \left\{ A \in L\mathcal{G}(R) \mid \nabla_{\mathbf{a}}(A) A^{-1} \in \frac{1}{(v+p)} L^+ \mathcal{M}(R) \right\}.$$

(We write $[M, N]$ for the usual Lie bracket on $L\mathcal{G}$, defined by $[M, N] \stackrel{\text{def}}{=} MN - NM$.)

We define the following closed subschemes of $U(\tilde{z}, \leq \eta)$:

- $U^{\text{nv}}(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}}) \stackrel{\text{def}}{=} U(\tilde{z}, \leq \eta) \cap L\mathcal{G}^{\nabla_{\mathbf{a}}}$; and
- $U(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}}) \subseteq U^{\text{nv}}(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}})$ as the p -flat closure of $U^{\text{nv}}(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}})_E$ inside $U(\tilde{z}, \leq \eta)$.

Thus $U(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}})$ is the closed subscheme $U(\tilde{z}, \leq \eta)$ whose ring of global functions is the quotient of $\mathcal{O}(U(\tilde{z}, \leq \eta))$ by the p -saturation of the ideal $I_{\nabla_{\mathbf{a}}}$ obtained by imposing that the universal matrix

$$A^{\text{univ}} \in U(\tilde{z}, \leq \eta)(\mathcal{O}(U(\tilde{z}, \leq \eta)))$$

satisfies

$$(\nabla_{\mathbf{a}}(A^{\text{univ}}))(A^{\text{univ}})^{-1} \in \frac{1}{(v+p)} L^+ \mathcal{M}(\mathcal{O}(U(\tilde{z}, \leq \eta))),$$

cf. [LLHLMb, Definition 7.1.8] for a list of generators of this ideal. Proposition 4.3.2 and Proposition 4.3.1 give results towards an explicit presentation for the affine scheme $U(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}})$ when $\tilde{z}$ has colength at most one.

We define analogously the closed subschemes $\tilde{U}(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}}) \subseteq \tilde{U}^{\text{nv}}(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}}) \subseteq \tilde{U}(\tilde{z}, \leq \eta)$, replacing $U(\tilde{z}, \leq \eta)$ by $\tilde{U}(\tilde{z}, \leq \eta)$ in the above items. Note that we also have

$$(4) \quad \tilde{U}(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}}) = T^{\vee} U(\tilde{z}, \leq \eta, \nabla_{\mathbf{a}}),$$

compatible with $\tilde{U}(\tilde{z}, \leq \eta) \stackrel{\text{def}}{=} T^{\vee} U(\tilde{z}, \leq \eta)$.

2.2.3. Monodromy condition and Galois representations. We follow the notation and terminology on tame inertial types and their lowest alcove presentations of [LLHLMb, §2.4]. In particular given $(s, \mu) = (s^{(j)}, \mu^{(j)})_{j \in \mathcal{J}} \in W^{\mathcal{J}} \times (X^*(T) \cap C_0)^{\mathcal{J}}$, [LLHLMb, Example 2.4.1, equations (5.2), (5.1)] produces a tame inertial type $\tau(s, \mu + \eta)$ and n -tuples $\mathbf{a}'^{(j')} \in \mathbb{Z}^n$ for $0 \leq j' \leq fr - 1$ (where r is the order of the element $s^{(0)} s^{(1)} \dots s^{(f-2)} s^{(f-1)} \in W$). If μ is 1-deep in $\underline{C}_0$, for each $0 \leq j' \leq fr - 1$ we define $s'_{\text{or}, j'}$ to be the element of W such that $(s'_{\text{or}, j'})^{-1}(\mathbf{a}'^{(j')}) \in \mathbb{Z}^n$ is dominant, and let

$$(5) \quad \mathbf{a}^{(j)} \stackrel{\text{def}}{=} (s'_{\text{or}})^{(j)}{}^{-1}(\mathbf{a}'^{(j)}) / (1 - p^{fr})$$

for $0 \leq j \leq f$.

Recall from [LLHLMb, §7.2] the p -adic formal algebraic stack $\mathcal{X}^{\leq \eta, \tau}$ over $\text{Spf} \mathcal{O}$. It is p -flat, equidimensional of dimension $(1 + [K : \mathbb{Q}_p] \binom{n}{2})$ and moreover if $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ is a continuous Galois representation, then $R_{\bar{\rho}}^{\leq \eta, \tau}$ is a versal ring for $\mathcal{X}^{\leq \eta, \tau}$ at its point corresponding to $\bar{\rho}$ (see [EG, §4.8]). Let $\tilde{z} = (\tilde{z}^{(j)})_{j \in \mathcal{J}} \in \widetilde{W}^{\vee, \mathcal{J}}$.

Let $N \geq n + 1$ and assume that μ is N -deep in alcove $\underline{C}_0$. From [LLHLMb, Theorem 7.2.3, Theorem 7.3.2] we have an open substack $\mathcal{X}^{\leq \eta, \tau}(\tilde{z}) \hookrightarrow \mathcal{X}^{\leq \eta, \tau}$, an $\mathcal{O}$ -flat closed p -adic formal subscheme $\tilde{U}(\tilde{z}, \leq \eta, \nabla_{\tau, \infty})$ of $\tilde{U}(\tilde{z}, \leq \eta)^{\wedge_p}$ (where $\tilde{U}(\tilde{z}, \leq \eta) \stackrel{\text{def}}{=} \prod_{j \in \mathcal{J}} \tilde{U}(\tilde{z}^{(j)}, \leq \eta)$ and the superscript $\wedge_p$ denotes the p -adic completion) and a formally smooth morphism

$$(6) \quad \tilde{U}(\tilde{z}, \leq \eta, \nabla_{\tau, \infty}) \rightarrow \mathcal{X}^{\leq \eta, \tau}(\tilde{z})$$

of relative dimension $n \# \mathcal{J}$.

Assume further that $N \geq 2n - 5$ and μ is N -deep in alcove $\underline{C}_0$. Then [LLHLMb, Proposition 7.1.10] shows that the formal scheme $\tilde{U}(\tilde{z}, \leq \eta, \nabla_{\tau, \infty})$ is the p -flat closure of a natural deformation of $\prod_{j \in \mathcal{J}} \tilde{U}(\tilde{z}^{(j)}, \leq \eta, \nabla_{\mathbf{a}^{(j)}})$ in the following sense: Letting $I_{\nabla_{\mathbf{a}^{(j)}}}^{(j)}$ be the ideal cutting out $\tilde{U}(\tilde{z}^{(j)}, \leq \eta, \nabla_{\mathbf{a}^{(j)}})$ in $\tilde{U}(\tilde{z}^{(j)}, \leq \eta)$, $\tilde{U}(\tilde{z}, \leq \eta, \nabla_{\tau, \infty})$ is cut out by the p -saturation of an ideal $I_{\nabla_{\tau, \infty}}$ which is obtained from $\sum_{j \in \mathcal{J}} I_{\nabla_{\mathbf{a}^{(j)}}}^{(j)}$ by adding a (specific but inexplicit) element divisible by p^{N-2n+5} to each element of its

natural set of generators (in particular, $(I_{\nabla_{\tau,\infty}}, p^{N-2n+5}) = \sum_j (I_{\nabla_{\mathbf{a}(j)}}^{(j)}, p^{N-2n+5})$ by definition). This property can be expressed in the following way: Given a p -adically complete Noetherian $\mathcal{O}$ -algebra R , if $\tilde{A} \in \tilde{U}(\tilde{z}, \leq \eta)^{\wedge p}(R)$ satisfies the equations in $I_{\nabla_{\tau,\infty}}$ then

$$(7) \quad (\nabla_{\mathbf{a}(j)}(\tilde{A}^{(j)}))((v+p)^{n-1}\tilde{A}^{(j)})^{-1} \in (v+p)^{n-2}\mathbf{M}_n(R[v]) + p^{N-2n+5}\mathbf{M}_n(R[[v]])$$

see [LLHLMb, Proposition 7.1.10 and its proof]. (We caution the reader that $I_{\nabla_{\tau,\infty}}$ is *different than* the ideal $\mathbf{I}_{\mathfrak{M},\nabla_{\infty}}$ appearing in [LLHLMb, §7.1], but the p -saturation of $I_{\nabla_{\tau,\infty}}$ equals $\mathbf{I}_{\mathfrak{M},\nabla_{\infty}}$ when $\mathfrak{M}$ is taken to be the universal Breuil–Kisin module over $\tilde{U}(\tilde{z}, \leq \eta)$.)

Finally let us recall the characterization of the Galois representations that contribute to $\mathcal{X}^{\leq \eta, \tau}(\tilde{z})$: given $\bar{\rho} \in \mathcal{X}^{\leq \eta, \tau}(\mathbb{F})$, then $\bar{\rho} \in \mathcal{X}^{\leq \eta, \tau}(\tilde{z})(\mathbb{F})$ if and only if its étale φ -module $\mathcal{M}_{\bar{\rho}} = \mathbb{V}_K^*(\bar{\rho}|_{G_{K_{\infty}}})$, cf. [LLHLMb, §5.5], admits a basis with respect to which the matrix of Frobenius belongs to $\tilde{U}(\tilde{z}, \leq \eta)(\mathbb{F}) \prod_{j \in \mathcal{J}} s_j^{-1} v^{\mu_j + \eta_j}$. We say that $\bar{\rho}$ has shape $\tilde{z}$ with respect to τ if additionally the matrix of Frobenius belongs to $\prod_{j \in \mathcal{J}} \mathcal{I}^{\tilde{z}(j)} \mathcal{I} s_j^{-1} v^{\mu_j + \eta_j}$. (Here $\mathcal{I}$ denotes the Iwahori subgroup of $\mathrm{GL}_n(\mathbb{F}((v)))$ corresponding to the Borel of upper triangular matrices.)

It follows from [LLHLMb, Proposition 5.4.7] that any $\bar{\rho} \in \mathcal{X}^{\leq \eta, \tau}(\mathbb{F})$ has a unique shape in $\mathrm{Adm}(\eta)$, which we denote by $\tilde{w}(\bar{\rho}, \tau)^*$.

3. FINITE HEIGHT CONDITIONS

In this section, we describe $U(\tilde{w}, \leq \eta)$ when $\tilde{w}^* \in \mathrm{Adm}(\eta)$ has colength one. Throughout this subsection R denotes a Noetherian $\mathcal{O}$ -flat $\mathcal{O}$ -algebra, and we use the notation $u_{\beta} : \mathbb{G}_a \hookrightarrow w_0 U w_0$ for the embedding to the β -entry for each $\beta \in \Phi^-$, where U is the subgroup of upper triangular unipotent matrices in GL_n . Moreover, given $A \in \mathrm{Mat}_n(R)$ and $\beta = \varepsilon_i - \varepsilon_j \in \Phi$, we write A_{β} and $A_{\beta, \ell}$ for A_{ij} and $a_{ik, \ell}$ respectively.

3.1. Finite height conditions: the first form. Let $\tilde{w}^* \in \mathrm{Adm}(\eta)$ be a colength one shape of the first form. Thus, by Proposition 2.1.2 and the definition of $\tilde{w} \mapsto \tilde{w}^*$, we have $\tilde{w} = w s_{\alpha} t_{\eta} w^{-1}$, where $w \in W$ satisfies $w(\alpha) > 0$ and $w(k)$ do not belong to the interval $(w(i_0), w(j_0))$ for all $i_0 < k < j_0$ if we set $\alpha = \alpha_{i_0 j_0} \in \Phi^+ \setminus \Delta$. For notational convenience we set $s \stackrel{\mathrm{def}}{=} s_{\alpha}$ and $w' \stackrel{\mathrm{def}}{=} w s$.

With an eye towards the monodromy conditions, we introduce the following:

Definition 3.1.1. We say that a negative root $\beta \in \Phi^-$ is *bad* (with respect to $\tilde{w}$) if

- β shares either the row or the column of α ;
- $\delta_{w(\beta) < 0} \neq \delta_{w'(\beta) < 0}$.

Proposition 3.1.2. Let $A \in U(\tilde{w}, \leq \eta)(R)$.

Then $A_{w(j_0)w(j_0)} = (w^{-1}Aw)_{j_0 j_0} = c(v+p)^{n-j_0}$ for some $c \in R$ and

$$(8) \quad s \cdot u_{-\alpha}(-c) \cdot (w^{-1}Aw) = (v+p)^{\eta} \cdot V,$$

where V is a lower triangular unipotent matrix whose entries are in $R[v]$.

Moreover, for each $\beta \in \Phi^-$ the entry V_{β} can be written as $v^{\kappa_{\beta}} \cdot f_{\beta}(v)$ for a polynomial $f_{\beta}(v) \in R[v]$ of the form $\sum_k c_{\beta, k}(v+p)^k$ where

- (1) if β is bad then $\kappa_{\beta} = \delta_{w(\beta) < 0} = 0$ and $\deg(f_{\beta}) < |\langle \eta, \beta^{\vee} \rangle| + 1$;

- (2) if β shares the row or the column of α but is not bad then $\kappa_\beta = \delta_{w(\beta) < 0}$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle|$;
- (3) if β shares the row of $-\alpha$ then $\kappa_\beta = \delta_{w'(\beta) < 0}$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle|$;
- (4) if β shares the column of $-\alpha$ with $\beta \neq -\alpha$ then $\kappa_\beta = \delta_{w(\beta) < 0}$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle| - \delta_{w(-\alpha) < w(\beta) < 0}$;
- (5) for all other roots $\beta \in \Phi^-$, $\kappa_\beta = \delta_{w(\beta) < 0}$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle|$.

Proof. It is easy to see that $(w^{-1}Aw)_{j_0 j_0}$ is a polynomial of degree less than $n - j_0 + 1$. Let $c \in R$ be the coefficient of v^{n-j_0} in $(w^{-1}Aw)_{j_0 j_0}$, and set $C \stackrel{\text{def}}{=} su_{-\alpha}(-c)(w^{-1}Aw)$ as in (8). One can readily observe the following degree bounds:

- $\deg(C_{ll}) \leq n - l$ for all $1 \leq l \leq n$;
- for $\beta = \alpha_{lm} \in \Phi$ with $l, m \notin \{i_0, j_0\}$, $C_\beta = v^{\delta_{w(\beta) < 0}} f_\beta(v)$ with $\deg(f_\beta) \leq n - m - 1$;
- for $\beta = \alpha_{i_0 m} \in \Phi$, $C_\beta = v^{\delta_{w(\beta) < 0}} f_\beta(v)$ with $\deg(f_\beta) \leq n - m - 1 + \delta_{w'(\beta) < 0 < w(\beta)}$;
- for $\beta = \alpha_{j_0 m} \in \Phi$, $C_\beta = v^{\delta_{w'(\beta) < 0}} f_\beta(v)$ with $\deg(f_\beta) \leq n - m - 1$;
- for $\beta = \alpha_{li_0} \in \Phi$ with $l \neq j_0$, $C_\beta = v^{\delta_{w(\beta) < 0}} f_\beta(v)$ with $\deg(f_\beta) \leq n - i_0 - \delta_{w(\beta) < 0} - \delta_{w(\beta) > w(-\alpha)}$;
- for $\beta = \alpha_{lj_0} \in \Phi$, $C_\beta = v^{\delta_{w(\beta) < 0}} f_\beta(v)$ with $\deg(f_\beta) \leq n - j_0 - \delta_{w(\beta) > w(\alpha)} - \delta_{w(\beta) < 0}$.

From the bullet points above it is now immediate to deduce the degree bounds (1)–(5).

We now prove the claim on the lower triangularity of V , and the divisibility by $(v + p)^\eta$. We will repeatedly use following inductive procedure (modeled on the height condition): Given an $\mathcal{O}$ -flat algebra R and $B \in \text{Mat}_n(R[v])$ such that

- (1) the right-bottom block of B of size $k \times k$ is lower triangular of the following form

$$\begin{pmatrix} B_{(n+1-k)(n+1-k)} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ B_{n(n+1-k)} & \cdots & B_{nn} \end{pmatrix},$$

and $(v + p)^{n-l} \mid B_{lm}$ if B_{lm} is in the $(k \times k)$ -block above;

- (2) for l and m with $1 \leq l, m \leq n - k$, the $(k+1) \times (k+1)$ -submatrix determined by B_{lm} and the $k \times k$ -block above is also lower-triangular;
- (3) $(v + p)^{\frac{k(k+1)}{2}} \mid ((k+1) \times (k+1)\text{-minors})$ for the $(k+1) \times (k+1)$ -submatrix above,

then we have $(v + p)^k \mid B_{lm}$. We apply the above procedure to $B = w^{-1}Aw$ and $B = C$.

First, imposing the finite height condition and inductively using items (1), (2), (3) above to $w^{-1}Aw$, we can assume without loss of generality that $(w^{-1}Aw)_\beta = 0$ for $\beta = \alpha_{lm}$ if $j_0 < m \leq n$ and $l < m$, and that $(v + p)^{n-l} \mid (w^{-1}Aw)_\beta$ for $\beta = \alpha_{lm}$ if $j_0 < l \leq n$ and $m \leq l$ (note that $(w^{-1}Aw)_\beta = 0$ for $\beta = \alpha_{ln}$ if $l < n$, by the degree bounds). Therefore, it is immediate that $C_\beta = 0$ for $\beta = \alpha_{lm}$ if $j_0 < m \leq n$ and $l < m$, and that $(v + p)^{n-l} \mid C_\beta$ for $\beta = \alpha_{lm}$ if $j_0 < l \leq n$ and $m \leq l$. Moreover, by a further application of items (1), (2), (3) above to C , we also conclude that the right-bottom $(n - i_0 + 1) \times (n - i_0 + 1)$ -block of C is as described in the statements, and that $(v + p)^{n-l} \mid C_{lm}$ if $i_0 \leq l \leq n$ and $m \leq l$.

We now check that $C_\beta = 0$ if $\beta = \alpha_{lm}$ with $1 \leq l < i_0$ and $l < m \leq j_0$. Note that we cannot, in general, conclude $C_{lj_0} = 0$ from the fact $(v+p)^{n-j_0} | C_{lj_0}$ induced from items (1), (2), (3) above for $1 \leq l < i_0$, since the degree bound of the polynomial C_{lj_0} is one higher than usual if $w(i_0) < w(l) < w(j_0)$. Let $\beta = \alpha_{lj_0}$ be such a root, and assume that the right-bottom $(n-l) \times (n-l)$ -block of C is as described in the statements. By items (1), (2), (3) and the degree bound, we may let $C_\beta = x(v+p)^{n-j_0}$ for $x \in R$. Consider $C' \stackrel{\text{def}}{=} u_\beta(-x)C$. Then we have $C'_\beta = 0$, and, moreover, the finite height condition together with the degree bounds implies that $C'_{lm} = 0$ for all $i_0 \leq m < j_0$. In particular, we have

$$C'_{li_0} = C_{li_0} - xC_{-\alpha} = 0.$$

By looking at the constant part of C'_{li_0} , which is the same as the constant part of $-xC_{-\alpha}$ as $v | C_{li_0}$, we have $p^{|\langle \eta, -\alpha^\vee \rangle| - 1} x c_{-\alpha} = 0$ in R where we let $p^{|\langle \eta, -\alpha^\vee \rangle| - 1} c_{-\alpha}$ be the constant part of $C_{-\alpha}$. On the other hand, by Lemma 3.1.6 $c_{-\alpha}$ is a unit in $R[\frac{1}{p}]$, so that we conclude that $x = 0$. (Note that to prove Lemma 3.1.6 we only used the fact that the right-bottom $(n-i_0+1) \times (n-i_0+1)$ -block of C is as described in the statements, which is already proved in the previous paragraph.) Now by repeatedly using items (1), (2), (3) above together with the degree bounds, we conclude that $C_{lm} = 0$ if $l < m \leq j_0$. Repeating this argument, we conclude that $C_\beta = 0$ if $\beta = \alpha_{lm}$ with $1 \leq l < i_0$ and $l < m \leq j_0$ and that $(v+p)^{n-l} | C_{lm}$ if $1 \leq m \leq l \leq n$.

Finally, we point out that we have $(w^{-1}Aw)_{j_0j_0} = c(v+p)^{n-j_0}$, which can be readily seen during the induction steps due to the degree bound. This completes the proof. $\square$

For the rest of this subsection, we observe some necessary properties of $\kappa_\beta \in \{0, 1\}$, defined in Proposition 3.1.2, as well as some identities of the coefficients of V from the finite height conditions. We first fix some notation:

- For $\beta \in \Phi^-$, we write $\mathfrak{D}_\beta$ for the set of the decompositions (β_1, β_2) of β into two negative roots with β_1 sharing the column of β (and β_2 sharing the row of β).
- For a bad root $\beta \in \Phi^-$, we write $\mathfrak{A}_\beta$ for the set $\{(\beta_1, \beta_2) \in \mathfrak{D}_\beta \mid \kappa_\beta = \kappa_{\beta_1} + \kappa_{\beta_2}\}$.

The following are immediate consequences of Proposition 3.1.2, which will be frequently used:

- if β shares the row of α then we have $\kappa_\beta = \kappa_{s(\beta)}$;
- if β is a bad root sharing the column of α then $\deg(v^{-\kappa_{s(\beta)}} V_{s(\beta)}) < |\langle \eta, s(\beta)^\vee \rangle| - 1$ and $\kappa_{s(\beta)} = 1 + \kappa_\beta$;
- if $(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}$ then $\kappa_{\beta_1} + \kappa_{\beta_2} = 1$;
- $\kappa_{-\alpha} = 0$.

Lemma 3.1.3. *Let $\beta = \alpha_{lm} \in \Phi^-$. Then if $l \leq j_0$ or $m > i_0$ then κ_β is subadditive, i.e., $\kappa_\beta \leq \kappa_{\beta_1} + \kappa_{\beta_2}$ for all $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$.*

Moreover, if β has a decomposition $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ with $\kappa_\beta > \kappa_{\beta_1} + \kappa_{\beta_2}$ then

- $l > j_0$ and $m \leq i_0$;
- $\kappa_\beta = 1$ and $\kappa_{\beta_1} = \kappa_{\beta_2} = 0$;
- β_1 shares the row of $-\alpha$ such that β_2 is bad and either $s(\beta_1)$ is bad or $\beta_1 = -\alpha$.

Proof. One can readily observe that

- if $\beta = \alpha_{lm}$ with $l \notin \{i_0, j_0\}$ then $\kappa_\beta = \delta_{w(\beta) < 0}$;
- if $\beta = \alpha_{i_0 m}$ then $\kappa_\beta = \delta_{w(\beta) < 0}$;
- if $\beta = \alpha_{j_0 m}$ then $\kappa_\beta = \delta_{w'(\beta) < 0}$.

The first part follows from combining these three observations together with the fact $\delta_{w(\beta) < 0} \leq \delta_{w(\beta_1) < 0} + \delta_{w(\beta_2) < 0}$ if $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$, case by case.

For the second part, if β has a decomposition $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ with $\kappa_\beta > \kappa_{\beta_1} + \kappa_{\beta_2}$ then it is easy to see that β_1 shares the row of $-\alpha$, and so if $\beta_1 \neq -\alpha$ then we have $w'(\beta_1) > 0$ and $w(\beta_1) < 0$ and so $s(\beta_1)$ is bad. β_2 is also bad, as $w'(\beta_2) < 0$. $\square$

Lemma 3.1.4. *Let $\beta \in \Phi^-$ be a bad root sharing the row of α . Then the map $(\beta_1, \beta_2) \mapsto (\beta_1, s(\beta_2))$ gives rise to a bijection between the following sets:*

- the set $\mathfrak{A}_\beta$;
- the set of elements (β_1, β_2) of $\mathfrak{A}_{s(\beta)}$ with $\beta_2 \neq -\alpha$.

Moreover, $\beta_2 \in \Phi^-$ is bad if $(\beta_1, \beta_2) \in \mathfrak{A}_\beta$.

Proof. For $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ it is easy to see that $\kappa_\beta = \kappa_{\beta_1} + \kappa_{\beta_2}$, if and only if $\kappa_{\beta_1} = 0 = \kappa_{\beta_2}$, if and only if $w(\beta_1) > 0$, $w(\beta_2) > 0$, and $w'(\beta_2) < 0$, if and only if $\kappa_{\beta_1} = 0 = \kappa_{s(\beta_2)}$, if and only if $\kappa_{s(\beta)} = \kappa_{\beta_1} + \kappa_{s(\beta_2)}$. Moreover, if $(\beta_1, \beta_2) \in \mathfrak{A}_\beta$ then $w(\beta_2) > 0$ and $w'(\beta_2) < 0$, and so β_2 is bad. This completes the proof. $\square$

Lemma 3.1.5. *Let $\beta \in \Phi^-$ be a root sharing the column of α . If β is bad then the map $(\beta_1, \beta_2) \mapsto (s(\beta_1), \beta_2)$ gives rise to a bijection between the following sets:*

- the set $\mathfrak{A}_\beta$;
- the set of elements (β_1, β_2) of $\mathfrak{A}_{s(\beta)}$ such that $s(\beta_1)$ is bad.

Moreover, if β is not bad and $(\beta_1, \beta_2) \in \mathfrak{A}_{s(\beta)}$ then $s(\beta_1)$ is not bad.

Proof. Assume β is bad. Then we have $\kappa_\beta = 0$ and $\kappa_{s(\beta)} = 1$. Also, for $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ it is easy to see that $\kappa_{s(\beta)} = \kappa_{\beta_1} + \kappa_{\beta_2}$ and $s(\beta_1)$ is bad, if and only if $\kappa_{\beta_1} = 1$, $\kappa_{s(\beta_1)} = 0$, and $\kappa_{\beta_2} = 0$, if and only if $\kappa_\beta = \kappa_{s(\beta_1)} + \kappa_{\beta_2}$. This completes the proof of the first part.

For the second part, it is clear if $\kappa_\beta = 0$. If $\kappa_\beta = 1$ then $\kappa_{s(\beta)} = 1$ as β is not bad, and so we have either $\kappa_{\beta_2} = 1$ or $\kappa_{\beta_1} = 1$. If $\kappa_{\beta_2} = 1$ then $\kappa_{\beta_1} = 0$ and so $s(\beta_1)$ is not bad. If $\kappa_{\beta_2} = 0$ then $\kappa_{\beta_1} = 1$ and $\kappa_{s(\beta_1)} = 1$, and so $s(\beta_1)$ is not bad. This completes the proof. $\square$

For each $\beta \in \Phi^-$, let c_β be the coefficient of v^{κ_β} in V_β . For each $\beta = \alpha_{lm} \in \Phi^-$ with either $l \leq j_0$ or $i_0 \leq m$, let c_β^i be the coefficient of $v^{\kappa_\beta - 1}$ (resp. v^{κ_β}) in V_β^i if $l > j_0$ and $m = i_0$ (resp. otherwise), where $V^i \stackrel{\text{def}}{=} V^{-1}$.

Lemma 3.1.6. *If β is a bad root sharing the row of α then*

$$c_\beta \cdot p^{|\langle \eta, \alpha^\vee \rangle|} = -c \cdot c_{s(\beta)}.$$

Moreover, we have $-c \cdot c_{-\alpha} = p^{|\langle \eta, \alpha^\vee \rangle|}$.

Proof. This is immediate from the elementary operations in (8) together with the finite height conditions. For instance, we have $C_{i_0, i_0} = v(v + p)^{n-i_0-1} - cC_{-\alpha} = (v + p)^{n-i_0}$ where C is defined in the proof of Proposition 3.1.2. Now, extracting the constant term of the identity we get $-c \cdot c_{-\alpha} = p^{|\langle \eta, \alpha^\vee \rangle|}$. $\square$

Lemma 3.1.7. *Let $\beta \in \Phi^-$ be a bad root.*

(1) *If $\beta \in \Phi^-$ is sharing the row of α then we have*

$$c_{s(\beta)} + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} c_{s(\beta_2)} c_{\beta_1}^i + c_{-\alpha} c_\beta^i + c_{s(\beta)}^i = 0.$$

(2) *If $\beta \in \Phi^-$ is sharing the column of α then we have*

$$c_\beta c_{-\alpha}^i + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} c_{\beta_2} c_{s(\beta_1)}^i + c_{s(\beta)}^i = 0.$$

Proof. (1) follows immediately from Lemma 3.1.4 and Lemma 3.1.3, by extracting the constant term in the equation $(V \cdot V^i)_{s(\beta)} = 0$.

It is immediate that if $\beta' \in \Phi^-$ with $\beta' \neq -\alpha$ shares the column of $-\alpha$ and $s(\beta')$ is not bad, then the coefficient of $v^{\kappa_{\beta'}-1}$ in $V_{\beta'}^i$ vanishes, as $\kappa_{\beta'}$ is subadditive if $s(\beta')$ is not bad. This result together with Lemma 3.1.5 and Lemma 3.1.3 implies (2), by extracting the constant term in the equation $(V \cdot V^i)_{s(\beta)} = 0$. $\square$

3.2. Finite height conditions: the second form. Let $\tilde{w}^* \in \text{Adm}(\eta)$ be a colength one shape of the second form. Write $\tilde{w} = ws_\alpha t_{\eta-\alpha} w^{-1}$ where $w \in W$ satisfies $w(\alpha) < 0$ and $w(j_0) < w(k) < w(i_0)$ for all $i_0 < k < j_0$ if we set $\alpha = \alpha_{i_0 j_0} \in \Phi^+ \setminus \Delta$. For notational convenience we set $w' \stackrel{\text{def}}{=} ws_\alpha$ and $s \stackrel{\text{def}}{=} s_\alpha$, so that we may write $\tilde{w} = w' t_{\eta-\alpha} s w'^{-1}$.

With an eye towards applying the monodromy conditions, we introduce the following:

Definition 3.2.1. We say that a negative root $\beta \in \Phi^-$ is *bad* (with respect to $\tilde{w}$) if

- β shares either the row or the column of α ;
- β satisfies $\delta_{w(\beta) < 0} = \delta_{w'(\beta) < 0}$.

Proposition 3.2.2. *Let $A \in U(\tilde{w}, \leq \eta)(R)$.*

Then $A_{w'(i_0)w'(i_0)} = (w'^{-1}Aw')_{i_0 i_0} = c(v+p)^{n-j_0}$ for some $c \in R$ and

$$(9) \quad u_\alpha(-c) \cdot v^{\varepsilon_{i_0}} \cdot (w'^{-1}Aw') \cdot s = (v+p)^\eta \cdot V',$$

where V' is a lower triangular matrix whose entries are in $R[v]$.

Moreover, $V \stackrel{\text{def}}{=} v^{-\varepsilon_{j_0}} \cdot V'$ is unipotent, and for each $\beta \in \Phi^-$ the entry V'_β can be written as $v^{\kappa'_\beta} \cdot f_\beta(v)$ for a polynomial $f_\beta(v) \in R[v]$ of the form $\sum_k c_{\beta,k}(v+p)^k$ where

- (1) *if β is bad then $\kappa'_\beta = \delta_{w'(\beta) < 0}$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle| + 1$;*
- (2) *if β shares the row or the column of α but is not bad then $\kappa'_\beta = 1 + \delta_{w'(\beta) < 0} = \delta_{w(\beta) < 0} = 1$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle|$;*
- (3) *if β shares the column of $-\alpha$ then $\kappa'_\beta = \delta_{w(\beta) < 0}$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle| - \delta_{w(\beta) > w'(\alpha)} - \delta_{w(\beta) < 0}$;*
- (4) *for all other roots $\beta \in \Phi^-$, $\kappa'_\beta = \delta_{w'(\beta) < 0}$ and $\deg(f_\beta) < |\langle \eta, \beta^\vee \rangle|$.*

Proof. It is easy to see that $(w'^{-1}Aw')_{i_0 i_0}$ is a polynomial of degree less than $n - j_0 + 1$. Let $c \in R$ be the coefficient of v^{n-j_0} in $(w'^{-1}Aw')_{i_0 i_0}$, and set $C \stackrel{\text{def}}{=} u_\alpha(-c)v^{\varepsilon_{i_0}}(w'^{-1}Aw')s$ as in (9). One can readily observe the following degree bound:

- $\deg(C_{ll}) \leq n - l$ if $l \neq j_0$, and $\deg(C_{j_0 j_0}) \leq n - l + 1$ and $C_{j_0 j_0} \in vR[v]$;

- for $\beta = \alpha_{lm} \in \Phi$ with $l \neq i_0$ and $m \notin \{i_0, j_0\}$, $C_\beta = v^{\delta_{w'(\beta)} < 0} f_\beta(v)$ with $\deg(f_\beta) \leq n - m - 1$;
- for $\beta = \alpha_{li_0} \in \Phi$, $C_\beta = v^{\delta_{w(\beta)} < 0} f_\beta(v)$ with $\deg(f_\beta) \leq n - i_0 - 1 - \delta_{w(\beta) > w'(\alpha)} - \delta_{w(\beta) < 0}$;
- for $\beta = \alpha_{lj_0} \in \Phi$ with $\beta \neq \alpha$, $C_\beta = v^{\delta_{w(\beta)} < 0} f_\beta(v)$ with $\deg(f_\beta) \leq n - j_0 + 1 - \delta_{w(\beta) > w'(-\alpha)} - \delta_{w(\beta) < 0}$;
- for $\beta = \alpha_{i_0 m} \in \Phi$, $C_\beta = v^{\delta_{w(\beta)} < 0} f_\beta(v)$ with $\deg(f_\beta) \leq n - m$ if $\beta \neq \alpha$, and $C_\alpha = v f_\beta(v)$ with $\deg(f_\beta) \leq n - j_0$.

Now, the rest of the proof is similar to Proposition 3.1.2, and we leave the details for the reader. $\square$

For the rest of this subsection, we observe some necessary properties of $\kappa'_\beta \in \{0, 1\}$, defined in Proposition 3.2.2, as well as some identities of the coefficients of V' (and so of V as well) from the finite height conditions. We first fix some notation:

- For $\beta \in \Phi^-$, we write $\mathfrak{D}_\beta$ for the set of the decompositions (β_1, β_2) of β into two negative roots with β_1 sharing the column of β (and β_2 sharing the row of β).
- For a bad root $\beta \in \Phi^-$, we write $\mathfrak{A}_\beta$ for the set $\{(\beta_1, \beta_2) \in \mathfrak{D}_\beta \mid \kappa'_\beta = \kappa'_{\beta_1} + \kappa'_{\beta_2}\}$.
- To describe the bottom degrees of each entry of V , we set $\kappa_\beta \stackrel{\text{def}}{=} \kappa'_\beta - 1$ if $\beta \in \Phi^-$ shares the row of $-\alpha$, and $\kappa_\beta \stackrel{\text{def}}{=} \kappa'_\beta$ otherwise.

The following are immediate consequences of Proposition 3.2.2, which will be frequently used:

- if β is bad then we have $\kappa'_\beta = \kappa'_{s(\beta)}$;
- if β is a bad root sharing the column of α then $\deg(v^{-\kappa'_{s(\beta)}} V'_{s(\beta)}) < |\langle \eta, s(\beta)^\vee \rangle| - 1$;
- if $\beta = \alpha_{j_0 m}$ with $i_0 < m \leq j_0$ then $\kappa'_\beta = 1$;
- if $\beta = \alpha_{li_0}$ with $i_0 < l \leq j_0$ then $\kappa'_\beta = 0$, and, in particular, $\kappa_{-\alpha} = -1$.

Lemma 3.2.3. *Let $\beta = \alpha_{lm} \in \Phi^-$. Then κ'_β is subadditive, i.e. $\kappa'_\beta \leq \kappa'_{\beta_1} + \kappa'_{\beta_2}$ for all $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$.*

Moreover, if β has a decomposition $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ with $\kappa_\beta > \kappa_{\beta_1} + \kappa_{\beta_2}$ then

- $l > j_0$ and $m \leq i_0$;
- β_1 shares the row of $-\alpha$ such that β_2 is bad and either $s(\beta_1)$ is bad or $\beta_1 = -\alpha$.

Proof. The proof is very similar to Lemma 3.1.3. Observe that

- if $\beta = \alpha_{lm}$ with $l \neq i_0$ and $m \notin \{i_0, j_0\}$ then $\kappa'_\beta = \delta_{w'(\beta) < 0}$;
- if $\beta = \alpha_{li_0}$ with $m \in \{i_0, j_0\}$ then $\kappa'_\beta = \delta_{w(\beta) < 0}$;
- if $\beta = \alpha_{i_0 m}$ then $\kappa'_\beta = \delta_{w(\beta) < 0}$.

The subadditivity of κ'_β follows from combining the three observations above together with the fact $\delta_{w'(\beta) < 0} \leq \delta_{w'(\beta_1) < 0} + \delta_{w'(\beta_2) < 0}$ if $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$. The subadditivity of κ_β when $l \leq j_0$ or $i_0 < m$ also easily follows from the subadditivity of κ'_β together with the colength one property of w .

Assume that β has a decomposition $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ with $\kappa_\beta > \kappa_{\beta_1} + \kappa_{\beta_2}$. Then as κ'_β is subadditive and κ_β is subadditive if $m > i_0$, we see that β_1 shares the row

of $-\alpha$ and so β_2 shares the column of α . It is also easy to see that β_2 is bad, and $s(\beta_1)$ is bad if $\beta_1 \neq -\alpha$. $\square$

Lemma 3.2.4. *Let $\beta \in \Phi^-$ be a bad root sharing the row of α . Then the map $(\beta_1, \beta_2) \mapsto (\beta_1, s(\beta_2))$ gives rise to a bijection between the following sets:*

- the set $\mathfrak{A}_\beta$;
- the set of elements (β_1, β_2) of $\mathfrak{A}_{s(\beta)}$ with $\beta_2 \neq -\alpha$.

Moreover, $\beta_2 \in \Phi^-$ is bad if $(\beta_1, \beta_2) \in \mathfrak{A}_\beta$.

Proof. The argument is very similar to Lemma 3.1.4. We leave the details for the reader. $\square$

Lemma 3.2.5. *Let $\beta \in \Phi^-$ be a root sharing the column of α . If β is bad then the map $(\beta_1, \beta_2) \mapsto (s(\beta_1), \beta_2)$ gives rise to a bijection between the following sets:*

- the set $\mathfrak{A}_\beta$;
- the set of elements (β_1, β_2) of $\mathfrak{A}_{s(\beta)}$ such that $s(\beta_1)$ is bad.

Moreover, if β is not bad and $(\beta_1, \beta_2) \in \mathfrak{A}_{s(\beta)}$ then $s(\beta_1)$ is not bad.

Proof. The argument is very similar to Lemma 3.1.5. We leave the details for the reader. $\square$

For each $\beta \in \Phi^-$, let c_β be the coefficient of $v^{\kappa'_\beta}$ in V'_β . For each $\beta = \alpha_{lm} \in \Phi^-$ with either $l \leq j_0$ or $i_0 \leq m$, let c_β^i be the coefficient of $v^{\kappa'_\beta - 1}$ (resp. $v^{\kappa'_\beta}$) in V_β^i if either $l = j_0$ or $l > j_0$ and $m = i_0$ (resp. otherwise), where $V^i \stackrel{\text{def}}{=} V^{-1}$.

Lemma 3.2.6. *If β is a bad root sharing the row of α then*

$$c_\beta \cdot p^{|\langle \eta, \alpha^\vee \rangle|} = -c \cdot c_{s(\beta)}.$$

Moreover, we have $-c \cdot c_{-\alpha} = p^{|\langle \eta, \alpha^\vee \rangle|}$.

Proof. The same argument as in Lemma 3.1.6 also works, using the elementary operations in (9) together with the finite height conditions. $\square$

Lemma 3.2.7. *Let $\beta \in \Phi^-$ be a bad root*

- (1) *If $\beta \in \Phi^-$ is sharing the row of α then we have*

$$c_{s(\beta)} + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} c_{s(\beta_2)} c_{\beta_1}^i + c_{-\alpha} c_\beta^i + c_{s(\beta)}^i = 0.$$

- (2) *If $\beta \in \Phi^-$ is sharing the column of α then we have*

$$c_\beta c_{-\alpha}^i + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} c_{\beta_2} c_{s(\beta_1)}^i + c_{s(\beta)}^i = 0.$$

Proof. The proof is similar to that of Lemma 3.1.7. The only difference is that we use Lemma 3.2.4, Lemma 3.2.3, and Lemma 3.2.5 instead of Lemma 3.1.4, Lemma 3.1.3, and Lemma 3.1.5, respectively. $\square$

3.3. Description of $U(\tilde{w}, \leq \eta)$ in colength one. Let V be the matrix introduced at Proposition 3.1.2 or at Proposition 3.2.2. For each $\beta \in \Phi^-$, we let $m_\beta \stackrel{\text{def}}{=} |\langle \eta, \beta^\vee \rangle| - 1$, and recall that the entry V_β is of the form

$$v^{\kappa_\beta} \sum_{k=0}^{m'_\beta} c_{\beta,k} (v+p)^k,$$

where

$$(10) \quad m'_\beta = \begin{cases} m_\beta + 1 & \text{if } \beta \text{ is bad;} \\ m_\beta - 1 & \text{if } \beta \text{ shares the column of } -\alpha \text{ such that } s(\beta) \in \Phi^- \text{ is bad;} \\ m_\beta & \text{otherwise.} \end{cases}$$

Proposition 3.3.1. *Let $\tilde{w}^* \in \text{Adm}(\eta)$ be a colength one shape. Then there is a closed immersion*

$$U(\tilde{w}, \leq \eta) \hookrightarrow \text{Spec} \mathcal{O}[\{c_{\beta,k} \mid \beta \in \Phi^- \text{ and } 0 \leq k \leq m'_\beta\} \cup \{c\}].$$

Proof. This is immediate from Propositions 3.1.2 and 3.2.2. $\square$

4. MONODROMY CONDITIONS

In this section, we induce certain identities and properties, that will be necessary to describe $U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}})$ for $\mathbf{a} \in \mathcal{O}^n$ when $\tilde{w}^* \in \text{Adm}(\eta)$ is of colength one. Throughout this section, by R we mean a Noetherian $\mathcal{O}$ -flat $\mathcal{O}$ -algebra. We keep the notation of §3.

We fix some notation. Set $\gamma_\beta \stackrel{\text{def}}{=} \kappa_\beta + m'_\beta$, where m'_β is defined in (10). For each $\beta \in \Phi^-$, we write $\mathbf{F}_{\geq \beta}$ (resp. $\mathbf{F}_{> \beta}$) for the free $\mathbb{Z}$ -module generated by the monomials $c_{\beta_1, k_1} \cdots c_{\beta_s, k_s}$ for all non-negative integers $k_1, \dots, k_s$ with $\sum_{i=1}^s k_i \leq \gamma_\beta + 1$ and for all negative roots $\beta_1, \dots, \beta_s$ with $\beta = \beta_1 + \cdots + \beta_s$ (resp. with $\beta = \beta_1 + \cdots + \beta_s$ and $s > 1$). For $\beta' \in \Phi^-$ with $\beta' > \beta$ we write $\mathbf{F}_{> \beta}^{\beta'}$ for the submodule of $\mathbf{F}_{> \beta}$ generated by the monomials $c_{\beta_1, k_1} \cdots c_{\beta_s, k_s}$ with $\beta_i \neq \beta'$ for all i . By abuse of notation, we will use the same notation for the images of all of these free $\mathbb{Z}$ -modules in the ring $\mathcal{O}(U(\tilde{w}, \leq \eta))$ via the closed immersion of Proposition 3.3.1. Finally, if $A \in U(\tilde{w}, \leq \eta)(R)$, we abuse notation again and write $\mathbf{F}_{\geq \beta}, \mathbf{F}_{> \beta}, \mathbf{F}_{> \beta}^{\beta'}$ for the image in $R[v]$ (via the map $\mathcal{O}(U(\tilde{w}, \leq \eta)) \rightarrow R$ corresponding to A) of the $\mathbb{Z}$ -modules above.

Let $A \in U(\tilde{w}, \leq \eta)(R)$, and let V be the corresponding matrix obtained by Proposition 3.1.2 (resp. by Proposition 3.2.2) if $\tilde{w}$ is of the first form (resp. if $\tilde{w}$ is of the second form). Then an elementary computation shows that $A \in U^{\text{nv}}(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}})(R)$ if and only if for each $\beta \in \Phi^-$

$$(11) \quad V_\beta^\# \stackrel{\text{def}}{=} (\nabla_{\mathbf{a}_w} V)_\beta + \sum_{(\beta_1, \beta_2) \in \mathfrak{D}_\beta} (\nabla_{\mathbf{a}_w} V)_{\beta_2} V_{\beta_1}^\beta \in (v+p)^{m_\beta} R[v],$$

where $\mathbf{a}_w \in \mathcal{O}^n$ satisfies $w(\mathbf{a}_w) = \mathbf{a}$. Throughout this subsection, we assume (11) holds for all $\beta \in \Phi^-$, and write ∇ for $\nabla_{\mathbf{a}_w}$ to lighten the notation.

4.1. Monodromy conditions: the first form. Let $\tilde{w}^* \in \text{Adm}(\eta)$ be a colength one shape of the first form (described in Proposition 2.1.2), and keep the notation of §3.1. In particular, we keep the notation of Proposition 3.1.2.

Lemma 4.1.1. *Let $\beta = \alpha_{lm} \in \Phi^-$ with $l \leq j_0$ or $i_0 \leq m$. Then we have*

(1) *if β is not bad then*

$$V_\beta^\sharp \in v^{\tilde{\kappa}_\beta}(v+p)^{m_\beta}(X_\beta + \mathbf{F}_{>\beta}),$$

where $X_\beta = (m_\beta + \kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m'_\beta}$ and

$$\tilde{\kappa}_\beta \stackrel{\text{def}}{=} \begin{cases} \kappa_\beta & \text{if either } l \leq j_0, i_0 < m, \text{ or } i_0 = m \text{ and } s(\beta) \in \Phi^- \text{ is not bad;} \\ \kappa_\beta - 1 & \text{if } i_0 = m \text{ and } s(\beta) \in \Phi^- \text{ is bad,} \end{cases}$$

(2) *if β is bad then*

$$V_\beta^\sharp \in v^{\kappa_\beta}(v+p)^{m_\beta}(Y_\beta + vX_\beta + \mathbf{F}_{>\beta}),$$

where

$$\begin{cases} X_\beta &= (m'_\beta + \kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m'_\beta}; \\ Y_\beta &= (m_\beta + \kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle + m_\beta + \kappa_\beta) c_{\beta, m_\beta} + p(\kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m'_\beta}. \end{cases}$$

Proof. Assume that $\beta = \alpha_{lm} \in \Phi^-$ is not bad. If either $l \leq j_0$, $i_0 < m$, or $i_0 = m$ and $s(\beta)$ is not bad then by Proposition 3.1.2 together with Lemma 3.1.3 it is routine to check the equation in (1). If $l > j_0$, $i_0 = m$, and $s(\beta) \in \Phi^-$ is bad then κ_β is not subadditive and $\deg(V_\beta/v^{\kappa_\beta}) < |\langle \eta, \beta^\vee \rangle| - 1$ by part (3) of Proposition 3.1.2, and so $v^{\kappa_\beta} \nmid \sum_{(\beta_1, \beta_2) \in \mathfrak{D}_\beta} (\nabla V)_{\beta_2} V_{\beta_1}^i$ while $v^{\kappa_\beta} \mid (\nabla V)_\beta$. Hence, in this case we get the equation in (1) with $\tilde{\kappa}_\beta = \kappa_\beta - 1$ in the equation.

Assume that β is bad. By part (1) of Proposition 3.1.2, $\deg V_\beta < |\langle \eta, \beta^\vee \rangle| + 1$ (as $\kappa_\beta = 0$) and κ_β is subadditive, and so we get the equation in (2). $\square$

For $\beta \in \Phi^-$ and for each integer $s \geq 1$, we set $I_{\beta, s}$ to be the set of the tuples of negative roots $(\beta_1, \beta_2, \dots, \beta_s)$ such that

- $\beta = \beta_1 + \beta_2 + \dots + \beta_s$ and β_1 shares the column of β ;
- $\beta_i + \beta_{i+1} \in \Phi^-$ for $i \in \{1, \dots, s-1\}$;
- $\sum_{i=1}^s \delta_{w(\beta_i) < 0} = s - \delta_{w(\beta) > 0}$.

Moreover, we set

$$I_\beta \stackrel{\text{def}}{=} \bigcup_{s \geq 1} I_{\beta, s}.$$

Lemma 4.1.2. *Let $\beta \in \Phi^-$ with $\beta \geq -\alpha$.*

(1) *If $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ then $\gamma_\beta \geq \gamma_{\beta_1} + \gamma_{\beta_2}$;*

(2) *$(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ satisfies $\gamma_\beta = \gamma_{\beta_1} + \gamma_{\beta_2}$ if and only if $\kappa_\beta < \kappa_{\beta_1} + \kappa_{\beta_2}$.*

In particular, if $(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}$ then $\gamma_{\beta_1} + \gamma_{\beta_2} = \gamma_{-\alpha}$, $\deg V_{\beta_1}^i = \gamma_{\beta_1} = m_{\beta_1} + \kappa_{\beta_1}$, and

$$\frac{1}{(\gamma_{\beta_1})!} \frac{d^{\gamma_{\beta_1}} V_{\beta_1}^i}{dv^{\gamma_{\beta_1}}} = \sum_{(\beta'_1, \dots, \beta'_s) \in I_{\beta_1}} (-1)^{\ell+j_0-s} c_{\beta'_1, m_{\beta'_1}} c_{\beta'_2, m_{\beta'_2}} \cdots c_{\beta'_s, m_{\beta'_s}},$$

where $\beta_1 = \alpha_{\ell i_0}$.

Proof. Recall that γ_β indicates the degree of V_β . By definition of w , $\kappa_\beta = \kappa_{s(\beta)}$ if either $\beta = \alpha_{li_0}$ with $i_0 < l < j_0$ or $\beta = \alpha_{i_0m}$ with $i_0 < m < j_0$. Now it is easy to check (1) and (2) case by case.

For the second part, let $(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}$. Then it is clear that $\gamma_{\beta_1} + \gamma_{\beta_2} = \gamma_{-\alpha}$ by (2), and it is also clear that $\deg V_{\beta_1}^i = \gamma_{\beta_1}$ by (1). Finally, the elements of I_β correspond to the monomials in $V_{\beta_1}^i$ which has the highest degree γ_{β_1} . This completes the proof. $\square$

Lemma 4.1.3. *We have*

$$p^{m_{-\alpha}+1} \cdot \langle \mathbf{a}_w, -\alpha^\vee \rangle = p^{m_{-\alpha}} \cdot c \cdot Z_{-\alpha},$$

where

$$\begin{aligned} Z_{-\alpha} &\stackrel{\text{def}}{=} (m_{-\alpha} - \langle \mathbf{a}_w, -\alpha^\vee \rangle) c_{-\alpha, m_{-\alpha}} \\ &+ \sum_{(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}} (m_{\beta_2} + \kappa_{\beta_2} - \langle \mathbf{a}_w, \beta_2^\vee \rangle) c_{\beta_2, m_{\beta_2}} \\ &\cdot \sum_{(\beta'_1, \dots, \beta'_s) \in I_{\beta_1}} (-1)^{\ell+j_0-s} c_{\beta'_1, m_{\beta'_1}} \cdots c_{\beta'_s, m_{\beta'_s}} \end{aligned}$$

if we write $\beta_1 = \alpha_{\ell i_0}$.

Proof. By Lemma 4.1.2, we see that $\deg((\nabla V)_{\beta_2} V_{\beta_1}^i) = \gamma_{\beta_2} + \gamma_{\beta_1} = \gamma_{-\alpha} = m_{-\alpha}$ for all $(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}$. Hence, if we apply the monodromy condition (11) to $V_{-\alpha}$ then we have

$$\begin{aligned} V_{-\alpha}^\# &= (m_{-\alpha} - \langle \mathbf{a}_w, -\alpha^\vee \rangle) c_{-\alpha, m_{-\alpha}} (v+p)^{m_{-\alpha}} \\ &+ \sum_{(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}} (m_{\beta_2} + \kappa_{\beta_2} - \langle \mathbf{a}_w, \beta_2^\vee \rangle) c_{\beta_2, m_{\beta_2}} \\ &\cdot \sum_{(\beta'_1, \dots, \beta'_s) \in I_{\beta_1}} (-1)^{\ell+j_0-s} c_{\beta'_1, m_{\beta'_1}} \cdots c_{\beta'_s, m_{\beta'_s}} (v+p)^{m_{-\alpha}} \end{aligned}$$

by applying the second part of Lemma 4.1.2. Since the constant term of $V_{-\alpha}^\#$ appears only at $(\nabla V)_{-\alpha}$ by Lemma 3.1.3, by extracting the constant term we have

$$-\langle \mathbf{a}_w, -\alpha^\vee \rangle c_{-\alpha} = p^{m_{-\alpha}} Z_{-\alpha}.$$

By the second part of Lemma 3.1.6, we get the desired identity. $\square$

We further eliminate the variables Y_β for bad roots $\beta \in \Phi^-$.

Lemma 4.1.4. *Let $\beta \in \Phi^-$ be a bad root.*

(1) *If β shares the row of α then we have*

$$p^{m_{s(\beta)}} \cdot (Y_\beta + c \cdot X_{s(\beta)}) \in p^{m_{s(\beta)}} \cdot \left(c \cdot c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{> \beta} + c \cdot \mathbf{F}_{> s(\beta)}^\beta \right).$$

(2) *If β shares the column of α then we have*

$$p^{m_{s(\beta)}} \cdot (Y_\beta - c \cdot X_{s(\beta)}) \in p^{m_{s(\beta)}} \cdot \left(c \cdot c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{> \beta} + c \cdot \mathbf{F}_{> s(\beta)}^\beta \right).$$

Proof. We first treat the case (1). By Lemma 3.1.4, extracting the constant term in the monodromy equation in part (1) of Lemma 4.1.1 for $s(\beta)$ gives rise to

$$(12) \quad \langle \mathbf{a}_w, s(\beta)^\vee \rangle c_{s(\beta)} + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} \langle \mathbf{a}_w, s(\beta_2)^\vee \rangle c_{s(\beta_2)} c_{\beta_1}^\flat + \langle \mathbf{a}_w, -\alpha^\vee \rangle c_{-\alpha} c_\beta^\flat \in -p^{m_{s(\beta)}} (X_{s(\beta)} + \mathbf{F}_{>s(\beta)}).$$

As $\deg(V_\beta V_{-\alpha}^\flat) = m'_\beta + m_{-\alpha} = m_\beta + 1 + m_{-\alpha} = m_{s(\beta)} = \tilde{\kappa}_{s(\beta)} + m_{s(\beta)}$, the quantity in (12), in fact, belongs to

$$-p^{m_{s(\beta)}} (X_{s(\beta)} + c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{>s(\beta)}^\beta).$$

By multiplying $-c$ and then applying Lemma 3.1.6 together with Lemma 3.1.4, we have

$$\begin{aligned} & p^{|\langle \eta, \alpha^\vee \rangle|} \langle \mathbf{a}_w, s(\beta)^\vee \rangle c_\beta \\ & + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} p^{|\langle \eta, \alpha^\vee \rangle|} \langle \mathbf{a}_w, s(\beta_2)^\vee \rangle c_{\beta_2} c_{\beta_1}^\flat + p^{|\langle \eta, \alpha^\vee \rangle|} \langle \mathbf{a}_w, -\alpha^\vee \rangle c_\beta^\flat \\ & \in c \cdot p^{m_{s(\beta)}} (X_{s(\beta)} + c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{>s(\beta)}^\beta). \end{aligned}$$

Applying the identity $c_\beta + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} c_{\beta_2} c_{\beta_1}^\flat + c_\beta^\flat = 0$, induced from $(V \cdot V^\flat)_\beta = 0$, together with $s(\beta) = \beta - \alpha$ and $s(\beta_2) = \beta_2 - \alpha$, we have

$$(13) \quad p^{|\langle \eta, \alpha^\vee \rangle|} \langle \mathbf{a}_w, \beta^\vee \rangle c_\beta + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} p^{|\langle \eta, \alpha^\vee \rangle|} \langle \mathbf{a}_w, \beta_2^\vee \rangle c_{\beta_2} c_{\beta_1}^\flat \in c \cdot p^{m_{s(\beta)}} (X_{s(\beta)} + c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{>s(\beta)}^\beta).$$

But by extracting the constant term in the monodromy equation in part (2) of Lemma 4.1.1 for β , the quantity in (13) also belongs to

$$-p^{|\langle \eta, -\alpha^\vee \rangle| + |\langle \eta, \beta^\vee \rangle| - 1} (Y_\beta + \mathbf{F}_{>\beta})$$

and so we have the desired result.

We now treat the case (2). By extracting the coefficient of v^{κ_β} in the equation $(V \cdot V^\flat)_\beta = 0$, we have $c_\beta + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} c_{\beta_2} c_{\beta_1}^\flat + c_\beta^\flat = 0$, which together with the equation in part (2) of Lemma 3.2.7 induces

$$\sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} c_{\beta_2} (c_{s(\beta_1)}^\flat - c_{-\alpha}^\flat c_{\beta_1}^\flat) + (c_{s(\beta)}^\flat - c_{-\alpha}^\flat c_\beta^\flat) = 0.$$

This equation together with Lemma 3.2.5 inductively induces that

$$(14) \quad c_{s(\beta)}^\flat - c_{-\alpha}^\flat c_\beta^\flat = 0$$

for any bad root β sharing the column of α .

By extracting the coefficient of v^{κ_β} in the monodromy equation in part (2) of Lemma 4.1.1 for β , we have

$$(\kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_\beta + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} (\kappa_{\beta_2} - \langle \mathbf{a}_w, \beta_2^\vee \rangle) c_{\beta_2} c_{\beta_1}^\flat \in p^{m_\beta} (Y_\beta + \mathbf{F}_{>\beta}).$$

Similarly, by extracting the coefficient of $v^{\kappa_{s(\beta)}-1}$ in the monodromy equation in part (1) of Lemma 4.1.1 for $s(\beta)$, we have

$$(15) \quad (\kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_\beta c_{-\alpha}^\iota + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_\beta} (\kappa_{\beta_2} - \langle \mathbf{a}_w, \beta_2^\vee \rangle) c_{\beta_2} c_{s(\beta_1)}^\iota \in p^{m_{s(\beta)}} (X_{s(\beta)} + \mathbf{F}_{>s(\beta)}).$$

As $\deg(V_\beta V_{-\alpha}^\iota) = \gamma_\beta + \gamma_{-\alpha} = m'_\beta + m_{-\alpha} = m_\beta + 1 + m_{-\alpha} = m_{s(\beta)} = \tilde{\kappa}_{s(\beta)} + m_{s(\beta)}$, the quantity in (15), in fact, belongs to

$$p^{m_{s(\beta)}} (X_{s(\beta)} + c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{>s(\beta)}^\beta).$$

Comparing these two equations via the identity (14), we have

$$p^{m_{s(\beta)}} X_{s(\beta)} - c_{-\alpha}^\iota p^{m_\beta} Y_\beta \in p^{m_{s(\beta)}} (c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{>s(\beta)}^\beta) + p^{m_\beta} \cdot c_{-\alpha}^\iota \cdot \mathbf{F}_{>\beta}.$$

As $c_{-\alpha} = -c_{-\alpha}^\iota$, by applying the second part of Lemma 3.1.6 we have

$$-cp^{m_{s(\beta)}} X_{s(\beta)} + p^{m_\beta + m_{-\alpha}} Y_\beta \in c \cdot p^{m_{s(\beta)}} (c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{>s(\beta)}^\beta) + p^{m_\beta + m_{-\alpha} + 1} \cdot \mathbf{F}_{>\beta}.$$

As $s(\beta) = \beta - \alpha$, we get the desired result. $\square$

We finally treat the case $\beta = \alpha_{lm} \in \Phi^-$ with $l > j_0$ and $m < i_0$. In this case, κ_β is not subadditive in general, so that Lemma 4.1.5 is not trivial.

Lemma 4.1.5. *For $\beta = \alpha_{lm} \in \Phi^-$ with $l > j_0$ and $m < i_0$,*

$$V_\beta^\sharp \in v^{\kappa_\beta} (v + p)^{|\langle \eta, \beta^\vee \rangle| - 1} (X_\beta + \mathbf{F}_{>\beta}),$$

where $X_\beta = (m_\beta + \kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m'_\beta}$.

Proof. We first claim that $V_{s(\beta_0)}^\iota \in vR[v]$ for a bad root $\beta_0 \in \Phi^-$ sharing the row of α . From part (1) of Lemma 3.1.7 together with Lemma 3.1.4 and Lemma 3.1.6, we have

$$p^{|\langle \eta, \alpha^\vee \rangle|} \left(c_{\beta_0} + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_{\beta_0}} c_{\beta_2} c_{\beta_1}^\iota + c_{\beta_0}^\iota \right) - cc_{s(\beta_0)}^\iota = 0.$$

As $c_{\beta_0} + \sum_{(\beta_1, \beta_2) \in \mathfrak{A}_{\beta_0}} c_{\beta_2} c_{\beta_1}^\iota + c_{\beta_0}^\iota = 0$ induced from extracting the constant term of $(V \cdot V^\iota)_{\beta_0} = 0$, we have $cc_{s(\beta_0)}^\iota = 0$. Hence, we conclude that $c_{s(\beta_0)}^\iota = 0$, as c is a unit in $R[\frac{1}{p}]$ by Lemma 3.1.6 and R is $\mathcal{O}$ -flat.

Let $\beta' = \alpha_{l'm'} \in \Phi^-$ with $m' < i_0$. We claim that $V_{\beta'}^\iota \in v^{\kappa_{\beta'}} R[v]$ if $l' \neq j_0$. It is clear that $V_{\beta'}^\iota \in v^{\kappa_{\beta'}} R[v]$ if $l' < j_0$, by Lemma 3.1.3. Assume $j_0 < l'$. Consider the following identity

$$0 = (V \cdot V^\iota)_{\beta'} = V_{\beta'} + \sum_{(\beta_1, \beta_2) \in \mathfrak{D}_{\beta'}} V_{\beta_2} \cdot V_{\beta_1}^\iota + V_{\beta'}^\iota,$$

and write $\beta_1 := \alpha_{km'}$. It is obvious that $V_{\beta'} \in v^{\kappa_{\beta'}} R[v]$, and that $V_{\beta_2} \cdot V_{\beta_1}^\iota \in v^{\kappa_{\beta'}} R[v]$ for $m' < k < j_0$ by Lemma 3.1.3, so that it is enough to check that $V_{\beta_2} \cdot V_{\beta_1}^\iota \in v^{\kappa_{\beta'}} R[v]$ for $j_0 \leq k < l'$.

Assume that $k = j_0$. If $s(\beta_1)$ is bad then $V_{\beta_1}^\iota \in vR[v]$ by the first claim, and so we conclude in this case that $V_{\beta_2} \cdot V_{\beta_1}^\iota \in v^{\kappa_{\beta'}} R[v]$. If $s(\beta_1)$ is not bad then it is clear that $V_{\beta_1}^\iota \in v^{\kappa_{\beta_1}} R[v]$, and so we also have $V_{\beta_2} \cdot V_{\beta_1}^\iota \in v^{\kappa_{\beta'}} R[v]$ by Lemma 3.1.3. Note that this also implies

$$(16) \quad (\nabla V)_{\beta_2} V_{\beta_1}^\iota \in v^{\kappa_{\beta'}} R[v]$$

in this case. Assume now that $j_0 < k < l'$. By induction hypothesis, we have $V_{\beta_1}^i \in v^{\kappa_{\beta_1}} R[v]$ and so we conclude that $V_{\beta_2} \cdot V_{\beta_1}^i \in v^{\kappa_{\beta'}} R[v]$ by Lemma 3.1.3, which completes the proof of the second claim.

Now, let $\beta = \alpha_{lm} \in \Phi^-$ with $l > j_0$ and $m < i_0$. For $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$, if we write $\beta_1 = \alpha_{km}$ then it is clear that $(\nabla V)_{\beta_2} V_{\beta_1}^i \in v^{\kappa_\beta} R[v]$ for $k \neq j_0$, by the claim above together with Lemma 3.1.3. If $k = j_0$, then we also have $(\nabla V)_{\beta_2} V_{\beta_1}^i \in v^{\kappa_\beta} R[v]$ by (16), which completes the proof. $\square$

4.2. Monodromy conditions: the second form. Let $\tilde{w}^* \in \text{Adm}(\eta)$ be a colength one shape of the second form (cf. Proposition 2.1.2), and keep the notation of §3.2. In particular, we keep the notation of Proposition 3.2.2.

Lemma 4.2.1. *Let $\beta = \alpha_{lm} \in \Phi^-$ with $l \leq j_0$ or $i_0 \leq m$. Then we have*

(1) *if β is not bad then*

$$V_\beta^\sharp \in v^{\tilde{\kappa}_\beta} (v + p)^{m_\beta} (X_\beta + \mathbf{F}_{>\beta}),$$

where $X_\beta = (m_\beta + \kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m'_\beta}$ and

$$\tilde{\kappa}_\beta \stackrel{\text{def}}{=} \begin{cases} \kappa_\beta & \text{if either } l < j_0, i_0 < m, \text{ or } i_0 = m \text{ and } s(\beta) \in \Phi^- \text{ is not bad;} \\ \kappa'_\beta - 1 & \text{if either } i_0 = m \text{ and } s(\beta) \in \Phi^- \text{ is bad or } l = j_0, \end{cases}$$

(2) *if β is bad then*

$$V_\beta^\sharp \in v^{\kappa_\beta} (v + p)^{m_\beta} (Y_\beta + vX_\beta + \mathbf{F}_{>\beta}),$$

where

$$\begin{cases} X_\beta &= (m'_\beta + \kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m'_\beta}; \\ Y_\beta &= (m_\beta + \kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m_\beta} + p(\kappa_\beta - \langle \mathbf{a}_w, \beta^\vee \rangle) c_{\beta, m'_\beta}. \end{cases}$$

Proof. The proof is similar to that of Lemma 4.1.1 using Proposition 3.2.2 and Lemma 3.2.3 instead of Proposition 3.1.2 and Lemma 3.1.3, respectively. We leave the details for the reader. $\square$

For $\beta \in \Phi^-$ and for each integer $s \geq 1$, we set $I_{\beta,s}$ to be the set of the tuples of negative roots $(\beta_1, \beta_2, \dots, \beta_s)$ such that

- $\beta = \beta_1 + \beta_2 + \dots + \beta_s$ and β_1 shares the column of β ;
- $\beta_i + \beta_{i+1} \in \Phi^-$ for $i \in \{1, \dots, s-1\}$;
- $w'(\beta_i) < 0$ for all $i \in \{1, \dots, s\}$.

Moreover, we set

$$I_\beta \stackrel{\text{def}}{=} \bigcup_{s \geq 1} I_{\beta,s}.$$

Lemma 4.2.2. *Let $\beta \in \Phi^-$ with $\beta \geq -\alpha$.*

(1) *If $(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ then $\gamma_\beta \geq \gamma_{\beta_1} + \gamma_{\beta_2}$;*

(2) *$(\beta_1, \beta_2) \in \mathfrak{D}_\beta$ satisfies $\gamma_\beta = \gamma_{\beta_1} + \gamma_{\beta_2}$ if and only if $\kappa_\beta < \kappa_{\beta_1} + \kappa_{\beta_2}$.*

In particular, if $(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}$ then $\gamma_{\beta_1} + \gamma_{\beta_2} = \gamma_{-\alpha}$, $\deg V_{\beta_1}^i = \gamma_{\beta_1} = m_{\beta_1}$, and

$$\frac{1}{(\gamma_{\beta_1})!} \frac{d^{\gamma_{\beta_1}} V_{\beta_1}^i}{dv^{\gamma_{\beta_1}}} = \sum_{(\beta'_1, \dots, \beta'_s) \in I_{\beta_1}} (-1)^{\ell + j_0 - s} c_{\beta'_1, m_{\beta'_1}} c_{\beta'_2, m_{\beta'_2}} \cdots c_{\beta'_s, m_{\beta'_s}},$$

where $\beta_1 = \alpha_{\ell i_0}$.

Proof. The proof is similar to that of Lemma 4.1.2. We leave the details for the reader. $\square$

Lemma 4.2.3. *We have*

$$p^{m_{-\alpha}+1} \cdot (\langle \mathbf{a}_w, -\alpha^\vee \rangle + 1) = p^{m_{-\alpha}} \cdot c \cdot Z_{-\alpha},$$

where

$$\begin{aligned} Z_{-\alpha} &\stackrel{\text{def}}{=} (m_{-\alpha} - 1 - \langle \mathbf{a}_w, -\alpha^\vee \rangle) c_{-\alpha, m_{-\alpha}} \\ &+ \sum_{(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}} (m_{\beta_2} - \langle \mathbf{a}_w, \beta_2^\vee \rangle) c_{\beta_2, m_{\beta_2}} \sum_{(\beta'_1, \dots, \beta'_s) \in I_{\beta_1}} (-1)^{\ell+j_0-s} c_{\beta'_1, m_{\beta'_1}} \cdots c_{\beta'_s, m_{\beta'_s}} \end{aligned}$$

if we write $\beta_1 = \alpha_{\ell i_0}$.

Proof. By Lemma 4.2.2, we see that $\deg((\nabla V)_{\beta_2} V_{\beta_1}^\vee) = \gamma_{\beta_2} + \gamma_{\beta_1} = m_{\beta_2} + m_{\beta_1} = m_{-\alpha} - 1$ for all $(\beta_1, \beta_2) \in \mathfrak{D}_{-\alpha}$. By the same argument as in Lemma 4.1.3, extracting the constant term of $v \cdot V_{-\alpha}^\#$ gives rise to the result. We leave the details for the reader. $\square$

We further eliminate the variables Y_β for bad roots $\beta \in \Phi^-$.

Lemma 4.2.4. *Let $\beta \in \Phi^-$ be a bad root.*

(1) *If β shares the row of α then we have*

$$p^{m_{s(\beta)}} \cdot (Y_\beta + c \cdot X_{s(\beta)}) \in p^{m_{s(\beta)}} \cdot \left(c \cdot c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{> \beta} + c \cdot \mathbf{F}_{> s(\beta)}^\beta \right).$$

(2) *If β shares the column of α then*

$$p^{m_{s(\beta)}} \cdot (Y_\beta - c \cdot X_{s(\beta)}) \in p^{m_{s(\beta)}} \cdot \left(c \cdot c_{\beta, m'_\beta} \cdot \mathbf{F}_{\geq -\alpha} + \mathbf{F}_{> \beta} + c \cdot \mathbf{F}_{> s(\beta)}^\beta \right).$$

Proof. The proof is similar to that of Lemma 4.1.4. The only difference is that we use Lemmas 3.2.6, 3.2.4, 4.2.1, 3.2.7, 3.2.5, and 3.2.6 instead of Lemmas 3.1.6, 3.1.4, 4.1.1, 3.1.7, 3.1.5, and 3.1.6, respectively. We leave the details for the reader. $\square$

Finally, we treat the case $\beta = \alpha_{lm} \in \Phi^-$ with $l > j_0$ and $m < i_0$. In this case, κ_β is not subadditive in general, so that Lemma 4.2.5 is not trivial.

Lemma 4.2.5. *Assume that $V_{s(\beta)}^\vee \in v^{\kappa_\beta} R[v]$ for a bad root $\beta \in \Phi^-$ sharing the row of α . For $\beta = \alpha_{lm} \in \Phi^-$ with $l > j_0$ and $m < i_0$, we have*

$$V_\beta^\# \in v^{\kappa_\beta} (v + p)^{m_\beta} (X_\beta + \mathbf{F}_{> \beta}),$$

where $X_\beta = (\langle \mathbf{a}_w, \beta^\vee \rangle + m_\beta + \kappa_\beta) c_{\beta, m'_\beta}$.

Proof. We first claim that $V_{s(\beta_0)}^\vee \in v^{\kappa_{\beta_0}} R[v]$ for each bad root $\beta_0 \in \Phi^-$ sharing the row of α , whose proof is almost identical to the first claim in the proof of Lemma 4.1.5. The rest of the proof also is similar to that of Lemma 4.1.5. The only difference is that we use Lemma 3.2.3 instead of Lemma 3.1.3. We leave the details for the reader. $\square$

4.3. Description of $U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}})$ in colength ≤ 1 . In this subsection we describe $U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}})$ when $\tilde{w}^* \in \widetilde{W}$ has colength ≤ 1 .

We first give an upper bound of $U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}})$ for $\tilde{w}^*$ of colength zero.

Proposition 4.3.1. *Let $\tilde{w}^* \in \text{Adm}(\eta)$ be a colength zero shape, and let $\mathbf{a} \in \mathcal{O}^n$. Assume that $\bar{\mathbf{a}}$ is n -generic ([LLHLMb, Definition 4.2.2]). Then there is a closed immersion*

$$U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}}) \hookrightarrow \text{Spec} \mathcal{O}[\{X_{\beta} \mid \beta \in \Phi^{-}\}].$$

Proof. This follows from the arguments in [LLHL19, §3.4], specifically the proof of [LLHL19, Proposition 3.4.12]. (Note that the argument in *loc. cit.* is written for complete local Noetherian $\mathcal{O}$ -algebras but is valid in our setting of $\mathcal{O}$ -flat Noetherian $\mathcal{O}$ -algebras.) $\square$

We now give an upper bound of $U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}})$ for $\tilde{w}^*$ of colength one. Recall that $Z_{-\alpha}$ is constructed in Lemma 4.1.3 (resp. in Lemma 4.2.3) if $\tilde{w}^*$ is of colength one of the first form (resp. of the second form).

Proposition 4.3.2. *Let $\tilde{w}^* \in \text{Adm}(\eta)$ be a colength one shape, and let $\mathbf{a} \in \mathcal{O}^n$. Assume that $\bar{\mathbf{a}}$ is n -generic (as defined in [LLHLMb, §4.2]). Then there is a closed immersion*

$$U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}}) \hookrightarrow \text{Spec} \frac{\mathcal{O}[\{X_{\beta} \mid \beta \in \Phi^{-}\} \cup \{c\}]}{(c \cdot Z_{-\alpha} - p)}.$$

Proof. By Proposition 3.3.1, $\mathcal{O}(U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}}))$ is generated by $c_{\beta,k}$ (with $\beta \in \Phi^{-}$ and $0 \leq k \leq m'_{\beta}$) and c . But Lemmas 4.1.1, 4.1.5, 4.2.1 and 4.2.5 show that for $k < m'_{\beta}$ and β not bad (resp. $k < m'_{\beta} - 1$ and β bad) the element $c_{\beta,k}$ is a $\mathbb{Z}$ -linear combination of X_{β} (resp. Y_{β}) and elements of $\mathbb{F}_{>\beta}$. We conclude, by induction on the poset Φ^{-} and change of variables, that we can also generate $\mathcal{O}(U(\tilde{w}, \leq \eta, \nabla_{\mathbf{a}}))$ using c , X_{β} ($\beta \in \Phi^{-}$) and Y_{β} (β bad). In turn, Lemmas 4.1.4 and 4.2.4 show that we can generate using just c and X_{β} . Finally Lemmas 4.1.3 and 4.2.3 (and p -flatness) give the relation $c \cdot Z_{-\alpha} - p$. $\square$

5. COLENGTH ONE DEFORMATION RINGS

In this section we apply the results of §3 and §4 to compute potentially crystalline deformation rings with Hodge–tate weights η , for sufficiently generic $\bar{\rho}$ and tame inertial types τ such that $\tilde{w}(\bar{\rho}, \tau)$ has colength at most one in each embedding.

5.1. Product structures and error terms. We first extend the technical results of §4.1 and §4.2 in a way which can be used to describe the closed immersion $\tilde{U}(\tilde{z}, \leq \eta, \nabla_{\tau, \infty}) \hookrightarrow \tilde{U}(\tilde{z}, \leq \eta)^{\wedge p}$, when $\tilde{z} = (\tilde{z}^{(j)})_{j \in \mathcal{J}}$ has colength at most one. This requires the modification of some of the previous formulas by allowing product structures, non-trivial diagonal entries, and an “error term” which takes into account the monodromy condition defined in §2.2.3.

Keep the notation of §3, and let R be a p -adically complete, topologically finite type, Noetherian $\mathcal{O}$ -algebra. Let $\tilde{w}^* \in \text{Adm}(\eta)$ be a colength one shape of the first form (resp. of the second form), and $\tilde{A} \in \tilde{U}(\tilde{w}, \leq \eta)(R)$ with its image $A \in U(\tilde{w}, \leq \eta)(R)$ under the natural morphism $\tilde{U}(\tilde{w}, \leq \eta) \rightarrow U(\tilde{w}, \leq \eta)$. We may write $w^{-1}\tilde{A}w = sD_0s \cdot w^{-1}Aw$ (resp. $w'^{-1}\tilde{A}w' = D_0 \cdot w'^{-1}Aw'$) for some $D_0 = \text{Diag}(a_1, \dots, a_n) \in T^{\vee}(R)$. Let $V \in \frac{1}{v}\text{Mat}_n(R[v])$ be the matrix obtained from Proposition 3.1.2 (resp. from Proposition 3.2.2) applied to $A \in U(\tilde{w}, \leq \eta)(R)$, and set $\tilde{V} \stackrel{\text{def}}{=} D_0 \cdot V$. Then

by the same argument as in Proposition 3.1.2 (resp. in Proposition 3.2.2) we may write $(w^{-1}\tilde{A}w)_{j_0j_0} = a_{j_0j_0}(v+p)^{n-j_0}$ (resp. $(w'^{-1}\tilde{A}w')_{i_0i_0} = a_{i_0i_0}(v+p)^{n-j_0}$) for some $a_{j_0j_0} \in R$ (resp. for some $a_{i_0i_0} \in R$), and we have the following identity:

$$(17) \quad s \cdot u_{-\alpha}(-\frac{a_{j_0j_0}}{a_{j_0}}) \cdot (w^{-1}\tilde{A}w) = (v+p)^\eta \cdot \tilde{V} \\ (\text{resp. } u_\alpha(-\frac{a_{i_0i_0}}{a_{j_0}}) \cdot v^{\varepsilon_{i_0}} \cdot (w'^{-1}\tilde{A}w') \cdot s = (v+p)^\eta \cdot v^{\varepsilon_{j_0}} \cdot \tilde{V}).$$

Note that we may identify c , defined in Proposition 3.1.2 (resp. in Proposition 3.2.2), with $a_{j_0j_0}/a_{i_0}$ (resp. with $a_{i_0i_0}/a_{i_0}$). We also note that the degree description of each entry of $\tilde{V}$ is exactly the same as that of V , as $\tilde{V} = D_0 \cdot V$.

We fix a tame inertial type τ with a N -generic lowest alcove presentation (s, μ) , together with an element $\tilde{z} \in \text{Adm}(\eta)^\vee$ which we write as $\tilde{z} = (\tilde{z}^{(j)})_{j \in \mathcal{J}}$.

Until the end of this subsection, assume that $j \in \mathcal{J}$ is such that $\ell(\tilde{z}^{(j)}) = \ell(t_\eta) - 1$, and let $A^{(j)} \in U(\tilde{z}^{(j)}, \leq \eta)(R)$ be the image of $\tilde{A}^{(j)} \in \tilde{U}(\tilde{z}^{(j)}, \leq \eta)(R)$. Let $\tilde{V}^{(j)} \in \frac{1}{v} \text{Mat}_n(R[v])$ be the matrix obtained from (17) (according to the two possible forms of $\tilde{z}^{(j)}$) applied to $\tilde{A}^{(j)} \in \tilde{U}(\tilde{z}^{(j)}, \leq \eta)(R)$. (We adapt the notation of Propositions 3.1.2 and 3.2.2 as well as condition (17) in our context by adding a superscript (j) , so that for instance a colength one shape of the second form has decomposition $w^{(j)}s_{\alpha^{(j)}}t_{\eta-\alpha^{(j)}}w^{(j)^{-1}}$.) It is easy to see that condition (11) shows that condition (7) has the form

$$(18) \quad \tilde{V}_{\beta^{(j)}}^{(j), \sharp} \stackrel{\text{def}}{=} \left((\nabla_{\mathbf{a}_w} \tilde{V}^{(j)}) \cdot \tilde{V}^{(j)} \right)_{\beta^{(j)}} \in (v+p)^{m_{\beta^{(j)}}} R[v] + p^{N-2n+5} R[v]$$

for all $\beta^{(j)} \in \Phi^-$, where $\mathbf{a}_w \in \mathbb{Z}^n$ is defined by $w^{(j)}(\mathbf{a}_w) = \mathbf{a}^{(j)}$.

As $\tilde{V}^{(j)} = D_0 \cdot V^{(j)}$ for some $D_0 \stackrel{\text{def}}{=} \text{Diag}(a_1, \dots, a_n) \in T^\vee(R)$, if we let $\beta^{(j)} = \alpha_{lm} \in \Phi^-$ then it is easy to see that

$$a_m \cdot \tilde{V}_{\beta^{(j)}}^{(j), \sharp} = a_l \cdot V_{\beta^{(j)}}^{(j), \sharp}.$$

Hence, condition (18) applied to $\tilde{V}^{(j)}$ induces all the relevant lemmas from §4.1 and §4.2 keeping track of the “error term” p^{N-2n+5} , as the diagonal entries a_k of D_0 are units in R . More precisely,

(i) in Lemma 4.1.1 (resp. in Lemma 4.2.1), we have

$$(1) \quad V_{\beta^{(j)}}^\sharp \in v^{\tilde{\kappa}_{\beta^{(j)}}} (v+p)^{m_{\beta^{(j)}}} (X_{\beta^{(j)}} + \mathbf{F}_{>\beta^{(j)}}) + p^{N-2n+5} R[v]; \\ (2) \quad V_{\beta^{(j)}}^\sharp \in v^{\kappa_{\beta^{(j)}}} (v+p)^{m_{\beta^{(j)}}} (Y_{\beta^{(j)}} + vX_{\beta^{(j)}} + \mathbf{F}_{>\beta^{(j)}}) + p^{N-2n+5} R[v],$$

(ii) in Lemma 4.1.3 (resp. in Lemma 4.2.3), we have

$$p^{m_{-\alpha^{(j)}}+1} \cdot \langle \mathbf{a}_w, -\alpha^{(j)\vee} \rangle + \kappa_{-\alpha^{(j)}} \in p^{m_{-\alpha^{(j)}}} \cdot c \cdot Z_{-\alpha^{(j)}} + p^{N-2n+5} R,$$

(iii) in Lemma 4.1.4 (resp. in Lemma 4.2.4), we have

$$p^{m_{s(\beta^{(j)})}} \cdot (Y_{\beta^{(j)}} \pm c \cdot X_{s(\beta^{(j)})}) \\ \in p^{m_{s(\beta^{(j)})}} \cdot \left(c \cdot c_{\beta^{(j)}, m'_{\beta^{(j)}}} \cdot \mathbf{F}_{\geq -\alpha^{(j)}} + \mathbf{F}_{>\beta^{(j)}} + c \cdot \mathbf{F}_{>s(\beta^{(j)})}^{(\beta^{(j)})} \right) + p^{N-2n+5} R,$$

(iv) in Lemma 4.1.5 (resp. in Lemma 4.2.5), we have

$$V_{\beta^{(j)}}^\sharp \in v^{\kappa_{\beta^{(j)}}} (v+p)^{m_{\beta^{(j)}}} (X_{\beta^{(j)}} + \mathbf{F}_{>\beta^{(j)}}) + p^{N-2n+5} R[v].$$

5.2. Potentially crystalline deformation rings. Fix $\bar{\rho} : G_K \rightarrow \mathrm{GL}_n(\mathbb{F})$, and let τ be a tame inertial type. We assume that τ has an N -generic lowest alcove presentation ([LLHLMb, Definition 2.4.3]) with $N > 3n - 6$. Assume that $R_{\bar{\rho}}^{\eta, \tau} \neq 0$ so that in particular $\tilde{w}(\bar{\rho}, \tau)$ is defined. Recall that when $\ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) = \ell(t_\eta) - 1$ for some $j \in \mathcal{J}$, then by Proposition 2.1.2 we have a positive root $\alpha^{(j)}$ and a corresponding element $Z_{-\alpha^{(j)}}$ constructed in Lemma 4.1.3 (resp. in Lemma 4.2.3).

Lemma 5.2.1. *Let τ be a tame inertial type with an N -generic lowest alcove presentation, where $N > 3n - 6$. Assume that $\tilde{w}(\bar{\rho}, \tau)$ satisfies $\ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) \geq \ell(t_\eta) - 1$ for each $j \in \mathcal{J}$. Then there is a closed immersion*

$$\tilde{U}((\tilde{w}(\bar{\rho}, \tau)^{(j)})^*, \leq_\eta, \nabla_{\mathbf{a}^{(j)}}) \hookrightarrow \mathrm{Spec} R^{(j)},$$

where $\mathbf{a}^{(j)} \in \mathbb{Z}_p$ are the constants defined in (5) and

$$R^{(j)} \stackrel{\text{def}}{=} \begin{cases} \frac{\mathcal{O}[\{X_\beta | \beta \in \Phi^-\} \cup \{c\}]}{(c \cdot Z_{-\alpha^{(j)}} - p)} \otimes_{\mathcal{O}} \left(\bigotimes_{i=1}^n \frac{\mathcal{O}[a_i, Y_i]}{(a_i \cdot Y_i - 1)} \right) & \text{if } \ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) = \ell(t_\eta) - 1; \\ \mathcal{O}[\{X_\beta | \beta \in \Phi^-\}] \otimes_{\mathcal{O}} \left(\bigotimes_{i=1}^n \frac{\mathcal{O}[a_i, Y_i]}{(a_i \cdot Y_i - 1)} \right) & \text{otherwise.} \end{cases}$$

Proof. The results follow immediately from Proposition 4.3.1 and Proposition 4.3.2 together with equation (4). $\square$

Proposition 5.2.2. *Let τ be a tame inertial type with an N -generic lowest alcove presentation, where $N > 3n - 6$. Assume that $\tilde{w}(\bar{\rho}, \tau)$ satisfies that $\ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) \geq \ell(t_\eta) - 1$ for each $j \in \mathcal{J}$. Then there is a closed immersion*

$$\tilde{U}(\tilde{w}(\bar{\rho}, \tau)^*, \leq_\eta, \nabla_{\tau, \infty}) \hookrightarrow \mathrm{Spf} \left(\bigotimes_{\mathcal{O}, j \in \mathcal{J}} R^{(j)} \right)^{\wedge_p},$$

where the rings $R^{(j)}$ have been defined in Lemma 5.2.1.

Proof. The proof goes very similar to the ones of Proposition 4.3.2 and Proposition 4.3.1 together with Lemma 5.2.1. The only difference is that we need to take care of the error terms. For instance, if $(\tilde{w}(\bar{\rho}, \tau)^{(j)})^*$ is of colength one of the first form, and if β is a bad root sharing the row of α , then there exist $F_\beta \in \mathbf{F}_{\geq -\alpha}$, $G_\beta \in \mathbf{F}_{> \beta}$, and $H_{s(\beta)} \in \mathbf{F}_{> s(\beta)}^\beta$ such that

$$Y_\beta = -c \cdot X_{s(\beta)} + \left(c \cdot c_{\beta, m'_\beta} \cdot F_\beta + G_\beta + c \cdot H_{s(\beta)} \right) + O(p^{N-2n+5-m_{s(\beta)}}),$$

by Lemma 4.1.4. The coefficient c_{β, m'_β} corresponds to X_β , and so due to our generic assumption, by scaling Y_β we can further eliminate the variable Y_β . (Note that since $N - 2n + 5 > m_\beta + 1$ for all negative roots β we can indeed eliminate Y_β .)

Repeating the same arguments, we conclude that there is a surjection

$$\left(\bigotimes_{\mathcal{O}, j \in \mathcal{J}} R^{(j)} \right)^{\wedge_p} \twoheadrightarrow \mathcal{O}(\tilde{U}(\tilde{w}(\bar{\rho}, \tau)^*, \leq_\eta, \nabla_{\tau, \infty})),$$

which completes the proof. $\square$

Let $\tilde{A} \in \tilde{U}(\tilde{w}(\bar{\rho}, \tau)^*, \leq_\eta)(\mathbb{F})$ correspond to $\bar{\rho}|_{G_{K^\infty}}$ as explained at the end of §2.2.3 and let A be the image of $\tilde{A}$ under the map $\tilde{U}(\tilde{w}(\bar{\rho}, \tau)^*, \leq_\eta)(\mathbb{F}) \rightarrow U(\tilde{w}(\bar{\rho}, \tau)^*, \leq_\eta)(\mathbb{F})$. Set

$$\mathcal{J}_{0,A} \stackrel{\text{def}}{=} \{j \in \mathcal{J} \mid \ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) = \ell(t_\eta) - 1 \text{ and } Z_{-\alpha^{(j)}} \equiv 0 \pmod{\varpi}\}$$

(where we consider the image of $Z_{-\alpha^{(j)}}$ in $\mathbb{F}$ via the map induced by A). Note that for all $j \in \mathcal{J}$ such that $\ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) = \ell(t_\eta) - 1$ the corresponding element $c \in R^{(j)}$ is sent to 0 via the map $R^{(j)} \rightarrow \mathbb{F}$ induced by $\tilde{A}$.

Theorem 5.2.3. *Let τ be a tame inertial type with an N -generic lowest alcove presentation, where $N > 3n - 6$. Assume that $R_{\bar{\rho}}^{\eta, \tau} \neq 0$, and the shape $\tilde{w}(\bar{\rho}, \tau)$ satisfies $\ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) \geq \ell(t_\eta) - 1$ for each $j \in \mathcal{J}$. Then $R_{\bar{\rho}}^{\eta, \tau}$ is formally smooth over*

$$(19) \quad \widehat{\bigotimes_{\mathcal{O}, j \in \mathcal{J}_{0,A}}} \frac{\mathcal{O}[[X, Y]]}{(XY - p)}.$$

Proof. First, notice that we may replace $Z_{-\alpha}$ with $X_{-\alpha}$, due to the equations of $Z_{-\alpha}$ in Lemma 4.1.3 and Lemma 4.2.3. Noting that the completion of $\bigotimes_{\mathcal{O}, j \in \mathcal{J}} R^{(j)}$ at $\tilde{A}$ is an integral domain, we see that the completion of $\mathcal{O}(\tilde{U}(\tilde{w}(\bar{\rho}, \tau)^*, \leq \eta, \nabla_{\tau, \infty}))$ at $\tilde{A}$ is formally smooth over the ring in (19), by dimension counting and Proposition 5.2.2. Moreover the deformation ring $R_{\bar{\rho}}^{\leq \eta, \tau}$ is also formally smooth over the ring in (19) by equation (6) and [Ham75, Theorem 5]. As the latter is irreducible, so is $R_{\bar{\rho}}^{\leq \eta, \tau}$, in particular $R_{\bar{\rho}}^{\leq \eta, \tau} = R_{\bar{\rho}}^{\eta, \tau}$ which completes the proof. $\square$

Remark 5.2.4. Under stronger genericity assumptions on τ we have $\tilde{U}(\tilde{z}, \leq \eta, \nabla_{\tau, \infty}) \neq \emptyset$ whenever $\tilde{z} \in \text{Adm}(\eta)^\vee$ ([LLHLMb, Lemma 7.3.5]).

6. APPLICATIONS

In this section we elaborate on how the explicit description of the potentially crystalline deformation rings from Theorem 5.2.3 can provide information on representation theory and automorphic forms through following the philosophy of the mod p local Langlands correspondence.

6.1. Subextremal weights. In this section we refine, in Definition 6.1.2, the notion of *defect* for Serre weights of $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ introduced in [LLHLMb, §8.6.1] and [LLHLMa, §3.7].

We fix once and for all a lowest alcove presentation $\tilde{w}(\bar{\rho}^{\text{ss}})$ for $\bar{\rho}$. All tame inertial types will be endowed with the unique lowest alcove presentation compatible with $\tilde{w}(\bar{\rho}^{\text{ss}})$. Throughout this subsection, we assume $S_p = \{v\}$ so that $F_p^+ = K$.

Recall from [CEG⁺16, Theorem 3.7] that given a tame inertial type τ for K there exists an irreducible smooth representation $\sigma(\tau)$ of $\text{GL}_n(\mathcal{O}_K)$ which satisfies properties towards the inertial local Langlands correspondence. By [LLHLMb, Theorem 2.5.3], if τ has a 1-generic lowest alcove presentation $(s, \mu - \eta)$ then $\sigma(\tau)$ can be taken to be $R_s(\mu)$.

Let $\bar{\rho}^{\text{sp}} : I_K \rightarrow \text{GL}_n(\mathbb{F})$ be a $2n$ -generic tame inertial $\mathbb{F}$ -type. The choice of a lowest alcove presentation $\tilde{w}(\bar{\rho}^{\text{sp}}) = t_{\mu+\eta}s$ for it gives a map

$$\begin{aligned} \tau_{\bar{\rho}^{\text{sp}}} : \text{Adm}^{\text{reg}}(\eta) &\longrightarrow \{\tau : I_K \rightarrow \text{GL}_n(E)\} \\ t_\nu w &\mapsto \tau(sw, \mu + \eta + s(\nu)). \end{aligned}$$

(Note that given $\tilde{w} \in \text{Adm}^{\text{reg}}(\eta)$ we have $\tilde{w}(\bar{\rho}^{\text{sp}}, \tau_{\bar{\rho}^{\text{sp}}}(\tilde{w})) = \tilde{w}$ by construction.)

Recall from [LLHLMb, §2.3.1] the background on Deligne–Lusztig representations and their lowest alcove presentations. In particular given a Deligne–Lusztig representation R with a $2(n-1)$ -generic lowest alcove presentation $\tilde{w}(R)$ we have a set $\text{JH}_{\text{out}}(\bar{R})$ of *outer weights* for R , which is a subset of $\text{JH}(\bar{R})$, the latter being the

set of the Jordan–Hölder factors of the mod ϖ -reduction of an arbitrary $\mathcal{O}$ -lattice in R . We also recall that we have a bijection

$$(20) \quad \left\{ (\tilde{w}_1, \tilde{w}_2) \in \left(\widetilde{W}^+ \times \widetilde{W}_1^+ \right) / X^0(\underline{T}) \mid \tilde{w}_1 \uparrow \tilde{w}_h^{-1} \tilde{w}_2 \right\} \longrightarrow \text{Adm}^{\text{reg}}(\eta)$$

$$(\tilde{w}_1, \tilde{w}_2) \longmapsto \tilde{w}_2^{-1} w_0 \tilde{w}_1$$

from [LLHLMb, Remark 2.1.8], and a bijection

$$\left\{ (\tilde{w}_1, \tilde{w}_2) \in \left(\widetilde{W}^+ \times \widetilde{W}_1^+ \right) / X^0(\underline{T}) \mid \tilde{w}_1 \uparrow \tilde{w}_2 \right\} \longrightarrow W^?(\bar{\rho}^{\text{sp}})$$

$$(\tilde{w}_1, \tilde{w}_2) \longmapsto F_{(\tilde{w}_2, \tilde{w}(\bar{\rho}^{\text{sp}})(\tilde{w}_1)^{-1}(0))}$$

from [LLHLMb, Proposition 2.6.2]. As multiplication by $\tilde{w}_h$ gives a self bijection on $\widetilde{W}_1^+$, we finally obtain a bijection

$$\sigma_{\bar{\rho}^{\text{sp}}} : \text{Adm}^{\text{reg}}(\eta) \longrightarrow W^?(\bar{\rho}^{\text{sp}}).$$

$$\tilde{w}_2^{-1} w_0 \tilde{w}_1 \longmapsto F_{(\tilde{w}_h^{-1} \tilde{w}_2, \tilde{w}(\bar{\rho}^{\text{sp}})(\tilde{w}_1)^{-1}(0))}.$$

We write $\tilde{w}_{\bar{\rho}^{\text{sp}}}$ for the inverse of $\sigma_{\bar{\rho}^{\text{sp}}}$.

Lemma 6.1.1. *Assume that $\tilde{w}(\bar{\rho}^{\text{sp}})$ is a $2n$ -generic lowest alcove presentation for $\bar{\rho}^{\text{sp}}$, and let $\tilde{w} \in \text{Adm}^{\text{reg}}(\eta)$. Then $\sigma_{\bar{\rho}^{\text{sp}}}(\tilde{w}) \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}_{\text{out}}(\overline{\sigma(\tau_{\bar{\rho}^{\text{sp}}}(\tilde{w})))}$ and it satisfies the following property: for any $\sigma' \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau_{\bar{\rho}^{\text{sp}}}(\tilde{w})))}$ we have*

$$(21) \quad \tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma')^{(j)} \geq \tilde{w}^{(j)}$$

for all $j \in \mathcal{J}$ with equality for all $j \in \mathcal{J}$ if and only if $\sigma' = \sigma$.

Proof. This is an immediate consequence the proof of [LLHLMb, Proposition 8.6.3] of which we employ here the notation and convention. In particular we let $\tau \stackrel{\text{def}}{=} \tau_{\bar{\rho}^{\text{sp}}}(\tilde{w})$ so that $\tilde{w} = \tilde{w}(\bar{\rho}, \tau) \in \text{Adm}^{\text{reg}}(\eta)$. Using the bijection (20) we decompose $\tilde{w}$ as $\tilde{w}_2^{-1} w_0 \tilde{w}_1$ where $(\tilde{w}_1, \tilde{w}_2) \in \widetilde{W}^+ \times \widetilde{W}_1^+$ satisfies $\tilde{w}_1 \uparrow \tilde{w}_h^{-1} \tilde{w}_2$.

We have $\sigma_{\bar{\rho}^{\text{sp}}}(\tilde{w}) = F_{(\tilde{w}_h^{-1} \tilde{w}_2, \tilde{w}(\bar{\rho}^{\text{sp}})(\tilde{w}_1)^{-1}(0))}$ by definition, and the latter Serre weight is the element $\kappa \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$ defined in [LLHLMb, Proposition 8.6.3]. As $\tilde{w}(\bar{\rho}^{\text{sp}})(\tilde{w}_1)^{-1}(0) = \tilde{w}(\tau) \tilde{w}_2^{-1}(0)$ by [LLHLMb, Proposition 2.6.4], the weight κ is in $\text{JH}_{\text{out}}(\overline{\sigma(\tau)})$, by definition of $\text{JH}_{\text{out}}(\overline{\sigma(\tau)})$ (see [LLHLMb, Proposition 2.3.7] and the beginning of [LLHLMb, §2.3.1]).

The fact that for any $\sigma' \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$ the inequality (21) holds is immediate from the proof of [LLHLMb, Proposition 8.6.3]. We provide the details: by [LLHLMb, Proposition 2.6.4] any $\sigma' \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$ is of the form $\sigma' = F_{(\tilde{w}', \tilde{w}(\tau) \tilde{s}_2^{-1}(0))}$ for some $\tilde{w}' \in \widetilde{W}_1^+$ and some pair $(\tilde{s}_1, \tilde{s}_2) \in \widetilde{W}^+ \times \widetilde{W}^+$ satisfying $\tilde{s}_1 \uparrow \tilde{w}' \uparrow \tilde{w}_h^{-1} \tilde{s}_2$ and $\tilde{s}_2^{-1} w \tilde{s}_1 = \tilde{w}(\bar{\rho}^{\text{sp}}, \tau)$ for some $w \in \underline{W}$. By Wang's Theorem (see [LLHL19, Theorem 4.1.1]) the condition $\tilde{s}_2 \uparrow \tilde{w}_h \tilde{w}'$, which is defined *embeddingwise*, gives $\tilde{s}_2^{(j)} \leq \tilde{w}_h^{(j)} \tilde{w}'^{(j)}$ for all $j \in \mathcal{J}$ and by [LLHL19, Lemma 4.1.9] we conclude that

$$(\tilde{w}_h^{(j)} \tilde{w}'^{(j)})^{-1} w_0 \tilde{s}_1^{(j)} \geq (\tilde{s}_2^{(j)})^{-1} w_0 \tilde{s}_1^{(j)} \geq (\tilde{s}_2^{(j)})^{-1} w \tilde{s}_1^{(j)} = \tilde{w}^{(j)}$$

since $(\tilde{s}_2^{(j)})^{-1} w_0 \tilde{s}_1^{(j)}$, $(\tilde{w}_h^{(j)} \tilde{w}'^{(j)})^{-1} w_0 \tilde{s}_1^{(j)}$ are reduced expressions for all $j \in \mathcal{J}$ and $w_0 \geq w$. As $\tilde{s}_1 \uparrow \tilde{w}' \in \widetilde{W}_1^+$ we see that $(\tilde{w}_h \tilde{w}', \tilde{s}_1)$ defines an element in the left hand side of (20) and hence $\tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma') = (\tilde{w}_h \tilde{w}')^{-1} w_0 \tilde{s}_1$, proving (21). The fact that

the equality holds for all $j \in \mathcal{J}$ if and only if $\sigma' = \sigma_{\bar{\rho}^{\text{sp}}}(\tilde{w})$ is immediate since $\sigma_{\bar{\rho}^{\text{sp}}}$ is a bijection. $\square$

The usual order on the set of non-negative integers $\mathbb{Z}_{\geq 0}$ induces the product partial order on $\mathbb{Z}_{\geq 0}^{\mathcal{J}}$, and for $\underline{h} = (h^{(j)})_{j \in \mathcal{J}} \in \mathbb{Z}_{\geq 0}^{\mathcal{J}}$ we define

$$W_{\leq \underline{h}}^?(\bar{\rho}^{\text{sp}}) \stackrel{\text{def}}{=} \left\{ \sigma \in W^?(\bar{\rho}^{\text{sp}}) \mid \ell(\tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma)^{(j)}) \geq \ell(t_{\eta}) - h^{(j)} \text{ for all } j \in \mathcal{J} \right\}.$$

Let $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ be a continuous Galois representation such that $\bar{\rho}^{\text{ss}}$ is 0-generic (so that $\tilde{w}(\bar{\rho}^{\text{ss}}) \in \widetilde{W}$ is defined). We say that a tame inertial $\mathbb{F}$ -parameter $\bar{\rho}^{\text{sp}} : I_K \rightarrow \text{GL}_n(\mathbb{F})$ is a *specialization* of $\bar{\rho}$, and write $\bar{\rho} \rightsquigarrow \bar{\rho}^{\text{sp}}$, if there exists an n -generic tame inertial type τ such that $\bar{\rho}$ is τ -admissible and $\tilde{w}(\bar{\rho}, \tau) = \tilde{w}(\bar{\rho}^{\text{sp}}, \tau)$. (In this definition, the lowest alcove presentations of τ and $\bar{\rho}^{\text{sp}}$ are always assumed to be compatible with a fixed lowest alcove presentation of $\bar{\rho}^{\text{ss}}$.)

Let $\mathcal{X}_{n,K}$ be the Noetherian formal algebraic stack over $\text{Spf } \mathcal{O}$ defined in [EG, Definition 3.2.1]. It has the property that $\mathcal{X}_{n,K}(\mathbb{F})$ is isomorphic to the groupoid of continuous representations of G_K over rank n vector spaces over $\mathbb{F}$. Moreover there is a bijection $\sigma \mapsto \mathcal{C}_{\sigma}$ between Serre weights of $G \stackrel{\text{def}}{=} \text{GL}_n(k)$ and irreducible components of the reduced special fiber of $\mathcal{X}_{n,K}$, described in [LLHLMb, §7.4]. (We refer the reader to [LLHLMb, §2.2] concerning Serre weights and their lowest alcove presentations.) This bijection is a renormalization of the bijection $\sigma \mapsto \mathcal{X}_{n,\text{red}}^{\sigma}$ of [EG, Theorem 6.5.1].

In particular, if $\bar{\rho} \in \mathcal{X}_{n,K}(\mathbb{F})$ we define the set of *geometric weights* of $\bar{\rho}$ as

$$W^g(\bar{\rho}) \stackrel{\text{def}}{=} \{ \sigma \mid \bar{\rho} \in \mathcal{C}_{\sigma}(\mathbb{F}) \}.$$

Definition 6.1.2. Let $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ be a continuous Galois representation such that $\bar{\rho}^{\text{ss}}$ is 0-generic. For $\underline{h} = (h^{(j)})_{j \in \mathcal{J}} \in \mathbb{Z}_{\geq 0}^{\mathcal{J}}$ define $W_{\leq \underline{h}}^g(\bar{\rho})$ to be

$$W^g(\bar{\rho}) \cap \left(\bigcup_{\bar{\rho} \rightsquigarrow \bar{\rho}^{\text{sp}}} W_{\leq \underline{h}}^?(\bar{\rho}^{\text{sp}}) \right).$$

We call the elements of $W_{\leq \underline{h}}^g(\bar{\rho})$ the *Serre weights of $\bar{\rho}$ of defect at most $\underline{h}$* .

In what follows we write $\underline{1}$ for the tuple of $\underline{h} \in \mathbb{Z}_{\geq 0}^{\mathcal{J}}$ satisfying $h^{(j)} = 1$ for all $j \in \mathcal{J}$, and similarly for $\underline{0}$. Note that $W_{\leq \underline{0}}^g(\bar{\rho}) = W_{\text{extr}}^g(\bar{\rho})$ is the set of *extremal weights* in [LLHLMa, Definition 3.7.1].

6.2. Application to patching functors. We introduce the formalism of patching functors following [LLHLMa, §5.2], giving applications to the results on the deformation rings in §5.

6.2.1. L -parameters. Recall that F_p^+ is a finite unramified étale $\mathbb{Q}_p$ -algebra, which we write as $\prod_{v \in S_p} F_v^+$ for a finite set S_p and finite unramified extensions $F_v^+/\mathbb{Q}_p$. We assume throughout that for any $v \in S_p$ the coefficient field E (resp. $\mathbb{F}$) contains the image of any homomorphism $F_v^+ \hookrightarrow \overline{\mathbb{Q}_p}$ (resp. $k_v \hookrightarrow \overline{\mathbb{F}_p}$, where k_v denotes the residue field of F_v^+). We let $\underline{G}^{\vee}$ denote the product $\prod_{F_p^+ \rightarrow E} \text{GL}_{n/\mathcal{O}}^{\vee}$ (the dual group of $\text{Res}_{F_p^+/\mathbb{Q}_p} \text{GL}_{n/\mathcal{O}_p}$) and ${}^L \underline{G}(\mathbb{F}) \stackrel{\text{def}}{=} \underline{G}^{\vee} \rtimes \text{Gal}(E/\mathbb{Q}_p)$, where $\text{Gal}(E/\mathbb{Q}_p)$ acts on the set $\{ \iota : F_p^+ \rightarrow E \}$ by post-composition. An L -homomorphism over $A \in \{E, \mathbb{F}\}$ is a continuous homomorphism $\bar{\tau}_p : G_{\mathbb{Q}_p} \rightarrow {}^L \underline{G}(A)$. An L -parameter is a $\underline{G}^{\vee}(E)$ -conjugacy class of an L -homomorphism. A tame inertial L -parameter

is a $\underline{G}^\vee(E)$ -conjugacy class of an homomorphism $\tau : I_{\mathbb{Q}_p} \rightarrow \underline{G}^\vee(E)$ which has open kernel and factors through the tame quotient of $I_{\mathbb{Q}_p}$, and which admits an extension to an L -homomorphism. By [GHS18, Lemma 9.4.1, Lemma 9.4.5], the datum of an L -parameters $\bar{\tau}_p$ (resp. a tame inertial L -parameter τ) is equivalent to the datum of a collection of continuous homomorphisms $\{\rho_v : G_{F_v^+} \rightarrow \mathrm{GL}_n(A)\}_{v \in S_p}$ (resp. tame inertial types $\{\tau_v : I_{F_v^+} \rightarrow \mathrm{GL}_n(A)\}_{v \in S_p}$). Via this bijection, we can therefore give the notion of lowest alcove presentations and genericity for L -parameters. Given a tame inertial L -parameter τ with corresponding collection $\{\tau_v : I_{F_v^+} \rightarrow \mathrm{GL}_n(A)\}_{v \in S_p}$ of tame inertial types, we let $\sigma(\tau)$ be the tame smooth irreducible representation of $\mathrm{GL}_n(\mathcal{O}_p)$ over E given by $\otimes_{v \in S_p, E} \sigma(\tau_v)$, where for each $v \in S_p$ we let $\sigma(\tau_v)$ be the tame smooth irreducible representation of $\mathrm{GL}_n(\mathcal{O}_{F_v^+})$ over E attached to τ_v via the inertial local Langlands correspondence of [LLHLMb, Proposition 2.5.5].

6.2.2. Patching functors and Serre weights. Let now $\bar{\tau}_p : G_{\mathbb{Q}_p} \rightarrow {}^L \underline{G}(\mathbb{F})$ be an L -homomorphism, with corresponding collection $\{\bar{\rho}_v : G_{F_v^+} \rightarrow \mathrm{GL}_n(\mathbb{F})\}_{v \in S_p}$.

We let R^p be a nonzero complete local Noetherian equidimensional flat $\mathcal{O}$ -algebra with residue field $\mathbb{F}$ such that each irreducible component of $\mathrm{Spec} R^p$ and of $\mathrm{Spec} \bar{R}^p$ is geometrically irreducible, and define

$$R_{\bar{\tau}_p} \stackrel{\mathrm{def}}{=} \widehat{\bigotimes_{v \in S_p, \mathcal{O}} R_{\bar{\rho}_v}^\square}, \quad R_\infty \stackrel{\mathrm{def}}{=} R^p \widehat{\otimes}_{\mathcal{O}} R_{\bar{\tau}_p}$$

(we suppress the dependence on R^p in the notation of R_∞).

Given a tame inertial L -homomorphism $\tau : I_{\mathbb{Q}_p} \rightarrow \underline{G}^\vee(E)$, with corresponding collection $\{\tau_v\}_{v \in S_p}$, we define

$$R_{\bar{\tau}_p}^{\eta, \tau} \stackrel{\mathrm{def}}{=} \widehat{\bigotimes_{v \in S_p} R_{\bar{\rho}_v}^{\eta, \tau_v}}$$

and $R_\infty(\tau) \stackrel{\mathrm{def}}{=} R_\infty \otimes_{R_{\bar{\tau}_p}} R_{\bar{\tau}_p}^{\eta, \tau}$, and write X_∞ , $X_\infty(\tau)$, $\bar{X}_\infty(\tau)$ for $\mathrm{Spec} R_\infty$, $\mathrm{Spec} R_\infty(\tau)$, and $\mathrm{Spec} \bar{R}_\infty(\tau)$ respectively.

We write $\mathrm{Mod}(X_\infty)$ be the category of coherent sheaves over X_∞ and $\mathrm{Rep}_{\mathcal{O}}(\mathrm{GL}_n(\mathcal{O}_p))$ be the category of topological $\mathcal{O}[\mathrm{GL}_n(\mathcal{O}_p)]$ -modules which are finitely generated over $\mathcal{O}$.

Definition 6.2.3. A *weak patching functor* for an L -homomorphism $\bar{\tau}_p : G_{\mathbb{Q}_p} \rightarrow {}^L \underline{G}(\mathbb{F})$ is a nonzero covariant exact functor $M_\infty : \mathrm{Rep}_{\mathcal{O}}(\mathrm{GL}_n(\mathcal{O}_p)) \rightarrow \mathrm{Mod}(X_\infty)$ such that for any tame inertial L -homomorphism τ and any $\mathcal{O}$ -lattice $\sigma(\tau)^\circ$ in $\sigma(\tau)$ one has:

- (1) $M_\infty(\sigma(\tau)^\circ)$ is a maximal Cohen–Macaulay sheaf on $X_\infty(\tau)$; and
- (2) for all $\sigma \in \mathrm{JH}(\overline{\sigma(\tau)})$, $M_\infty(\sigma)$ is either zero or a maximal Cohen–Macaulay sheaf on $\bar{X}_\infty(\tau)$.

Given a weak patching functor for an L -homomorphism $\bar{\tau}_p$ we thus define

$$(22) \quad W_{M_\infty}(\bar{\tau}_p) \stackrel{\mathrm{def}}{=} \{\sigma \mid \sigma \text{ is a } 3(n-1)\text{-deep Serre weight of } G \text{ and } M_\infty(\sigma) \neq 0\}.$$

By [LLHLMb, Proposition 2.4.5] and [LLHLMa, Theorem 5.1.1, Proposition 5.4.1] we see that if $\bar{\tau}_p$ is $6(n-1)$ -generic then condition that σ is $3(n-1)$ -deep in the right hand side of (22) is automatically satisfied.

6.2.4. *Modularity of weights of defect at most one.* We now assume $S_p = \{v\}$ and $F_v^+ = K$. In particular a continuous homomorphism $\bar{\rho} : G_K \rightarrow \mathrm{GL}_n(\mathbb{F})$ can be seen as an L -parameter, and have weak patching functors associated to it.

Proposition 6.2.5. *Let $\bar{\rho} : G_K \rightarrow \mathrm{GL}_n(\mathbb{F})$ be $6(n-1)$ -generic and let M_∞ be a weak patching functor for $\bar{\rho}$. Then the following are equivalent:*

- (1) $W_{\leq 0}^g(\bar{\rho}) \cap W_{M_\infty}(\bar{\rho}) \neq \emptyset$; and
- (2) $W_{\leq 1}^g(\bar{\rho}) \subseteq W_{M_\infty}(\bar{\rho})$.

Before the proof, we record Lemma 6.2.6, which will be also used in the proof of [PQ22, Conjecture 5.3.1] in §6.3.

Lemma 6.2.6. *Let $\bar{\rho}^{\mathrm{sp}} : G_K \rightarrow \mathrm{GL}_n(\mathbb{F})$ be $6(n-1)$ -generic and let $\tilde{w} \in \mathrm{Adm}^{\mathrm{reg}}(\eta)$ satisfy $\ell(t_\eta) - \ell(\tilde{w}^{(j)}) \leq 1$ for all $j \in \mathcal{J}$. Let $\tau \stackrel{\mathrm{def}}{=} \tau_{\bar{\rho}^{\mathrm{sp}}}(\tilde{w})$ and write $\tilde{w} = \tilde{w}_2^{-1} w_0 t_\nu \tilde{w}_1$ where $\tilde{w}_1, \tilde{w}_2 \in \widetilde{W}_1^+$ and $\nu \in X^+(\underline{T})$ are uniquely determined up to $X^0(\underline{T})$. Then:*

$$W^?(\bar{\rho}^{\mathrm{sp}}) \cap \mathrm{JH}(\overline{\sigma(\tau)}) = \left\{ \sigma_{\bar{\rho}^{\mathrm{sp}}}(\tilde{w}') \mid \tilde{w}'^{(j)} \in S^{(j)} \text{ for all } j \in \mathcal{J} \right\},$$

where $S^{(j)} \stackrel{\mathrm{def}}{=} \left\{ (\tilde{w}_2^{(j)})^{-1} w_0 t_{\nu^{(j)}} \tilde{w}_1^{(j)}, (\tilde{w}_h^{(j)} \tilde{w}_1^{(j)})^{-1} w_0 \tilde{w}_1^{(j)} \right\}$. Moreover, $\#S^{(j)} = 1$ if and only if $\tilde{w}_1^{(j)} = \tilde{w}_h^{-1} \tilde{w}_2^{(j)}$ and $\nu^{(j)} \in X^0(T)$, if and only if $\ell(\tilde{w}^{(j)}) = \ell(t_\eta)$.

Note that the existence and uniqueness of the decomposition of $\tilde{w}$ in the lemma is guaranteed by [LLHLMb, Proposition 2.1.5].

Proof. The relation $F_{(\tilde{x}, \omega)} \in W^?(\bar{\rho}^{\mathrm{sp}}) \cap \mathrm{JH}(\overline{\sigma(\tau)})$ is equivalent, by [LLHLMb, Proposition 2.6.4], to the existence of a factorization $\tilde{w} = (\tilde{x}_2)^{-1} s \tilde{x}_1$ with $s \in \underline{W}$, $\tilde{x}_1, \tilde{x}_2 \in \widetilde{W}_1^+$, $\tilde{x}_1 \uparrow \tilde{x} \uparrow \tilde{w}_h^{-1} \tilde{x}_2$ and $\omega = \tilde{w}(\bar{\rho}^{\mathrm{sp}})(\tilde{x}_1)^{-1}(0)$. By Lemma 2.1.4 applied to each $\tilde{w}^{(j)}$ we have $s = w_0$ and for all $j \in \mathcal{J}$ either $\tilde{x}^{(j)} = \tilde{x}_1^{(j)} \in \tilde{w}_1^{(j)} X^0(T)$ (in which case, by the uniqueness of the factorization in [LLHLMb, Proposition 2.1.5] we further have $\tilde{x}_2^{(j)} = t_{-w_0(\nu^{(j)})} \tilde{w}_2^{(j)}$) or $\tilde{x}^{(j)} = (\tilde{w}_h^{(j)})^{-1} \tilde{x}_2^{(j)} \in (\tilde{w}_h^{(j)})^{-1} \tilde{w}_2^{(j)} X^0(T)$ (in which case, by the uniqueness of the factorization in [LLHLMb, Proposition 2.1.5] we further have $\tilde{x}_1^{(j)} = t_{\nu^{(j)}} \tilde{w}_1^{(j)}$). The conclusion follows now from the definition of the map $\sigma_{\bar{\rho}^{\mathrm{sp}}}$. $\square$

Proof of Proposition 6.2.5. The proof is by induction on the following quantity $\delta_{\bar{\rho}}(\sigma)$ attached to a Serre weight $\sigma \in W_{\leq 1}^g(\bar{\rho})$:

$$(23) \quad \delta_{\bar{\rho}}(\sigma) \stackrel{\mathrm{def}}{=} \min \left\{ \sum_{j \in \mathcal{J}} \ell(t_\eta) - \ell(\tilde{w}_{\bar{\rho}^{\mathrm{sp}}}(\sigma)^{(j)}) \mid \bar{\rho} \rightsquigarrow \bar{\rho}^{\mathrm{sp}} \right\}.$$

We fix throughout the proof a choice of an algebraic central character ζ ; all lowest alcove presentations below will be chosen to be compatible with ζ .

By definition of $W_{\leq 1}^g(\bar{\rho})$ and Lemma 6.1.1, for each $\sigma \in W_{\leq 1}^g(\bar{\rho})$ there exist $\bar{\rho} \rightsquigarrow \bar{\rho}^{\mathrm{sp}}$ and a tame inertial type τ (depending on $\bar{\rho}^{\mathrm{sp}}$) such that

- (1) $\sigma \in W^g(\bar{\rho}) \cap W_{\leq 1}^?(\bar{\rho}^{\mathrm{sp}}) \cap \mathrm{JH}_{\mathrm{out}}(\overline{\sigma(\tau)})$;
- (2) $\tilde{w}_{\bar{\rho}^{\mathrm{sp}}}(\sigma) = \tilde{w}(\bar{\rho}^{\mathrm{sp}}, \tau) \in \mathrm{Adm}^{\mathrm{reg}}(\eta)$; and
- (3) $\tilde{w}_{\bar{\rho}^{\mathrm{sp}}}(\sigma')^{(j)} \geq \tilde{w}_{\bar{\rho}^{\mathrm{sp}}}(\sigma)^{(j)}$ for all $j \in \mathcal{J}$ and all $\sigma' \in W^?(\bar{\rho}^{\mathrm{sp}}) \cap \mathrm{JH}(\overline{\sigma(\tau)})$.

(Note that condition (2) determines τ uniquely, and we thus let $\tau = \tau_{\bar{\rho}^{\mathrm{sp}}}(\tilde{w}_{\bar{\rho}^{\mathrm{sp}}}(\sigma))$ in the notation of Lemma 6.1.1.)

For each $\sigma \in W_{\leq 1}^g(\bar{\rho})$ with corresponding $\bar{\rho} \rightsquigarrow \bar{\rho}^{\text{sp}}$ and τ as above, we now claim that

$$(24) \quad W^g(\bar{\rho}) \cap \text{JH}(\overline{\sigma(\tau)}) = W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$$

or, equivalently by [LLHLMa, Theorem 5.1.1], that any $\sigma' \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$ is in $W^g(\bar{\rho})$. Indeed by Lemma 6.2.6 there exists $\sigma_0 \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$ such that

(1) $\delta_{\bar{\rho}}(\sigma_0) = 0$; and

(2) for all $\sigma' \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$ we have $\tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma')^{(j)} \in \{\tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma_0)^{(j)}, \tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma)^{(j)}\}$.

By item (1) and [LLHLMb, Theorem 7.4.2] (see also [LLHLMa, Remark 3.9.1]) we have $\sigma_0 \in W^g(\bar{\rho})$ and thus $\bar{\rho} \in \mathcal{C}_\sigma \cap \mathcal{C}_{\sigma_0}$. Recall from [LLHLMb, equation (4.11) and Definition 4.6.1] the variety $\tilde{C}_\sigma^\zeta$, with its decomposition $\prod_{j \in \mathcal{J}} \tilde{C}_{\sigma'}^{\zeta, (j)}$. Each $\tilde{C}_{\sigma'}^{\zeta, (j)}$ is determined explicitly by $\tilde{w}(\bar{\rho}^{\text{sp}})^{(j)}$ and $\tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma')^{(j)}$ (see *loc. cit.* Definition 4.3.2, Theorem 4.3.9) so that $\tilde{C}_{\sigma'}^{\zeta, (j)} = \tilde{C}_\sigma^{\zeta, (j)}$ if $\tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma')^{(j)} = \tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma)^{(j)}$, and $\tilde{C}_{\sigma'}^{\zeta, (j)} = \tilde{C}_{\sigma_0}^{\zeta, (j)}$ if $\tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma')^{(j)} = \tilde{w}_{\bar{\rho}^{\text{sp}}}(\sigma_0)^{(j)}$. Thus, by item (2), for all $\sigma' \in W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)})$ we have

$$\tilde{C}_\sigma^\zeta \cap \tilde{C}_{\sigma_0}^\zeta = \prod_{j \in \mathcal{J}} \tilde{C}_\sigma^{\zeta, (j)} \cap \tilde{C}_{\sigma_0}^{\zeta, (j)} \subseteq \prod_{j \in \mathcal{J}} \tilde{C}_{\sigma'}^{\zeta, (j)} = \tilde{C}_{\sigma'}^\zeta$$

which together with [LLHLMb, Theorem 7.4.2] implies $\mathcal{C}_\sigma \cap \mathcal{C}_{\sigma_0} \subseteq \mathcal{C}_{\sigma'}$. This proves (24).

We now proceed to the inductive argument. We freely use the notation for cycles from patching functors introduced in [LLHLMa, §5.3], in particular we write $\overline{\text{pr}}$ to indicate the projection map from cycles over the reduced union $\cup_{\tau \in \mathcal{T}} \overline{X}_\infty(\tau)$ (for a set of generic tame inertial L -parameters $\mathcal{T}$) to cycles over the special fiber of the multi-type deformation ring associated to the set $\mathcal{T}$. The set $\mathcal{T}$ can be fixed to satisfy condition (ii) in [LLHLMa, §5.3] since all the tame inertial types τ involved in this proof satisfy $\ell(\tilde{w}(\bar{\rho}, \tau)^{(j)}) \geq \ell(t_\eta) - 1$ for all $j \in \mathcal{J}$, hence Theorem 5.2.3 applies. Furthermore, given $\sigma \in W^g(\bar{\rho})$ we will write $\mathcal{C}_\sigma(\bar{\rho})$ to denote the pullback of the component $\mathcal{C}_\sigma$ to the versal ring of $\mathcal{X}_{n,K}$ at $\bar{\rho} \in \mathcal{X}_{n,K}(\mathbb{F})$.

For $\sigma \in W_{\leq 1}^g(\bar{\rho})$, we prove by induction on $\delta_{\bar{\rho}}(\sigma)$ that the support of $\overline{\text{pr}} \circ Z(M_\infty(\sigma))$ contains $\mathcal{C}_\sigma(\bar{\rho})$, and that any *other* component in the support of $\overline{\text{pr}} \circ Z(M_\infty(\sigma))$ is of the form $\mathcal{C}_\kappa(\bar{\rho})$ with $\kappa \in W^g(\bar{\rho})$, $\delta_{\bar{\rho}}(\kappa) < \delta_{\bar{\rho}}(\sigma)$.

If $\delta_{\bar{\rho}}(\sigma) = 0$ then $\sigma \in W_{\leq 0}^g(\bar{\rho})$, and $W_{\leq 0}^g(\bar{\rho}) \cap W_{M_\infty}(\bar{\rho}) \neq \emptyset$ implies $W_{\leq 0}^g(\bar{\rho}) \subseteq W_{M_\infty}(\bar{\rho})$ by the main result of [LLHLMa]. As $W^g(\bar{\rho}) \cap \text{JH}(\overline{\sigma(\tau)}) = W^?(\bar{\rho}^{\text{sp}}) \cap \text{JH}(\overline{\sigma(\tau)}) = \{\sigma\}$ by Lemma 6.2.6 and $M_\infty(\overline{\sigma(\tau)}^\circ)$ has full support over (a formally smooth modification of) $\overline{R}_\rho^{\eta, \tau}$, we conclude that the cycle $\overline{\text{pr}} \circ Z(M_\infty(\overline{\sigma(\tau)}^\circ))$ is supported on $\mathcal{C}_\sigma(\bar{\rho})$. (Here and below we write $\overline{\sigma(\tau)}^\circ$ to denote the mod ϖ -reduction of any $\mathcal{O}$ -lattice $\sigma(\tau)^\circ$ in $\sigma(\tau)$.)

Assume now that $\delta_{\bar{\rho}}(\sigma) > 0$ and that for any $\sigma' \in W_{\leq 1}^g(\bar{\rho})$ with $\delta_{\bar{\rho}}(\sigma') < \delta_{\bar{\rho}}(\sigma)$ the cycle $\overline{\text{pr}} \circ Z(M_\infty(\sigma'))$ is supported on $\mathcal{C}_{\sigma'}(\bar{\rho})$ and possibly other components of the form $\mathcal{C}_\kappa(\bar{\rho})$ where $\kappa \in W^g(\bar{\rho})$ and $\delta_{\bar{\rho}}(\kappa) < \delta_{\bar{\rho}}(\sigma')$. We have

$$\begin{aligned} (25) \quad \text{Supp}(Z(\overline{R}_\rho^{\eta, \tau})) &= \text{Supp}(\text{red}(Z(R_\rho^{\eta, \tau})[1/p])) \\ &= \text{Supp}(\text{red} \circ \text{pr}(Z(M_\infty(\sigma(\tau)^\circ)[1/p]))) \\ &= \text{Supp}(\overline{\text{pr}} \circ \text{red}(Z(M_\infty(\sigma(\tau)^\circ)[1/p]))) \\ &= \text{Supp}(\overline{\text{pr}} \circ Z(M_\infty(\overline{\sigma(\tau)}^\circ))), \end{aligned}$$

where the first and last equality follow from [LLHLMa, Proposition 5.3.1], the second from the fact that $R_{\bar{\rho}}^{\eta, \tau}$ is geometrically integral (Theorem 5.2.3), the third from $\text{red} \circ \text{pr} = \overline{\text{pr}} \circ \text{red}$ (see [LLHLMa, §5.3]). Since $R_{\bar{\rho}}^{\eta, \tau}$ is geometrically integral by Theorem 5.2.3, by exactness of M_{∞} and [LLHLMa, Theorem 5.1.1 and Proposition 5.4.1] we have

$$\begin{aligned} \overline{\text{pr}} \circ Z(M_{\infty}(\overline{\sigma(\tau)})) &= \sum_{\sigma' \in W^?(\overline{\rho^{\text{sp}}}) \cap \text{JH}(\overline{\sigma(\tau)})} \overline{\text{pr}} \circ Z(M_{\infty}(\sigma')) \\ &= \overline{\text{pr}} \circ Z(M_{\infty}(\sigma)) + \sum_{\sigma' \in W^?(\overline{\rho^{\text{sp}}}) \cap \text{JH}(\overline{\sigma(\tau)}) \setminus \{\sigma\}} \overline{\text{pr}} \circ Z(M_{\infty}(\sigma')). \end{aligned}$$

From the definition of $\delta_{\bar{\rho}}$ together with Lemma 6.1.1, we easily check that $\delta_{\bar{\rho}}(\sigma') < \delta_{\bar{\rho}}(\sigma)$ for all $\sigma' \in W^?(\overline{\rho^{\text{sp}}}) \cap \text{JH}(\overline{\sigma(\tau)}) \setminus \{\sigma\}$. We thus deduce, using the inductive hypothesis and (24), that

$$\bigcup_{\sigma' \in W^?(\overline{\rho^{\text{sp}}}) \cap \text{JH}(\overline{\sigma(\tau)}) \setminus \{\sigma\}} \text{Supp}(\overline{\text{pr}} \circ Z(M_{\infty}(\sigma'))) = \bigcup_{\sigma' \in W^?(\overline{\rho^{\text{sp}}}) \cap \text{JH}(\overline{\sigma(\tau)}) \setminus \{\sigma\}} \mathcal{C}_{\sigma'}(\bar{\rho}).$$

On the other hand, we have

$$\text{Supp}(Z(\overline{R}_{\bar{\rho}}^{\eta, \tau})) = \bigcup_{\sigma' \in W^g(\bar{\rho}) \cap \text{JH}(\overline{\sigma(\tau)})} \mathcal{C}_{\sigma'}(\bar{\rho})$$

by [LLHLMb, Theorem 7.4.2] (and [Sta19, Lemma 0DRB, Lemma 0DRD and Definition 0DRA]) and hence $\overline{\text{pr}} \circ Z(M_{\infty}(\sigma))$ is necessarily supported on $\mathcal{C}_{\sigma}(\bar{\rho})$, and possibly on other components $\mathcal{C}_{\sigma'}(\bar{\rho})$ with $\sigma' \in W^g(\bar{\rho})$ and $\delta_{\bar{\rho}}(\sigma') < \delta_{\bar{\rho}}(\sigma)$. $\square$

Remark 6.2.7. If, in the statement of Proposition 6.2.5, we furthermore assume that weak patching functor M_{∞} is *minimal* ([LLHLMa, Definition 5.2.1]), then equation (25) can be replaced with the stronger statement

$$e(R_{\bar{\rho}}^{\eta, \tau}) = e(M_{\infty}(\overline{\sigma(\tau)}))$$

(where $e(\cdot)$ denotes the Hilbert–Samuel multiplicity), which forces $e(M_{\infty}(\sigma)) = 1$ for all $\sigma \in W^?(\overline{\rho^{\text{sp}}}) \cap \text{JH}(\overline{\sigma(\tau)})$.

Remark 6.2.8. Let $\bar{\tau}_p : G_{\mathbb{Q}_p} \rightarrow {}^L\underline{G}(\mathbb{F})$ be an L -parameter, with corresponding collection $\{\bar{\rho}_v : G_{F_v^+} \rightarrow \text{GL}_n(\mathbb{F})\}_{v \in S_p}$. For each $v \in S_p$ let $\mathcal{J}_v$ denote the set of ring homomorphisms $\{k_v \hookrightarrow \mathbb{F}\}$, so that $\mathcal{J}_p = \prod_{v \in S_p} \mathcal{J}_v$. Given $\underline{h}_v \in \mathbb{Z}_{\geq 0}^{\mathcal{J}_v}$ for each $v \in S_p$ we then have a collection $\{W_{\leq \underline{h}_v}^g(\bar{\rho}_v)\}_{v \in S_p}$ whose elements are Serre weights for G by taking tensor products over $v \in S_p$. In particular, given $\underline{h} \in \mathbb{Z}_{\geq 0}^{\mathcal{J}_p}$ we can define the set $W_{\leq \underline{h}}^g(\bar{\tau}_p)$ for an L -parameter $\bar{\tau}_p : G_{\mathbb{Q}_p} \rightarrow {}^L\underline{G}(\mathbb{F})$. The proofs of Proposition 6.2.5 and Lemma 6.2.6 go through, *mutatis mutandis*, replacing $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ with an L -homomorphism $\bar{\tau}_p : G_{\mathbb{Q}_p} \rightarrow {}^L\underline{G}(\mathbb{F})$ (for these kind of passages from $\#S_p = 1$ to $\#S_p > 1$ see also [LLHLMb, Remark 7.3.4]).

6.3. Global applications and the conjecture of [PQ22]. We apply the results above to obtain our main global applications. We follow the setup and notation of [LLHLMa, §5.5.1, 5.5.2, 5.5.3]. In particular, we have a totally real field $F^+/\mathbb{Q}$ not equal to $\mathbb{Q}$, and F/F^+ a CM extension. We assume from now on that all places of F^+ above p are unramified over $\mathbb{Q}_p$, and that they are further totally split in F . Given a reductive group $G_{/F^+}$ which is an outer form for GL_n , split over F ,

and such that $G(F^+ \otimes_{\mathbb{Q}} \mathbb{R}) = U_n(F^+ \otimes_{\mathbb{Q}} \mathbb{R})$, we define for a continuous Galois representation $\bar{\tau} : G_{F^+} \rightarrow \mathcal{G}_n(\mathbb{F})$ the notion of being *automorphic* (relatively to G_{F^+}) as in [LLHLMa, Definition 5.5.1], as well as the set $W(\bar{\tau})$ of modular Serre weights of $\bar{\tau}$ (*loc. cit.* Definition 5.5.2). Here, $\mathcal{G}_n$ is the group scheme defined in [CHT08, §2]. Finally, let $\bar{\tau}_p$ be the L -homomorphism induced from the collection of continuous representations $\bar{\tau}|_{G_{F_v^+}} : G_{F_v^+} \rightarrow \mathrm{GL}_n(\mathbb{F})$ for $v|p$. In particular, we denote by S_p the set of the finite places of F^+ above p .

Let $\bar{\tau} : G_{F^+} \rightarrow \mathcal{G}_n(\mathbb{F})$ be automorphic and such that $\bar{\tau}(G_{F(\zeta_p)}) \subseteq \mathrm{GL}_n(\mathbb{F})$ is adequate. Then by [LLHLMa, Lemma 5.5.4] we can and do fix a weak patching functor M_{∞} for the L -homomorphism $\bar{\tau}_p$ such that for any Serre weight σ of $G = \prod_{v \in S_p} \mathrm{GL}_n(k_v)$ we have

$$(26) \quad M_{\infty}(\sigma) \neq 0 \iff \sigma \in W(\bar{\tau}).$$

Theorem 6.3.1 (Modularity of weights of defect at most one). *Let $\bar{\tau} : G_{F^+} \rightarrow \mathcal{G}(\mathbb{F})$ be an automorphic representation such that*

- $\bar{\tau}(G_{F(\zeta_p)}) \subseteq \mathrm{GL}_n(\mathbb{F})$ is adequate; and
- $\bar{\tau}_p$ is $6(n-1)$ -generic.

Then the following are equivalent:

- (1) $W_{\leq 0}^g(\bar{\tau}_p) \cap W(\bar{\tau}) \neq \emptyset$; and
- (2) $W_{\leq 1}^g(\bar{\tau}_p) \subset W(\bar{\tau})$.

Proof. This follows from Proposition 6.2.5 and Remark 6.2.8 using (26). $\square$

Remark 6.3.2. Keep the setup and notation of Theorem 6.3.1 and Remark 6.2.8. If $\sigma \in W_{\leq 1}^g(\bar{\tau}_p^{\mathrm{sp}})$ for some $\bar{\tau}_p \rightsquigarrow \bar{\tau}_p^{\mathrm{sp}}$, then $\sigma \in W_{M_{\infty}}(\bar{\tau})$ if and only if $\sigma \in W^g(\bar{\tau}_p)$ if and only if $\sigma \in W_{\leq 1}^g(\bar{\tau}_p)$. For each such σ , there exists τ such that

$$(27) \quad \tilde{w}(\bar{\tau}_p, \tau) = \tilde{w}(\bar{\tau}_p^{\mathrm{sp}}, \tau) \quad \text{and} \quad \ell(\tilde{w}(\bar{\tau}_p^{\mathrm{sp}}, \tau)^{(j)}) \geq \ell(t_{\eta}) - 1$$

for each $j \in \mathcal{J}_p$ and $\sigma \in W^?(\bar{\tau}_p^{\mathrm{sp}}) \cap \mathrm{JH}(\bar{\sigma}(\tau))$, with $\sigma \in W^g(\bar{\tau}_p)$ if and only if $\bar{\tau}_p$ satisfies $Z_{-\alpha(j)} = 0$ for each $j \in \mathcal{J}_p$ for which the inequality in (27) is an equality.

We now recall the setup of the local–global compatibility result of [PQ22]. Assume that p is totally split in F and fix a place $w|p$ of F . Assume that $\bar{\tau}_w \stackrel{\mathrm{def}}{=} \bar{\tau}|_{G_{F_w}}$ is Fontaine–Laffaille of niveau one, and that satisfies a geometric genericity condition dictated by its position in the moduli of Fontaine–Laffaille modules (see [PQ22, Definition 3.2.5]). In particular we have a lowest alcove presentation $(1, \mu)$ for $\bar{\tau}_w^{\mathrm{ss}}$ (with $\mu = (c_{n-1}, c_{n-2}, \dots, c_1, c_0)$ in the notation of §1 in *loc. cit.*) and a niveau one tame inertial type τ with lowest alcove presentation $(1, s_{\alpha}((\mu^{\square, i_1, j_1})^{\vee}))$ (with $\mu^{\square, i_1, j_1} \stackrel{\mathrm{def}}{=} \mu^{\vee} + \langle \langle \eta, \alpha^{\vee} \rangle + 1 \rangle \alpha$ where $0 \leq i_1, j_1 \leq n-1$ corresponds to a positive root $\alpha = \alpha_{i_1+1, j_1+1}$). Then by the proof of [PQ22, Lemma 3.4.1] (namely, from the expression of $\mathrm{Mat}_{e''}(\phi)$ in *loc. cit.*) we see that $\tilde{w}(\bar{\tau}_w, \tau) = t_{\eta-\alpha} s_{\alpha}$, which is a regular colength one shape. In particular $\bar{\tau}_w \rightsquigarrow \bar{\tau}^{\mathrm{sp}}$ with $\tilde{w}(\bar{\tau}^{\mathrm{sp}}, \tau) = t_{\eta-\alpha} s_{\alpha}$. By Lemma 6.2.6 we obtain:

Theorem 6.3.3 (Conjecture 5.3.1 [PQ22]). *Let $w|p$. Assume that $\bar{\tau}_w$ is Fontaine–Laffaille of niveau one, that $(\bar{\tau}_w)_{n-i_0, n-j_0}$ is Fontaine–Laffaille generic in the sense of [PQ22, Definition 3.2.5], and moreover that $\bar{\tau}_w^{\mathrm{ss}}$ is $3(n-1)$ -generic. Then*

$$W_w(\bar{\tau}) \cap \mathrm{JH}(\bar{\sigma}(\tau)) \subseteq \{F(\mu)^{\vee}, F(\mu^{\square, i_1, j_1})^{\vee}\},$$

where $W_w(\bar{\tau})$ denotes the set of modular weights for $\bar{\tau}$ at w (as defined in the paragraph just below [PQ22, Definition 5.2.2]).

Proof. By [LLHLMa, Theorem 5.1.1], $W_w(\bar{\tau}) \cap \mathrm{JH}(\bar{\sigma}(\tau)) \subseteq W^?(\bar{\rho}^{\mathrm{sp}}) \cap \mathrm{JH}(\bar{\sigma}(\tau))$. Now, it follows immediately from Lemma 6.2.6, as $\tilde{w}(\bar{\rho}^{\mathrm{sp}}, \tau) = s_\alpha t_{\eta-\alpha}$. $\square$

The above result removes the weight elimination condition of [PQ22, Theorem 5.6.2]. However the fact that the deformation ring $R_{\bar{\rho}}^{\eta, \tau}$ is formally smooth over $\mathcal{O}$ when $Z_{-\alpha} \not\equiv 0$ modulo ϖ makes the argument of [PQ22, Theorem 5.6.2] more direct, since the modules of algebraic automorphic forms are in this case free over the Hecke algebra by patching arguments (cf. [PQ22, Remark 5.4.6]).

ACKNOWLEDGMENT

Part of the work was carried out during a visit at the Università degli Studi di Padova (2019), which we would like to heartily thank for the excellent working conditions which provided to us.

REFERENCES

- [BD14] Christophe Breuil and Fred Diamond, *Formes modulaires de Hilbert modulo p et valeurs d'extensions entre caractères galoisiens* (French, with English and French summaries), Ann. Sci. Éc. Norm. Supér. (4) **47** (2014), no. 5, 905–974, DOI 10.24033/asens.2230. MR3294620
- [BHH⁺a] Christophe Breuil, Florian Herzig, Yongquan Hu, Stefano Morra, and Benjamin Schraen, *Conjectures and results on modular representations of $\mathrm{GL}_n(k)$ for a p -adic field k* , Preprint, [arXiv:2102.06188](#), 2021.
- [BHH⁺b] Christophe Breuil, Florian Herzig, Yongquan Hu, Stefano Morra, and Benjamin Schraen, *Gelfand–Kirillov dimension and mod p cohomology for GL_2* , Preprint, [arXiv:2009.03127](#), 2020.
- [CDT99] Brian Conrad, Fred Diamond, and Richard Taylor, *Modularity of certain potentially Barsotti–Tate Galois representations*, J. Amer. Math. Soc. **12** (1999), no. 2, 521–567, DOI 10.1090/S0894-0347-99-00287-8. MR1639612
- [CEG⁺16] Ana Caraiani, Matthew Emerton, Toby Gee, David Geraghty, Vytas Paškūnas, and Sug Woo Shin, *Patching and the p -adic local Langlands correspondence*, Camb. J. Math. **4** (2016), no. 2, 197–287, DOI 10.4310/CJM.2016.v4.n2.a2. MR3529394
- [CHT08] Laurent Clozel, Michael Harris, and Richard Taylor, *Automorphy for some l -adic lifts of automorphic mod l Galois representations*, Publ. Math. Inst. Hautes Études Sci. **108** (2008), 1–181, DOI 10.1007/s10240-008-0016-1. With Appendix A, summarizing unpublished work of Russ Mann, and Appendix B by Marie-France Vignéras. MR2470687
- [DL21] Andrea Dotto and Daniel Le, *Diagrams in the mod p cohomology of Shimura curves*, Compos. Math. **157** (2021), no. 8, 1653–1723, DOI 10.1112/s0010437x21007375. MR4283560
- [EG] Matthew Emerton and Toby Gee, *Moduli stacks of étale (φ, Γ) -modules and the existence of crystalline lifts*, Annals of Mathematics Studies, vol. 215, Princeton University Press, Princeton, NJ, [2023] ©2023, DOI 10.1515/9780691241364. MR4529886
- [EGS15] Matthew Emerton, Toby Gee, and David Savitt, *Lattices in the cohomology of Shimura curves*, Invent. Math. **200** (2015), no. 1, 1–96, DOI 10.1007/s00222-014-0517-0. MR3323575
- [EL] John Enns and Heejons Lee, *Mod p local–global compatibility for $\mathrm{GSp}_4(\mathbf{Q}_p)$ in the ordinary case*, Preprint, [arXiv:2112.12256](#), 2021.
- [GHS18] Toby Gee, Florian Herzig, and David Savitt, *General Serre weight conjectures*, J. Eur. Math. Soc. (JEMS) **20** (2018), no. 12, 2859–2949, DOI 10.4171/JEMS/826. MR3871496
- [Ham75] Eloise Hamann, *On power-invariance*, Pacific J. Math. **61** (1975), no. 1, 153–159. MR407009

- [HC02] Thomas J. Haines and Ngô Bao Châu, *Alcoves associated to special fibers of local models*, Amer. J. Math. **124** (2002), no. 6, 1125–1152. MR1939783
- [He16] Xuhua He, *Hecke algebras and p -adic groups*, Current developments in mathematics 2015, Int. Press, Somerville, MA, 2016, pp. 73–135. MR3642544
- [Her09] Florian Herzig, *The weight in a Serre-type conjecture for tame n -dimensional Galois representations*, Duke Math. J. **149** (2009), no. 1, 37–116, DOI 10.1215/00127094-2009-036. MR2541127
- [HH17] Thomas J. Haines and Xuhua He, *Vertexwise criteria for admissibility of alcoves*, Amer. J. Math. **139** (2017), no. 3, 769–784, DOI 10.1353/ajm.2017.0020. MR3650232
- [HLM17] Florian Herzig, Daniel Le, and Stefano Morra, *On mod p local-global compatibility for GL_3 in the ordinary case*, Compos. Math. **153** (2017), no. 11, 2215–2286, DOI 10.1112/S0010437X17007357. MR3705291
- [HW22] Yongquan Hu and Haoran Wang, *On the mod p cohomology for GL_2 : the non-semisimple case*, Camb. J. Math. **10** (2022), no. 2, 261–431. MR4461834
- [Jan03] Jens Carsten Jantzen, *Representations of algebraic groups*, 2nd ed., Mathematical Surveys and Monographs, vol. 107, American Mathematical Society, Providence, RI, 2003. MR2015057
- [LLHL19] Daniel Le, Bao V. Le Hung, and Brandon Levin, *Weight elimination in Serre-type conjectures*, Duke Math. J. **168** (2019), no. 13, 2433–2506, DOI 10.1215/00127094-2019-0015. MR4007598
- [LLHLMa] Daniel Le, Bao Viet Le Hung, Brandon Levin, and Stefano Morra, *Extremal weights and a tameness criterion for mod p Galois representations*, Preprint, [arXiv:2206.06442](https://arxiv.org/abs/2206.06442), 2022.
- [LLHLMb] Daniel Le, Bao Viet Le Hung, Brandon Levin, and Stefano Morra, *Local models for galois deformation rings and applications*, Inv. Math. , to appear.
- [LLHLMc] Daniel Le, Bao Viet Le Hung, Brandon Levin, and Stefano Morra, *Serre weights for three-dimensional wildly ramified Galois representations*, Preprint, [arXiv:2202.03303](https://arxiv.org/abs/2202.03303), 2021.
- [LLHLM18] Daniel Le, Bao V. Le Hung, Brandon Levin, and Stefano Morra, *Potentially crystalline deformation rings and Serre weight conjectures: shapes and shadows*, Invent. Math. **212** (2018), no. 1, 1–107, DOI 10.1007/s00222-017-0762-0. MR3773788
- [LLHLM20] Daniel Le, Bao Viet Le Hung, Brandon Levin, and Stefano Morra, *Serre weights and Breuil’s lattice conjectures in dimension three*, Forum Math. Pi **8** (2020), e5, 135. MR4079756
- [LLHM⁺] Daniel Le, Bao Viet Le Hung, Stefano Morra, Chol Park, and Zicheng Qian, *Moduli of Fontaine–Laffaille representations and a mod- p compatibility result*, Preprint, [arXiv:2109.02720](https://arxiv.org/abs/2109.02720), 2021.
- [LMP18] Daniel Le, Stefano Morra, and Chol Park, *On mod p local-global compatibility for $\mathrm{GL}_3(\mathbb{Q}_p)$ in the non-ordinary case*, Proc. Lond. Math. Soc. (3) **117** (2018), no. 4, 790–848, DOI 10.1112/plms.12148. MR3873135
- [MP17] Stefano Morra and Chol Park, *Serre weights for three-dimensional ordinary Galois representations*, J. Lond. Math. Soc. (2) **96** (2017), no. 2, 394–424, DOI 10.1112/jlms.12064. MR3708956
- [PQ22] Chol Park and Zicheng Qian, *On mod p local-global compatibility for $\mathrm{GL}_n(\mathbb{Q}_p)$ in the ordinary case* (English, with English and French summaries), Mém. Soc. Math. Fr. (N.S.) **173** (2022), vi+150, DOI 10.24033/msmf.48. MR4439201
- [Sta19] The Stacks Project Authors, *Stacks Project*, <https://stacks.math.columbia.edu>, 2019.
- [Wan87] Jian Pan Wang, *Partial orderings on affine Weyl groups* (Chinese, with English summary), J. East China Norm. Univ. Natur. Sci. Ed. **4** (1987), 15–25. MR980127

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY STREET, WEST LAFAYETTE, INDIANA 47907

Email address: `ledt@purdue.edu`

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, 2033 SHERIDAN ROAD, EVANSTON, ILLINOIS 60208

Email address: `lhvietbao@googlemail.com`

UNIVERSITÉ PARIS 8, LABORATOIRE D'ANALYSE, GÉOMÉTRIE ET APPLICATIONS, LAGA, UNIVERSITÉ SORBONNE PARIS NORD, CNRS, UMR 7539, F-93430, VILLETANEUSE, FRANCE

Email address: `morra@math.univ-paris13.fr`

DEPARTMENT OF MATHEMATICAL SCIENCES, ULSAN NATIONAL INSTITUTE OF SCIENCE AND TECHNOLOGY, UNIST-GIL 50, ULSAN 44919, SOUTH KOREA

Email address: `cholpark@unist.ac.kr`

MORNINGSIDE CENTER OF MATHEMATICS, NO.55, ZHONGGUANCUN EAST ROAD, BEIJING 100190, PEOPLE'S REPUBLIC OF CHINA

Email address: `qianzicheng@amss.ac.cn`