

https://doi.org/10.1093/jmammal/gyae084 Advance access publication 16 August 2024 Research Article

## Research Article

# The allometry of brain size in Euarchontoglires: clade-specific patterns and their impact on encephalization quotients

Sergi López-Torres<sup>1,2,\*</sup>, Ornella C. Bertrand<sup>3,4</sup>, Łucja Fostowicz-Frelik<sup>5,6,7</sup>, Madlen M. Lang<sup>8</sup>, Chris J. Law<sup>2,9,10</sup>, Gabriela San Martin-Flores<sup>8</sup>, Michael A. Schillaci<sup>8</sup>, Mary T. Silcox<sup>8</sup>,

Associate Editor was Deyan Ge

#### Abstract

The timing and nature of evolutionary shifts in the relative brain size of Primates have been extensively studied. Less is known, however, about the scaling of the brain-to-body size in their closest living relatives, i.e., among other members of Euarchontoglires (Dermoptera, Scandentia, Lagomorpha, Rodentia). Ordinary least squares (OLS), reduced major axis (RMA), and phylogenetic generalized least squares (PGLS) regressions were fitted to the largest euarchontogliran data set of brain and body mass, comprising 715 species. Contrary to previous inferences, lagomorph brain sizes (PGLS slope = 0.465; OLS slope = 0.593) scale relative to body mass similarly to rodents (PGLS = 0.526; OLS = 0.638), and differently than primates (PGLS = 0.607; OLS = 0.794). There is a shift in the pattern of the scaling of the brain in Primates, with Strepsirrhini occupying an intermediate stage similar to Scandentia but different from Rodentia and Lagomorpha, while Haplorhini differ from all other groups in the OLS and RMA analyses. The unique brain-body scaling relationship of Primates among Euarchontoglires illustrates the need for clade-specific metrics for relative brain size (i.e., encephalization quotients; EQs) for more restricted taxonomic entities than Mammalia. We created clade-specific regular and phylogenetically adjusted EQ equations at superordinal, ordinal, and subordinal levels. When using fossils as test cases, our results show that generalized mammalian equations underestimate the encephalization of the stem lagomorph *Megalagus turgidus* in the context of lagomorphs, overestimate the encephalization of the stem primate *Microsyops annectens* and the early euprimate *Necrolemur antiquus*, but provide similar EQ values as our new strepsirrhine-specific EQ when applied to the early euprimate *Nadapis parisiensis*.

Key words: biological scaling, body size, brain size, comparative anatomy, endocasts, neurobiology.

One of the defining characteristics of modern primates is their relatively large brain size compared to other mammalian orders (Martin 1990). Primates have historically received significant attention with respect to the question of how their brain size scales in relation to body size (e.g., Count 1947; Gould 1975; Lande 1979; Clutton-Brock and Harvey 1980; Bronson 1981; Shea 1983; Armstrong 1985; Martin and Harvey 1985; Marino 1998; Sherwood et al. 2008; Boddy et al. 2012; Smaers et al. 2012, 2021; Holekamp et al. 2013; Burger et al. 2019). Within the order Primates, hominoids (Hominoidea) are

particularly encephalized, both in the larger context of Mammalia and compared to the much less encephalized and smaller bodied strepsirrhines (Strepsirrhini). This association between body size and degree of encephalization leads to Primates having the highest slope value in ordinary least square (OLS) regression analyses of brain mass against body mass among any of the mammalian orders (Burger et al. 2019). However, to fully understand evolutionary trends that define the primate brain, it is necessary to place the scaling of the primate brain within a phylogenetic context.

<sup>&</sup>lt;sup>1</sup>University of Warsaw, Faculty of Biology, Biological and Chemical Research Centre, Institute of Evolutionary Biology, Żwirki i Wigury 101, 02-089 Warsaw, Poland

Division of Paleontology, American Museum of Natural History, 79th Street and Central Park West, New York, NY 10024-5192, United States

<sup>&</sup>lt;sup>3</sup>Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, Edifici ICTA-ICP, c/ Columnes s/n, Campus de la UAB, 08193 Cerdanyola del Vallès, Spain

<sup>&</sup>lt;sup>4</sup>School of Geosciences, The University of Edinburgh, Grant Institute, Edinburgh EH9 3FE, United Kingdom

Department of Organismal Biology and Anatomy, The University of Chicago, 1027 East 57th Street, Chicago, IL 60637, United States

<sup>&</sup>lt;sup>6</sup>Key Laboratory of Evolutionary Systematics of Vertebrates, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, 142 Xi-Zhi-Men-Wai Street, Beijing 100044, People's Republic of China

<sup>&</sup>lt;sup>7</sup>Institute of Paleobiology, Polish Academy of Sciences, Twarda 51/55, 00-818 Warsaw, Poland

<sup>\*</sup>Department of Anthropology, University of Toronto Scarborough, 1265 Military Trail, Toronto, ON M1C 1A4, Canada

Department of Biology and Burke Museum of Natural History and Culture, University of Washington, Seattle, WA 98105, United States

<sup>&</sup>lt;sup>10</sup>Department of Integrative Biology, University of Texas, 2415 Speedway #C0930, Austin, TX 78712, United States

<sup>&#</sup>x27;Corresponding author: University of Warsaw, Faculty of Biology, Biological and Chemical Research Centre, Institute of Evolutionary Biology, Żwirki i Wigury 101, 02-089 Warsaw, Poland. Email: s.lopez-torres@uw.edu.pl

This exercise is 2-fold: on the one hand, it is important to explore how brain-to-body scaling relationships differ at a subordinal level within Primates, while on the other hand, it is important to put primate data into the larger context of their close, non-primate relatives.

While there is a long history of the study of relative brain size in primates, the consensus around the broader phylogenetic context within which these data should be assessed has changed relatively recently (Murphy et al. 2001). Literature from only a few decades ago treated "insectivorans" as a good model for primitive states for the primate brain—this was true, for example, with respect to the classic and oft-reanalyzed compilation of quantitative data on primate brains by Stephan et al. (1970, 1981). However, modern phylogenetic studies (e.g., Foley et al. 2023) position eulipotyphlan "insectivorans" as relatively distant to primates. Instead, Primates are broadly considered to be members of the superorder Euarchonta (together with the orders Dermoptera and Scandentia; Waddell et al. 1999), and Euarchonta is understood to be most closely related to the superorder Glires (Rodentia + Lagomorpha) within the greater grouping of Euarchontoglires (Murphy et al. 2001; Foley et al. 2023).

Recent molecular analyses have supported both Dermoptera (Mason et al. 2016; Zhang et al. 2019; Foley et al. 2023) and Sundatheria (Dermoptera + Scandentia; Olson et al. 2005; Upham et al. 2019) as the sister group of Primates. However, dermopterans are only known from 2 species, which makes it problematic to delve into the relationship between brain mass and body mass at an ordinal level. In the case of Scandentia, the present paper is the first detailed consideration of allometric relationships between brain and body mass for this group. Smaers et al. (2021) inferred a value for Scandentia from an ancestral state reconstruction analysis, but that was based on only 5 taxa, all of them tupaiines. Relative brain size has been more intensively studied in rodents (e.g., Count 1947; Mace et al. 1981; Mace and Eisenberg 1982; Pilleri et al. 1984; Hafner and Hafner 1984; Towe and Mann 1992; Bernard and Nurton 1993; Mann and Towe 2003; Matějů et al. 2016; Burger et al. 2019; Bertrand et al. 2021), but much less for lagomorphs. Burger et al. (2019) reported that the regression slope values for rodent and lagomorph regressions of brain mass on body mass were notably different. In their analysis, rodents had a slope value of 0.64, but lagomorphs were reported to have a slope value of 0.75 (Burger et al. 2019). This result implies that the brains in lagomorphs would scale more similarly to primates (slope value of 0.79; Burger et al. 2019) than they would to rodents, which would be surprising for a group that is not generally thought of as having comparatively encephalized larger

The calculation of allometric regressions describing brain scaling for the various euarchontogliran groups is also of particular relevance for the calculation of encephalization quotients (EQs; Jerison 1973; Supplementary Data SD1). Although sometimes criticized (e.g., Deacon 1990; Begun and Kordos 2004; Schoenemann 2006; Deaner et al. 2007; Gilbert and Jungers 2017), the calculation of metrics related to EQs for fossil taxa remains a critically important part of assessing brain size evolution through time in Primates and other orders (e.g., Boddy et al. 2012; Ni et al. 2019; Bertrand et al. 2022; Silcox et al. 2023). To date, the most commonly used equations for calculation of EQ in fossil Euarchontoglires are based on general mammalian samples (i.e., Jerison 1973; Eisenberg 1981). However, some authors have taken the approach of studying relative brain size evolution using EQs calculated from more restricted samples, which might provide a better framework for studying variation and evolution of relative brain size within the context of a particular group. For example, Pilleri et al. (1984) developed a rodent-specific equation that has recently been used for the calculation of EQs in many fossil rodent studies (Bertrand and Silcox 2016; Bertrand et al. 2016, 2017, 2018, 2019a). More recently, Grabowski et al. (2016) and Ni et al. (2019) developed primate-specific equations based on PGLS regressions of endocranial volume versus body mass, and Ni et al. (2019) coined the term "phylogenetic encephalization quotient" (PEQ). Further, Bertrand et al. (2021) developed a PEQ equation for their own sample of sciuroid rodents and extinct closest relatives, Püschel et al. (2021) also used their own equation for PEQ using extant and extinct hominoids, and Bertrand et al. (2022) conceived their own equation of PEQ for a sample of extinct placental mammals.

In the present study we aim to: (1) analyze the allometry of brain mass to body mass across Euarchontoglires and at lower taxonomic levels; (2) examine the patterns in relative brain size variation among and within major evolutionary lineages; and (3) generate clade-specific EQ equations. The goal of generating clade-specific EQ equations is to minimize error and allow authors to tailor their research questions to narrower taxonomic frameworks. The current work therefore considers patterns of allometric scaling in all euarchontogliran orders and probes the question of how best to use that information to study the evolution of relative brain size through

# Materials and methods. The studied sample.

We acquired estimates of brain size and body size for 715 species of Euarchontoglires (Supplementary Data SD2, SD4). In our analyses, we used data from previously published literature (Bertrand and Silcox 2016; Bertrand et al. 2017, 2018, 2019a, 2021; Burger et al. 2019; López-Torres et al. 2020; Smaers et al. 2021; Lang et al. 2022). In all cases, when data originated from endocranial volumes (cm3), they were converted to brain mass (g) by dividing the endocranial volume by 1.036 (Stephan et al. 1981). For details about original sources of data, see Supplementary Data SD2. In the process of combining data sets, we were careful to avoid duplicating entries. In some cases, new data for particular species were combined with data from previous literature to generate new average values. For rodents, we combined the data sets from Burger et al. (2019), Bertrand and Silcox (2016), Bertrand et al. (2018, 2021), and Lang et al. (2022). For lagomorphs, we combined the data sets from Burger et al. (2019), López-Torres et al. (2020), and Smaers et al. (2021). For primates, we combined Burger et al.'s (2019) and Lang et al.'s (2022) data sets. For scandentians, we combined Burger et al.'s (2019) and San Martin-Flores et al.'s (2018) data sets. The new scandentian sample (n = 14) allows us to generate the first regression lines for brain versus body mass for this order of mammals. The brain size data for dermopterans were obtained from Lang et al. (2022) and include both extant species; the body mass data were taken from Stafford and Szalay (2000). Although it is fundamentally uninformative to generate a regression line from only 2 points, the dermopteran species were still included in the calculations for the regression lines for Euarchonta and Euarchontoglires.

Following the total number of species in different mammalian orders by Burgin et al. (2018), the present study samples 100% of currently recognized dermopteran species (2/2), 58% of currently recognized scandentian species (14/24), 47.88% of currently recognized primate species (248/518), 31.63% of currently recognized lagomorph species (31/98), and 16.46% of currently recognized rodent species (420/2,552). Only extant taxa were included in this analysis. The main reason for the exclusion of extinct taxa is that uncertainty in body mass estimates for fossils would lead to very large amounts of error around the calculated regression lines, making it difficult to interpret any differences as being a product of relative brain size versus uncertainty in the underlying data. Also, because there is a temporal effect in brain size in several mammalian orders (Jerison 1973; Silcox et al. 2010; Orliac and Gilissen 2012; Yao et al. 2012; Bertrand et al. 2019a, 2022), simultaneously analyzing fossils with extant taxa has the potential of conflating temporal patterns with scaling relationships that exist within particular groups at a given moment in evolutionary time. However, it is a fair concern that these temporal shifts could make the scaling relationships calculated here an inaccurate representation of those that would have applied at other points in the evolution of euarchontoglirans. Investigating these complexities is beyond the scope of the current analysis.

No research was conducted on live animals.

## **Analysis**

The scaling of brain size with body size has typically been characterized by a power law (Snell 1892; Dubois 1898; Jerison 1973),

$$(Brain\ mass) = \alpha (Body\ mass)^{\beta}$$

and  $\alpha$  and  $\beta$  are constants representing the intercept and slope, respectively. This relationship becomes linear by log transforming both sides of the equation:

$$log(Brain \ mass) = log(\alpha) + \beta \cdot log(Body \ mass)$$

OLS, reduced major axis (RMA), and phylogenetic generalized least squares (PGLS) regressions were fitted to our data. We used all 3 approaches to allow for comparison to previous analyses and to enable the calculation of both regular EQs and PEQs. Bootstrapped estimates of OLS and RMA regression slopes and intercepts were generated for each taxonomic grouping based on 10,000 random resampling iterations using the Statistics101 software package (http://www.statistics101.net/). The PGLS analysis was conducted by incorporating an extensively sampled, time-scaled mammalian phylogeny (Upham et al. 2019); of the total of 715 taxa, 696 taxa could be included based on their presence in the Upham et al. (2019) analysis. All regression parameters were simultaneously estimated with phylogenetic signal in the residual error as Pagel's lambda (Pagel 1999; Revell 2010) using the phylolm function in the R package "phylolm" (Tung Ho and Ané 2014). Regression lines of the 3 types were calculated at multiple taxonomic levels: supraordinal (Euarchonta, Glires, and Euarchontoglires), ordinal (Primates, Scandentia, Rodentia, and Lagomorpha), and subordinal (Strepsirrhini and Haplorhini).

## Calculation of significance.

P-values are strongly dependent on sample size and relate to an arbitrarily chosen alpha level. In this study, rather than making determinations of statistical significance using standard P-values consistent with traditional hypothesis testing, we chose to estimate what the difference in slope and intercept values is likely to be between taxonomic groups (see discussion in Smith 2020). To do this we used random resampling with replacement to generate resampling distributions of the difference between taxonomic groupings in slope and intercept values, also based on 10,000 resampling iterations. We used these resampling distributions to generate 95% confidence intervals (CIs), which is the interval that comprises 95% of the resampled differences between the 2 taxonomic groupings compared. Standard deviations and 95% CIs for the slopes, intercepts, and differences between taxonomic groupings were generated from the resampling distributions. We use standard deviation of the resampled distribution as the standard error of the bootstrap estimate (i.e., slope, intercept, or difference).

**Table 1.** Formulae to calculate the expected brain mass (E<sub>c</sub>) for a given body mass (BM) for specific taxonomic groups. The EQ and PEQ are calculated by dividing the actual brain mass by the E given in this table. The equations are formulated the following way: 10intercept \* (body mass)slope.

	E <sub>c</sub> for EQ	E <sub>c</sub> for PEQ
Euarchontoglires	0.045 * BM <sup>0.809</sup>	0.214 * BM <sup>0.540</sup>
Euarchonta	0.067 * BM <sup>0.805</sup>	0.189 * BM <sup>0.603</sup>
Glires	0.090 * BM <sup>0.635</sup>	0.168 * BM <sup>0.524</sup>
Rodentia	See Pilleri et al. (1984)	0.166 * BM <sup>0.526</sup>
Lagomorpha	0.117 * BM <sup>0.594</sup>	0.262 * BM <sup>0.465</sup>
Scandentia	0.151 * BM <sup>0.604</sup>	0.167 * BM <sup>0.580</sup>
Primates	0.074 * BM <sup>0.794</sup>	0.247 * BM <sup>0.607</sup>
Strepsirrhini	0.117 * BM <sup>0.679</sup>	0.206 * BM <sup>0.614</sup>
Haplorhini	0.146 * BM <sup>0.723</sup>	0.341 * BM <sup>0.578</sup>

#### New EQ equations.

In this study, we propose new clade-specific EQs (Table 1), following the model suggested by Pilleri et al. (1984) specifically for rodents, which has been previously used in studies focusing on fossil rodent endocasts (Bertrand et al. 2016; Bertrand and Silcox 2016, 2017). These equations are based on more constrained samples, instead of general mammalian samples as in Jerison (1973) or Eisenberg (1981). The new equations are derived from the OLS and the PGLS regression equations. Additionally, in this study we are using a few fossil specimens as test cases for the new EQs. In particular, we use the stem lagomorph Megalagus turgidus (FMNH UC 1642; López-Torres et al. 2020), the stem primate Microsyops annectens (UW 14559; Silcox et al. 2010), the adapoid primate Adapis parisiensis (NHM M 1345; Harrington et al. 2016), and the omomyoid primate Necrolemur antiquus (MaPhQ 289; Harrington et al. 2020) to allow us to consider the value of taxonomically constrained EQ equations. Brain mass and body mass data for these species are provided in Table 2.

## Institutional abbreviations.

FMNH UC-University of Chicago collection, Field Museum of Natural History, Chicago, Illinois, United States; MaPhQ—Muséum d'Histoire Naturelle Victor Brun, Montauban, France; NHM—Natural History Museum, London, United Kingdom; UW-University of Wyoming, Laramie, Wyoming, United States.

#### Results

Results for the OLS analysis are shown in Tables 3 and 4; the results for the PGLS analysis are shown in Tables 5 and 6; and the results for the RMA analysis are shown in the Supplementary Data SD3. The calculated regression lines are shown at the supraordinal level (Euarchonta, Glires, and Euarchontoglires; Fig. 1), ordinal level (Primates, Scandentia, Rodentia, and Lagomorpha; Fig. 2), and subordinal level (Strepsirrhini and Haplorhini; Fig. 3). Figure 4 combines different regression lines in 1 graph to better visualize the comparison between slopes and intercepts.

#### OLS analysis.

Our results show that the brains of rodents, lagomorphs, and scandentians scale similarly with respect to body mass (i.e., have similar slopes; 0.638, 0.594, and 0.604, respectively; Fig. 2B-D). The 95% CIs for each of the comparisons among these 3 taxonomic groupings included zero. The bootstrap estimates of the differences in slope for the 3 comparisons ranged from -0.034 to 0.044 (Table 4). Our

Table 2. Brain mass and body mass data (in grams) for selected taxa of fossil Euarchontoglires.

Order	Family	Species	Specimen number	Brain mass	Body mass	Source
Lagomorpha	Megalagidae	Megalagus turgidus	FMNH UC 1642	6.72	2,325.01	López-Torres et al. (2020)
Primates	Microsyopidae	Microsyops annectens	UW 12362	5.62	1,358 to 2,568	Silcox et al. (2010)
Primates	Adapidae	Adapis parisiensis	NHM M1345	8.39	1,103	Harrington et al. (2016)
Primates	Microchoeridae	Necrolemur antiquus	MaPhQ 289	2.36	144	Harrington et al. (2020)

reanalysis of the Burger et al. (2019) data set (i.e., excluding the new data used in the present analysis) failed to recover their results with respect to lagomorphs (i.e., a slope of 0.75), suggesting that the appearance of a difference is based on an error in their analysis of their data.

When assessed at an ordinal level, the brains of primates (slope = 0.794; Fig. 2A) do show evidence of scaling differently from other euarchontogliran orders (Fig. 4A). None of the 95% CIs for any of the comparisons with Primates included zero. This same pattern was observed for Haplorhini, which exhibit a high slope value (slope = 0.723; Fig. 3A). Interestingly, the 95% CIs for comparisons of Strepsirrhini with both Rodentia and Lagomorpha excluded zero, while the comparisons with Haplorhini and Scandentia included zero. This pattern of results suggests that while the slope value for haplorhines is likely higher than other members of Euarchontoglires, strepsirrhines (slope = 0.679) occupy an intermediate scaling pattern between non-primate Euarchontoglires and haplorhines. These findings were largely mirrored by the results of RMA regression (Supplementary Data SD3), with the exception of the Strepsirrhini-Rodentia comparison.

The intercepts for these various regressions (Table 3) show that hypothetically small haplorhines would have larger brains than similarly small strepsirrhines (Fig. 3). However, the overall primate regression intersects with the y axis at a lower value (-1.133) than the haplorhine (-0.834) and the strepsirrhine (-0.932) regressions do—a byproduct of having large-brained, large-bodied anthropoids in the primate sample, which tilt the regression line in a way that increases the slope and decreases the intercept when strepsirrhines and haplorrhines are combined (Fig. 2A). Small rodents would also be expected to have smaller brains than small strepsirrhines and small haplorhines, based on the estimate of the rodent intercept (-1.050; Fig. 2C). But when looking at the general primate regression, small rodents would have larger brains than small primates (Fig. 4A)—again, is a byproduct of the high slope of the overall primate regression. The highest intercept is observed in scandentians (-0.821), with the hypothetical small scandentians having larger brains than even small haplorhines. This high intercept is probably due to the fact that scandentians have both a shallow slope (0.604; similar to the lagomorph slope, 0.594) and that they have slightly larger brains than lagomorphs of similar body mass (Fig. 4A).

#### PGLS analysis.

The PGLS analysis showed a strong phylogenetic signal (i.e.,  $\lambda \ge$ 0.875 for all groupings). The main difference between the values calculated in the OLS and PGLS analyses is that PGLS gives, in the case of this study, systematically lower slope values and higher intercept values than OLS (Table 5). This contrast is particularly pronounced in primates and lagomorphs, because in these 2 groups there is a strong clade-specific pattern in body mass—for example, among haplorhines, the different major clades (i.e., tarsioids, platyrrhines, cercopithecoids, hominoids) are each fairly constrained in their size range, with only a small degree of overlap between these groups. Because this pattern is also related to relative brain mass, controlling for this pattern has the effect of substantially decreasing the slope in the regression lines generated by PGLS.

Only the comparisons between Euarchonta and Glires, Primates and Glires, and Primates and Lagomorpha excluded zero in their 95% CIs. Within primates, the slope of the regression line analyzed ordinally (0.607) is intermediate in value between the strespirrhine (0.614) and haplorrhine (0.578) slopes. Scandentians are more similar to primates in having a relatively higher slope value (0.580) than rodents (0.526) or lagomorphs (0.465; Fig. 4A and B). In comparing the PGLS and OLS slope values for the various orders the greatest similarity is observed for Scandentia (0.580 vs. 0.604, respectively), likely as a consequence of the fact that there is little phylogenetic effect on body mass in Scandentia (Fig. 2). Both the largest and smallest species of scandentians are tupaiines (Tupaia everetti, 249.5 g; Dendrogale murina, 45 g; Sargis 2002). The slope for Glires (0.524) is intermediate between the rodent (0.526) and lagomorph (0.465) slopes, although it is very similar to the rodents. It is likely that the slope for Glires is so similar to rodents because the sample of Glires is mostly composed of rodents (421 rodents vs. 31 lagomorphs). Also, the range of body mass of lagomorphs is contained within the range of body mass for rodents (Fig. 1C), so it is not expected that lagomorphs would shift the numbers substantially, as they might have if lagomorphs were clustered at one extreme or the other. The euarchontan slope (0.603) is also intermediate between the primate (0.607) and the scandentian (0.580) slopes but very close in value to the primate slope. That is a similar situation to the one seen with Glires; the euarchontan sample is largely made up of primates (248 primates, 14 scandentians, 2 dermopterans), and the scandentian and dermopteran ranges of body mass are within that of strepsirrhines (Fig. 1B). Finally, the slope for Euarchontoglires (0.540) has a value similar to those calculated for non-primate Euarchontoglires.

As in OLS, the intercepts of the PGLS regressions show that hypothetically small haplorhines would have larger brains than small strepsirrhines. However, in the PGLS analysis, the overall primate regression intersects with the y axis between the haplorhine and the strepsirrhine intercepts. Contrary to the finding in the OLS analysis, the highest intercept in PGLS, at an ordinal level, is observed for lagomorphs instead of scandentians (Fig. 4B).

Another important difference between the OLS and PGLS regressions is that in the OLS analysis there are more groups that could be considered different based on the 95% CI values in terms of scaling patterns, but PGLS fails to find most of these differences. In pairwise comparisons PGLS only finds clear differences in scaling between Euarchonta and Glires, between Primates and Glires, and between Primates and Rodentia (Table 6).

#### **Discussion**

Our results differ from the conclusion of Burger et al. (2019) that the lagomorph brain scales more similarly to primates—instead our results indicate that lagomorphs show a scaling relationship closer to that seen in rodents with respect to body mass (Fig. 4A and B).

Downloaded from https://academic.oup.com/jmammal/article/105/6/1430/7734701 by University of California Library - Berkeley Library user on 07 July 2025

Allometries for log10 brain size versus log10 body size across different taxonomic levels within Euarchontoglires based on bootstrap estimates of OLS regression parameters. n = number of sampled species; r = correlation coefficient; SD = standard deviation

Taxon	и	۲	Slope	SD	ID %56		Intercept	SD	12 % GI	
					Lower end	Upper end			Lower end	Upper end
Euarchontoglires	715	0.959	0.809	600.0	0.791	0.826	-1.344	0.021	-1.386	-1.303
Euarchonta	264	0.963	0.805	0.011	0.783	0.827	-1.177	0.037	-1.251	-1.105
Glires	451	0.962	0.635	0.008	0.619	0.652	-1.044	0.019	-1.082	-1.008
Rodentia	420	0.959	0.638	0.010	0.619	0.657	-1.050	0.021	-1.089	-1.008
Lagomorpha	31	0.955	0.594	0.037	0.517	0.668	-0.931	0.117	-1.160	-0.693
Scandentia	14	0.964	0.604	0.048	0.502	0.692	-0.821	660.0	-1.004	-0.608
Primates	248	0.959	0.794	0.014	0.769	0.822	-1.133	0.047	-1.228	-1.044
Strepsirrhini	63	0.958	0.679	0.019	0.645	0.719	-0.932	0.053	-1.045	-0.837
Haplorhini	185	0.954	0.723	0.018	0.687	0.759	-0.834	0.066	-0.961	0.702

This correction highlights the uniqueness of primates in a euarchontogliran context. This is further supported in the OLS analysis, where primates are the only order of Euarchontoglires with the 95% CIs for differences in slope that do not include zero.

Strepsirrhines have similar OLS slopes to both haplorhines and scandentians, differing from those of rodents and lagomorphs. However, the scandentian slope differs from haplorhines, making strepsirrhines an apparently intermediate stage between other primates and non-primate euarchontoglirans. On the other hand, the CIs for the slope in the PGLS analysis overlap among many more ordinal groupings. However, primates stand out even in the PGLS analysis for being the only order with a CI for their slope that does not overlap with rodents. Consequently, the fact that primates appear to have a unique brain-body scaling relationship in the context of Euarchontoglires (supported by both the OLS and the PGLS analyses) further illustrates the necessity for clade-specific regular EQs and phylogenetic EQs for more restricted taxonomic entities than Mammalia.

A recent analysis (Smaers et al. 2021) examined allometric scaling relationships across Mammalia, identifying particular points in the mammalian tree where they inferred grade shifts had occurred. In their analysis, several groups analyzed here were reconstructed as part of the ancestral mammalian grade (Scandentia, Lagomorpha, Dermoptera, squirrel-related clade, Tarsioidea) based on their allometric scaling relationships not being found to have differed from the relationship calculated for the common mammalian ancestor, with a primitive PGLS slope of 0.51. However, this slope lies outside the 95% CI for Euarchontoglires, Euarchonta, Primates, and Strepsirrhini calculated here (Table 5). As such, our analysis would suggest that a grade shift might have happened earlier in the evolution of this clade than their analysis suggests. In particular, the contrast between Primates and Glires found in all analyses suggests an ordinal level shift for Primates, contrasting with inferred changes occurring only within primate subgroups in the result of Smaers et al. (2021). Possible explanations for this contrast are the notably stronger sampling of both Scandentia and Rodentia here, which allowed for a refinement of previous estimates, but also the direct inclusion of fossils in Smaers et al. (2021). However, these inferences require testing using fossil specimens that allow for a more direct window into such grade shifts.

## Encephalization quotients.

The EQ is a widely used index of brain size scaled to body size (Jerison 1985). As noted above, some of the most commonly used EQ equations in the field of paleoprimatology are Jerison's (1973) and Eisenberg's (1981), even though they are based on generalized mammalian samples. However, the use of EQs has been criticized for poorly modeling brain scaling relationships in fossil primates (Gilbert and Jungers 2017), as well as for not being a good predictor of cognitive abilities (Deaner et al. 2007). Although imperfect, no alternative has been suggested to EQs for comparing brain sizes in animals of different body masses that is not also problematic. For example, taking a narrow allometric approach (as suggested by Gilbert and Jungers 2017) is very prone to error being introduced by inconsistencies in body mass estimation, and does not offer a clear solution for making comparisons to fossil taxa outside the modern range of variation (White et al. 2023). While we agree with Smaers et al. (2021) that making comparisons between particular brain regions offers a much richer understanding of brain size evolution than looking at relative overall size (see, for example, Bertrand et al. 2019b, 2021), such an approach is limited in dealing with fossils because only certain brain regions can be isolated on endocasts. We would also argue that, in light of the high physiological cost of

Table 4. Pairwise comparisons of bootstrap estimates of differences among Euarchontoglires in OLS regression parameters. SD = standard deviation. Asterisks = likely to be different.

Taxon	Slope	SD	95% CI		Intercept	SD	95% CI	
	bootstrap difference		Lower end	Upper end	bootstrap difference		Lower end	Upper end
Euarchonta-Glires	0.169*	0.042	0.143	0.197	-0.133*	0.042	-0.215	-0.051
Rodentia–Lagomorpha	0.044	0.039	-0.031	0.123	-0.119	0.120	-0.364	0.114
Rodentia-Scandentia	-0.034	0.102	-0.138	0.055	0.228*	0.102	0.044	0.445
Lagomorpha–Scandentia	0.010	0.153	-0.111	0.126	0.110	0.153	-0.187	0.416
Primates-Glires	0.158*	0.050	0.128	0.191	-0.088	0.050	-0.190	0.007
Primates-Rodentia	0.156*	0.050	0.125	0.189	-0.084	0.050	-0.185	0.011
Primates–Lagomorpha	0.200*	0.128	0.123	0.283	-0.204	0.128	-0.468	0.042
Primates–Scandentia	0.190*	0.108	0.103	0.294	-0.313*	0.108	-0.542	-0.114
Strepsirrhini-Haplorhini	0.044	0.085	-0.010	0.094	0.097	0.085	-0.060	0.272
Strepsirrhini-Scandentia	0.074	0.113	-0.018	0.182	-0.111	0.113	-0.348	0.093
Strepsirrhini-Rodentia	0.040*	0.056	0.002	0.083	0.117	0.056	-0.002	0.218
Strepsirrhini–Lagomorpha	0.085*	0.130	0.003	0.170	-0.002	0.130	-0.267	0.246
Haplorhini–Scandentia	0.112*	0.138	0.010	0.248	0.003	0.138	-0.302	0.246
Haplorhini–Rodentia	0.085*	0.069	0.042	0.125	0.215	0.069	0.083	0.356
Haplorhini–Lagomorpha	0.130*	0.137	0.048	0.216	0.093	0.137	-0.184	0.357

maintaining brain tissue, considering relative brain size has fundamental merit in discussions of brain evolution, whether or not it is an accurate proxy for measures of cognition. As such, EQs continue to serve as useful tools, and merit further work to overcome identified problems. The growing availability of brain and body mass data for Euarchontoglires allows us to elaborate subordinal, ordinal, and supraordinal EQ equations, which may allow for more meaningful comparisons relevant to some evolutionary questions than are possible with general mammalian equations, and may solve some of the problems that have been identified with EQs such as a lack of independence from body mass (Begun and Kordos 2004; Gilbert and Jungers 2017). Our results show that in the OLS regression analyses the slopes generally increase and the intercepts decrease the higher we go in taxonomic level within Euarchontoglires (see also Martin 1990). Therefore, in order to avoid overestimation and underestimation of EQs in animals with extreme body masses, it is preferable to use the most taxonomically specific EQ possible. In fact, Pilleri et al. (1984) already elaborated an OLS-based EQ equation specific to rodents that has been used in recent studies on the brain evolution of rodents (Bertrand and Silcox 2016; Bertrand et al. 2016, 2017, 2018, 2019a). Other groups within Euarchontoglires are worth exploring with respect to this approach.

López-Torres et al. (2020) described the first virtual endocast of a fossil lagomorph and provided EQs for both extant lagomorphs and a specimen of M. turgidus (FMNH UC 1642), a stem lagomorph. The body mass of M. turgidus was calculated by López-Torres et al. (2020) using the width of the occipital condyles, which is argued to be one of the most reliable measures of body mass estimation in lagomorphs ( $r^2 = 0.957$ ; Moncunill-Solé et al. 2015). Because of the lack of a lagomorph-specific EQ, the authors used Jerison's (1973) and Eisenberg's (1981) equations. The present OLS-based lagomorph-specific equation (Table 1) provides a higher EQ estimate (0.53) than those calculated with generalized mammalian equations (Jerison's EQ = 0.32; Eisenberg's EQ = 0.40). This equation also provides higher estimates for living lagomorphs than previously calculated (i.e., López-Torres et al. 2020). Here, the critical point is that calculating the group-specific value makes it clear that in the

lagomorph context Megalagus was more encephalized than previously thought (Supplementary Data SD8), with a brain only approximately half the size expected rather than approximately one-third. Our Glires-specific EQ for Megalagus gives a very similar result (0.55) to the lagomorph-specific EQ (Supplementary Data SD10). However, the Euarchontoglires-specific EQ for Megalagus drops to 0.29, even lower than the generalized mammalian EQs of Jerison (1973) and Eisenberg (1981). This is most certainly due to the effect of adding primates, and in particular highly encephalized anthropoids (Supplementary Data SD11). These differing values highlight the importance of making a choice of EQ equation that is relevant to the evolutionary question being asked, and that EQ calculations need to be put in a specific taxonomic framework.

With respect to primates, Grabowski et al. (2016) pioneered using PGLS regressions to calculate EQs instead of using the traditional OLS. They developed a primate-specific EQ, although based on endocranial volume rather than brain mass. Recently, Ni et al. (2019) expanded upon analyses of Grabowski et al. (2016) by publishing EQ equations specific for anthropoids, platyrrhines, and catarrhines based on PGLS regressions (i.e., PEQ; Ni et al. 2019). Here we complement these equations with primate-, strepsirrhine-, and haplorhine-specific equations based on OLS (regular EQ) and PGLS (phylogenetic EQ or PEQ) using brain mass (see Table 1 for the specific EQ and PEQ equations). It is worth noting that EQs and PEQs are not directly comparable, so future studies may benefit from calculating both. For example, whereas an EQ = 1 always means that the brain size of an animal is exactly the brain size expected for an animal of its size, this is not necessarily true when PEQ = 1 because the PGLS analysis corrects the position of the regression line based on phylogenetic effect (if there is one). As such, PGLS estimates have the benefit of including a metric (EQ) that allows for some degree of comparability with past analyses, while also providing information within the context of a particular analysis that reflects our understanding of phylogeny (PEQ).

Here we use 3 early primates (a plesiadapiform, an adapoid, and an omomyoid) as test cases for our new primate equations. Microsyops annectens is a microsyopid plesiadapiform from the

**Table 5.** Allometries for  $\log_{10}$  brain size versus  $\log_{10}$  body size across different taxonomic levels within Euarchontoglires based on bootstrap estimates of PGLS regression parameters. n = n

Taxon	и	$\mathbb{R}^2$	~	Slope	SD	95% CI		Intercept	SD	95% CI	
						Lower end	Upper end			Lower end	Upper end
Euarchontoglires	969	0.766	0.898	0.540	0.011	0.518	0.561	-0.670	0.067	-0.811	-0.540
Euarchonta	254	0.779	0.882	0.603	0.022	0.561	0.644	-0.724	960.0	-0.911	-0.538
Glires	442	0.779	0.882	0.524	0.013	0.499	0.549	-0.774	0.088	-0.949	-0.611
Rodentia	413	0.784	0.875	0.526	0.013	0.501	0.550	-0.779	0.097	-0.967	-0.593
Lagomorpha	29	0.784	0.875	0.465	0.084	0.295	0.628	-0.581	0.272	-1.092	-0.083
Scandentia	14	0.784	0.875	0.580	0.114	0.364	0.803	-0.778	0.250	-1.262	-0.313
Primates	238	0.784	0.875	0.607	0.022	0.566	0.650	-0.607	0.114	-0.833	-0.385
Strepsirrhini	09	0.725	0.915	0.614	0.044	0.527	0.701	-0.687	0.193	-1.053	-0.316
Haplorhini	178	0.725	0.915	0.578	0.029	0.518	0.635	-0.468	0.162	-0.793	-0.154

middle Eocene of North America known from a substantively complete cranium (UW 12362; Silcox et al. 2010, 2020) with a range of EQs between 0.25 and 0.38 using Jerison's (1973) equation, and between 0.31 and 0.49 using Eisenberg's (1981) equation. The range of EQs is due to the fact that Silcox et al. (2010) calculated several body mass estimates, including a cranial length insectivoran equation (Thewissen and Gingerich 1989), a cranial length horizontal primate PGLS equation (Silcox et al. 2009), a cranial length generic primate equation (Martin 1990), and an upper molar area equation (Gingerich et al. 1982). Using our OLS-based primatespecific equation, Microsyops yields a lower range of EQ estimates (0.15 to 0.25) than those resulting from Jerison's (1973) and Eisenberg's (1981) equations. The euarchontan-specific equation gives the same EQ values as the primate-specific equation. The Euarchontoglires-specific equation provides more similar values to those of Jerison's (1973) equation, with EQs in the range of 0.22 to 0.37. These results accentuate the fact that M. annectens had an extremely small brain for a primate (Supplementary Data SD7, SD9) but was quite similar to the stem lagomorph M. turgidus when the EQs of both are calculated using a Euarchontoglires-specific equation (Supplementary Data SD11).

Adapis parisiensis is an adapid adapoid from the late Eocene of Europe. Jerison's (1973) equation provides an EQ of 0.65 and Eisenberg's (1981) an EQ of 0.86 based on a very complete cranium (NHM M1345; Harrington et al. 2016). Harrington et al. (2016) calculated the body mass of A. parisiensis using the euarchontan ectal facet equation (Yapuncich et al. 2015). The relationships of adapoids are somewhat controversial, with conflicting views about whether adapoids are haplorhines (Wortman 1903; Gingerich 1973, 1984, 2012, 2015; Gingerich and Schoeninger 1977; Rasmussen and Simons 1988, 1992, 1994; Simons 1989; Simons and Rasmusen 1989; Rasmussen 1990, 1994; Bloch et al. 1997; Franzen et al. 2009) or stepsirrhines (Gregory 1920; Hoffstetter 1977; Beard et al. 1988; Dagosto 1988; Kay et al. 1997; Seiffert et al. 2009; Williams et al. 2010; Gilbert and Maiolino 2015). Current large-scale cladistic analyses tend to place adapoids in the strepsirrhine side of the primate tree (Ni et al. 2016; Gunnell et al. 2018; Seiffert et al. 2018), so the consensus leans toward considering them as most likely stem strepsirrhines. Here, we follow the current consensus and consider adapoids stem strepsirrhines and consequently apply the strepsirrhine-specific EQ equation to Adapis (Supplementary Data SD5). This equation yields an EQ of 0.62, similar to that derived from Jerison's (1973) equation. The primate and the euarchontan-specific equations give similar results (EQ $_{Primates}$  = 0.44; EQ $_{Euarchonta}$  = 0.45; Supplementary Data SD7, SD9), and the Euarchontoglires-specific equation (Supplementary Data SD11) provides a similar EQ to the strepsirrhine-specific and Jerison's (1973) equations (0.65).

Harrington et al. (2020) published a virtual endocast of the omomyoid, N. antiquus, a late Eocene European microchoerid, based on a very complete cranium (MaPhQ 289). Jerison's (1973) equation yields an EQ of 0.68 for Necrolemur, and Eisenberg's (1981) an EQ of 1.04. Harrington et al. (2020) calculated the body mass of N. antiquus using a cranial length equation (Martin 1990). Unlike adapoids, the phylogenetic relationships of omomyoids are much less controversial and they are mostly regarded as haplorhines, with a closer relationship to modern tarsiers (e.g., Ni et al. 2016). The haplorhinespecific equation gives a much lower EQ (0.43; Supplementary Data SD6) than that provided by the generalized mammalian equations, whereas the primate and euarchontan equations give more intermediate values (EQ<sub>Primates</sub> = 0.59; EQ<sub>Euarchonta</sub> = 0.62; Supplementary Data SD7, SD9). The Euarchontoglires-specific EQ yields a more similar value (0.91) to that given using Eisenberg's (1981) equation. The results from both A. parisiensis and N. antiquus highlight

Downloaded from https://academic.oup.com/jmammal/article/105/6/1430/7734701 by University of California Library - Berkeley Library user on 07 July 2025

**Table 6.** Pairwise comparisons of bootstrap estimates of the differences in PGLS regression parameters among Euarchontoglires. SD = standard deviation. Asterisks = likely to be different.

Taxon	Slope	SD	95% CI		Intercept	SD	95% CI	
	bootstrap difference		Lower end	Upper end	bootstrap difference		Lower end	Upper end
Euarchonta-Glires	*620.0	0.025	0.031	0.128	0.050	0.126	-0.197	0.286
Rodentia–Lagomorpha	-0.060	0.085	-0.226	0.106	0.198	0.272	-0.301	0.708
Rodentia–Scandentia	-0.054	0.115	-0.279	0.161	0.000	0.269	-0.543	0.522
Lagomorpha–Scandentia	-0.115	0.142	-0.386	0.168	0.197	0.366	-0.484	0.901
Primates–Glires	0.083*	0.026	0.031	0.132	0.167	0.144	-0.133	0.441
Primates–Rodentia	0.082*	0.026	0.034	0.134	0.172	0.147	-0.105	0.450
Primates–Lagomorpha	-0.142	0.088	-0.309	0.027	0.026	0.296	-0.534	0.616
Primates–Scandentia	0.027	0.116	-0.211	0.247	0.172	0.276	-0.365	0.708
Strepsirrhini–Haplorhini	-0.036	0.053	-0.143	0.067	0.219	0.230	-0.234	999.0
Strepsirrhini-Scandentia	-0.043	0.123	-0.281	0.197	-0.080	0.323	-0.720	0.578
Strepsirrhini–Rodentia	0.088	0.046	-0.002	0.184	0.092	0.214	-0.309	0.511
Strepsirrhini–Lagomorpha	0.149	0.091	-0.020	0.330	-0.106	0.316	-0.704	0.509
Haplorhini–Scandentia	0.007	0.118	-0.213	0.245	0.299	0.310	-0.313	0.888
Haplorhini–Rodentia	0.052	0.032	-0.012	0.117	0.311	0.188	-0.056	0.668
Haplorhini–Lagomorpha	0.113	0.090	-0.067	0.288	0.114	0.319	-0.513	0.741




Fig. 1. Bivariate plots of log brain mass versus log body mass for euarchontogliran species. Points are color-coded by order, except for Primates, which are divided into the suborders Haplorhini and Strepsirrhini. Best fit lines are shown for the OLS (dashed) and PGLS (solid) regressions. (A) Data for the entire sample, with the best fit line for Euarchontoglires (OLS slope = 0.809, intercept = -1.344; PGLS slope = 0.540, intercept = -0.670); (B) data for members of Euarchonta (Primates, Dermoptera, Scandentia; OLS slope = 0.805, intercept = -1.177; PGLS slope = 0.603, intercept = -0.724); (C) data for members of Glires (Rodentia, Lagomorpha; OLS slope = 0.635, intercept = -1.044; PGLS slope = 0.524, intercept = -0.774).




Fig. 2. Bivariate plots of log brain mass versus log body mass for individual euarchontogliran orders. Best fit lines are shown for the OLS (dashed) and PGLS (solid) regressions. (A) Data for Primates, with points color-coded by suborder. The best fit lines are for all Primates (OLS slope = 0.794, intercept = -1.133; PGLS slope = 0.607, intercept = -0.607); (B) data for Scandentia (OLS slope = 0.604, intercept = -0.821; PGLS slope = 0.580, intercept = -0.778); (C) data for Rodentia (OLS slope = 0.638, intercept = -0.931; PGLS slope = 0.594, intercept = -0.931; PGLS slope = 0.931; PGLS slope = 0.93 slope = 0.465, intercept = -0.581).




Fig. 3. Bivariate plots of log brain mass versus log body mass for Primates by suborder. Best fit lines are shown for the OLS (dashed) and PGLS (solid) regressions. (A) Data for Haplorhini (OLS slope = 0.723, intercept = -0.834; PGLS slope = 0.578, intercept = -0.468); (B) data for Strepsirrhini (OLS slope = 0.679; intercept = -0.932; PGLS slope = 0.614, intercept = -0.687).

⋖

the importance of selecting an EQ equation based on the research question being asked. In particular, for N. antiquus, the very low haplorhine EQ value emphasizes the contrast in relative brain size between this fossil taxon and living members of Haplorhini. So, N. antiquus was quite encephalized for an early Tertiary primate, but actually not very encephalized for a modern haplorrhine, as might be expected for a stem taxon.

To conclude, the present study explores scaling relationships between body mass and brain mass in Euarchontoglires. To our knowledge, this is the first instance that the scaling relationships between brain mass and body mass have been reported for Scandentia (Fig. 2B), a group that has been underrepresented in previous studies for lack of brain data. Our results also show that the brain of lagomorphs scales with respect to body mass similarly to rodents, and very differently than Primates (Fig. 4A and B; contra Burger et al. 2019).

Generating clade-specific EQs provides some insight into issues that have been raised with the general mammalian equations of Jerison (1973) and Eisenberg (1981). As shown by Gilbert and Jungers (2017); see also Begun and Kordos (2004), these EQ equations do an imperfect job at controlling for body mass. A critical reason for this issue is that the brain allometries in the various subgroups of mammals vary greatly, making clade-specific EQs necessary. In other words, the relationship between EQ and body mass observed by those authors stems from using a scaling equation (either Jerison's or Eisenberg's) that does not correctly control for body mass because it does not accurately reflect scaling relationships of the mammalian subsample of interest (i.e., Primates, Rodentia, Lagomorpha, etc.). Therefore, clade-specific EQs will be more appropriate when the research questions being asked concern a more taxonomically narrow sample. The use of Jerison's (1973) and Eisenberg's (1981) equations might be appropriate for research questions that concern a broad sample of mammals, particularly distantly related mammals. Choosing Jerison's (1973) or Eisenberg's (1981) equations for a taxonomically narrow sample (like Rodentia or Primates) will maximize the overestimations and underestimations of relative brain size for members with more extreme body sizes (either small or large, respectively). Eisenberg's (1981) equation gives closer EQ values than Jerison's (1973) does for rodents to those yielded by Pilleri et al.'s (1984) equation, which is a rodent-specific OLS-based equation. This makes Eisenberg's (1981) equation a more appropriate choice for rodents over Jerison's (1973). However, Jerison's (1973) equation yields similar EQ values to those given by the strepsirrhine-specific EQ, making Jerison's (1973) equation more appropriate for strepsirrhines over Eisenberg's (1981). This contrast highlights the reality that neither is a generally preferred option across all scales of comparisons. In sum, clade-specific equations (Pilleri et al. 1984; Grabowski et al. 2016; Ni et al. 2019; Bertrand et al. 2021; this paper) should be given preference whenever the research question focuses on a particular mammalian group, and generalized mammalian equations (Jerison 1973; Eisenberg 1981) are only appropriate for broad comparative mammalian samples.

# Supplementary data

Supplementary data are available at Journal of Mammalogy online.

Supplementary Data SD1. Text explaining how encephalization quotients are calculated.

Supplementary Data SD2. Body mass and brain mass data for 715 euarchontogliran species, sorted by order. Sample sizes are included as specifically as the literature allows, sometimes including ranges. Data have been extracted from the "Direct




Fig. 4. Bivariate plots of log brain mass versus log body mass for all 715 euarchontoglirans. (A) Data are color-coded by order, best fit lines are shown for the OLS regressions for each order except for Dermoptera. Primates slope = 0.794, intercept = -1.133; Scandentia slope = 0.604, intercept = -0.821; Rodentia slope = 0.638, intercept = -1.050; Lagomorpha slope = 0.594, intercept = -0.931. (B) Data are color-coded by order, best fit lines are shown for the PGLS regressions for each order except for Dermoptera. Primates slope = 0.607, intercept = -0.607; Scandentia slope = 0.580, intercept = -0.778; Rodentia slope = 0.526, intercept = -0.779; Lagomorpha slope = 0.465, intercept = -0.581. (C) Data are color-coded by superorder, best fit lines are shown for the OLS (dashed) and PGLS (solid) regressions for Euarchonta (Primates, Dermoptera, Scandentia; OLS slope = 0.805, intercept = -1.177; PGLS slope = 0.603, intercept = -0.724); and Glires (Rodentia, Lagomorpha; OLS slope = 0.635, intercept = -1.044; PGLS slope = 0.524, intercept = -0.774).

source." Direct sources often include original sources where the data came from in the respective manuscripts. When a direct source is listed as "Combined," it means that data from several original sources (sometimes including newly generated data) have been combined.

Supplementary Data SD3. Results from the reduced major axis (RMA) analysis. This file includes slope and intercept data for several taxonomic levels in Euarchontoglires using RMA and pairwise comparisons between regression parameters of different taxonomic groups of Euarchontoglires. Values in red denote coverage probabilities that do not include zero. N = number of species; SD = standard deviation; CI = confidence interval; CP = coverage probability.

Supplementary Data SD4. References used for tables in Supplementary Data SD2.

Supplementary Data SD5. Bivariate plots of log brain mass versus log body mass for Strepsirrhini (Fig. 3B) with Adapis parisiensis indicated with a black star. OLS regression represented as a dashed line; PGLS regression represented as a solid line. Note that the body mass and brain mass values for A. parisiensis have not been taken into account when calculating the regression lines.

Supplementary Data SD6. Bivariate plots of log brain mass versus log body mass for Strepsirrhini (Fig. 3A) with Necrolemur antiquus indicated with a black star. OLS regression represented as a dashed line; PGLS regression represented as a solid line. Note that the body mass and brain mass values for N. antiquus have not been taken into account when calculating the regression lines.

Supplementary Data SD7. Bivariate plots of log brain mass versus log body mass for Primates (Fig. 2A) with Microsyops annectens, Adapis parisiensis, and Necrolemur antiquus indicated with black stars. OLS regression represented as a dashed line; PGLS regression represented as a solid line. Note that the body mass and brain mass values for M. annectens, A. parisiensis, and N. antiquus have not been taken into account when calculating the regression lines. BM1, high body mass estimate; BMJ, low body mass estimate.

Supplementary Data SD8. Bivariate plots of log brain mass versus log body mass for Lagomorpha (Fig. 2D) with Megalagus turgidus indicated with a black star. OLS regression represented as a dashed line; PGLS regression represented as a solid line. Note that the body mass and brain mass values for M. turgidus have not been taken into account when calculating the regression lines.

Supplementary Data SD9. Bivariate plots of log brain mass versus log body mass for Euarchonta (Fig. 1B) with Microsyops annectens, Adapis parisiensis, and Necrolemur antiquus indicated with black stars. OLS regression represented as a dashed line; PGLS regression represented as a solid line. Note that the body mass and brain mass values for M. annectens, A. parisiensis, and N. antiquus have not been taken into account when calculating the regression lines. BM1, high body mass estimate; BMJ, low body mass estimate.

Supplementary Data SD10. Bivariate plots of log brain mass versus log body mass for Glires (Fig. 1C) with Megalagus turgidus indicated with a black star. OLS regression represented as a dashed line; PGLS regression represented as a solid line. Note that the body mass and brain mass values for M. turgidus have not been taken into account when calculating the regression lines.

Supplementary Data SD11. Bivariate plots of log brain mass versus log body mass for Euarchontoglires (Fig. 1A) with Microsyops annectens, Adapis parisiensis, Necrolemur antiquus, and Megalagus turgidus indicated with black stars. OLS regression represented as a dashed line; PGLS regression represented as a solid line. Note that the body mass and brain mass values for M. annectens, A. parisiensis, N. antiquus, and M. turgidus have not been taken into account when calculating the regression lines. BM1, high body mass estimate; BM1, low body mass estimate.

## Acknowledgments

We thank A. Billings for helping the authors with a literature search; her work was funded by the University of Toronto Work Study program. We thank Deyan Ge, Carrie Mongle, Brett Riddle, and an anonymous reviewer for helpful comments that improved this paper. The figures in the manuscript include some silhouettes from phylopic.org, specifically of Cebus (by Sarah Werning; green; CC BY 3.0), Galeopterus variegatus (by Yan Wong; flipped horizontally, red; CC0 1.0), and Lemur catta (by Geoff Shaw; flipped horizontally, yellow; CC BY-NC 3.0).

## **Author contributions**

SL-T and MTS contributed to conceptualization and project administration; SL-T, OCB, ŁF-F, MML, GSM-F, and MTS to data curation; CJL and MAS to formal analysis, methodology, software, and validation; SL-T, OCB, ŁF-F, MML, CJL, and MTS to funding acquisition; MTS to supervision; SL-T and CJL to visualization; SL-T to writing-original draft; all authors contributed to investigation, resources, and writing-review & editing.

## **Funding**

This study was supported by a National Science Centre (Cracow, Poland) grant (number 2022/45/NZ8/03585) to SL-T, a CERCA Programme/Generalitat de Catalunya, a Marie Skłodowska-Curie Actions: Individual Fellowship (H2020-MSCA-IF-2018-2020; no. 792611), and a Beatriu de Pinós 2021 BP 00042 to OCB; a National Science Centre (Cracow, Poland) grant (number 2015/18/NZ8/00637) to ŁF-F, an NSERC CGS grant to MML; National Science Foundation (DBI-1906248, DBI-2128146), a Gerstner Family Foundation Scholars Postdoctoral Fellowship, and a University of Texas Early Career Provost Fellowship to CJL; and an NSERC Discovery Grant to MTS.

#### **Conflict of interest**

None declared.

## Data availability

All data are available in the Supplementary Data.

#### References

Armstrong E. 1985. Allometric considerations of the adult mammalian brain, with special emphasis on primates. In: Jungers WL, editor. Size and scaling in primate biology. New York (NY, USA): Springer Science+Business Media; p. 115-146.

Beard KC, Dagosto M, Gebo DL, Godinot M. 1988. Interrelationships among primate higher taxa. Nature 331(6158):712-714. https:// doi.org/10.1038/331712a0

Begun DR, Kordos L. 2004. Cranial evidence of the evolution of intelligence in fossil apes. In: Russon AE, Begun DR, editors. The evolution of thought: evolutionary origins of great ape intelligence. Cambridge (UK): Cambridge University Press; p. 260-279.

Bernard RTF, Nurton J. 1993. Ecological correlates of relative brain size in some South American rodents. South African Journal of Zoology 28(2):95-98. https://doi.org/10.1080/02541858.1993.11448300

Bertrand OC, Amador-Mughal F, Lang MM, Silcox MT. 2018. Virtual endocasts of fossil Sciuroidea: brain size reduction in the evolution of fossoriality. Palaeontology 61(6):919-948. https://doi. org/10.1111/pala.12378

- Bertrand OC, Amador-Mughal F, Lang MM, Silcox MT. 2019a. New virtual endocasts of Eocene Ischyromyidae and their relevance in evaluating neurological changes occurring through time in Rodentia. Journal of Mammalian Evolution 26(3):345–371. https:// doi.org/10.1007/s10914-017-9425-6.
- Bertrand OC, Amador-Mughal F, Silcox MT. 2016. Virtual endocasts of Eocene Paramys (Paramyinae): oldest endocranial record for Rodentia and early brain evolution in Euarchontoglires. Proceedings of the Royal Society of London, B: Biological Sciences 283(1823):20152316. https://doi.org/10.1098/rspb.2015.2316
- Bertrand OC, Amador-Mughal F, Silcox MT. 2017. Virtual endocast of the early Oligocene Cedromus wilsoni (Cedromurinae) and brain evolution in squirrels. Journal of Anatomy 230(1):128-151. https:// doi.org/10.1111/joa.12537
- Bertrand OC, Püschel HP, Schwab JA, Silcox MT, Brusatte SL. 2021. The impact of locomotion on the brain evolution of squirrels and close relatives. Communications Biology 4(1):460. https://doi. org/10.1038/s42003-021-01887-8
- Bertrand OC, San Martin-Flores G, Silcox MT. 2019b. Endocranial shape variation in the squirrel-related clade and their fossil relatives using 3D geometric morphometrics: contributions of locomotion and phylogeny to brain shape. Journal of Zoology 308(3):197-211. https://doi.org/10.1111/jzo.12665
- Bertrand OC, Shelley SL, Williamson TE, Wible JR, Chester SGB, Flynn JJ, Holbrook LT, Lyson TR, Meng J, Miller IM, et al. 2022. Brawn before brains in placental mammals after the end-Cretaceous extinction. Science 376(6588):80-85. https://doi.org/10.1126/science.abl5584
- Bertrand OC, Silcox MT. 2016. First virtual endocasts of a fossil rodent: Ischyromys typus (Ischyromyidae, Oligocene) and brain evolution in rodents. Journal of Vertebrate Paleontology 36(3):e1095762. https://doi.org/10.1080/02724634.2016.1095762
- Bloch JI, Fisher DC, Gingerich PD, Gunnell GF, Simons EL, Uhen MD. 1997. Cladistic analysis and anthropoid origins. Science 278(5346):2134-2136. https://doi.org/10.1126/science.278.5346.2134
- Boddy AM, McGowen MM, Sherwood CC, Grossman LI, Goodman M, Wildman DE. 2012. Comparative analysis of encephalization in mammals reveals relaxed constraints on anthropoid primate and cetacean brain scaling. Journal of Evolutionary Biology 25(5):981-994. https://doi.org/10.1111/j.1420-9101.2012.02491.x
- Bronson RT. 1981. Brain weight-body weight relationships in 12 species of nonhuman primates. American Journal of Physical Anthropology 56(1):77-81. https://doi.org/10.1002/ajpa.1330560109
- Burger JP, George MA Jr, Leadbetter C, Shaikh F. 2019. The allometry of brain size in mammals. Journal of Mammalogy 100(2):276-283. https://doi.org/10.1093/jmammal/gyz043
- Burgin CH, Colella JP, Kahn PL, Upham NS. 2018. How many species of mammals are there? Journal of Mammalogy 99(1):1-14. https:// doi.org/10.1093/jmammal/gyx147
- Clutton-Brock TH, Harvey PH. 1980. Primates, brains and ecology. Journal of Zoology 190(3):309-323. org/10.1111/j.1469-7998.1980.tb01430.x
- Count EW. 1947. Brain and body weight in man: their antecedents in growth and evolution. A study in dynamic somatometry. Annals of the New York Academy of Sciences 46(10):993-1122. https://doi. org/10.1111/j.1749-6632.1947.tb36165.x
- Dagosto M. 1988. Implications of postcranial evidence for the origin of euprimates. Journal of Human Evolution 17(1-2):35-56. https:// doi.org/10.1016/0047-2484(88)90048-6
- Deacon TW. 1990. Fallacies of progression in theories of brain-size evolution. International Journal of Primatology 11(3):193-236. https:// doi.org/10.1007/bf02192869
- Deaner RO, Isler K, Burkart J, van Schaik C. 2007. Overall brain size, and not encephalization quotient, best predicts cognitive ability across

- non-human primates. Brain Behavior and Evolution 70(2):115-124. https://doi.org/10.1159/000102973
- Dubois E. 1898. Ueber die Abhängigkeit des Hirngewichtes von der Körpergrosse bei den Säugethieren. Archiv fur Anthropologie 25:1-28.
- Eisenberg JF. 1981. The mammalian radiations: an analysis of trends in evolution, adaptation and behavior. Chicago (IL, USA): University of Chicago Press.
- Foley NM, Mason VC, Harris AJ, Bredmeyer KR, Damas J, Lewin HA, Eizirik E, Gatesy J, Karlsson EK, Lindbald-Toh K, et al. 2023. A genomic timescale for placental mammal evolution. Science 380(6643):eab18189. https://doi.org/10.1126/science.abl8189
- Franzen JL, Gingerich PD, Habersetzer J, Hurum JH, von Koenigswald W, Smith BH. 2009. Complete primate skeleton from the middle Eocene of Messel in Germany: morphology and paleobiology. PLoS One 4(5):e5723. https://doi.org/10.1371/journal.pone.0005723
- Gilbert CC, Jungers WL. 2017. Comment on relative brain size in early primates and the use of encephalization quotients in primate evolution. Journal of Human Evolution 109:79-87. https://doi. org/10.1016/j.jhevol.2017.04.007
- Gilbert CC, Maiolino SA. 2015. Comment to 'Primates in the Eocene' by Gingerich (2012). Palaeobiodiversity and Palaeoenvironments 95(2):237-241. https://doi.org/10.1007/s12549-015-0184-1
- Gingerich PD. 1973. Anatomy of the temporal bone in the Oligocene anthropoid Apidium and the origin of Anthropoidea. Folia Primatologica 19(5):329–337. https://doi.org/10.1159/000155549
- Gingerich PD. 1984. Primate evolution: evidence from the fossil record, comparative morphology, and molecular biology. American Journal of Physical Anthropology 27(S5):57-72. https://doi.org/10.1002/ ajpa.1330270504
- Gingerich PD. 2012. Primates in the Eocene. In: Lehman T, Schaal SFK, editors. The world at the time of Messel: puzzles in palaeobiology, palaeoenvironment, and the history of early primates. Frankfurt am Main (Germany): Senckenberg Gesellschaft für Naturforschung; p. 67-68.
- Gingerich PD. 2015. Reply to comment on "Primates in the Eocene" by Gilbert and Maiolino (2015). Palaeobiodiversity and Palaeoenvironments 95(2):243-246. https://doi.org/10.1007/ s12549-015-0194-z
- Gingerich PD, Schoeninger M. 1977. The fossil record and primate phylogeny. Journal of Human Evolution 6(5):483-505. https://doi. org/10.1016/s0047-2484(77)80059-6
- Gingerich PD, Smith BH, Rosenberg KR. 1982. Allometric scaling in the dentition of primates and prediction of body weight from tooth size in fossils. American Journal of Physical Anthropology 58(1):81–100. https://doi.org/10.1002/ajpa.1330580110
- Gould SJ. 1975. Allometry in primates, with emphasis on scaling and evolution of the brain. Contributions to Primatology 5:244-292.
- Grabowski M, Voje KL, Hansen TF. 2016. Evolutionary modeling and correcting for observation error support a 3/5 brain-body allometry for primates. Journal of Human Evolution 94:106–116. https:// doi.org/10.1016/j.jhevol.2016.03.001
- Gregory WK. 1920. On the structure and relations of Notharctus, an American Eocene primate. Memoirs of the American Museum of Natural History 3(2):49–243.
- Gunnell GF, Boyer DM, Friscia AR, Heritage S, Manthi FK, Miller ER, Sallam HM, Simmons NB, Stevens NJ, Seiffert ER. 2018. Fossil lemurs from Egypt and Kenya suggest an African origin for Madagascar's aye-aye. Nature Communications 9(1):3193. https:// doi.org/10.1038/s41467-018-05648-w
- Hafner MS, Hafner JC. 1984. Brain size, adaptation and heterochrony in geomyoid rodents. Evolution 38(5):1088–1098. https://doi. org/10.1111/j.1558-5646.1984.tb00378.x

- Harrington AR, Silcox MT, Yapuncich GS, Boyer DM, Bloch JI. 2016. First virtual endocasts of adapiform primates. Journal of Human Evolution 99:52-78. https://doi.org/10.1016/j.jhevol.2016.06.005
- Harrington AR, Yapuncich GS, Boyer DM. 2020. The digital endocast of Necrolemur antiquus. Palaeovertebrata 43(2):e1. https://doi. org/10.18563/pv.43.2.e1
- Hoffstetter R. 1977. Phylogénie des primates: confrontation des résultats obtenus per les diverses voies d'approche du problème. Bulletins et Mémoires de la Société d'Anthropologie de Paris 4(4):327-346. https://doi.org/10.3406/bmsap.1977.1886
- Holekamp KE, Swanson EM, Van Meter PE. 2013. Developmental constraints on behavioural flexibility. Philosophical Transactions of the Royal Society of London, B: Biological Sciences 368(1618):20120350. https://doi.org/10.1098/rstb.2012.0350
- Jerison JH. 1973. Evolution of the brain and intelligence. New York (NY, USA): Academic Press.
- Jerison JH. 1985. Animal intelligence as encephalization. Philosophical Transactions of the Royal Society of London, B: Biological Sciences 308(1135):21-35. https://doi.org/10.1098/rstb.1985.0007
- Kay RF, Ross CF, Williams BA. 1997. Anthropoid origins. Science 275(5301):797-804. https://doi.org/10.1126/science.275.5301.797
- Lande R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain: body size allometry. Evolution 33(1):402-416. https://doi.org/10.1111/j.1558-5646.1979.tb04694.x
- Lang MM, Bertrand OC, San Martin-Flores G, Law CJ, Abdul-Sater J, Spakowski S, Silcox MT. 2022. Scaling patterns of cerebellar petrosal lobules in Euarchontoglires: impacts of ecology and phylogeny. Anatomical Record 305(12):3472-3503. https://doi. org/10.1002/ar.24929
- López-Torres S, Bertrand OC, Lang MM, Silcox MT, Fostowicz-Frelik Ł. 2020. Cranial endocast of the stem lagomorph Megalagus and brain structure of basal Euarchontoglires. Proceedings of the Royal Society of London, B: Biological Sciences 287(1929):20200665. https://doi.org/10.1098/rspb.2020.0665
- Mace GM, Eisenberg JF. 1982. Competition, niche specialization and the evolution of brain size in the genus Peromyscus. Biological Journal of the Linnean Society 17(3):243-257. https://doi. org/10.1111/j.1095-8312.1982.tb02019.x
- Mace GM, Harvey PH, Clutton-Brock TH. 1981. Brain size and ecology in small mammals. Journal of Zoology 193(3):333-354. https://doi. org/10.1111/j.1469-7998.1981.tb03449.x
- Mann MD, Towe AL. 2003. Brain-body size relations in grasshopper mice. Brain, Behavior and Evolution 62(1):13-18. https://doi. org/10.1159/000071956
- Marino L. 1998. A comparison of encephalization between odontocete cetaceans and anthropoid primates. Brain, Behavior and Evolution 51(4):230-238. https://doi.org/10.1159/000006540
- Martin RD. 1990. Primate origins and evolution: a phylogenetic reconstruction. London (UK): Chapman & Hall.
- Martin RD, Harvey PH. 1985. Brain size allometry: Ontogeny and phylogeny. In: Jungers WL, editor. Size and scaling in primate biology. New York (NY, USA): Springer Science+Business Media; p. 147-174.
- Mason VC, Li G, Minx P, Schmitz J, Churakov G, Doronina L, Melin AD, Dominy NJ, Lim NT-L, Springer MS, et al. 2016. Genomic analysis reveals hidden biodiversity within colugos, the sister group to primates. Science Advances 2(8):e1600633. https://doi.org/10.1126/ sciadv.1600633
- Matějů JL, Kratochvíl L, Pavelková Z, Řičánková VP, Vohralík V, Němec P. 2016. Absolute, not relative brain size correlates with sociality in ground squirrels. Proceedings of the Royal Society of London, B: Biological Sciences 283(1827):20152725. https://doi.org/10.1098/ rspb.2015.2725

- Moncunill-Solé B, Quintana J, Jordana X, Engelbrektsson P, Köhler M. 2015. The weight of fossil leporids and ochotonids: body mass estimation models for the order Lagomorpha. Journal of Zoology 295(4):269-278. https://doi.org/10.1111/jzo.12209
- Murphy WJ, Eizirik E, Johnson WE, Wang YP, Ryder OA, O'Brien SJ. 2001. Molecular phylogenetics and the origins of placental mammals. Nature 409:614–618. https://doi.org/10.1038/35054550
- Ni X, Flynn JJ, Wyss AR, Zhang C. 2019. Cranial endocast of a stem platyrrhine primate and ancestral brain conditions in anthropoids. Science Advances 5(8):eaav7913. https://doi.org/10.1126/sciadv.aav7913
- Ni X, Li Q, Li L, Beard KC. 2016. Oligocene primates from China reveal divergence between African and Asian primate evolution. Science 352(6286):673-677. https://doi.org/10.1126/science.aaf2107
- Olson LE, Sargis EJ, Martin RD. 2005. Intraordinal phylogenetics of treeshrews (Mammalia: Scandentia) based on evidence from the mitochondrial 12S rRNA gene. Molecular Phylogenetics and Evolution 35(3):656-673. https://doi.org/10.1016/j.ympev.2005.01.005
- Orliac MJ, Gilissen E. 2012. Virtual endocranial cast of earliest Eocene Diacodexis (Artiodactyla, Mammalia) and morphological diversity of early artiodactyl brains. Proceedings of the Royal Society of London, B: Biological Sciences 279(1743):3670-3677. https://doi. org/10.1098/rspb.2012.1156
- Pagel M. 1999. Inferring the historical patterns of biological evolution. Nature 401(6756):877-884. https://doi.org/10.1038/44766
- Pilleri G, Gihr M, Kraus C. 1984. Cephalization in rodents with particular reference to the Canadian beaver (Castor canadensis). In: Pilleri G, editor. Investigations on beavers. Bern, Switzerland: Brain Anatomy Institute; p. 11-102.
- Püschel HP, Bertrand OC, O'Reilly JE, Bobe R, Püschel TA. 2021. Divergence-time estimates for hominins provide insights into encephalization and body mass trends in human evolution. Nature Ecology & Evolution 5:808-819. https://doi.org/10.1038/ s41559-021-01431-1
- Rasmussen DT. 1990. The phylogenetic position of Mahgarita stevensi: protoanthropoid or lemuroid? International Journal of Primatology 11(5):439-469. https://doi.org/10.1007/bf02196131
- Rasmussen DT. 1994. The different meanings of a tarsioid-anthropoid clade and a new model of anthropoid origin. In: Fleagle JG, Kay RF, editors. Anthropoid origins. New York (NY, USA): Plenum; p. 335-360.
- Rasmussen DT, Simons EL. 1988. New specimens of Oligopithecus savage, early Oligocene primate from the Fayum, Egypt. Folia Primatologica 51(4):182–208. https://doi.org/10.1159/000156372
- Rasmussen DT, Simons EL. 1992. Paleobiology of oligopithecines, the earliest known anthropoid primate. International Journal of Primatology 13(5):477–508. https://doi.org/10.1007/bf02547829
- Rasmussen DT, Simons EL. 1994. A whole new world of ancestors: Eocene anthropoideans from Africa. Evolutionary Anthropology 3(4):128–139. https://doi.org/10.1002/evan.1360030407
- Revell LJ. 2010. Phylogenetic signal and linear regression on species data. Methods in Ecology and Evolution 1(4):319-329. https://doi. org/10.1111/j.2041-210x.2010.00044.x
- San Martin-Flores GA, Nagendran L, Silcox MT. 2018. Insights into the primitive brain of primates: treeshrew cranial endocasts analyzed using geometric morphometrics. American Journal of Physical Anthropology 165(S66):237. https://doi.org/10.1002/ajpa.23489
- Sargis EJ. 2002. A multivariate analysis of the postcranium of tree shrews (Scandentia, Tupaiidae) and its taxonomic impli-66(4):579-598. cations. Mammalia https://doi.org/10.1515/ mamm.2002.66.4.579
- Schoenemann PT. 2006. Evolution of the size and functional areas of the human brain. Annual Review of Anthropology 35(1):379-406. https://doi.org/10.1146/annurev.anthro.35.081705.123210

- Seiffert ER, Boyer DM, Fleagle JG, Gunnell GF, Heesy CP, Perry JMG, Sallam HM. 2018. New adaptform primate fossils from the late Eocene of Egypt. Historical Biology 30(1-2):204-226. https://doi.org /10.1080/08912963.2017.1306522
- Seiffert ER, Perry JMG, Simons EL, Boyer DM. 2009. Convergent evolution of anthropoid-like adaptations in Eocene adapiform primates. Nature 461(7267):1118–1121. https://doi.org/10.1038/nature08429
- Shea BT. 1983. Phyletic size change and brain/body allometry: a consideration based on African pongids and other primates. International Journal of Primatology 4(1):33-62. https://doi. org/10.1007/bf02739359
- Sherwood CC, Subiaul F, Zawidzki TW. 2008. A natural history of the human mind: tracing evolutionary changes in brain and cognition. Journal of Anatomy 212(4):426-454. https://doi. org/10.1111/j.1469-7580.2008.00868.x
- Silcox MT, Benham AE, Bloch JI. 2010. Endocasts of Microsyops (Microsyopidae, Primates) and the evolution of the brain in primitive primates. Journal of Human Evolution 58(6):505-521. https:// doi.org/10.1016/j.jhevol.2010.03.008
- Silcox MT, Bertrand OC, Harrington AR, Lang MM, San Martin-Flores G, López-Torres S. 2023. Early evolution of the brain in primates and their close kin. In: Dozo MT, Paulina-Carabajal A, Macrini TE, Walsh S, editors. Paleoneurology of amniotes: new directions in the study of fossil endocasts. Switzerland: Springer Cham; p. 457-506.
- Silcox MT, Bloch JI, Boyer DM, Godinot M, Ryan TM, Spoor F, Walker A. 2009. Semicircular canal system in early primates. Journal of Human Evolution 56(3):315-327. https://doi.org/10.1016/j. jhevol.2008.10.007
- Silcox MT, Gunnell GF, Bloch JI. 2020. Cranial anatomy of Microsyops annectens (Microsyopidae, Euarchonta, Mammalia) from the middle Eocene of Northwestern Wyoming. Journal of Paleontology 94(5):979-1006. https://doi.org/10.1017/jpa.2020.24
- Simons EL. 1989. Description of two genera and species of late Eocene Anthropoidea of Egypt. Proceedings of the National Academy of Sciences of the United States of America 86(24):9956-9960. https:// doi.org/10.1073/pnas.86.24.9956
- Simons EL, Rasmussen DT. 1989. Cranial morphology of Aegyptopithecus and Tarsius and the question of the tarsier-anthropoidean clade. American Journal of Physical Anthropology 79(1):1–23. https://doi. org/10.1002/ajpa.1330790103
- Smaers JB, Dechmann DKN, Goswami A, Soligo C, Safi K. 2012. Comparative analyses of evolutionary rates reveal different pathways to encephalization in bats, carnivorans, and primates. Proceedings of the National Academy of Sciences of the United States of America 109(44):18006-18011. https://doi.org/10.1073/ pnas.1212181109
- Smaers JB, Rothman RS, Hudson DR, Balanoff AM, Beatty B, Dechmann DKN, DeVries D, Dunn JC, Fleagle JG, Gilbert CC, et al. 2021. The evolution of mammalian brain size. Science Advances 7(18):eabe2101. https://doi.org/10.1126/sciadv.abe2101
- Smith RJ. 2020. P > .05: the incorrect interpretation of "not significant" results is a significant problem. American Journal of Physical Anthropology 172(4):521–527. https://doi.org/10.1002/ajpa.24092

- Snell O. 1892. Die Abhangigkeit des Hirngewitches von dem Körpergewicht un den geistigen Fähigkeiten. Archiv für Psychiatrie un Nervenkrankheiten 23:436-446.
- Stafford BJ, Szalay FS. 2000. Craniodental functional morphology and taxonomy of dermopterans. Journal of Mammalogy 81(2):360-385. https://doi.org/10.1093/jmammal/81.2.360
- Stephan H, Frahm H, Baron G. 1981. New and revised data on volumes of brain structures in insectivore and primates. Folia Primatologica 35(1):1-29. https://doi.org/10.1159/000155963
- Stephan HR, Bauchot R, Andy OJ. 1970. Data on size of the brain and of various brain parts in insectivores and primates. In: Noback CR, Montagna W, editors. The primate brain. New York (NY, USA): Appleton-Century-Crofts; p. 289–297.
- Thewissen JGM, Gingerich PD. 1989. Skull and endocranial cast of Eoryctes melanus, a new palaeoryctid (Mammalia: Insectivora) from the early Eocene of western North America. Journal of Vertebrate Paleontology 9(4):459-470. https://doi.org/10.1080/02724634.1989. 10011778
- Towe AL, Mann MD. 1992. Brain size/body length relations among myomorph rodents. Brain, Behavior and Evolution 39(1):17-23. https:// doi.org/10.1159/000114101
- Tung Ho LS, Ané C. 2014. A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. Systematic Biology 63(3):397-408. https://doi.org/10.1093/sysbio/syu005
- Upham NS, Esselstyn JA, Jetz W. 2019. Inferring the mammal tree: species-level sets of phylogenies for questions in ecology, evolution, and conservation. PLoS Biology 17(12):e3000494. https://doi. org/10.1371/journal.pbio.3000494
- Waddell PJ, Okada N, Hasegawa M. 1999. Towards resolving the interordinal relationships of placental mammals. Systematic Biology 48(1):1-5.
- White CL, Bloch JI, Morse PE, Silcox MT. 2023. Virtual endocast of late Paleocene Niptomomys (Microsyopidae, Euarchonta) and early primate brain evolution. Journal of Human Evolution 175:103303. https://doi.org/10.1016/j.jhevol.2022.103303
- Williams BA, Kay RF, Kirk EC, Ross CF. 2010. Darwinius masillae is a strepsirrhine—a reply to Franzen et al. (2009). Journal of Human Evolution 59(5):567–573. https://doi.org/10.1016/j. jhevol.2010.01.003
- Wortman JL. 1903. Studies of Eocene Mammalia in the Marsh collection, Peabody Museum. Part III, Primates. American Journal of Science 15:163-176, 399-414, 419-436; 16:345-368; 17:23-33, 133-140, 203-214.
- Yao L, Brown JO, Stampanoni M, Marone F, Isler K, Martin RD. 2012. Evolutionary change in the brain size of bats. Brain, Behavior & Evolution 80(1):15-25. https://doi.org/10.1159/000338324
- Yapuncich GS, Gladman JT, Boyer DM. 2015. Predicting euarchontan body mass: a comparison of tarsal and dental variables. American Journal of Physical Anthropology 157(3):472-506. https://doi. org/10.1002/ajpa.22735
- Zhang M-L, Li M-L, Ayoola AO, Murphy WJ, Wu D-D, Shao Y. 2019. Conserved sequences identify the closest living relatives of primates. Zoological Research 40(6):532-540. https://doi. org/10.24272/j.issn.2095-8137.2019.057