

SARS-CoV-2 small viral RNA suppresses gene expression via complementary binding to mRNA 3' UTR

Haley A Delcher¹, Jeffrey D DeMeis¹, Nicole Ghobar¹, Noel L Godang¹, Sierra L Knight¹, Shahem Y Alqudah¹, Kevin N Nguyen¹, Brianna C Watters¹, Glen M Borchert^{1,2§}

¹Department of Pharmacology, College of Medicine, University of South Alabama, Mobile, AL

²Department of Biology, College of Arts and Sciences, University of South Alabama, Mobile, AL

§To whom correspondence should be addressed: borchert@southalabama.edu

Abstract

SARS-CoV-2 (SC2) has been intensely studied since its emergence. However, the mechanisms of host immune dysregulation triggered by SC2 remain poorly understood. That said, it is well established that many prominent viral families encode microRNAs (miRNAs) or related small viral RNAs (svRNAs) capable of regulating human genes involved in immune function. Importantly, recent reports have shown that SC2 encodes its own svRNAs. In this study, we have identified 12 svRNAs expressed during SC2 infection and show that one of these svRNAs can regulate target gene expression via complementary binding to mRNA 3' untranslated regions (3'UTRs) much like human microRNAs.

A

Average (start-end)	svRNA Sequence	RPM	4hpi-A	4hpi-B	12hpi-A	12hpi-B	24hpi-A	24hpi-B
			SRR11550029	SRR11550030	SRR11550025	SRR11550026	SRR11550027	SRR11550028
228 - 259	CACAUCAUGGUUUGUCGGGGUGUGACCGAAA	0	0	115	134	202	120	
1606 - 1636	UGACACCUUCUUGAAUACUCCAAAAGAG	0	0	0	0	0	92	
1787 - 1811	AAUUUUAAAAGUUACAAAAGGAAAG	0	0	0	0	0	99	
2451 - 2474	UCAUGCCUCUAAAAGCCCCAAAG	0	0	0	0	278	207	
3177 - 3196	AAGAAGGACAAGAAGAGAU	0	0	0	0	281	209	
3837 - 3857	AAAUGAAGAGUGAAAAGCAAG	0	0	0	0	138	105	
27267 - 27290	GAUUAUCAUAAAACCUCUAAU	0	0	199	242	0	0	
27696 - 27730	UUUUAUACACUUUGCUUCACUCAAAAGAAAGAC	0	0	140	176	0	0	
28535 - 28554	GUGGGGGUGACGGUAAAUG	0	0	124	138	0	0	
28655 - 28684	CUGAGGGAGCCUUGAAUACACAAAAGAAC	0	0	112	0	0	0	
29094 - 29121	GCAGACGUGGUCCAGAACAAACCCAAG	0	0	135	168	278	167	
29331 - 29357	CAAAACAUUCCACCAACAGAGCCUA	53	0	2021	2533	956	578	
29821 - 29846	GAUUUUAAAAGCUUCUUAGGAGAAU	45	0	703	903	309	149	




Figure 1. SARS-CoV-2 svRNA-29094 mimic suppresses target gene in PC3 cells:

(A) 12 svRNA sequences expressed in SARS-CoV-2 infected Calu-3 cells. Average start-end (position in the SC2 NC_045512 genome) alongside corresponding svRNA sequence (+ strand) are shown. Expression (in Reads Per Million) of individual svRNA sequences are indicated under the corresponding SRR Identifiers (Wyler et al. 2021). (B) mFold predicted secondary structure of svRNA-29094 hairpin (Zuker 2003). The excised svRNA is highlighted in green and included below the hairpin. (C) Alignment of putative target mRNA 3'UTRs with the svRNA-29094 sequence. (D) PC3 cell lysates were collected at 24-

1/18/2024 - Open Access

and 48-hours after co-transfection of a small RNA mimic and a psiCHECK-2 reporter expressing Renilla luciferase mRNA harboring the SLC5A9 target site (shown in C) in its 3'UTR. Graph depicts Relative Light Units (RLU) of Renilla luciferase normalized to firefly luciferase (internal control independently expressed from psiCHECK-2). 29094 mimic, commercially synthesized small RNA identical to the svRNA 29094 sequence depicted in B. Ctl mimic, standard Dharmacon negative control microRNA mimic. Error bars indicate standard deviation (n=3). P-values were obtained using a standard t-test.

Description

Over the past 4 years, SARS-CoV-2 (SC2) has killed millions of people and has upended the global economy. SC2 belongs to the Coronaviridae family of viruses and has been classified as a positive sense, single-stranded RNA virus. Although asymptomatic in some patients, SC2 infection can lead to the development of coronavirus disease 19 (COVID-19), a life-threatening inflammatory illness that primarily affects the lungs (Zhang et al. 2020). The severity of COVID-19 is largely reliant upon the ability of SC2 to replicate efficiently while simultaneously shutting down the host immune system (Chang et al. 2021). Despite being intensely studied since its discovery, much is still unknown about the elements of the virus that enable it to dysregulate the host immune response in severe cases of COVID-19. That said, it is well established that many prominent viral families such as the Herpes and Papilloma virus families encode microRNAs (miRNAs) or related short viral RNAs (svRNAs) capable of regulating human genes involved in immune function (Samols et al. 2007; Sadri Nahand et al. 2020). MiRNAs are small (~22nt) noncoding RNA that participate in post-transcriptional gene regulation by complementary binding to the 3'UTR of their respective target mRNAs with some miRNAs possessing the capacity to regulate dozens of target genes in a single controllable event (Lee et al. 1993). Importantly, recent reports have now indicated that SC2 encodes its own svRNAs, and two other groups have already shown that the principle svRNA examined in this work, svRNA-29094, is specifically excised and highly expressed during SC2 infection (Meng et al. 2021; Neeb et al. 2022).

Analysis of small RNA transcriptomes corresponding to Calu-3 cells at 1-, 12-, and 24-hours post-infection (hpi) with SC2 (Wyler et al. 2021) found ~95% of resultant small RNA sequences mapped to the human genome while < 0.4% aligned to the SC2 genome. Strikingly, we observed that roughly 92% of these viral small RNA reads mapped to one of 12 specific genomic positions, are ~20 nt in length and encoded on the positive strand and had peak expressions ranging between ~100 and 2,000 RPM (**Figure 1A**). We selected svRNA-29094 (**Figure 1B**) for further analysis primarily based on (1) its predicted target sequence alignments (**Figure 1C**) and (2) previous reports confirming its expression in SC2-infected cell lines and patients (Meng et al. 2021; Neeb et al. 2022). To confirm the ability of svRNA-29094 to regulate its predicted targets, we performed co-transfections of small RNA mimics and a psiCHECK-2 reporter expressing Renilla luciferase mRNA harboring the SLC5A9 target site (shown in Figure 1C) in its 3'UTR. Luciferase expression was measured at 24- and 48-hours post-transfection and ~27% and ~79% decreases (respectively) in Renilla activity were observed in cells treated with svRNA-29094 mimic as compared to cells transfected with mimic controls (**Figure 1D**).

Viruses such as SC2 remodel host gene expression through a variety of mechanisms including post-transcriptional gene regulation. SvRNAs have now been implicated in the regulation of a variety of different biological pathways related to infection and host immune evasion. Additionally, svRNAs have recently been identified to play important roles in viral propagation in SARS-CoV-1, influenza, and hepatitis infections (Meng et al. 2021). For example, much like with SC2, SARS-CoV-1 infection frequently results in an exaggerated inflammatory response, significant lung pathology, and is associated with a high rate of mortality (Channappanavar and Perlman 2017). The relevance of specific svRNAs in SARS-CoV-1 pathology was previously established, when ten ~20 nt svRNAs were discovered in deep sequenced RNAs from infected mice lungs. Importantly, antagomiR-based inhibition of one of these, svRNA-N, can significantly reduce pro-inflammatory cytokine expression and lung pathology *in vivo* (Morales et al. 2017). These results demonstrate that (at least) this svRNA could directly contribute to SARS-CoV-1 pathogenesis, and also suggests that antagomiRs targeting other coronavirus svRNAs could potentially prove of value as a novel form of potent antiviral.

That said, multiple groups have now independently reported the expression of SC2 svRNAs, although their cellular functions remain almost entirely unresolved (Meng et al. 2021). In this study, we have identified 12 small viral RNAs (svRNAs) expressed during SC2 infection and find at least one of these svRNAs can regulate target gene expression via complementary binding to mRNA 3' untranslated regions. In short, our data confirms the ability of a svRNA-29094 mimic to suppress the expression of a mRNA bearing a 3'UTR target site, and as such, to our knowledge, this study is the first to confirm miRNA-like target mRNA repression by SC2 svRNAs. Furthermore, although we only directly confirmed the ability of svRNA-29094 to repress the intestinal sodium/glucose cotransporter SLC5A9 target site (Tazawa et al. 2005), similarly strong alignments were identified with KPNA6 (known to facilitate nuclear translocation of interferon regulatory factor 3 and inhibited by SC2 membrane protein (Zhang et al. 2021)), ERI1 (known to regulate aspects of Influenza A mRNA transcription (Declercq et al. 2020)), DDX19A (known to detect and bind viral RNA to activate the NLRP3 inflammasome (Li et al. 2015)), and DDX19B (a component of the nuclear pore complex known to interact with Influenza A viral polymerase (Munier et al. 2013)) (**Figure**

1C) further supporting a role for svRNA-29094 in remodeling host gene expression related to infection and host immune evasion.

In conclusion, we suggest (1) the ability of svRNA-29094 to suppress the expression of a mRNA bearing a complementary 3'UTR target site, (2) the presence of putative svRNA-29094 target sites in several host genes related to infection and host immune evasion, and (3) the identification of svRNA-29094 expression during SC2 infection in both cell culture models and patient samples by three independent laboratories (Meng et al. 2021; Neeb et al. 2022), strongly supports further characterization of svRNA-29094 and evaluation of its potential to serve as a novel therapeutic target.

Methods

Sequence analyses

To identify svRNAs expressed during SC2 infection, we obtained next-generation small RNA deep sequencing libraries corresponding to Calu-3 cells infected with an SC2 patient isolate 1, 12, and 24 hours post-infection (Devadoss et al. 2021 May 18; Wyler et al. 2021; Devadoss et al. 2022). Alignments between the SC2 genome and individual small RNA-seq reads were performed on the Alabama Supercomputer Center SGI UV 2000 and DMC cluster and obtained via Basic Local Alignment Search Tool (BLAST+) using the following parameters: 100% identity, word size = 6, un-gapped, and evalue = 0.001 (Camacho et al. 2009). All accepted BLAST+ alignments were restricted to perfect matches (100% identity) between 16 and 32 nts. The frequency of alignments to each position across the full-length genome was calculated by counting reads defined as ≥ 16 nts, perfect matches (100% identity) and svRNAs called as previously defined and employed (Roberts et al. 2013; Huang et al. 2014; Roberts et al. 2014; Amin et al. 2016; Huang, Eilbeck, et al. 2016; Cardin and Borchert 2017; Barnhill et al. 2019; Houserova et al. 2021; Coley et al. 2022; Godang et al. 2023). Publicly available next-generation small RNA deep sequencing libraries were obtained from the NCBI Sequence Read Archive (SRA) (www.ncbi.nlm.nih.gov/sra/). These included SRR1155025, SRR1155026, SRR1155027, SRR1155028, SRR1155029, and SRR1155030 (Wyler et al. 2021). Putative human 3'UTR targets were identified for svRNAs using an in-house BLAST-based pipeline previously defined and routinely employed by our lab (Borchert et al. 2009; Borchert et al. 2011; Filshtein et al. 2012; Huang, Gutierrez, et al. 2016; Chevalier and Borchert 2017; Patterson et al. 2017; Roberts and Borchert 2017; Chen et al. 2018; Roberts et al. 2018; Alexander B. Coley et al. 2022).

Luciferase assays

Antisense reporters were constructed by oligonucleotide primer extension PCR in 50 μ l reactions at standard concentrations (1.5 mM MgCl₂, 0.2 mM dNTP, 1x Biolase PCR buffer, 2.5 U Taq (Bioline USA, Inc., Randolph, MA), 0.1 μ M each primer) and using standard cycling parameters (95°C - 2 min, (95°C - 40 sec, 58°C - 40 sec, 72°C - 40 sec) x 35 cycles, 72°C - 2 min) with primers containing 5' Xho-I and 3' Not-I restriction enzyme sites. Primer sequences were:

Forward: ACTCGAGAACGTAGCACACAAAGCCAAAAACTGGGTTGTCAGGACCACCCAGAAGAAG
AAGAGCAAGAAGAAGATTG

Reverse: TCGGGCCGCAACCAGCCTCATCCACGCACAATTGGTGCCACTCTGCTGCTCCAATCTT
CTTCTTGCTCTTCTTCTTCTGGG

Following PCR amplification and subsequent digestion, amplicons were ligated into the Renilla luciferase 3'UTR of psiCHECK-2 (Promega) vector linearized with Xho-I and Not-I. The prostate cancer PC3 cell line was obtained from GenLantis (San Diego, CA) and cultured in a humidified atmosphere with 5% CO₂ at 37°C in MEM (Mediatech, Herndon, VA) supplemented with 10% fetal bovine serum (Hyclone, Logan, UT), 25 mg/ml streptomycin and 25 I.U. penicillin (Mediatech). PC3 cells were seeded in a 96-well plate (10,000 cells per well) and allowed to adhere for 4 hours. Then cells were co-transfected with 500 ng of reporter plasmid and 10 pmol of svRNA mimic following lipofectamine 2000 (Invitrogen) protocol. Cells were lysed in the plate according to standard Dual-Glo Luciferase Reporter System (Promega, Madison, WI) at 24h and 48h post-transfection. Firefly and Renilla luciferase were measured sequentially at each time point using a 96-well plate luminometer (SpectraMax iD5 Multi-mode microplate reader (Molecular Devices) and normalized to an empty vector.

Acknowledgements: We thank the University of South Alabama College of Medicine Department of Pharmacology for their ongoing support.

References

Amin SV, Roberts JT, Patterson DG, Coley AB, Allred JA, Denner JM, et al., Borchert GM. 2016. Novel small RNA (sRNA) landscape of the starvation-stress response transcriptome of *Salmonella enterica* serovar *typhimurium*. *RNA Biol* 13: 331-42. PubMed ID: [26853797](https://pubmed.ncbi.nlm.nih.gov/26853797/)

1/18/2024 - Open Access

Barnhill EC, Crucello A, Houserova D, King VM, Amin SV, Roberts JT, et al., Borchert GM. 2019. Characterization of novel small RNAs (sRNAs) contributing to the desiccation response of *Salmonella enterica* serovar *Typhimurium*. *RNA Biol* 16: 1643-1657. PubMed ID: [31390935](#)

Borchert GM, Gilmore BL, Spengler RM, Xing Y, Lanier W, Bhattacharya D, Davidson BL. 2009. Adenosine deamination in human transcripts generates novel microRNA binding sites. *Hum Mol Genet* 18: 4801-7. PubMed ID: [19776031](#)

Borchert GM, Holton NW, Larson ED. 2011. Repression of human activation induced cytidine deaminase by miR-93 and miR-155. *BMC Cancer* 11: 347. PubMed ID: [21831295](#)

Cardin SE, Borchert GM. 2017. Viral MicroRNAs, Host MicroRNAs Regulating Viruses, and Bacterial MicroRNA-Like RNAs. *Methods Mol Biol* 1617: 39-56. PubMed ID: [28540675](#)

Chang R, Mamun A, Dominic A, Le NT. 2020. SARS-CoV-2 Mediated Endothelial Dysfunction: The Potential Role of Chronic Oxidative Stress. *Front Physiol* 11: 605908. PubMed ID: [33519510](#)

Channappanavar R, Perlman S. 2017. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Semin Immunopathol* 39: 529-539. PubMed ID: [28466096](#)

Chen H, Zhang D, Zhang G, Li X, Liang Y, Kasukurthi MV, et al., Huang J. 2018. A semantics-oriented computational approach to investigate microRNA regulation on glucocorticoid resistance in pediatric acute lymphoblastic leukemia. *BMC Med Inform Decis Mak* 18: 57. PubMed ID: [30066657](#)

Chevalier D, Borchert GM. 2017. Genome-Wide Analysis of MicroRNA-Regulated Transcripts. *Methods Mol Biol* 1617: 93-107. PubMed ID: [28540679](#)

Coley AB, DeMeis JD, Chaudhary NY, Borchert GM. 2022. Small Nucleolar Derived RNAs as Regulators of Human Cancer. *Biomedicines* 10: . PubMed ID: [36009366](#)

Coley AB, Stahly AN, Kasukurthi MV, Barchie AA, Hutcheson SB, Houserova D, et al., Borchert GM. 2022. MicroRNA-like snoRNA-Derived RNAs (sdRNAs) Promote Castration-Resistant Prostate Cancer. *Cells* 11: . PubMed ID: [35455981](#)

Declercq M, Biquand E, Karim M, Pietrosemoli N, Jacob Y, Demeret C, Barbezange C, van der Werf S. 2020. Influenza A virus co-opts ERI1 exonuclease bound to histone mRNA to promote viral transcription. *Nucleic Acids Res* 48: 10428-10440. PubMed ID: [32960265](#)

Devadoss D, Acharya A, Manevski M, Houserova D, Cioffi MD, Pandey K, et al., Chand HS. 2022. Immunomodulatory LncRNA on antisense strand of ICAM-1 augments SARS-CoV-2 infection-associated airway mucoinflammatory phenotype. *iScience* 25: 104685. PubMed ID: [35789750](#)

Devadoss D, Acharya A, Manevski M, Pandey K, Borchert GM, Nair M, et al., Chand HS. 2021. Distinct Mucoinflammatory Phenotype and the Immunomodulatory Long Noncoding Transcripts Associated with SARS-CoV-2 Airway Infection. *medRxiv* : . PubMed ID: [34031668](#)

Filshtein TJ, Mackenzie CO, Dale MD, Dela-Cruz PS, Ernst DM, Frankenberger EA, et al., Larson ED. 2012. OrbId: Origin-based identification of microRNA targets. *Mob Genet Elements* 2: 184-192. PubMed ID: [23087843](#)

Godang NL, DeMeis JD, Houserova D, Chaudhary NY, Salter CJ, Xi Y, McDonald OG, Borchert GM. 2023. Global Switch from DICER-dependent MicroRNA to DICER-independent SnoRNA-derived RNA Biogenesis in Malignancy. *MicroPubl Biol* 2023: . PubMed ID: [36818311](#)

Houserova D, Dahmer DJ, Amin SV, King VM, Barnhill EC, Zambrano ME, et al., Borchert GM. 2021. Characterization of 475 Novel, Putative Small RNAs (sRNAs) in Carbon-Starved *Salmonella enterica* Serovar *Typhimurium*. *Antibiotics (Basel)* 10: . PubMed ID: [33809610](#)

Huang J, Dang J, Borchert GM, Eilbeck K, Zhang H, Xiong M, et al., Tan M. 2014. OMIT: dynamic, semi-automated ontology development for the microRNA domain. *PLoS One* 9: e100855. PubMed ID: [25025130](#)

Huang J, Eilbeck K, Smith B, Blake JA, Dou D, Huang W, et al., Tan M. 2016. The Non-Coding RNA Ontology (NCRO): a comprehensive resource for the unification of non-coding RNA biology. *J Biomed Semantics* 7: 24. PubMed ID: [27152146](#)

Huang J, Gutierrez F, Strachan HJ, Dou D, Huang W, Smith B, et al., Ruttenberg A. 2016. OmniSearch: a semantic search system based on the Ontology for MiCroRNA Target (OMIT) for microRNA-target gene interaction data. *J Biomed Semantics* 7: 25. PubMed ID: [27175225](#)

Lee RC, Feinbaum RL, Ambros V. 1993. The *C. elegans* heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. *Cell* 75: 843-54. PubMed ID: [8252621](#)

1/18/2024 - Open Access

Li J, Hu L, Liu Y, Huang L, Mu Y, Cai X, Weng C. 2015. DDX19A Senses Viral RNA and Mediates NLRP3-Dependent Inflammasome Activation. *J Immunol* 195: 5732-49. PubMed ID: [26538395](#)

Meng F, Siu GK, Mok BW, Sun J, Fung KSC, Lam JY, et al., Huang CL. 2021. Viral MicroRNAs Encoded by Nucleocapsid Gene of SARS-CoV-2 Are Detected during Infection, and Targeting Metabolic Pathways in Host Cells. *Cells* 10: . PubMed ID: [34359932](#)

Morales L, Oliveros JC, Fernandez-Delgado R, tenOever BR, Enjuanes L, Sola I. 2017. SARS-CoV-Encoded Small RNAs Contribute to Infection-Associated Lung Pathology. *Cell Host Microbe* 21: 344-355. PubMed ID: [28216251](#)

Munier S, Rolland T, Diot C, Jacob Y, Naffakh N. 2013. Exploration of binary virus-host interactions using an infectious protein complementation assay. *Mol Cell Proteomics* 12: 2845-55. PubMed ID: [23816991](#)

Neeb ZT, Ritter AJ, Chauhan LV, Katzman S, Lipkin WI, Mishra N, Sanford JR. 2022. A potential role for SARS-CoV-2 small viral RNAs in targeting host microRNAs and modulating gene expression. *Sci Rep* 12: 21694. PubMed ID: [36522444](#)

Patterson DG, Roberts JT, King VM, Houserova D, Barnhill EC, Crucello A, et al., Borchert GM. 2017. Human snoRNA-93 is processed into a microRNA-like RNA that promotes breast cancer cell invasion. *NPJ Breast Cancer* 3: 25. PubMed ID: [28702505](#)

Roberts JT, Borchert GM. 2017. Computational Prediction of MicroRNA Target Genes, Target Prediction Databases, and Web Resources. *Methods Mol Biol* 1617: 109-122. PubMed ID: [28540680](#)

Roberts JT, Cardin SE, Borchert GM. 2014. Burgeoning evidence indicates that microRNAs were initially formed from transposable element sequences. *Mob Genet Elements* 4: e29255. PubMed ID: [25054081](#)

Roberts JT, Cooper EA, Favreau CJ, Howell JS, Lane LG, Mills JE, et al., Borchert GM. 2013. Continuing analysis of microRNA origins: Formation from transposable element insertions and noncoding RNA mutations. *Mob Genet Elements* 3: e27755. PubMed ID: [24475369](#)

Roberts JT, Patterson DG, King VM, Amin SV, Polska CJ, Houserova D, et al., Borchert GM. 2018. ADAR Mediated RNA Editing Modulates MicroRNA Targeting in Human Breast Cancer. *Processes (Basel)* 6: . PubMed ID: [30197877](#)

Sadri Nahand J, Moghoofei M, Salmaninejad A, Bahmanpour Z, Karimzadeh M, Nasiri M, et al., Hamblin MR. 2020. Pathogenic role of exosomes and microRNAs in HPV-mediated inflammation and cervical cancer: A review. *Int J Cancer* 146: 305-320. PubMed ID: [31566705](#)

Samols MA, Skalsky RL, Maldonado AM, Riva A, Lopez MC, Baker HV, Renne R. 2007. Identification of cellular genes targeted by KSHV-encoded microRNAs. *PLoS Pathog* 3: e65. PubMed ID: [17500590](#)

Tazawa S, Yamato T, Fujikura H, Hiratuchi M, Itoh F, Tomae M, et al., Isaji M. 2005. SLC5A9/SGLT4, a new Na⁺-dependent glucose transporter, is an essential transporter for mannose, 1,5-anhydro-D-glucitol, and fructose. *Life Sci* 76: 1039-50. PubMed ID: [15607332](#)

Wyler E, Mösbauer K, Franke V, Diag A, Gottula LT, Arsiè R, et al., Landthaler M. 2021. Transcriptomic profiling of SARS-CoV-2 infected human cell lines identifies HSP90 as target for COVID-19 therapy. *iScience* 24: 102151. PubMed ID: [33585804](#)

Zhang JJ, Dong X, Cao YY, Yuan YD, Yang YB, Yan YQ, Akdis CA, Gao YD. 2020. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* 75: 1730-1741. PubMed ID: [32077115](#)

Zhang Q, Chen Z, Huang C, Sun J, Xue M, Feng T, et al., Dai J. 2021. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Membrane (M) and Spike (S) Proteins Antagonize Host Type I Interferon Response. *Front Cell Infect Microbiol* 11: 766922. PubMed ID: [34950606](#)

Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* 31: 3406-15. PubMed ID: [12824337](#)

Funding: Funding was provided in part by NSF grant NSF2030080 (GMB) and NSF CAREER grant 1350064 (with co-funding provided by the NSF EPSCoR program) (GMB) both awarded by the Division of Molecular and Cellular Biosciences. Graduate funding was also provided in part by Alabama Commission on Higher Education ALEPSCoR grant 210471 (JDD). The project used an instrument funded, in part, by the National Science Foundation MRI Grant No. CNS-1726069. Research reported in this publication was also supported by the National Center for Advancing Translational Research of the National Institutes of Health under award number UL1TR001417 (GMB).

1/18/2024 - Open Access

Author Contributions: Haley A Delcher: writing - original draft, writing - review editing. Jeffrey D DeMeis: writing - review editing, formal analysis. Nicole Ghobar: formal analysis, investigation. Noel L Godang: writing - review editing. Sierra L Knight: formal analysis, investigation. Shahem Y Alqudah: writing - review editing. Kevin N Nguyen: writing - review editing. Brianna C Watters: formal analysis. Glen M Borchert: conceptualization, methodology, supervision, writing - review editing.

Reviewed By: Anonymous

History: Received March 4, 2023 **Revision Received** January 12, 2024 **Accepted** January 17, 2024 **Published Online** January 18, 2024 **Indexed** February 1, 2024

Copyright: © 2024 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Delcher, HA; DeMeis, JD; Ghobar, N; Godang, NL; Knight, SL; Alqudah, SY; et al.; Borchert, GM (2024). SARS-CoV-2 small viral RNA suppresses gene expression via complementary binding to mRNA 3' UTR. *microPublication Biology*. [10.17912/micropub.biology.000790](https://doi.org/10.17912/micropub.biology.000790)