

RECEIVED: September 6, 2024

REVISED: October 30, 2024

ACCEPTED: November 20, 2024

PUBLISHED: January 9, 2025

Measurement of CP violation in $B^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays



The LHCb collaboration

E-mail: louis.gerken@cern.ch

ABSTRACT: A time-dependent, flavour-tagged measurement of CP violation is performed with $B^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays, using data collected by the LHCb detector in proton-proton collisions at a centre-of-mass energy of 13 TeV corresponding to an integrated luminosity of 6 fb^{-1} . In $B^0 \rightarrow D^+ D^-$ decays the CP -violation parameters are measured to be

$$S_{D^+ D^-} = -0.552 \pm 0.100 (\text{stat}) \pm 0.010 (\text{syst}),$$

$$C_{D^+ D^-} = 0.128 \pm 0.103 (\text{stat}) \pm 0.010 (\text{syst}).$$

In $B_s^0 \rightarrow D_s^+ D_s^-$ decays the CP -violating parameter formulation in terms of ϕ_s and $|\lambda|$ results in

$$\phi_s = -0.086 \pm 0.106 (\text{stat}) \pm 0.028 (\text{syst}) \text{ rad},$$

$$|\lambda_{D_s^+ D_s^-}| = 1.145 \pm 0.126 (\text{stat}) \pm 0.031 (\text{syst}).$$

These results represent the most precise single measurement of the CP -violation parameters in their respective channels. For the first time in a single measurement, CP symmetry is observed to be violated in $B^0 \rightarrow D^+ D^-$ decays with a significance exceeding six standard deviations.

KEYWORDS: CP Violation, Hadron-Hadron Scattering

ARXIV EPRINT: [2409.03009](https://arxiv.org/abs/2409.03009)

Contents

1	Introduction	1
2	Detector and simulation	3
3	Selection	4
4	Mass fit	6
5	Flavour tagging	6
6	Decay-time fit	8
7	Systematic uncertainties and cross-checks	10
8	Results and interpretation	12
A	Results and correlations of external parameters	14
	The LHCb collaboration	19

1 Introduction

Measurements of CP violation in $B_{(s)}^0$ mesons play a crucial role in the search for physics beyond the Standard Model (SM). With the increase in experimental precision, control over hadronic matrix elements becomes more important, which is a major challenge in most decay modes. In decays of beauty mesons to two charmed mesons $B \rightarrow DD$, this can be achieved by employing U-spin flavour symmetry and constraining the hadronic contributions by relating different CP -violation and branching fraction measurements [1–5].

The $B \rightarrow DD$ system gives access to a variety of interesting observables that probe elements of the Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [6, 7]. In $B^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays, the CP -violating weak phases β and β_s can be measured, respectively. The phases arise in the interference between the B^0 – $\bar{B}^0$ (B_s^0 – $\bar{B}_s^0$) mixing and the tree-level decay amplitudes to the $D^+ D^-$ ($D_s^+ D_s^-$) final state, leading to time-dependent CP asymmetries. The decays can also proceed through several other diagrams, as shown in figure 1. The CP asymmetries may arise from both SM contributions and new physics effects, if present.

In $B^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays, the same final state is accessible from both $B_{(s)}^0$ and $\bar{B}_{(s)}^0$ states. The partial decay rate as a function of the decay time t is given by

$$\frac{d\Gamma(t, d)}{dt} \propto e^{-t/\tau_{B_{(s)}^0}} \left(\cosh \frac{\Delta\Gamma_q t}{2} + D_f \sinh \frac{\Delta\Gamma_q t}{2} + d C_f \cos \Delta m_q t - d S_f \sin \Delta m_q t \right), \quad (1.1)$$




Figure 1. Dominant Feynman diagrams contributing to the $B^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays. The (top left) tree-level, (bottom left) exchange, (top right) penguin and (bottom right) penguin annihilation diagrams are shown.

where $\Delta\Gamma_q = \Gamma_{qL} - \Gamma_{qH}$ and $\Delta m_q = m_{qH} - m_{qL}$ are the decay-width difference and mass difference of the heavy and light B^0 ($q = d$) or B_s^0 ($q = s$) mass eigenstates, $\tau_{B(s)^0}$ is the mean lifetime of the $B(s)^0$ meson and the tag d represents the flavour at production taking the value $+1$ for a $B(s)^0$ meson and -1 for a $\bar{B}(s)^0$ meson. The CP -violation parameters are defined as

$$D_f = -\frac{2|\lambda_f| \cos \phi_q}{1 + |\lambda_f|^2}, \quad C_f = \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2}, \quad S_f = -\frac{2|\lambda_f| \sin \phi_q}{1 + |\lambda_f|^2}, \quad (1.2)$$

$$\lambda_f = \frac{q}{p} \frac{\bar{A}_f}{A_f} \quad \text{and} \quad \phi_q = -\arg \lambda_f,$$

where A_f and $\bar{A}_f$ are the decay amplitudes of $B(s)^0$ and $\bar{B}(s)^0$ to the common final state f and the ratio q/p describes mixing of the $B(s)^0$ mesons. The parameter D_f cannot be measured in B^0 decays because, at the current experimental precision, $\Delta\Gamma_d$ is compatible with zero. Thus, the decay rates for $B^0 \rightarrow D^+ D^-$ can be simplified to

$$\frac{d\Gamma(t, d)}{dt} \propto e^{-t/\tau_{B^0}} (1 + d C_{D^+ D^-} \cos \Delta m_d t - d S_{D^+ D^-} \sin \Delta m_d t). \quad (1.3)$$

If only tree-level contributions in $B^0 \rightarrow D^+ D^-$ decays are considered, direct CP violation vanishes resulting in $C_{D^+ D^-} = 0$ and $S_{D^+ D^-} = -\sin \phi_d = -\sin 2\beta$. This assumption is valid within the current experimental precision for $B^0 \rightarrow J/\psi K_S^0$ decays, where β can be

measured with high precision as recently reported by LHCb [8]. However, in $B^0 \rightarrow D^+ D^-$ measurements the loop-mediated penguin contributions shown in figure 1 cannot be neglected and an additional phase shift is measured via $\sin(2\beta + \Delta\phi_d) = -S_{D^+ D^-} / \sqrt{1 - C_{D^+ D^-}^2}$. This measurement enables higher-order corrections to the measurement of ϕ_s in $B_s^0 \rightarrow D_s^+ D_s^-$ decays to be constrained, under the assumption of U-spin flavour symmetry.

Due to the similarities of the two decay channels, a parallel measurement of the CP -violation parameters in $B^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays is performed. Both decays have been previously studied by LHCb [9, 10], while measurements of the CP -violation parameters in $B^0 \rightarrow D^+ D^-$ decays have been performed by BaBar [11] and Belle [12]. The Belle result lies outside the physically allowed region and shows a small tension with the other measurements.

This analysis uses proton-proton (pp) collision data collected by the LHCb experiment during the years 2015 to 2018 corresponding to an integrated luminosity of 6 fb^{-1} . The $B^0 \rightarrow D^+ D^-$ candidates are reconstructed through the decays $D^+ \rightarrow K^- \pi^+ \pi^+$ and $D^+ \rightarrow K^- K^+ \pi^+$.¹ These decays have the highest branching fractions into charged kaons and pions. Candidates where both $D^\pm$ mesons decay via $D^+ \rightarrow K^- K^+ \pi^+$ are not considered due to the smaller branching fraction of this mode. Similarly, one of the $D_s^\pm$ mesons from the $B_s^0 \rightarrow D_s^+ D_s^-$ candidates is always reconstructed through the decay $D_s^+ \rightarrow K^- K^+ \pi^+$ and the other is reconstructed through the decays $D_s^+ \rightarrow K^- K^+ \pi^+$, $D_s^+ \rightarrow \pi^- K^+ \pi^+$ or $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$.

Both signal channels and a dedicated $B^0 \rightarrow D_s^+ D^-$ control channel are selected by similar criteria with only minor differences as described in section 3. A mass fit is performed separately for each final state to statistically subtract the remaining background as described in section 4. The knowledge of the initial flavour of the candidates is crucial for measurements of time-dependent asymmetries in neutral B -meson decays. In section 5 the algorithms used to determine the initial flavour of the $B_{(s)}^0$ mesons are described. The decay-time fit to measure the CP -violation parameters is described in section 6 and the systematic uncertainties are discussed in section 7. In section 8 the results are presented from both this analysis and in combination with previous LHCb measurements.

2 Detector and simulation

The LHCb detector [13, 14] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [15], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 T m, and three stations of silicon-strip detectors and straw drift tubes [16] placed downstream of the magnet. The tracking system provides a measurement of the momentum, p , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/ c . The minimum distance of a track to a primary pp collision vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the

¹If not stated otherwise, charge-conjugated decays are implied.

beam, in GeV/ c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [17]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [18].

Simulation is required to model the effects of the detector acceptance and the imposed selection requirements. Samples of signal decays are used to determine the parameterisation of the signal mass distributions and decay-time resolution model. In the simulation, pp collisions are generated using PYTHIA [19] with a specific LHCb configuration [21]. Decays of unstable particles are described by EVTGEN [22], in which final-state radiation is generated using PHOTOS [23]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [24] as described in ref. [26]. The underlying pp interaction is reused multiple times, with an independently generated signal decay for each [27]. To account for differences between the distributions of particle identification (PID) variables in simulation and data, the PIDCalib package [28] is used to reweight the distributions in the simulation.

3 Selection

The online event selection is performed by a trigger [29], which consists of a hardware stage based on information from the calorimeter and muon systems, followed by a software stage which applies a full event reconstruction. At the hardware trigger stage, events are required to have a muon with high p_T or a hadron, photon or electron with high transverse energy in the calorimeters. The software trigger requires a two-, three- or four-track secondary vertex with a significant displacement from any primary pp interaction vertex. At least one charged particle must have a transverse momentum $p_T > 1.6$ GeV/ c and be inconsistent with originating from a PV. A multivariate algorithm [30, 31] is used for the identification of secondary vertices consistent with the decay of a b hadron.

In the offline selection, $D^\pm$ and $D_s^\pm$ candidates are reconstructed through their decays into the selected final-state particles, which are required to satisfy loose selection criteria on their momentum, transverse momentum and PID variables, and be inconsistent with originating from any PV. The $D^\pm$ and $D_s^\pm$ candidates should form vertices with a good fit quality and the scalar sum of transverse momenta of their three final-state particles should be greater than 1800 MeV/ c . All possible combinations of tracks forming a common vertex should have a distance of closest approach smaller than 0.5 mm. The $B_{(s)}^0$ candidates are reconstructed from two $D^\pm$ or $D_s^\pm$ candidates with opposite charges that form a good-quality vertex. The momentum vector of the $B_{(s)}^0$ candidates should point from the PV to the secondary vertex. The scalar sum of the transverse momenta of all six final-state particles is required to be greater than 5000 MeV/ c . The invariant masses of the $D^\pm$ and $D_s^\pm$ candidates are required to be within a window of ± 45 MeV/ c^2 around their known values [32]. This requirement, of about ± 4 times the mass resolution, retains almost all candidates while separating the $D^\pm$ from the $D_s^\pm$ mass region. To suppress single-charm decays of the form $B_{(s)}^0 \rightarrow D_{(s)}^+ h^+ h^- h^-$, both $D_{(s)}^+$ candidates are required to have a significant flight distance from the $B_{(s)}^0$ decay vertex ($> 5\sigma$).

In the reconstruction of the $D_{(s)}^+$ candidates, background contributions can arise from the misidentification of the final-state particles. Misidentification from a pion, kaon or proton is considered. The three-body invariant masses are recomputed to identify background decays from D^+ , D_s^+ and Λ_c^+ states. The masses for potential two-body background contributions arising from intermediate ϕ and D^0 decays are similarly computed. These background sources are suppressed by PID requirements within the mass windows of the known particle masses.

A particularly challenging background arises from the misidentification between $D^+ \rightarrow K^- \pi^+ \pi^+$ and $D_s^+ \rightarrow K^- K^+ \pi^+$ decays. The $\pi^+ \leftrightarrow K^+$ misidentification shifts the mass region of the reconstructed D^+ candidates to that of the D_s^+ or vice versa. In this case, a simple PID requirement does not provide the necessary rejection of the particularly large background contribution from $B^0 \rightarrow D_s^+ D^-$ decays. To distinguish between the two decays a boosted decision tree (BDT) algorithm is trained utilising the `xgboost` module from the `scikit-learn` package [33]. Simulated D^+ and D_s^+ decays from the $B^0 \rightarrow D^+ D^-$, $B^0 \rightarrow D_s^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ samples are used to train the BDT classifier. A k -folding procedure with $k = 5$ is used to avoid overtraining [34]. Various two- and three-body invariant masses, recomputed with different final-state particle hypotheses, are used in the training. Additionally, the flight distance of the $D_{(s)}^+$ candidates, and the PID variables of those particles that are potentially misidentified, are used. The requirements on the BDT-classifier output are chosen to suppress the D_s^+ candidates in the $B^0 \rightarrow D^+ D^-$ channel and D^+ candidates in the $B_s^0 \rightarrow D_s^+ D_s^-$ channel to negligible levels. This is verified by applying the requirements to the simulated samples, which results in the rejection of more than 99% of the respective candidates.

A second BDT classifier is trained to suppress combinatorial background. As a signal proxy, all available simulated $B^0 \rightarrow D^+ D^-$, $B^0 \rightarrow D_s^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ samples are used while the background proxy is taken from the upper-mass sideband of the data, which is defined as $m_{D_{(s)}^+ D_{(s)}^-} > 5600 \text{ MeV}/c^2$, beyond the $B_{(s)}^0$ -candidate mass fit region. The variables used in the training are all transverse momenta of intermediate and final-state particles; the flight distance and the difference in invariant mass from the known value [32] of the $D_{(s)}^+$ candidates; the angle between the $D_{(s)}^+$ flight direction and each of the decay products; the χ_{IP}^2 of the $B_{(s)}^0$ and $D_{(s)}^+$ candidates, which is the difference in the χ^2 value of the PV fit with and without the particle being considered in the calculation. Similar to the strategy used in ref. [35], the requirement on the BDT-classifier output is chosen to minimise the uncertainties on the CP -violation parameters.

The invariant mass used in the mass fits is computed from a kinematic fit to the decay chain with constraints on all charm-meson masses to improve the invariant-mass resolution of the $B_{(s)}^0$ candidates [36]. For calculation of the decay time, a constraint on the PV is used in the kinematic fit. To avoid correlations between the decay time and the invariant mass, no constraints on the charm-masses are used.

Contributions from partially reconstructed backgrounds are reduced to negligible levels by restricting the invariant mass of the B^0 candidates to lie within the range 5240–5540 MeV/c^2 . The decay-time range is chosen to be 0.3–10.3 ps, where the lower boundary is set to reduce background originating from the PV. For B_s^0 candidates the same decay-time range is chosen, but the invariant-mass range is 5300–5600 MeV/c^2 .

After the selection, multiple candidates are found in about 1% of the events. Usually, these candidates differ in just one track or PID assignment. Since it is very unlikely to find two genuine candidates in one event, only one of the candidates is chosen in a reproducible random way.

4 Mass fit

An extended unbinned maximum-likelihood fit to the invariant mass of the $B_{(s)}^0$ candidates is performed to extract per-event weights via the *sPlot* technique [37]. These weights are used in the decay-time fit to statistically subtract the background. Pseudoexperiment studies indicate that any residual correlation between the decay time and the mass introduces no meaningful bias into the CP -violation measurement.

The mass model in the $B^0 \rightarrow D^+ D^-$ channel consists of a signal component and two background components to model $B_s^0 \rightarrow D^+ D^-$ decays and the combinatorial background. A double-sided Hypatia probability density function (PDF) [38] is used to model the signal component. The shape parameters are determined by a fit to simulated $B^0 \rightarrow D^+ D^-$ decays and fixed in the fit to data, while the peak position and width of the distribution are allowed to vary. The same model is used for the $B_s^0 \rightarrow D^+ D^-$ component with a shift of the peak position by the known mass difference between the B^0 and B_s^0 mesons [32]. An exponential PDF is used to model the combinatorial background.

The mass model in the $B_s^0 \rightarrow D_s^+ D_s^-$ channel consists only of a signal component and a combinatorial background component, which are parameterised as in the $B^0 \rightarrow D^+ D^-$ fit. Mass fits are performed separately for each final state. Figures 2 and 3 show the results of the fits to all $B^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$ final states, respectively. The fits yield an overall number of $5\,695 \pm 100$ $B^0 \rightarrow D^+ D^-$ and $13\,313 \pm 135$ $B_s^0 \rightarrow D_s^+ D_s^-$ signal decays.

5 Flavour tagging

For time-dependent CP violation measurements of neutral B mesons, the flavour of the meson at production is required. At LHCb the method used to determine the initial flavour is called flavour tagging. These algorithms exploit the fact that in pp collisions, b and $\bar{b}$ quarks are almost exclusively produced in pairs. When the b quark forms a $\bar{B}$ meson (and similarly the $\bar{b}$ quark forms a B meson), additional particles are produced in the fragmentation process. From the charges and types of these particles, the flavour of the signal B meson at production can be inferred. The tagging algorithm that uses charged pions or protons from the fragmentation process of the b quark that leads to the $\bar{B}^0$ signal is called the same-side (SS) tagger [39]. In the case of signal $\bar{B}_s^0$ mesons, charged kaons are used by the SS tagger [40]. The opposite-side (OS) tagger uses information from electrons and muons from semileptonic b decays, kaons from the $b \rightarrow c \rightarrow s$ decay chain, secondary charm hadrons and the charges of tracks from the secondary vertex of the other b -hadron decay [41, 42]. Each algorithm i provides individual tag decisions, d_i , and a predicted mistag, η_i , which is an estimate of the probability that the tag decision is wrong. The tag decision takes the values -1 for a $\bar{B}$ meson, 1 for a B meson and 0 if no tag decision can be made. The predicted mistag ranges from 0 to 0.5 and takes the value of 0.5 for untagged events. Each predicted mistag




Figure 2. Invariant-mass distribution of $B^0 \rightarrow D^+ D^-$ decays. The data are shown as points and the full PDF is shown as a solid-blue line for (left) both $D^\pm$ candidates decaying through $D^+ \rightarrow K^- \pi^+ \pi^+$ and (right) one $D^\pm$ candidate decaying through $D^+ \rightarrow K^- K^+ \pi^+$.




Figure 3. Invariant-mass distribution of $B_s^0 \rightarrow D_s^+ D_s^-$ decays. The data are shown as points and the full PDF is shown as a solid-blue line for (left) both $D_s^\pm$ candidates decaying through $D_s^+ \rightarrow K^- K^+ \pi^+$, (right) one $D_s^\pm$ candidate decaying through $D_s^+ \rightarrow \pi^- K^+ \pi^+$ and (bottom) one $D_s^\pm$ candidate decaying through $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$.

distribution is given by the output of a BDT that is trained on flavour-specific decays [43] and has to be calibrated to represent the mistag probability, $\omega_i(\eta_i)$, in the signal decay. Flavour-specific control channels with kinematics similar to the signal are used to obtain

a calibration curve. This is found to be well-described by a linear function. Following calibration, the individual taggers are combined separately for OS and SS cases, and the resulting mistag distributions are recalibrated. These calibrations are used in the decay-time fit to determine the CP -violation parameters to which the uncertainties on the calibration parameters are propagated through means of a Gaussian constraint.

To calibrate the SS and OS taggers of the $B^0 \rightarrow D^+ D^-$ channel, as well as the OS tagger of the $B_s^0 \rightarrow D_s^+ D_s^-$ channel, $B^0 \rightarrow D_s^+ D^-$ decays are used. These have very similar kinematics to the signal decays and the selection is very similar, as described in section 3. The SS kaon tagger used for $B_s^0 \rightarrow D_s^+ D_s^-$ decays is calibrated with the $B_s^0 \rightarrow D_s^- \pi^+$ channel. A reweighting process is applied to ensure the calibration sample matches the distributions of the signal channel in the transverse momentum of the B_s^0 meson, the pseudorapidity, the number of tracks and the number of PVs. Additionally, the compatibility of the calibration between $B_s^0 \rightarrow D_s^- \pi^+$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays is verified by comparing the calibration parameters determined using simulation.

The performance of the tagging algorithms is measured by the tagging power $\epsilon_{\text{tag}} D^2$, where ϵ_{tag} is the fraction of tagged candidates and $D = 1 - 2\omega$ is the dilution factor introduced by the mistag probability, ω . The tagging power is a statistical dilution factor due to imperfect tagging, equivalent to an efficiency with respect to a sample with perfect tagging. Overall tagging powers of $(6.28 \pm 0.11)\%$ in $B^0 \rightarrow D^+ D^-$ and $(5.60 \pm 0.07)\%$ in $B_s^0 \rightarrow D_s^+ D_s^-$ decays are achieved.

6 Decay-time fit

An unbinned maximum-likelihood fit to the signal-weighted decay-time distribution is performed to determine the CP -violation parameters. In order to avoid experimenter bias, the values of the CP -violation parameters were not examined until the full procedure had been finalised.

The measured decay-time distribution of the $B_{(s)}^0$ candidates given the tag decisions $\vec{d} = (d_{\text{OS}}, d_{\text{SS}})$ and predicted mistags $\vec{\eta} = (\eta_{\text{OS}}, \eta_{\text{SS}})$ is described by the PDF

$$\mathcal{P}(t, \vec{d} | \vec{\eta}) = \epsilon(t) \cdot \left(\mathcal{B}(t', \vec{d} | \vec{\eta}) \otimes \mathcal{R}(t - t') \right), \quad (6.1)$$

where $\mathcal{B}(t', \vec{d} | \vec{\eta})$ describes the distribution of the true decay time t' , which is convolved with the decay-time resolution function $\mathcal{R}(t - t')$, and the acceptance function $\epsilon(t)$ describes the total efficiency as a function of the reconstructed decay time. The PDF describing the decay-time distribution can be deduced from eq. (1.1) and takes the general form

$$\begin{aligned} \mathcal{B}(t', \vec{d} | \vec{\eta}) \propto e^{-t'/\tau} & \left(C_{\cosh}^{\text{eff}}(\vec{d} | \vec{\eta}) \cosh \frac{\Delta\Gamma_q t'}{2} + C_{\sinh}^{\text{eff}}(\vec{d} | \vec{\eta}) \sinh \frac{\Delta\Gamma_q t'}{2} \right. \\ & \left. - C_{\cos}^{\text{eff}}(\vec{d} | \vec{\eta}) \cos \Delta m_q t' + C_{\sin}^{\text{eff}}(\vec{d} | \vec{\eta}) \sin \Delta m_q t' \right). \end{aligned} \quad (6.2)$$

The effective coefficients are given by

$$\begin{aligned} C_{\cosh}^{\text{eff}} &= \Sigma(\vec{d} | \vec{\eta}) + A_{\text{prod}} \Delta(\vec{d} | \vec{\eta}), & C_{\cos}^{\text{eff}} &= C_f \left(\Delta(\vec{d} | \vec{\eta}) + A_{\text{prod}} \Sigma(\vec{d} | \vec{\eta}) \right), \\ C_{\sinh}^{\text{eff}} &= D_f \left(\Sigma(\vec{d} | \vec{\eta}) + A_{\text{prod}} \Delta(\vec{d} | \vec{\eta}) \right), & C_{\sin}^{\text{eff}} &= S_f \left(\Delta(\vec{d} | \vec{\eta}) + A_{\text{prod}} \Sigma(\vec{d} | \vec{\eta}) \right), \end{aligned} \quad (6.3)$$




Figure 4. Decay-time distribution of (left) $B^0 \rightarrow D^+ D^-$ and (right) $B_s^0 \rightarrow D_s^+ D_s^-$ candidates. The background-subtracted data are shown as points and the projection of the PDF is shown as a solid blue line.

where the production asymmetry $A_{\text{prod}} = (N_{\bar{B}_{(s)}} - N_{B_{(s)}})/(N_{\bar{B}_{(s)}} + N_{B_{(s)}})$ represents the difference in the production rates of $\bar{B}_{(s)}^0$ and $B_{(s)}^0$ mesons. The functions

$$\begin{aligned} \Sigma(\vec{d}, \vec{\eta}) &= P(\vec{d}, \vec{\eta} | \bar{B}_{(s)}^0) + P(\vec{d}, \vec{\eta} | B_{(s)}^0) \quad \text{and} \\ \Delta(\vec{d}, \vec{\eta}) &= P(\vec{d}, \vec{\eta} | \bar{B}_{(s)}^0) - P(\vec{d}, \vec{\eta} | B_{(s)}^0) \end{aligned} \quad (6.4)$$

are dependent on the tagging calibration parameters, where $P(\vec{d}, \vec{\eta} | B_{(s)}^0)$ and $P(\vec{d}, \vec{\eta} | \bar{B}_{(s)}^0)$ are the probabilities of observing the tagging decisions $\vec{d}$ and the predicted mistags $\vec{\eta}$, given the true flavour $B_{(s)}^0$ or $\bar{B}_{(s)}^0$, respectively.

$B^0 \rightarrow D^+ D^-$

The decay-time fit of $B^0 \rightarrow D^+ D^-$ decays is insensitive to $C_{\text{sinh}}^{\text{eff}}$ under the assumption that $\Delta\Gamma_d$ is zero. Moreover, due to the long oscillation period of the B^0 mesons, the decay-time resolution of around 52 fs has a very small impact on the CP -violation parameters. The decay-time resolution model consists of three Gaussian functions that have a common mean and different widths. The parameters of the model are determined from simulation and fixed in the fit to data.

The selection and reconstruction efficiency depends on the B^0 decay time due to displacement requirements made on the final-state particles and a decrease in the reconstruction efficiency for tracks with large impact parameter with respect to the beamline [44]. The decay-time dependent efficiency is modeled by cubic-spline functions [45] with five knots at (0.3, 0.5, 2.7, 6.3, 10.3) ps, whose positions were determined using simulation. The spline coefficients are free to vary in the fit.

Gaussian constraints are used to account for the uncertainties on the tagging calibration parameters, the B^0 lifetime, the oscillation frequency, Δm_d , and the production asymmetry. The world-average values are used for the external parameters [46], while the production asymmetry is taken from a similar time-dependent analysis of $B^0 \rightarrow D^{*\pm} D^\mp$ decays [47]. The tagging efficiencies are free to vary in the decay-time fit. Figure 4 (left) shows the results of the decay-time fit for this channel.

$$B_s^0 \rightarrow D_s^+ D_s^-$$

In the decay-time fit of $B_s^0 \rightarrow D_s^+ D_s^-$ decays, the hyperbolic terms of eq. (6.2) can be measured provided that $\Delta\Gamma_s$ is not zero. Moreover, the definitions from eq. (1.2) are used to directly determine the parameters ϕ_s and $|\lambda|$. The acceptance function, the tagging parameters and external parameters are treated in the same way as for the $B^0 \rightarrow D^+ D^-$ decays. In addition to the lifetime and the oscillation frequency, Δm_s , the decay-width difference $\Delta\Gamma_s$ is constrained in the fit to the world-average value [46]. The value of the production asymmetry is taken from the control channel $B_s^0 \rightarrow D_s^- \pi^+$ as described in ref. [48].

Due to the high oscillation frequency of the B_s^0 meson, the decay-time resolution plays an important role. A per-event decay-time resolution is determined based on the per-event decay-time uncertainty estimated from the vertex fit, which is calibrated using a sample of $D_s^- \pi^+$ candidates, with $D_s^- \rightarrow \phi(K^+ K^-) \pi^-$, and additional requirements imposed to suppress candidates produced in B decays to negligible levels. The measured decay time of the remaining candidates, which originate from the PV, is consistent with zero, and their distribution is used to assess resolution and bias effects. A linear fit to the measured and predicted decay-time resolution is performed. A scale factor is then applied to translate the resulting calibration to the signal $B_s^0 \rightarrow D_s^+ D_s^-$ mode. It is determined by comparing the decay-time resolution of $B_s^0 \rightarrow D_s^- \pi^+$ and $B_s^0 \rightarrow D_s^+ D_s^-$ decays in simulation. Figure 4 (right) shows the results of the decay-time fit for this channel.

The decay-time-dependent CP asymmetry and the projection of the PDF are shown in figure 5 for (left) $B^0 \rightarrow D^+ D^-$ and (right) $B_s^0 \rightarrow D_s^+ D_s^-$ decays. The CP asymmetry in each decay-time bin is given by $A^{CP} = -\left(\sum_j w_j d_j D_j\right) / \left(\sum_j w_j D_j^2\right)$ with the tagging decision d_j , the tagging dilution D_j and the signal weight w_j obtained by the *sPlot* method [8], for each candidate j .

7 Systematic uncertainties and cross-checks

A variety of cross-checks are performed and potential sources of systematic uncertainties are considered.

The decay-time fit is performed on a simulated $B^0 \rightarrow D^+ D^-$ sample using the same strategy for the tagging calibration as for the fit to data. A second fit is performed where instead of the reconstructed tagging, the truth information of the initial flavour of the B^0 mesons is used. Both results of the CP -violation parameters agree with the generated values.

The decay-time fit is performed on several subsets of the data to test the consistency of the results. The data subdivision is done according to the final state, magnet polarity, years of data taking and tagging information (OS only or SS only). Consistent results are found in all cases.

A bootstrapping procedure [49] is used to cross-check the statistical uncertainty from the decay-time fit to data. A data set is created by randomly drawing candidates from the original sample until a certain number of candidates is reached that itself is drawn from a Poisson distribution with the expected number of candidates matching the original data sample. This entails that the same candidate can be drawn multiple times. The mass and decay-time fits are performed on this data set to first statistically subtract the background




Figure 5. Decay-time-dependent CP asymmetry of (left) $B^0 \rightarrow D^+ D^-$ and (right) $B_s^0 \rightarrow D_s^+ D_s^-$ candidates. The asymmetry in the background-subtracted data is shown as points and the projection of the PDF is shown as a solid blue line. Due to the high oscillation frequency of the B_s^0 mesons, the corresponding distribution is folded onto one oscillation period.

and then determine the CP -violation parameters. The residual of the fit result with respect to the baseline fit is stored and the whole procedure is repeated until the distribution of the residuals is not significantly affected by statistical fluctuations. The statistical uncertainties from the fits to data are shown to be accurate as they are consistent with the standard deviations of the residuals, and the correlation coefficients lie within expectations.

A decay-time fit with a different set of knots for the acceptance function is performed. The difference in the results with respect to the baseline fit is assigned as a systematic uncertainty.

To test the fit strategy, pseudoexperiments are performed. In each pseudoexperiment, the mass and decay time are generated using the results of the baseline fit to data. The background contributions are generated with a specific time dependence, assuming CP symmetry for the $B_s^0 \rightarrow D^+ D^-$ background. Similar to the bootstrapping procedure, the baseline fitting procedure is performed on the pseudoexperiments and the residuals are collected. For $B^0 \rightarrow D^+ D^-$ decays, the mean values of the results are found to be consistent with the input values within the statistical uncertainties, while the fits to the $B_s^0 \rightarrow D_s^+ D_s^-$ pseudoexperiments show a small bias of -0.002 in ϕ_s and 0.008 in $|\lambda_{D_s^+ D_s^-}|$. This is of the order of a few percent of the statistical uncertainty and is subtracted from the biases found in the following studies.

The following systematic uncertainties are determined using the same procedure, with the only difference being that an alternative model is used to generate pseudoexperiments in each case. A bias in the distribution of the residuals is assigned as a systematic uncertainty.

The sum of two Crystal Ball functions [50], with parameters obtained from a fit to simulation, is used in the pseudoexperiments to test the choice of the signal mass model.

Since $\Delta\Gamma_d$ is fixed to zero in the decay-time fit of $B^0 \rightarrow D^+ D^-$ decays, a systematic uncertainty is assigned for this assumption. The value of $\Delta\Gamma_d$ is varied in the pseudoexperiments from the assumed value of zero by $\pm 1\sigma$, where σ is the uncertainty of the world average value of $\Delta\Gamma_d$ [32]. The value of $D_{D^+ D^-}$ is calculated from the normalisation

Source	$S_{D^+D^-}$	$C_{D^+D^-}$	ϕ_s [rad]	$ \lambda_{D_s^+D_s^-} $
Mass model	0.001	0.005	0.003	0.005
$\Delta\Gamma$	0.010	0.005	—	—
Decay-time resolution	0.002	0.007	0.011	0.027
Decay-time bias	—	—	0.026	0.014
Acceptance function	0.001	0.001	< 0.001	0.001
Total	0.010	0.010	0.028	0.031

Table 1. Systematic uncertainties for the $B^0 \rightarrow D^+D^-$ and $B_s^0 \rightarrow D_s^+D_s^-$ channel. A dash (—) is used to denote that a systematic has not been evaluated. The total systematic uncertainty is the quadratic sum of the individual uncertainties.

condition $D_{D^+D^-} = \pm\sqrt{1 - S_{D^+D^-}^2 - C_{D^+D^-}^2}$ and the largest deviation is assigned as the systematic uncertainty.

In the $B^0 \rightarrow D^+D^-$ channel the decay-time-resolution model is determined on simulation. Due to differences between simulation and data the resolution could be underestimated. The effect of underestimating the resolution is tested by increasing the width of the resolution function by 10% in the pseudoexperiments, which corresponds to the level measured in the B_s^0 system. It is found to be small and no further studies are considered.

In the $B_s^0 \rightarrow D_s^+D_s^-$ channel, $D_s^-\pi^+$ candidates originating from the PV are used to determine a per-event resolution calibration. Only $D_s^- \rightarrow \phi(K^+K^-)\pi^-$ decays are used and assumed to represent the resolution of the whole sample. A second calibration is obtained using a sample of $D_s^- \rightarrow K^+K^-\pi^-$ decays without specific requirements on the intermediate decays and used in the pseudoexperiments to assign a systematic uncertainty.

A decay-time bias caused by the misalignment of the vertex detector was observed in other LHCb analyses of data taken during the same period [8, 48] and confirmed in the present analysis. Due to the low oscillation frequency of B^0 mesons, this has a negligible effect on the measurement of the CP -violation parameters, as shown in ref. [48] and so is not evaluated here. However, in B_s^0 decays, this bias could have a significant impact on the measurement. To evaluate the effect, the mean of the resolution function in the generation of the pseudoexperiments is set to the largest observed bias.

The individual systematic uncertainties on the CP -violation parameters are reported in table 1 and summed in quadrature. Compared to the previous LHCb results [9, 10], the systematic uncertainties in the $B^0 \rightarrow D^+D^-$ measurement are significantly reduced, while they are comparable for the $B_s^0 \rightarrow D_s^+D_s^-$ measurement. The reduction in the first measurement is achieved by the improved suppression of backgrounds from single-charm, misidentified and partially reconstructed decays. They are the leading sources of systematic uncertainties in the previous measurement and are reduced to negligible levels in this analysis.

8 Results and interpretation

A flavour-tagged time-dependent analysis of $B^0 \rightarrow D^+D^-$ and $B_s^0 \rightarrow D_s^+D_s^-$ decays is performed using proton-proton collision data collected by the LHCb experiment during the

years 2015 to 2018, corresponding to an integrated luminosity of 6 fb^{-1} . Approximately 5 700 $B^0 \rightarrow D^+ D^-$ signal candidates are observed. A fit to their decay-time distribution, including evaluation of systematic uncertainties, gives the final results

$$\begin{aligned} S_{D^+ D^-} &= -0.552 \pm 0.100 \text{ (stat)} \pm 0.010 \text{ (syst)}, \\ C_{D^+ D^-} &= 0.128 \pm 0.103 \text{ (stat)} \pm 0.010 \text{ (syst)}, \end{aligned}$$

with a statistical correlation between the two parameters of $\rho(S_{D^+ D^-}, C_{D^+ D^-}) = 0.472$. The results and correlations of the external parameters from the decay-time fit are presented in appendix A. Wilks' theorem [51] is used to determine the significance of the result, excluding systematic uncertainties. The hypothesis of CP symmetry, corresponding to $S_{D^+ D^-} = C_{D^+ D^-} = 0$, can be rejected by more than six standard deviations. The values are consistent with previous results from LHCb and BaBar [11], which correspond to a small contribution from higher-order SM corrections. Thus, this measurement will move the world average further away from the Belle measurement, which lies outside the physical region [12].

The result is combined with the previous LHCb measurement in this channel [9] using the **GammaCombo** package [52]. Due to the small effect of the external parameters on the result, the two measurements are assumed to be uncorrelated and the combined values are

$$\begin{aligned} S_{D^+ D^-} &= -0.549 \pm 0.085 \text{ (stat)} \pm 0.015 \text{ (syst)}, \\ C_{D^+ D^-} &= 0.162 \pm 0.088 \text{ (stat)} \pm 0.009 \text{ (syst)}, \end{aligned}$$

with a statistical correlation between the two parameters of $\rho(S_{D^+ D^-}, C_{D^+ D^-}) = 0.474$.

Approximately 13 000 $B_s^0 \rightarrow D_s^+ D_s^-$ signal candidates are observed and the final results of the decay-time fit and the systematic uncertainties are

$$\begin{aligned} \phi_s &= -0.086 \pm 0.106 \text{ (stat)} \pm 0.028 \text{ (syst)} \text{ rad}, \\ |\lambda_{D_s^+ D_s^-}| &= 1.145 \pm 0.126 \text{ (stat)} \pm 0.031 \text{ (syst)}, \end{aligned}$$

with a statistical correlation between the two parameters of $\rho(\phi_s, |\lambda_{D_s^+ D_s^-}|) = -0.007$. Further information on the results of the decay-time fit is shown in appendix A. This result is consistent with, and more precise than, the previous LHCb measurement [10]. The combination with the previous LHCb measurement, following the same strategy as for the $B^0 \rightarrow D^+ D^-$ decays, yields the values

$$\begin{aligned} \phi_s &= -0.055 \pm 0.090 \text{ (stat)} \pm 0.021 \text{ (syst)} \text{ rad}, \\ |\lambda_{D_s^+ D_s^-}| &= 1.054 \pm 0.099 \text{ (stat)} \pm 0.020 \text{ (syst)}, \end{aligned}$$

with a statistical correlation between the two parameters of $\rho(\phi_s, |\lambda_{D_s^+ D_s^-}|) = 0.005$. The values are consistent with CP symmetry in the $B_s^0 \rightarrow D_s^+ D_s^-$ channel.

These results can be used in combination with other $B \rightarrow DD$ measurements to perform a global analysis and extract SM parameters as has previously been performed in ref. [5]. They represent the most precise single measurements of the CP -violation parameters in their respective channels and the combined results supersede the previous LHCb measurements. For the first time, CP symmetry can be excluded by more than six standard deviations in a single measurement of $B^0 \rightarrow D^+ D^-$ decays.

Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (The Netherlands); MNiSW and NCN (Poland); MCID/IFA (Romania); MICIU and AEI (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (U.S.A.). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), and Polish WLCG (Poland). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); Minciencias (Colombia); EPLANET, Marie Skłodowska-Curie Actions, ERC and NextGenerationEU (European Union); A*MiDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); AvH Foundation (Germany); ICSC (Italy); Severo Ochoa and María de Maeztu Units of Excellence, GVA, XuntaGal, GENCAT, InTalent-Inditex and Prog. Atracción Talento CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

A Results and correlations of external parameters

Parameter	Input value	Fit result
$\Delta m_d [\text{ps}^{-1}]$	0.5065 ± 0.0019	0.5065 ± 0.0019
$\tau_{B^0} [\text{ps}]$	1.519 ± 0.004	1.519 ± 0.004

Table 2. Results of the external parameters from the decay-time fit to $B^0 \rightarrow D^+ D^-$ data.

	$S_{D^+ D^-}$	$C_{D^+ D^-}$	Δm_d	τ_{B^0}
$S_{D^+ D^-}$	1.000	0.472	-0.014	< 0.001
$C_{D^+ D^-}$		1.000	-0.022	< 0.001
Δm_d			1.000	< 0.001
τ_{B^0}				1.000

Table 3. Correlation matrix of the CP parameters and the external parameters from the decay-time fit to $B^0 \rightarrow D^+ D^-$ data.

Parameter	Input value	Fit result
$\Delta\Gamma_s$ [ps ⁻¹]	0.083 ± 0.005	0.083 ± 0.005
Δm_s [ps ⁻¹]	17.765 ± 0.006	17.765 ± 0.006
$\tau_{B_s^0}$ [ps]	1.521 ± 0.005	1.521 ± 0.005

Table 4. Results of the external parameters from the decay-time fit to $B_s^0 \rightarrow D_s^+ D_s^-$ data.

	ϕ_s	$\lambda_{D_s^+ D_s^-}$	Δm_s	$\Delta\Gamma_s$	$\tau_{B_s^0}$
ϕ_s	1.000	-0.007	-0.010	0.001	< 0.001
$\lambda_{D_s^+ D_s^-}$		1.000	-0.018	-0.010	< 0.001
Δm_s			1.000	< 0.001	< 0.001
$\Delta\Gamma_s$				1.000	< 0.001
$\tau_{B_s^0}$					1.000

Table 5. Correlation matrix of the CP parameters and the external parameters from the decay-time fit to $B_s^0 \rightarrow D_s^+ D_s^-$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] R. Fleischer, *Exploring CP violation and penguin effects through $B_d^0 \rightarrow D^+ D^-$ and $B_s^0 \rightarrow D_s^+ D_s^-$* , *Eur. Phys. J. C* **51** (2007) 849 [[arXiv:0705.4421](https://arxiv.org/abs/0705.4421)] [[INSPIRE](#)].
- [2] R. Fleischer, *Extracting γ from $B_{s(d)} \rightarrow J/\psi K_S$ and $B_{d(s)} \rightarrow D_{d(s)}^+ D_{d(s)}^-$* , *Eur. Phys. J. C* **10** (1999) 299 [[hep-ph/9903455](https://arxiv.org/abs/hep-ph/9903455)] [[INSPIRE](#)].
- [3] M. Jung and S. Schacht, *Standard model predictions and new physics sensitivity in $B \rightarrow DD$ decays*, *Phys. Rev. D* **91** (2015) 034027 [[arXiv:1410.8396](https://arxiv.org/abs/1410.8396)] [[INSPIRE](#)].
- [4] L. Bel, K. De Bruyn, R. Fleischer, M. Mulder and N. Tuning, *Anatomy of $B \rightarrow D\bar{D}$ decays*, *JHEP* **07** (2015) 108 [[arXiv:1505.01361](https://arxiv.org/abs/1505.01361)] [[INSPIRE](#)].
- [5] J. Davies, M. Jung and S. Schacht, *$\bar{B} \rightarrow \bar{D}D$ decays and the extraction of f_d/f_u at hadron colliders*, *JHEP* **01** (2024) 191 [[arXiv:2311.16952](https://arxiv.org/abs/2311.16952)] [[INSPIRE](#)].
- [6] N. Cabibbo, *Unitary Symmetry and Leptonic Decays*, *Phys. Rev. Lett.* **10** (1963) 531 [[INSPIRE](#)].
- [7] M. Kobayashi and T. Maskawa, *CP Violation in the Renormalizable Theory of Weak Interaction*, *Prog. Theor. Phys.* **49** (1973) 652 [[INSPIRE](#)].
- [8] LHCb collaboration, *Measurement of CP Violation in $B^0 \rightarrow \psi(\rightarrow \ell^+ \ell^-) K_S^0(\rightarrow \pi^+ \pi^-)$ Decays*, *Phys. Rev. Lett.* **132** (2024) 021801 [LHCb-PAPER-2023-013] [CERN-EP-2023-177] [[arXiv:2309.09728](https://arxiv.org/abs/2309.09728)] [[INSPIRE](#)].
- [9] LHCb collaboration, *Measurement of CP violation in $B^0 \rightarrow D^+ D^-$ decays*, *Phys. Rev. Lett.* **117** (2016) 261801 [LHCb-PAPER-2016-037] [CERN-EP-2016-203] [[arXiv:1608.06620](https://arxiv.org/abs/1608.06620)] [[INSPIRE](#)].

- [10] LHCb collaboration, *Measurement of the CP-violating phase ϕ_s in $\bar{B}_s^0 \rightarrow D_s^+ D_s^-$ decays*, *Phys. Rev. Lett.* **113** (2014) 211801 [LHCb-PAPER-2014-051] [CERN-PH-EP-2014-223] [[arXiv:1409.4619](#)] [[INSPIRE](#)].
- [11] BABAR collaboration, *Measurements of time-dependent CP asymmetries in $B^0 \rightarrow D^{(*)+} D^{(*)-}$ decays*, *Phys. Rev. D* **79** (2009) 032002 [[arXiv:0808.1866](#)] [[INSPIRE](#)].
- [12] BELLE collaboration, *Measurements of Branching Fractions and Time-dependent CP Violating Asymmetries in $B^0 \rightarrow D^{(*)\pm} D^\mp$ Decays*, *Phys. Rev. D* **85** (2012) 091106 [[arXiv:1203.6647](#)] [[INSPIRE](#)].
- [13] LHCb collaboration, *The LHCb Detector at the LHC*, *2008 JINST* **3** S08005 [[INSPIRE](#)].
- [14] LHCb collaboration, *LHCb Detector Performance*, *Int. J. Mod. Phys. A* **30** (2015) 1530022 [LHCb-DP-2014-002] [CERN-PH-EP-2014-290] [[arXiv:1412.6352](#)] [[INSPIRE](#)].
- [15] R. Aaij et al., *Performance of the LHCb Vertex Locator*, *2014 JINST* **9** P09007 [LHCb-DP-2014-001] [[arXiv:1405.7808](#)] [[INSPIRE](#)].
- [16] P. d'Argent et al., *Improved performance of the LHCb Outer Tracker in LHC Run 2*, *2017 JINST* **12** P11016 [LHCb-DP-2017-001] [[arXiv:1708.00819](#)] [[INSPIRE](#)].
- [17] LHCb RICH collaboration, *Performance of the LHCb RICH detector at the LHC*, *Eur. Phys. J. C* **73** (2013) 2431 [LHCb-DP-2012-003] [[arXiv:1211.6759](#)] [[INSPIRE](#)].
- [18] A.A. Alves Jr. et al., *Performance of the LHCb muon system*, *2013 JINST* **8** P02022 [LHCb-DP-2012-002] [[arXiv:1211.1346](#)] [[INSPIRE](#)].
- [19] T. Sjöstrand, S. Mrenna and P.Z. Skands, *A Brief Introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852 [[arXiv:0710.3820](#)] [[INSPIRE](#)].
- [20] T. Sjöstrand, S. Mrenna and P.Z. Skands, *PYTHIA 6.4 Physics and Manual*, *JHEP* **05** (2006) 026 [[hep-ph/0603175](#)] [[INSPIRE](#)].
- [21] I. Belyaev et al., *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, *J. Phys. Conf. Ser.* **331** (2011) 032047 [[INSPIRE](#)].
- [22] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [23] N. Davidson, T. Przydzinski and Z. Was, *PHOTOS interface in C++: Technical and Physics Documentation*, *Comput. Phys. Commun.* **199** (2016) 86 [[arXiv:1011.0937](#)] [[INSPIRE](#)].
- [24] J. Allison et al., *Geant4 developments and applications*, *IEEE Trans. Nucl. Sci.* **53** (2006) 270 [[INSPIRE](#)].
- [25] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [26] M. Clemencic et al., *The LHCb simulation application, Gauss: Design, evolution and experience*, *J. Phys. Conf. Ser.* **331** (2011) 032023 [[INSPIRE](#)].
- [27] D. Müller, M. Clemencic, G. Corti and M. Gersabeck, *ReDecay: A novel approach to speed up the simulation at LHCb*, *Eur. Phys. J. C* **78** (2018) 1009 [LHCb-DP-2018-004] [[arXiv:1810.10362](#)] [[INSPIRE](#)].
- [28] R. Aaij et al., *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2*, *Eur. Phys. J. Tech. Instrum.* **6** (2019) 1 [LHCb-DP-2018-001] [[arXiv:1803.00824](#)] [[INSPIRE](#)].

- [29] R. Aaij et al., *The LHCb Trigger and its Performance in 2011*, [2013 JINST 8 P04022](#) [LHCb-DP-2012-004] [[arXiv:1211.3055](#)] [[INSPIRE](#)].
- [30] V.V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, [2013 JINST 8 P02013](#) [[arXiv:1210.6861](#)] [[INSPIRE](#)].
- [31] T. Likhomanenko et al., *LHCb Topological Trigger Reoptimization*, *J. Phys. Conf. Ser.* **664** (2015) 082025 [[arXiv:1510.00572](#)] [[INSPIRE](#)].
- [32] PARTICLE DATA collaboration, *Review of particle physics*, *Phys. Rev. D* **110** (2024) 030001 [[INSPIRE](#)].
- [33] F. Pedregosa et al., *Scikit-learn: Machine Learning in Python*, *J. Mach. Learn. Res.* **12** (2011) 2825 [[arXiv:1201.0490](#)] [[INSPIRE](#)].
- [34] A. Blum, A. Kalai and J. Langford, *Beating the hold-out: bounds for K-fold and progressive cross-validation*, in the proceedings of the *Twelfth Annual Conference on Computational Learning Theory (COLT '99)*, Santa Cruz, CA, U.S.A., 7–9 July 1999, Association for Computing Machinery, New York, NY, U.S.A. (1999), pp. 203–208 [[DOI:10.1145/307400.307439](#)].
- [35] LHCb collaboration, *First study of the CP-violating phase and decay-width difference in $B_s^0 \rightarrow \psi(2S)\phi$ decays*, *Phys. Lett. B* **762** (2016) 253 [LHCb-PAPER-2016-027] [CERN-EP-2016-192] [[arXiv:1608.04855](#)] [[INSPIRE](#)].
- [36] W.D. Hulsbergen, *Decay chain fitting with a Kalman filter*, *Nucl. Instrum. Meth. A* **552** (2005) 566 [[physics/0503191](#)] [[INSPIRE](#)].
- [37] M. Pivk and F.R. Le Diberder, *SPlot: A Statistical tool to unfold data distributions*, *Nucl. Instrum. Meth. A* **555** (2005) 356 [[physics/0402083](#)] [[INSPIRE](#)].
- [38] D. Martínez Santos and F. Dupertuis, *Mass distributions marginalized over per-event errors*, *Nucl. Instrum. Meth. A* **764** (2014) 150 [[arXiv:1312.5000](#)] [[INSPIRE](#)].
- [39] LHCb collaboration, *New algorithms for identifying the flavour of B^0 mesons using pions and protons*, *Eur. Phys. J. C* **77** (2017) 238 [LHCb-PAPER-2016-039] [CERN-EP-2016-251] [[arXiv:1610.06019](#)] [[INSPIRE](#)].
- [40] LHCb collaboration, *A new algorithm for identifying the flavour of B_s^0 mesons at LHCb*, [2016 JINST 11 P05010](#) [LHCb-PAPER-2015-056] [CERN-EP-2016-030] [[arXiv:1602.07252](#)] [[INSPIRE](#)].
- [41] LHCb collaboration, *Opposite-side flavour tagging of B mesons at the LHCb experiment*, *Eur. Phys. J. C* **72** (2012) 2022 [CERN-PH-EP-2012-034] [LHCb-PAPER-2011-027] [[arXiv:1202.4979](#)] [[INSPIRE](#)].
- [42] LHCb collaboration, *B flavour tagging using charm decays at the LHCb experiment*, [2015 JINST 10 P10005](#) [LHCb-PAPER-2015-027] [CERN-PH-EP-2015-193] [[arXiv:1507.07892](#)] [[INSPIRE](#)].
- [43] LHCb collaboration, *Flavour Tagging in the LHCb experiment*, in the proceedings of the *6th Large Hadron Collider Physics Conference (LHCP 2018)*, Bologna, Italy, 4–9 June 2018, *PoS LHCP2018* (2018) 230 [[INSPIRE](#)].
- [44] LHCb collaboration, *Measurements of the B^+ , B^0 , B_s^0 meson and Λ_b^0 baryon lifetimes*, *JHEP* **04** (2014) 114 [LHCb-PAPER-2013-065] [CERN-PH-EP-2014-018] [[arXiv:1402.2554](#)] [[INSPIRE](#)].
- [45] T.M. Karbach, G. Raven and M. Schiller, *Decay time integrals in neutral meson mixing and their efficient evaluation*, [arXiv:1407.0748](#) [[INSPIRE](#)].
- [46] PARTICLE DATA collaboration, *Review of particle physics*, *Prog. Theor. Exp. Phys.* **2022** (2022) 083C01 [[INSPIRE](#)].

- [47] LHCb collaboration, *Measurement of CP violation in $B^0 \rightarrow D^{*\pm} D^\mp$ decays*, *JHEP* **03** (2020) 147 [LHCb-PAPER-2019-036] [CERN-EP-2019-264] [[arXiv:1912.03723](#)] [[INSPIRE](#)].
- [48] LHCb collaboration, *Precise determination of the $B_s^0 - \bar{B}_s^0$ oscillation frequency*, *Nat. Phys.* **18** (2022) 1 [LHCb-PAPER-2019-036] [CERN-EP-2019-264] [[arXiv:2104.04421](#)] [[INSPIRE](#)].
- [49] B. Efron, *Bootstrap Methods: Another Look at the Jackknife*, *Ann. Statist.* **7** (1979) 1 [[INSPIRE](#)].
- [50] T. Skwarnicki, *A study of the radiative CASCADE transitions between the Upsilon-Prime and Upsilon resonances*, Ph.D. Thesis, Institute of Nuclear Physics, Krakow, Poland (1986) [[INSPIRE](#)].
- [51] S.S. Wilks, *The Large-Sample Distribution of the Likelihood Ratio for Testing Composite Hypotheses*, *Ann. Math. Stat.* **9** (1938) 60 [[INSPIRE](#)].
- [52] M. Kenzie, M. Karbach, T. Mombächer, M. Schlupp and K. Schubert, *GammaCombo: A statistical analysis framework for combining measurements, fitting datasets and producing confidence intervals*, (2021) [[DOI:10.5281/zenodo.3371421](#)].
- [53] LHCb collaboration, *Measurement of the CKM angle γ from a combination of LHCb results*, *JHEP* **12** (2016) 087 [LHCb-PAPER-2016-032] [CERN-EP-2016-270] [[arXiv:1611.03076](#)] [[INSPIRE](#)].

The LHCb collaboration

R. Aaij³⁷, A.S.W. Abdelmotteleb⁵⁶, C. Abellan Beteta⁵⁰, F. Abudinén⁵⁶, T. Ackernley⁶⁰, A.A. Adefisoye⁶⁸, B. Adeva⁴⁶, M. Adinolfi⁵⁴, P. Adlarson⁸¹, C. Agapopoulou¹⁴, C.A. Aidala⁸², Z. Ajaltouni¹¹, S. Akar⁶⁵, K. Akiba³⁷, P. Albicocco²⁷, J. Albrecht^{19,f}, F. Alessio⁴⁸, M. Alexander⁵⁹, Z. Aliouche⁶², P. Alvarez Cartelle⁵⁵, R. Amalric¹⁶, S. Amato³, J.L. Amey⁵⁴, Y. Amhis^{14,48}, L. An⁶, L. Anderlini²⁶, M. Andersson⁵⁰, A. Andreianov⁴³, P. Andreola⁵⁰, M. Andreotti²⁵, D. Andreou⁶⁸, A. Anelli^{30,o}, D. Ao⁷, F. Archilli^{36,u}, M. Argenton²⁵, S. Arguedas Cuendis^{9,48}, A. Artamonov⁴³, M. Artuso⁶⁸, E. Aslanides¹³, R. Ataíde Da Silva⁴⁹, M. Atzeni⁶⁴, B. Audurier¹², D. Bacher⁶³, I. Bachiller Perea¹⁰, S. Bachmann²¹, M. Bachmayer⁴⁹, J.J. Back⁵⁶, P. Baladron Rodriguez⁴⁶, V. Balagura¹⁵, W. Baldini²⁵, L. Balzani¹⁹, H. Bao⁷, J. Baptista de Souza Leite⁶⁰, C. Barbero Pretel^{46,12}, M. Barbetti²⁶, I.R. Barbosa⁶⁹, R.J. Barlow⁶², M. Barnyakov²⁴, S. Barsuk¹⁴, W. Barter⁵⁸, M. Bartolini⁵⁵, J. Bartz⁶⁸, J.M. Basels¹⁷, S. Bashir³⁹, G. Bassi^{34,r}, B. Batsukh⁵, P.B. Battista¹⁴, A. Bay⁴⁹, A. Beck⁵⁶, M. Becker¹⁹, F. Bedeschi³⁴, I.B. Bediaga², N.A. Behling¹⁹, S. Belin⁴⁶, V. Bellee⁵⁰, K. Belous⁴³, I. Belov²⁸, I. Belyaev³⁵, G. Benane¹³, G. Bencivenni²⁷, E. Ben-Haim¹⁶, A. Berezhnoy⁴³, R. Bernet⁵⁰, S. Bernet Andres⁴⁴, A. Bertolin³², C. Betancourt⁵⁰, F. Betti⁵⁸, J. Bex⁵⁵, Ia. Bezshyiko⁵⁰, J. Bhom⁴⁰, M.S. Bieker¹⁹, N.V. Biesuz²⁵, P. Billoir¹⁶, A. Biolchini³⁷, M. Birch⁶¹, F.C.R. Bishop¹⁰, A. Bitadze⁶², A. Bizzeti¹⁶, T. Blake⁵⁶, F. Blanc⁴⁹, J.E. Blank¹⁹, S. Blusk⁶⁸, V. Bocharnikov⁴³, J.A. Boelhauve¹⁹, O. Boente Garcia¹⁵, T. Boettcher⁶⁵, A. Bohare⁵⁸, A. Boldyrev⁴³, C.S. Bolognani⁷⁸, R. Bolzonella^{25,l}, N. Bondar⁴³, A. Bordelius⁴⁸, F. Borgato^{32,p}, S. Borghi⁶², M. Borsato^{30,o}, J.T. Borsuk⁴⁰, S.A. Bouchiba⁴⁹, M. Bovill⁶³, T.J.V. Bowcock⁶⁰, A. Boyer⁴⁸, C. Bozzi²⁵, A. Brea Rodriguez⁴⁹, N. Breer¹⁹, J. Brodzicka⁴⁰, A. Brossa Gonzalo^{46,56,45,†}, J. Brown⁶⁰, D. Brundu³¹, E. Buchanan⁵⁸, A. Buonauro⁵⁰, L. Buonincontri^{32,p}, A.T. Burke⁶², C. Burr⁴⁸, J.S. Butter⁵⁵, J. Buytaert⁴⁸, W. Byczynski⁴⁸, S. Cadeddu³¹, H. Cai⁷³, A.C. Caillet¹⁶, R. Calabrese^{25,l}, S. Calderon Ramirez⁹, L. Calefice⁴⁵, S. Cali²⁷, M. Calvi^{30,o}, M. Calvo Gomez⁴⁴, P. Camargo Magalhaes^{2,y}, J.I. Cambon Bouzas⁴⁶, P. Campana²⁷, D.H. Campora Perez⁷⁸, A.F. Campoverde Quezada⁷, S. Capelli³⁰, L. Capriotti²⁵, R. Caravaca-Mora⁹, A. Carbone^{24,j}, L. Carcedo Salgado⁴⁶, R. Cardinale^{28,m}, A. Cardini³¹, P. Carniti^{30,o}, L. Carus²¹, A. Casais Vidal⁶⁴, R. Caspary²¹, G. Casse⁶⁰, J. Castro Godinez⁹, M. Cattaneo⁴⁸, G. Cavallero^{25,48}, V. Cavallini^{25,l}, S. Celani²¹, D. Cervenkov⁶³, S. Cesare^{29,n}, A.J. Chadwick⁶⁰, I. Chahrouh⁸², M. Charles¹⁶, Ph. Charpentier⁴⁸, E. Chatzianagnostou³⁷, M. Chefdeville¹⁰, C. Chen¹³, S. Chen⁵, Z. Chen⁷, A. Chernov⁴⁰, S. Chernyshenko⁵², X. Chiotopoulos⁷⁸, V. Chobanova⁸⁰, S. Cholak⁴⁹, M. Chrzaszcz⁴⁰, A. Chubykin⁴³, V. Chulikov⁴³, P. Ciambone²⁷, X. Cid Vidal⁴⁶, G. Ciezarek⁴⁸, P. Cifra⁴⁸, P.E.L. Clarke⁵⁸, M. Clemencic⁴⁸, H.V. Cliff⁵⁵, J. Closier⁴⁸, C. Cocha Toapaxi²¹, V. Coco⁴⁸, J. Cogan¹³, E. Cogneras¹¹, L. Cojocariu⁴², P. Collins⁴⁸, T. Colombo⁴⁸, M. Colonna¹⁹, A. Comerma-Montells⁴⁵, L. Congedo²³, A. Contu³¹, N. Cooke⁵⁹, I. Corredoira⁴⁶, A. Correia¹⁶, G. Corti⁴⁸, J. Cottee Meldrum⁵⁴, B. Couturier⁴⁸, D.C. Craik⁵⁰, M. Cruz Torres^{2,g}, E. Curras Rivera⁴⁹, R. Currie⁵⁸, C.L. Da Silva⁶⁷, S. Dadabaev⁴³, L. Dai⁷⁰, X. Dai⁶, E. Dall’Occo¹⁹, J. Dalseno⁴⁶,

C. D'Ambrosio ⁴⁸, J. Daniel ¹¹, A. Danilina ⁴³, P. d'Argent ²³, A. Davidson ⁵⁶,
J.E. Davies ⁶², A. Davis ⁶², O. De Aguiar Francisco ⁶², C. De Angelis ^{31,k}, F. De Benedetti ⁴⁸,
J. de Boer ³⁷, K. De Bruyn ⁷⁷, S. De Capua ⁶², M. De Cian ^{21,48},
U. De Freitas Carneiro Da Graca ^{2,a}, E. De Lucia ²⁷, J.M. De Miranda ², L. De Paula ³,
M. De Serio ^{23,h}, P. De Simone ²⁷, F. De Vellis ¹⁹, J.A. de Vries ⁷⁸, F. Debernardis ²³,
D. Decamp ¹⁰, V. Dedu ¹³, S. Dekkers ¹, L. Del Buono ¹⁶, B. Delaney ⁶⁴, H.-P. Dembinski ¹⁹,
J. Deng ⁸, V. Denysenko ⁵⁰, O. Deschamps ¹¹, F. Dettori ^{31,k}, B. Dey ⁷⁶, P. Di Nezza ²⁷,
I. Diachkov ⁴³, S. Didenko ⁴³, S. Ding ⁶⁸, L. Dittmann ²¹, V. Dobishuk ⁵², A.D. Docheva ⁵⁹,
C. Dong ^{4,b}, A.M. Donohoe ²², F. Dordei ³¹, A.C. dos Reis ², A.D. Dowling ⁶⁸, W. Duan ⁷¹,
P. Duda ⁷⁹, M.W. Dudek ⁴⁰, L. Dufour ⁴⁸, V. Duk ³³, P. Durante ⁴⁸, M.M. Duras ⁷⁹,
J.M. Durham ⁶⁷, O.D. Durmus ⁷⁶, A. Dziurda ⁴⁰, A. Dzyuba ⁴³, S. Easo ⁵⁷, E. Eckstein ¹⁸,
U. Egede ¹, A. Egorychev ⁴³, V. Egorychev ⁴³, S. Eisenhardt ⁵⁸, E. Ejopu ⁶², L. Eklund ⁸¹,
M. Elashri ⁶⁵, J. Ellbracht ¹⁹, S. Ely ⁶¹, A. Ene ⁴², E. Eppe ⁶⁵, J. Eschle ⁶⁸, S. Esen ²¹,
T. Evans ⁶², F. Fabiano ^{31,k}, L.N. Falcao ², Y. Fan ⁷, B. Fang ⁷³, L. Fantini ^{33,q,48},
M. Faria ⁴⁹, K. Farmer ⁵⁸, D. Fazzini ^{30,o}, L. Felkowski ⁷⁹, M. Feng ^{5,7}, M. Feo ^{19,48},
A. Fernandez Casani ⁴⁷, M. Fernandez Gomez ⁴⁶, A.D. Ferez ⁶⁶, F. Ferrari ^{24,j},
F. Ferreira Rodrigues ³, M. Ferrillo ⁵⁰, M. Ferro-Luzzi ⁴⁸, S. Filippov ⁴³, R.A. Fini ²³,
M. Fiorini ^{25,l}, M. Firlej ³⁹, K.L. Fischer ⁶³, D.S. Fitzgerald ⁸², C. Fitzpatrick ⁶²,
T. Fiutowski ³⁹, F. Fleuret ¹⁵, M. Fontana ²⁴, L.F. Foreman ⁶², R. Forty ⁴⁸,
D. Foulds-Holt ⁵⁵, V. Franco Lima ³, M. Franco Sevilla ⁶⁶, M. Frank ⁴⁸, E. Franzoso ^{25,l},
G. Frau ⁶², C. Frei ⁴⁸, D.A. Friday ⁶², J. Fu ⁷, Q. Fühling ^{19,f,55}, Y. Fujii ¹, T. Fulghesu ¹⁶,
E. Gabriel ³⁷, G. Galati ²³, M.D. Galati ³⁷, A. Gallas Torreira ⁴⁶, D. Galli ^{24,j},
S. Gambetta ⁵⁸, M. Gandelman ³, P. Gandini ²⁹, B. Ganie ⁶², H. Gao ⁷, R. Gao ⁶³,
T.Q. Gao ⁵⁵, Y. Gao ⁸, Y. Gao ⁶, Y. Gao ⁸, M. Garau ^{31,k}, L.M. Garcia Martin ⁴⁹,
P. Garcia Moreno ⁴⁵, J. García Pardiñas ⁴⁸, K.G. Garg ⁸, L. Garrido ⁴⁵, C. Gaspar ⁴⁸,
R.E. Geertsema ³⁷, L.L. Gerken ¹⁹, E. Gersabeck ⁶², M. Gersabeck ⁶², T. Gershon ⁵⁶,
S. Ghizzo ^{28,m}, Z. Ghorbanimoghaddam ⁵⁴, L. Giambastiani ^{32,p}, F.I. Giasemis ^{16,e},
V. Gibson ⁵⁵, H.K. Gienza ⁴¹, A.L. Gilman ⁶³, M. Giovannetti ²⁷, A. Gioventù ⁴⁵,
L. Girardey ⁶², P. Gironella Gironell ⁴⁵, C. Giugliano ^{25,l}, M.A. Giza ⁴⁰, E.L. Gkougkousis ⁶¹,
F.C. Glaser ^{14,21}, V.V. Gligorov ^{16,48}, C. Göbel ⁶⁹, E. Golobardes ⁴⁴, D. Golubkov ⁴³,
A. Golutvin ^{61,48,43}, S. Gomez Fernandez ⁴⁵, F. Goncalves Abrantes ⁶³, M. Goncerz ⁴⁰,
G. Gong ^{4,b}, J.A. Gooding ¹⁹, I.V. Gorelov ⁴³, C. Gotti ³⁰, J.P. Grabowski ¹⁸,
L.A. Granado Cardoso ⁴⁸, E. Graugés ⁴⁵, E. Graverini ^{49,s}, L. Grazette ⁵⁶, G. Graziani ⁶,
A.T. Grecu ⁴², L.M. Greeven ³⁷, N.A. Grieser ⁶⁵, L. Grillo ⁵⁹, S. Gromov ⁴³, C. Gu ¹⁵,
M. Guarise ²⁵, L. Guerry ¹¹, M. Guittiere ¹⁴, V. Guliaeva ⁴³, P.A. Günther ²¹,
A.-K. Guseinov ⁴⁹, E. Gushchin ⁴³, Y. Guz ^{6,48,43}, T. Gys ⁴⁸, K. Habermann ¹⁸,
T. Hadavizadeh ¹, C. Hadjivasiliou ⁶⁶, G. Haefeli ⁴⁹, C. Haen ⁴⁸, J. Haimberger ⁴⁸,
M. Hajheidari ⁴⁸, G. Hallett ⁵⁶, M.M. Halvorsen ⁴⁸, P.M. Hamilton ⁶⁶, J. Hammerich ⁶⁰,
Q. Han ⁸, X. Han ²¹, S. Hansmann-Menzemer ²¹, L. Hao ⁷, N. Harnew ⁶³, M. Hartmann ¹⁴,
S. Hashmi ³⁹, J. He ^{7,c}, F. Hemmer ⁴⁸, C. Henderson ⁶⁵, R.D.L. Henderson ^{1,56},
A.M. Hennequin ⁴⁸, K. Hennessy ⁶⁰, L. Henry ⁴⁹, J. Herd ⁶¹, P. Herrero Gascon ²¹,
J. Heuel ¹⁷, A. Hicheur ³, G. Hijano Mendizabal ⁵⁰, D. Hill ⁴⁹, S.E. Hollitt ¹⁹, J. Horswill ⁶²,
R. Hou ⁸, Y. Hou ¹¹, N. Howarth ⁶⁰, J. Hu ²¹, J. Hu ⁷¹, W. Hu ⁶, X. Hu ^{4,b}, W. Huang ⁷,
W. Hulsbergen ³⁷, R.J. Hunter ⁵⁶, M. Hushchyn ⁴³, D. Hutchcroft ⁶⁰, M. Idzik ³⁹, D. Ilin ⁴³,

P. Ilten ⁶⁵, A. Inglessi ⁴³, A. Iniukhin ⁴³, A. Ishteev ⁴³, K. Ivshin ⁴³, R. Jacobsson ⁴⁸,
H. Jage ¹⁷, S.J. Jaimes Elles ^{47,74}, S. Jakobsen ⁴⁸, E. Jans ³⁷, B.K. Jashal ⁴⁷,
A. Jawahery ^{66,48}, V. Jevtic ¹⁹, E. Jiang ⁶⁶, X. Jiang ^{5,7}, Y. Jiang ⁷, Y.J. Jiang ⁶,
M. John ⁶³, A. John Rubesh Rajan ²², D. Johnson ⁵³, C.R. Jones ⁵⁵, T.P. Jones ⁵⁶,
S. Joshi ⁴¹, B. Jost ⁴⁸, J. Juan Castella ⁵⁵, N. Jurik ⁴⁸, I. Juszcak ⁴⁰, D. Kaminaris ⁴⁹,
S. Kandybei ⁵¹, M. Kane ⁵⁸, Y. Kang ^{4,b}, C. Kar ¹¹, M. Karacson ⁴⁸, D. Karpenkov ⁴³,
A. Kauniskangas ⁴⁹, J.W. Kautz ⁶⁵, M.K. Kazanecki ⁴⁰, F. Keizer ⁴⁸, M. Kenzie ⁵⁵,
T. Ketel ³⁷, B. Khanji ⁶⁸, A. Kharisova ⁴³, S. Kholodenko ^{34,48}, G. Khreich ¹⁴, T. Kirn ¹⁷,
V.S. Kirsebom ^{30,o}, O. Kitouni ⁶⁴, S. Klaver ³⁸, N. Kleijne ^{34,r}, K. Klimaszewski ⁴¹,
M.R. Kmiec ⁴¹, S. Koliiev ⁵², L. Kolk ¹⁹, A. Konoplyannikov ⁴³, P. Kopciwicz ^{39,48},
P. Koppenburg ³⁷, M. Korolev ⁴³, I. Kostiuk ³⁷, O. Kot ⁵², S. Kotriakhova ⁴³, A. Kozachuk ⁴³,
P. Kravchenko ⁴³, L. Kravchuk ⁴³, M. Kreps ⁵⁶, P. Krokovny ⁴³, W. Krupa ⁶⁸,
W. Krzemien ⁴¹, O. Kshyvanskyi ⁵², S. Kubis ⁷⁹, M. Kucharczyk ⁴⁰, V. Kudryavtsev ⁴³,
E. Kulikova ⁴³, A. Kupsc ⁸¹, B.K. Kutsenko ¹³, D. Lacarrere ⁴⁸, P. Laguarda Gonzalez ⁴⁵,
A. Lai ³¹, A. Lampis ³¹, D. Lancierini ⁵⁵, C. Landesa Gomez ⁴⁶, J.J. Lane ¹, R. Lane ⁵⁴,
G. Lanfranchi ²⁷, C. Langenbruch ²¹, J. Langer ¹⁹, O. Lantwin ⁴³, T. Latham ⁵⁶,
F. Lazzari ^{34,s}, C. Lazzeroni ⁵³, R. Le Gac ¹³, H. Lee ⁶⁰, R. Lefèvre ¹¹, A. Leflat ⁴³,
S. Legotin ⁴³, M. Lehuraux ⁵⁶, E. Lemos Cid ⁴⁸, O. Leroy ¹³, T. Lesiak ⁴⁰, E.D. Lesser ⁴⁸,
B. Leverington ²¹, A. Li ^{4,b}, C. Li ¹³, H. Li ⁷¹, K. Li ⁸, L. Li ⁶², M. Li ⁸, P. Li ⁷,
P.-R. Li ⁷², Q. Li ^{5,7}, S. Li ⁸, T. Li ^{5,d}, T. Li ⁷¹, Y. Li ⁸, Y. Li ⁵, Z. Lian ^{4,b}, X. Liang ⁶⁸,
S. Libralon ⁴⁷, C. Lin ⁷, T. Lin ⁵⁷, R. Lindner ⁴⁸, V. Lisovskyi ⁴⁹, R. Litvinov ^{31,48},
F.L. Liu ¹, G. Liu ⁷¹, K. Liu ⁷², S. Liu ^{5,7}, W. Liu ⁸, Y. Liu ⁵⁸, Y. Liu ⁷², Y.L. Liu ⁶¹,
A. Lobo Salvia ⁴⁵, A. Loi ³¹, J. Lomba Castro ⁴⁶, T. Long ⁵⁵, J.H. Lopes ³,
A. Lopez Huertas ⁴⁵, S. López Soliño ⁴⁶, Q. Lu ¹⁵, C. Lucarelli ²⁶, D. Lucchesi ^{32,p},
M. Lucio Martinez ⁷⁸, V. Lukashenko ^{37,52}, Y. Luo ⁶, A. Lupato ^{32,i}, E. Luppi ^{25,l},
K. Lynch ²², X.-R. Lyu ⁷, G.M. Ma ^{4,b}, R. Ma ⁷, S. Maccolini ¹⁹, F. Machefert ¹⁴,
F. Maciuc ⁴², B. Mack ⁶⁸, I. Mackay ⁶³, L.M. Mackey ⁶⁸, L.R. Madhan Mohan ⁵⁵,
M.J. Madurai ⁵³, A. Maevskiy ⁴³, D. Magdalinski ³⁷, D. Maisuzenko ⁴³, M.W. Majewski ³⁹,
J.J. Malczewski ⁴⁰, S. Malde ⁶³, L. Malentacca ⁴⁸, A. Malinin ⁴³, T. Maltsev ⁴³,
G. Manca ^{31,k}, G. Mancinelli ¹³, C. Mancuso ^{29,14,n}, R. Manera Escalero ⁴⁵, D. Manuzzi ²⁴,
D. Marangotto ^{29,n}, J.F. Marchand ¹⁰, R. Marchevski ⁴⁹, U. Marconi ²⁴, E. Mariani ¹⁶,
S. Mariani ⁴⁸, C. Marin Benito ⁴⁵, J. Marks ²¹, A.M. Marshall ⁵⁴, L. Martel ⁶³,
G. Martelli ^{33,q}, G. Martellotti ³⁵, L. Martinazzoli ⁴⁸, M. Martinelli ^{30,o},
D. Martinez Santos ⁴⁶, F. Martinez Vidal ⁴⁷, A. Massafferri ², R. Matev ⁴⁸, A. Mathad ⁴⁸,
V. Matiunin ⁴³, C. Matteuzzi ⁶⁸, K.R. Mattioli ¹⁵, A. Mauri ⁶¹, E. Maurice ¹⁵, J. Mauricio ⁴⁵,
P. Mayencourt ⁴⁹, J. Mazorra de Cos ⁴⁷, M. Mazurek ⁴¹, M. McCann ⁶¹, L. McConnell ²²,
T.H. McGrath ⁶², N.T. McHugh ⁵⁹, A. McNab ⁶², R. McNulty ²², B. Meadows ⁶⁵,
G. Meier ¹⁹, D. Melnychuk ⁴¹, F.M. Meng ^{4,b}, M. Merk ^{37,78}, A. Merli ⁴⁹, L. Meyer Garcia ⁶⁶,
D. Miao ^{5,7}, H. Miao ⁷, M. Mikhasenko ⁷⁵, D.A. Milanes ⁷⁴, A. Minotti ^{30,o}, E. Minucci ⁶⁸,
T. Miralles ¹¹, B. Mitreska ¹⁹, D.S. Mitzel ¹⁹, A. Modak ⁵⁷, R.A. Mohammed ⁶³,
R.D. Moise ¹⁷, S. Mokhnenko ⁴³, E.F. Molina Cardenas ⁸², T. Mombächer ⁴⁸, M. Monk ^{56,1},
S. Monteil ¹¹, A. Morcillo Gomez ⁴⁶, G. Morello ²⁷, M.J. Morello ^{34,r}, M.P. Morgenthaler ²¹,
J. Moron ³⁹, A.B. Morris ⁴⁸, A.G. Morris ¹³, R. Mountain ⁶⁸, H. Mu ^{4,b}, Z.M. Mu ⁶,
E. Muhammad ⁵⁶, F. Muheim ⁵⁸, M. Mulder ⁷⁷, K. Müller ⁵⁰, F. Muñoz-Rojas ⁹,

R. Murta ⁶¹, P. Naik ⁶⁰, T. Nakada ⁴⁹, R. Nandakumar ⁵⁷, T. Nanut ⁴⁸, I. Nasteva ³,
M. Needham ⁵⁸, N. Neri ^{29,n}, S. Neubert ¹⁸, N. Neufeld ⁴⁸, P. Neustroev ⁴³, J. Nicolini ^{19,14},
D. Nicotra ⁷⁸, E.M. Niel ⁴⁹, N. Nikitin ⁴³, Q. Niu ⁷², P. Nogarolli ³, P. Nogga ¹⁸,
C. Normand ⁵⁴, J. Novoa Fernandez ⁴⁶, G. Nowak ⁶⁵, C. Nunez ⁸², H.N. Nur ⁵⁹,
A. Oblakowska-Mucha ³⁹, V. Obraztsov ⁴³, T. Oeser ¹⁷, S. Okamura ^{25,l}, A. Okhotnikov ⁴³,
O. Okhrimenko ⁵², R. Oldeman ^{31,k}, F. Oliva ⁵⁸, M. Olocco ¹⁹, C.J.G. Onderwater ⁷⁸,
R.H. O’Neil ⁵⁸, D. Osthues ¹⁹, J.M. Otalora Goicochea ³, P. Owen ⁵⁰, A. Oyanguren ⁴⁷,
O. Ozcelik ⁵⁸, F. Paciolla ^{34,v}, A. Padee ⁴¹, K.O. Padeken ¹⁸, B. Pagare ⁵⁶, P.R. Pais ²¹,
T. Pajero ⁴⁸, A. Palano ²³, M. Palutan ²⁷, G. Panshin ⁴³, L. Paolucci ⁵⁶, A. Papanestis ^{57,48},
M. Pappagallo ^{23,h}, L.L. Pappalardo ^{25,l}, C. Pappenheimer ⁶⁵, C. Parkes ⁶², B. Passalacqua ²⁵,
G. Passaleva ²⁶, D. Passaro ^{34,r}, A. Pastore ²³, M. Patel ⁶¹, J. Patoc ⁶³, C. Patrignani ^{24,j},
A. Paul ⁶⁸, C.J. Pawley ⁷⁸, A. Pellegrino ³⁷, J. Peng ^{5,7}, M. Pepe Altarelli ²⁷, S. Perazzini ²⁴,
D. Pereima ⁴³, H. Pereira Da Costa ⁶⁷, A. Pereiro Castro ⁴⁶, P. Perret ¹¹, A. Perro ^{48,13},
K. Petridis ⁵⁴, A. Petrolini ^{28,m}, J.P. Pfaller ⁶⁵, H. Pham ⁶⁸, L. Pica ^{34,r}, M. Piccini ³³,
L. Piccolo ³¹, B. Pietrzyk ¹⁰, G. Pietrzyk ¹⁴, D. Pinci ³⁵, F. Pisani ⁴⁸, M. Pizzichemi ^{30,o,48},
V. Placinta ⁴², M. Plo Casasus ⁴⁶, T. Poeschl ⁴⁸, F. Polci ^{16,48}, M. Poli Lener ²⁷,
A. Poluektov ¹³, N. Polukhina ⁴³, I. Polyakov ⁴³, E. Polycarpo ³, S. Ponce ⁴⁸, D. Popov ⁷,
S. Poslavskii ⁴³, K. Prasanth ⁵⁸, C. Prouve ⁴⁶, D. Provenzano ^{31,k}, V. Pugatch ⁵²,
G. Punzi ^{34,s}, S. Qasim ⁵⁰, Q.Q. Qian ⁶, W. Qian ⁷, N. Qin ^{4,b}, S. Qu ^{4,b}, R. Quagliani ⁴⁸,
R.I. Rabadan Trejo ⁵⁶, J.H. Rademacker ⁵⁴, M. Rama ³⁴, M. Ramírez García ⁸²,
V. Ramos De Oliveira ⁶⁹, M. Ramos Pernas ⁵⁶, M.S. Rangel ³, F. Ratnikov ⁴³, G. Raven ³⁸,
M. Rebollo De Miguel ⁴⁷, F. Redi ^{29,i}, J. Reich ⁵⁴, F. Reiss ⁶², Z. Ren ⁷, P.K. Resmi ⁶³,
R. Ribatti ⁴⁹, G.R. Ricart ^{15,12}, D. Riccardi ^{34,r}, S. Ricciardi ⁵⁷, K. Richardson ⁶⁴,
M. Richardson-Slipper ⁵⁸, K. Rinnert ⁶⁰, P. Robbe ¹⁴, G. Robertson ⁵⁹, E. Rodrigues ⁶⁰,
E. Rodriguez Fernandez ⁴⁶, J.A. Rodriguez Lopez ⁷⁴, E. Rodriguez Rodriguez ⁴⁶, J. Roensch ¹⁹,
A. Rogachev ⁴³, A. Rogovskiy ⁵⁷, D.L. Rolf ⁴⁸, P. Roloff ⁴⁸, V. Romanovskiy ⁶⁵,
M. Romero Lamas ⁴⁶, A. Romero Vidal ⁴⁶, G. Romolini ²⁵, F. Ronchetti ⁴⁹, T. Rong ⁶,
M. Rotondo ²⁷, S.R. Roy ²¹, M.S. Rudolph ⁶⁸, M. Ruiz Diaz ²¹, R.A. Ruiz Fernandez ⁴⁶,
J. Ruiz Vidal ^{81,z}, A. Ryzhikov ⁴³, J. Ryzka ³⁹, J.J. Saavedra-Arias ⁹, J.J. Saborido Silva ⁴⁶,
R. Sadek ¹⁵, N. Sagidova ⁴³, D. Sahoo ⁷⁶, N. Sahoo ⁵³, B. Saitta ^{31,k}, M. Salomoni ^{30,48,o},
I. Sanderswood ⁴⁷, R. Santacesaria ³⁵, C. Santamarina Rios ⁴⁶, M. Santimaria ^{27,48},
L. Santoro ², E. Santovetti ³⁶, A. Saputi ^{25,48}, D. Saranin ⁴³, A. Sarnatskiy ⁷⁷, G. Sarpis ⁵⁸,
M. Sarpis ⁶², C. Satriano ^{35,t}, A. Satta ³⁶, M. Saur ⁶, D. Savrina ⁴³, H. Sazak ¹⁷,
F. Sborzacchi ^{48,27}, L.G. Scantlebury Smead ⁶³, A. Scarabotto ¹⁹, S. Schael ¹⁷, S. Scherl ⁶⁰,
M. Schiller ⁵⁹, H. Schindler ⁴⁸, M. Schmelling ²⁰, B. Schmidt ⁴⁸, S. Schmitt ¹⁷, H. Schmitz ¹⁸,
O. Schneider ⁴⁹, A. Schopper ⁴⁸, N. Schulte ¹⁹, S. Schulte ⁴⁹, M.H. Schune ¹⁴,
R. Schwemmer ⁴⁸, G. Schwering ¹⁷, B. Sciascia ²⁷, A. Sciuccati ⁴⁸, S. Sellam ⁴⁶,
A. Semennikov ⁴³, T. Senger ⁵⁰, M. Senghi Soares ³⁸, A. Sergi ^{28,m,48}, N. Serra ⁵⁰,
L. Sestini ³², A. Seuthe ¹⁹, Y. Shang ⁶, D.M. Shangase ⁸², M. Shapkin ⁴³, R.S. Sharma ⁶⁸,
I. Shchemerov ⁴³, L. Shchutska ⁴⁹, T. Shears ⁶⁰, L. Shekhtman ⁴³, Z. Shen ⁶, S. Sheng ^{5,7},
V. Shevchenko ⁴³, B. Shi ⁷, Q. Shi ⁷, Y. Shimizu ¹⁴, E. Shmanin ²⁴, R. Shorkin ⁴³,
J.D. Shupperd ⁶⁸, R. Silva Coutinho ⁶⁸, G. Simi ^{32,p}, S. Simone ^{23,h}, N. Skidmore ⁵⁶,
T. Skwarnicki ⁶⁸, M.W. Slater ⁵³, J.C. Smallwood ⁶³, E. Smith ⁶⁴, K. Smith ⁶⁷, M. Smith ⁶¹,
A. Snoch ³⁷, L. Soares Lavra ⁵⁸, M.D. Sokoloff ⁶⁵, F.J.P. Soler ⁵⁹, A. Solomin ^{43,54},

A. Solovov ⁴³, I. Solov'yev ⁴³, R. Song ¹, Y. Song ⁴⁹, Y. Song ^{4,b}, Y.S. Song ⁶,
 F.L. Souza De Almeida ⁶⁸, B. Souza De Paula ³, E. Spadaro Norella ^{28,m}, E. Spedicato ²⁴,
 J.G. Speer ¹⁹, E. Spiridenkov ⁴³, P. Spradlin ⁵⁹, V. Sriskaran ⁴⁸, F. Stagni ⁴⁸, M. Stahl ⁴⁸,
 S. Stahl ⁴⁸, S. Stanislaus ⁶³, E.N. Stein ⁴⁸, O. Steinkamp ⁵⁰, O. Stenyakin ⁴³, H. Stevens ¹⁹,
 D. Strekalina ⁴³, Y. Su ⁷, F. Suljik ⁶³, J. Sun ³¹, L. Sun ⁷³, Y. Sun ⁶⁶, D. Sundfeld ²,
 W. Sutcliffe ⁵⁰, P.N. Swallow ⁵³, K. Swientek ³⁹, F. Swystun ⁵⁵, A. Szabelski ⁴¹, T. Szumlak ³⁹,
 Y. Tan ^{4,b}, M.D. Tat ⁶³, A. Terentev ⁴³, F. Terzuoli ^{34,v,48}, F. Teubert ⁴⁸, E. Thomas ⁴⁸,
 D.J.D. Thompson ⁵³, H. Tilquin ⁶¹, V. Tisserand ¹¹, S. T'Jampens ¹⁰, M. Tobin ^{5,48},
 L. Tomassetti ^{25,l}, G. Tonani ^{29,n,48}, X. Tong ⁶, D. Torres Machado ², L. Toscano ¹⁹,
 D.Y. Tou ^{4,b}, C. Tripp ⁴⁴, G. Tuci ²¹, N. Tuning ³⁷, L.H. Uecker ²¹, A. Ukleja ³⁹,
 D.J. Unverzagt ²¹, E. Ursov ⁴³, A. Usachov ³⁸, A. Ustyuzhanin ⁴³, U. Uwer ²¹,
 V. Vagnoni ²⁴, V. Valcarce Cadenas ⁴⁶, G. Valenti ²⁴, N. Valls Canudas ⁴⁸, H. Van Hecke ⁶⁷,
 E. van Herwijnen ⁶¹, C.B. Van Hulse ^{46,x}, R. Van Laak ⁴⁹, M. van Veghel ³⁷, G. Vasquez ⁵⁰,
 R. Vazquez Gomez ⁴⁵, P. Vazquez Regueiro ⁴⁶, C. Vázquez Sierra ⁴⁶, S. Vecchi ²⁵,
 J.J. Velthuis ⁵⁴, M. Veltri ^{26,w}, A. Venkateswaran ⁴⁹, M. Verdoglia ³¹, M. Vesterinen ⁵⁶,
 D. Vico Benet ⁶³, P. Vidrier Villalba ⁴⁵, M. Vieites Diaz ⁴⁸, X. Vilasis-Cardona ⁴⁴,
 E. Vilella Figueras ⁶⁰, A. Villa ²⁴, P. Vincent ¹⁶, F.C. Volle ⁵³, D. vom Bruch ¹³,
 N. Voropaev ⁴³, K. Vos ⁷⁸, G. Vouters ¹⁰, C. Vrahas ⁵⁸, J. Wagner ¹⁹, J. Walsh ³⁴,
 E.J. Walton ^{1,56}, G. Wan ⁶, C. Wang ²¹, G. Wang ⁸, H. Wang ⁷², J. Wang ⁶, J. Wang ⁵,
 J. Wang ^{4,b}, J. Wang ⁷³, M. Wang ²⁹, N.W. Wang ⁷, R. Wang ⁵⁴, X. Wang ⁸, X. Wang ⁷¹,
 X.W. Wang ⁶¹, Y. Wang ⁶, Y.W. Wang ⁷², Z. Wang ¹⁴, Z. Wang ^{4,b}, Z. Wang ²⁹,
 J.A. Ward ^{56,1}, M. Waterlaan ⁴⁸, N.K. Watson ⁵³, D. Websdale ⁶¹, Y. Wei ⁶, J. Wendel ⁸⁰,
 B.D.C. Westhenry ⁵⁴, C. White ⁵⁵, M. Whitehead ⁵⁹, E. Whiter ⁵³, A.R. Wiederhold ⁶²,
 D. Wiedner ¹⁹, G. Wilkinson ⁶³, M.K. Wilkinson ⁶⁵, M. Williams ⁶⁴, M.R.J. Williams ⁵⁸,
 R. Williams ⁵⁵, Z. Williams ⁵⁴, F.F. Wilson ⁵⁷, M. Winn ¹², W. Wislicki ⁴¹, M. Witek ⁴⁰,
 L. Witola ²¹, G. Wormser ¹⁴, S.A. Wotton ⁵⁵, H. Wu ⁶⁸, J. Wu ⁸, Y. Wu ⁶, Z. Wu ⁷,
 K. Wyllie ⁴⁸, S. Xian ⁷¹, Z. Xiang ⁵, Y. Xie ⁸, A. Xu ³⁴, J. Xu ⁷, L. Xu ^{4,b}, L. Xu ^{4,b},
 M. Xu ⁵⁶, Z. Xu ⁴⁸, Z. Xu ⁷, Z. Xu ⁵, D. Yang ^{4,b}, K. Yang ⁶¹, S. Yang ⁷, X. Yang ⁶,
 Y. Yang ^{28,m}, Z. Yang ⁶, Z. Yang ⁶⁶, V. Yeroshenko ¹⁴, H. Yeung ⁶², H. Yin ⁸, X. Yin ⁷,
 C.Y. Yu ⁶, J. Yu ⁷⁰, X. Yuan ⁵, Y. Yuan ^{5,7}, E. Zaffaroni ⁴⁹, M. Zavertyaev ²⁰,
 M. Zdybal ⁴⁰, F. Zenesini ^{24,j}, C. Zeng ^{5,7}, M. Zeng ^{4,b}, C. Zhang ⁶, D. Zhang ⁸,
 J. Zhang ⁷, L. Zhang ^{4,b}, S. Zhang ⁷⁰, S. Zhang ⁶³, Y. Zhang ⁶, Y.Z. Zhang ^{4,b}, Y. Zhao ²¹,
 A. Zharkova ⁴³, A. Zhelezov ²¹, S.Z. Zheng ⁶, X.Z. Zheng ^{4,b}, Y. Zheng ⁷, T. Zhou ⁶,
 X. Zhou ⁸, Y. Zhou ⁷, V. Zhovkovska ⁵⁶, L.Z. Zhu ⁷, X. Zhu ^{4,b}, X. Zhu ⁸, V. Zhukov ¹⁷,
 J. Zhuo ⁴⁷, Q. Zou ^{5,7}, D. Zuliani ^{32,p}, G. Zunica ⁴⁹

¹ School of Physics and Astronomy, Monash University, Melbourne, Australia

² Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

³ Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

⁴ Department of Engineering Physics, Tsinghua University, Beijing, China

⁵ Institute Of High Energy Physics (IHEP), Beijing, China

⁶ School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁷ University of Chinese Academy of Sciences, Beijing, China

⁸ Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁹ Consejo Nacional de Rectores (CONARE), San Jose, Costa Rica

- ¹⁰ *Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France*
- ¹¹ *Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*
- ¹² *Université Paris-Saclay, Centre d'Etudes de Saclay (CEA), IRFU, Saclay, France, Gif-Sur-Yvette, France*
- ¹³ *Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France*
- ¹⁴ *Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*
- ¹⁵ *Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France*
- ¹⁶ *LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*
- ¹⁷ *I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ¹⁸ *Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany*
- ¹⁹ *Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ²⁰ *Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ²¹ *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ²² *School of Physics, University College Dublin, Dublin, Ireland*
- ²³ *INFN Sezione di Bari, Bari, Italy*
- ²⁴ *INFN Sezione di Bologna, Bologna, Italy*
- ²⁵ *INFN Sezione di Ferrara, Ferrara, Italy*
- ²⁶ *INFN Sezione di Firenze, Firenze, Italy*
- ²⁷ *INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ²⁸ *INFN Sezione di Genova, Genova, Italy*
- ²⁹ *INFN Sezione di Milano, Milano, Italy*
- ³⁰ *INFN Sezione di Milano-Bicocca, Milano, Italy*
- ³¹ *INFN Sezione di Cagliari, Monserrato, Italy*
- ³² *INFN Sezione di Padova, Padova, Italy*
- ³³ *INFN Sezione di Perugia, Perugia, Italy*
- ³⁴ *INFN Sezione di Pisa, Pisa, Italy*
- ³⁵ *INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³⁶ *INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³⁷ *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ³⁸ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ³⁹ *AGH - University of Krakow, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ⁴⁰ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ⁴¹ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ⁴² *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ⁴³ *Affiliated with an institute covered by a cooperation agreement with CERN*
- ⁴⁴ *DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁴⁵ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁶ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁷ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- ⁴⁸ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴⁹ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁵⁰ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁵¹ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁵² *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁵³ *School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*
- ⁵⁴ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁵⁵ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁶ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁷ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵⁸ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁹ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁶⁰ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*

- ⁶¹ Imperial College London, London, United Kingdom
⁶² Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom
⁶³ Department of Physics, University of Oxford, Oxford, United Kingdom
⁶⁴ Massachusetts Institute of Technology, Cambridge, MA, United States
⁶⁵ University of Cincinnati, Cincinnati, OH, United States
⁶⁶ University of Maryland, College Park, MD, United States
⁶⁷ Los Alamos National Laboratory (LANL), Los Alamos, NM, United States
⁶⁸ Syracuse University, Syracuse, NY, United States
⁶⁹ Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to³
⁷⁰ School of Physics and Electronics, Hunan University, Changsha City, China, associated to⁸
⁷¹ Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to⁴
⁷² Lanzhou University, Lanzhou, China, associated to⁵
⁷³ School of Physics and Technology, Wuhan University, Wuhan, China, associated to⁴
⁷⁴ Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to¹⁶
⁷⁵ Ruhr Universitaet Bochum, Fakultät f. Physik und Astronomie, Bochum, Germany, associated to¹⁹
⁷⁶ Eotvos Lorand University, Budapest, Hungary, associated to⁴⁸
⁷⁷ Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to³⁷
⁷⁸ Universiteit Maastricht, Maastricht, The Netherlands, associated to³⁷
⁷⁹ Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland, associated to⁴⁰
⁸⁰ Universidade da Coruña, A Coruña, Spain, associated to⁴⁴
⁸¹ Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, associated to⁵⁹
⁸² University of Michigan, Ann Arbor, MI, United States, associated to⁶⁸

^a Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Rio De Janeiro, Brazil

^b Center for High Energy Physics, Tsinghua University, Beijing, China

^c Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China

^d School of Physics and Electronics, Henan University, Kaifeng, China

^e LIP6, Sorbonne Université, Paris, France

^f Lamarr Institute for Machine Learning and Artificial Intelligence, Dortmund, Germany

^g Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras

^h Università di Bari, Bari, Italy

ⁱ Università di Bergamo, Bergamo, Italy

^j Università di Bologna, Bologna, Italy

^k Università di Cagliari, Cagliari, Italy

^l Università di Ferrara, Ferrara, Italy

^m Università di Genova, Genova, Italy

ⁿ Università degli Studi di Milano, Milano, Italy

^o Università degli Studi di Milano-Bicocca, Milano, Italy

^p Università di Padova, Padova, Italy

^q Università di Perugia, Perugia, Italy

^r Scuola Normale Superiore, Pisa, Italy

^s Università di Pisa, Pisa, Italy

^t Università della Basilicata, Potenza, Italy

^u Università di Roma Tor Vergata, Roma, Italy

^v Università di Siena, Siena, Italy

^w Università di Urbino, Urbino, Italy

^x Universidad de Alcalá, Alcalá de Henares, Spain

^y Facultad de Ciencias Físicas, Madrid, Spain

^z Department of Physics/Division of Particle Physics, Lund, Sweden

[†] Deceased