



The value of visualization in improving compound flood hazard communication: A new perspective through a Euclidean Geometry lens

5 Soheil Radfar^{1,2*}, Georgios Boumis^{1,2}, Hamed R. Moftakhari^{1,2}, Wanyun Shao³, Larisa Lee⁴, Alison N. Rellinger^{4,5}

¹ Center for Complex Hydrosystems Research, Tuscaloosa, AL, USA

² Department of Civil, Construction, and Environmental Engineering, University of Alabama, Tuscaloosa, AL, USA

³ Department of Geography & the Environment, University of Alabama, Tuscaloosa, AL, USA

10 ⁴ Coastal Research and Extension Center, Mississippi State University, Biloxi, MS, USA

⁵ Mississippi-Alabama Sea Grant Consortium, Ocean Springs, MS, USA

Correspondence to: Soheil Radfar (sradfar@ua.edu)

Abstract. Compound flooding, caused by the sequence/co-occurrence of flood drivers (i.e. river discharge and elevated sea
15 level) can lead to devastating consequences for society. Weak and insufficient progress toward sustainable development and disaster risk reduction are likely to exacerbate the catastrophic impacts of these events on vulnerable communities. For this reason, it is indispensable to develop new perspectives on evaluating compound flooding dependence and communicating the associated risks to meet UN Sustainable Development Goals (SDGs) related to climate action, sustainable cities, and sustainable coastal communities. An indispensable first step for studies examining the dependence between these bivariate
20 extremes is plotting the data in the variable space, i.e., visualizing a scatterplot, where each axis represents a variable of interest, then computing a form of correlation between them. This paper introduces the Angles method, based on Euclidean geometry of the so-called “*subject space*,” for visualizing the dependence structure of compound flooding drivers. Here, we evaluate, for the first time, the utility of this geometric space in computing and visualizing the dependence structure of compound flooding drivers. To assess the effectiveness of this method as a risk communication tool, we conducted a survey
25 with a diverse group of end-users, including academic and non-academic respondents. The survey results provide insights into the perceptions of applicability of the Angles method and highlight its potential as an intuitive alternative to scatterplots in depicting the evolution of dependence in the non-stationary environment. This study emphasizes the importance of innovative visualization techniques in bridging the gap between scientific insights and practical applications, supporting more effective compound flood hazard communication in a warming climate.



30 1 Introduction

Compound flooding from terrestrial (i.e. river discharge) and coastal (i.e. storm surge) drivers due to long-lasting (extra)tropical cyclones can have severe social and economic impacts for coastal communities around the globe (Zscheischler et al., 2018). Compound flooding risks are becoming more prevalent due to climate change, presenting a major sustainability challenge worldwide (Chan et al., 2024; Lai et al., 2021). A comprehensive risk communication strategy is essential to engaging stakeholders and informing decision-making and mitigation efforts (Khan and Mishra, 2022), as well as supporting the UN Sustainable Development Goals (SDGs), specifically, SDG Target 11.5, which calls for reducing the adverse effects of natural disasters. The primary barrier to anticipating and responding to compound flood events is the lack of effective communication (Kruczkiewicz et al., 2021).

To date, many researchers have extensively explored the likelihood of co-occurrence of anomalously large river discharges and high sea water levels, at both local (Kim et al., 2023) and global (Couasnon et al., 2020) scales over the years (Radfar et al., 2024; Green et al., 2024). This type of analysis is well-grounded in the scientific literature and can be done by analysing the dependence structure of coinciding extremal samples of the variables of interest.

The first key step of bivariate dependence analysis is to exhibit the variables of interest graphically in the variable space, via a scatterplot, where each axis represents a variable, and then calculate the correlation coefficient between them, e.g., the linear Pearson's r or the non-linear Kendall's τ or Spearman's ρ . This kind of visualization and computation of the dependence is prevalent in current scientific literature. To name a few examples, Robins et al. (Robins et al., 2021) plot coinciding extremes of river discharge and skew surge from two estuaries in the UK using a scatterplot and then calculate the Kendall's τ , while Jane et al. (Jane et al., 2022) use the variable space for depicting the relationship between concurrent extremal values of storm surge and river discharge for three sites along the Texas Gulf Coast and subsequently compute the Kendall's τ . Nasr et al. (Nasr et al., 2021) also follow Kendall's τ approach for quantifying dependence among different pairs of environmental extremes, including river discharge and storm surge across 36 coastal sites in the US. The variable space, however, does pose a limitation to studying the dependence structure, in the sense that it strongly places emphasis on the individual observations (subjects) themselves, which are denoted by points on the scatterplot, rather than the two variables for which inference is sought as generic entities. However, a deeper understanding of multivariate statistics and particularly of bivariate dependence requires an effective and intuitive way of visualizing the relationship between variables with minimal focus on individual subjects. This is particularly important when the dimensionality of the problem increases, e.g., when an additional time dimension is introduced, to examine potential non-stationarities in the dependence structure of two variables. To overcome these limitations, we propose the Angles method, which uses Euclidean geometry to visualize the relationship between flood drivers in an intuitive way.


In reality, multivariate statistics have a strong flavor of Euclidean geometry (Farnsworth, 2000; Friendly et al., 2013), which in turn can be an aid to unraveling the relationship between compound flooding drivers. Unlike the variable space and the scatterplot, if we think about the data in the “*subject space*” instead, where each subject (observation) of coinciding



extremal pairs defines an axis (dimension), then the two variables can be represented as two points inside that space (Wickens, 2014). The idea of the subject space, although long known in statistical scientific literature, has rarely been explored in environmental sciences, let alone in studies concerning environmental extremes that may lead to flooding. In this note, we demonstrate how the use of this geometric space provides an alternative way of studying the dependence structure between environmental bivariate extremes, specifically river discharge and storm surge. What sets our work apart is its application in a multivariate non-stationary context, where it enhances risk communication by providing an insightful means of visualizing evolving dependencies. Effective risk communication is a critical component in disaster risk reduction (Fakhruddin et al., 2020; Pile et al., 2018) as it helps to inform, engage and educate vulnerable communities and stakeholders about the risks associated with natural hazards (Auermüller, 2019). This is of paramount importance to improve resilience against compound flooding, which is becoming an increasing threat to coastal communities in the changing climate (Bevacqua et al., 2020; Ghanbari et al., 2021). In this regard, the present study evaluates the effectiveness of the Angles method in visualizing evolving dependencies in compound flooding, emphasizing its potential for enhanced risk communication.

2. Materials and Methods

For our analysis, we first used still water level data, composed of mean sea level, astronomical tide, and non-tidal residual, from tide gauges at Washington, DC, and Baltimore, MD, extracted from the GESLA3 database (Haigh et al., 2023). To extract the non-tidal residual, i.e., the storm surge, we performed tidal harmonic analysis on a rolling-year basis involving 60 major tidal constituents. Additionally, we utilized discharge data from rivers that drain to the respective tidal river outlet of each city, originating from the Global Runoff Data Center (GRDC) (Recknagel et al., 2023). Figure 1 illustrates the pairs of annual maximum discharge (Q) and respective maximum surge (S) within (+/-) a day of maximum-discharge timing (i.e., coinciding extremes) for the two coastal cities. In the case of the freshwater-influenced tide gauge at Washington, DC, the scatterplot is constructed from all available 83 years of measurements and the linear Pearson's r correlation coefficient is found to be 0.96, while the non-linear Spearman's ρ correlation coefficient is 0.84. On the other end, the scatterplot at Baltimore, MD, tide gauge is produced from available records of 54 years and the correlation coefficients are found to be weaker with Pearson's r and Spearman's ρ being 0.41 and 0.52, respectively (see also (Nasr et al., 2021) for Kendall's τ). For both tide gauges, years that have more than 20% of missing data were not considered for the illustration of scatterplots and the calculation of correlation coefficients.



90 **Figure 1. Scatterplots of discharge (Q) annual maxima and surge (S) maxima within ± 1 day of the maximum-discharge timing for: (a) Washington, DC; (b) Baltimore, MD.**

In addition to traditional correlation analyses, the Angles method is applied to assess and visualize dependencies, offering a new perspective on compound flood dynamics. In the subject space of the data at Washington, DC, which consists of 83 axes (dimensions), equal to the number of pairs (subjects), discharge and storm surge can be defined by the two points:

$$Q = (4615, 3436, 3398, \dots, 3086) \text{ and } S = (1.03, 0.88, 0.57, \dots, 0.61) \quad (1)$$

95 Likewise, the data at Baltimore, MD, can be thought of as belonging to a 54-dimensional subject space where discharge and storm surge are simply two points:

$$Q = (2449, 5748, 2973, \dots, 7673) \text{ and } S = (0.26, 0.20, -0.01, \dots, 0.27) \quad (2)$$

Picturing variables Q and S in such high-dimensional spaces is obviously an impossible task for the human mind. Despite this limitation, the two points (Q and S), together with the origin (O) of the subject space, form a 2-D plane which is easy to grasp, and thus discharge and storm surge can be plotted as two vectors (starting from the origin and extending to the respective point). For the sake of simplicity, we can center the two variables by subtracting the respective mean value of each variable so that the origin of the 2-D plane becomes zero, while their correlation and variances remain unchanged:

$$q = Q - \bar{Q} \text{ and } s = S - \bar{S} \quad (3)$$

From Euclidean geometry, we know that the length of a vector, e.g., the discharge vector ($\bar{q}$) is given by the following formula:



$$|\bar{q}| = \sqrt{q_1^2 + q_2^2 + q_3^2 + \dots + q_N^2} \quad (4)$$

The squared length of $\bar{q}$ is then equivalent to the sum of squared deviations from the zero mean:

$$|\bar{q}|^2 = \sum_{i=1}^N q_i^2 \quad (5)$$

105 Hence, the length of vector $\bar{q}$ is directly related to the unbiased estimator of the standard deviation of discharge population:

$$\sigma_q = \frac{|\bar{q}|}{\sqrt{N-1}} \quad (6)$$

Correspondingly, it follows the same for the standard deviation of surge population:

$$\sigma_s = \frac{|\bar{s}|}{\sqrt{N-1}} \quad (7)$$

Euclidean geometry, and particularly trigonometry, indicates also that the cosine of the angle between two vectors is equal to their dot product ($\cdot$) over the product of their lengths, as shown below:

$$\cos(\theta) = \frac{\bar{q} \cdot \bar{s}}{|\bar{q}| |\bar{s}|} \quad (8)$$

110 where $\theta = \angle(\bar{q}, \bar{s})$, and $\bar{q} \cdot \bar{s} = q_1 s_1 + q_2 s_2 + q_3 s_3 + \dots + q_N s_N$. It is now easy to see that the expression in Equation 8 matches that of Pearson's r correlation coefficient:

$$r = \cos(\theta) = \frac{\bar{q} \cdot \bar{s}}{|\bar{q}| |\bar{s}|} = \frac{\sum_{i=1}^N q_i s_i}{\sqrt{\left(\sum_{i=1}^N q_i^2\right) \left(\sum_{i=1}^N s_i^2\right)}} \quad (9)$$

115 In the subject space, uncorrelated discharge and surge variables are displayed as perpendicular vectors ($\theta = 90^\circ$), whereas correlated discharge and surge variables are displayed as collinear vectors ($\theta = 0^\circ$ or $\theta = 180^\circ$). Many times, in multivariate statistics, variables are not only centered around zero but also scaled by dividing them with their standard deviation. In such an instance, the standard deviation of each variable becomes then one and thus from Equations 6 and 7 it follows that the vectors $\bar{q}$ and $\bar{s}$ have the same length, only dependent on N . For convenience, one may choose to work with vectors of unit length $|\bar{q}| = |\bar{s}| = 1$ and hence the constant $\sqrt{N-1}$ can be neglected – then, the only characteristic of the



two vectors that truly matters is the angle between them. Consequently, a greater angle θ , i.e., a smaller Pearson's r will lead to a bigger parallelogram area between the two vectors since Euclidean geometry suggests that:

$$\text{Area}_{\text{parallelogram}} = |\vec{q}||\vec{s}|\sin(\theta) = \sin(\theta) = \sqrt{1 - (\cos[\theta])^2} = \sqrt{1 - r^2} \quad (10)$$

3. Results and discussion

3.1. Application of the Angles method for visualizing CCF dependencies

Figure 2 shows the two variables (Q and S) represented as unit-length vectors ($\vec{q}$ and $\vec{s}$) on a 2-D plane of the subject space with zero origin. As opposed to scatterplots, these graphs show the two variables as general entities rather than individual observations. The angle between the two vectors is proportional to how dependent they are, as shown in Equations 9 and 10.

Observe the small and large angles (or, equivalently, parallelogram areas) that the two vectors form at Washington, DC, and Baltimore, MD, respectively. In fact, for the two cases, computing the vectors' lengths as well as the dot product of the discharge and surge vectors and then plugging them in Equation 8, yields $\cos(\theta) = r = 0.96 \Rightarrow \theta \sim 16^\circ$ and $\cos(\theta) = r = 0.41 \Rightarrow \theta \sim 66^\circ$, respectively. In Washington, DC, the smaller angle between the unit vectors indicates a tighter interplay between river discharge and storm surge. This suggests a higher correlation and potential for severe compound flooding events. Conversely, in Baltimore, MD, the larger angle suggests a lower degree of correlation. This indicates that while both factors are relevant, they may not coincide as frequently to create severe compound events.




Figure 2. Discharge (Q) and surge (S) variables represented as unit-length vectors in the subject space for: (a) Washington, DC where $\cos[\theta] = 0.96$; (b) Baltimore, MD with $\cos[\theta] = 0.41$.

The subject space can be of greater value when dealing with more than two variables, e.g., multi-driver compound flooding from discharge, surge, precipitation, and wind waves. It is inherently difficult to illustrate 4-dimensional



scatterplots, and the interactions of multiple flooding drivers cannot be visually captured by such a plot. Euclidean geometry can, nevertheless, be a helpful tool for such an analysis since all variables can be projected as vectors on a 2-D plane and their interrelations can be visualized as explained herein. In addition, plots like those of Figure 2 can be a great tool for visualization of the change of dependence structure over time; if human-induced climate change is making the co-occurrence of flood drivers more likely (Wahl et al., 2015), this can be visualized by a frame with a shrinking angle θ . For example, Figure 3 illustrates a scatterplot of bivariate sampling where the y axis shows annual maxima still water levels at Galveston Pier 21, TX, while the x axis represents co-occurring (± 5 days) maxima of discharge at Buffalo Bayou which drains into Galveston Bay – data from different time periods are highlighted with different colors. From Figure 3 alone, it is not evident if the dependence between the two flooding drivers is getting stronger with time. Specifically, it appears rather hard to determine (by a mere visualization) whether the correlation coefficient from 1972-1996 is greater than that of the period from 1997-2022. In many times, the scatterplot fails to reveal evolving patterns of dependence.




Figure 3. Scatterplot of annual maxima sea levels and discharge maxima within ± 5 days of the maximum-sea-level timing for Galveston Bay, TX. Pairs are colored based on period of observation.

On the contrary, visualizing the variables as unit-length centred vectors again, where the pair-wise angle is the only key characteristic between them, allows us to infer that the dependence between extreme sea levels and river discharge at Galveston Bay has been increasing over time (observe the shrinking angle θ in Figure 4). From Figure 4, it is clear that the correlation coefficient of the period 1997-2022 is greater than that of 1972-1996 since θ is smaller (thus, the cosine is greater). This evolving trend is a sign of non-stationarity in dependence structure, which is per se a difficult concept to communicate to a variety of stakeholder groups. Presenting this simple graph underscores the importance of considering temporal changes in dependence structure when planning and implementing flood risk management strategies. This dynamic understanding aligns with adaptive management principles in coastal engineering. It necessitates continuous monitoring and



re-assessment of flood risks considering potential non-stationarity in hydrodynamic, hydrological and meteorological relationships.




Figure 4. Subject space showing a stronger dependence between sea levels and discharge over non-overlapping time periods (end year shown in red), i.e., a shrinking angle θ between the two vectors at Galveston Bay, TX.

Another important aspect within compound flooding framework is non-stationarities of the dependence structure among flood drivers. In the context of a warming climate, depending solely on stationarity assumptions could be insufficient (Milly et al., 2008), as it does not consider increasing changes in variation of flood drivers due to climate change (Kim et al., 2018). Natural climatic variability and anthropogenic climate change are among the most important drivers of non-stationarity (Galiatsatou and Prinos, 2011). Ignoring this effect can easily lead to misinterpretation of results in a multivariate framework (Radfar and Galiatsatou, 2023; Corbella and Stretch, 2012). Non-stationarity also influences the dependence structures among compound flood drivers over time (Naseri and Hummel, 2022). Using non-stationary dynamic copulas is a reliable solution for better understanding of variability and long-term trends (Pirani and Najafi, 2023). However, due to complexities in using this approach, the existing literature commonly relies on moving window approach or simplifying the assumption of stationarity of the dependence structure (Radfar et al., 2023). Public perception of this impact is even more challenging. The expected annual economic losses due to compound flooding damage amount to billions of dollars. Yet, the knowledge about non-stationarity in compound flood drivers is still very limited among practitioners and stakeholders and this could hinder proper preparedness and mitigation efforts against this increasing risk to coastal communities. To disseminate information about changing dependence structures to the target audience, it would be necessary to adopt effective communication approaches. Figure 5 illustrates how non-stationarity in the dependence of the two variables over multiple, possibly overlapping time periods, can be effectively visualized with the use of the subject space. Observe, for example, how θ shrinks from an obtuse angle in 1950-1991 (past) to an acute angle in 1982-2021 (present), indicating that the negative correlation between discharge and sea level extremes has gradually evolved into a strong positive dependence over time. A unique characteristic of semi-circular representation of Figure 5 is its capability to encompass equal, unit-length vectors to



clearly depict an evolving correlation among flood drivers over time. This easy-to-follow visualization technique could help overcome challenges in communicating with non-experts, aiding in their better understanding of the shifting dependence between multiple flood drivers, and ultimately, motivating them about compound flood risk mitigation efforts. It is expected that such simple visualization efforts will better reflect climate change effects and emphasize the need for resilient infrastructure and adaptive measures to safeguard against flood risks. Ultimately, this enables vulnerable coastal communities to remain resilient and sustainable in the face of a warming climate, which is an overarching objective of SDGs 11 (Sustainable Cities and Communities) and 13 (Climate Action).




Figure 5. Subject space showing a stronger dependence between sea levels and discharge across multiple overlapping time periods, i.e., a shrinking angle θ between the two vectors at Galveston Bay, TX. Observe how an obtuse angle, i.e., a negative correlation in the past, gradually transforms into an acute angle indicating strong positive correlation.


3.2. Evaluating stakeholder perceptions of the Angles method for CCF risk communication

To assess the end users' perceptions of the effectiveness of the Angles method and the subject space for visualizing CCF dependencies, we conducted a survey targeting a diverse group of end users. The survey was administered online from July 10 to September 10, utilizing various distribution channels to reach a broad audience. These channels included email lists, members of the working Group 4 of the Cooperative Institute for Research to Operations in Hydrology (CIROH) institution which is responsible for impact-based decision-making research, stakeholders from the NOAA project "Coastal Nature-Based Solutions to Mitigate Flood Impacts and Enhance Resilience," and the network of the Program for Local Adaptation to Climate Effects (PLACE). This distribution strategy allowed us to gather input from a wide range of respondents, including members from the academia, industry, non-governmental organizations (NGOs), and state, federal, and local government agencies.



205 The survey consisted of questions designed to gauge the respondents' familiarity with CCF dependencies, the clarity of
 non-stationarity concepts, and the effectiveness of the Angles method in communicating risk. Likert scale questions were
 used to capture the degree of agreement or disagreement on various aspects of the Angles method, including its
 understandability, applicability, and perceived usefulness in CCF risk communication. The survey collected 91 complete
 responses. The top panel in Figure 6 shows a world map highlighting the global reach of the survey, with participants spread
 across multiple continents. Respondents were primarily concentrated in the United States, which accounted for most
 210 responses (64), followed by the United Kingdom (5), India (4), France (3), the Netherlands, Spain, Australia (2), and nine
 other countries with one respondent. This distribution reflects the diverse global engagement with the survey, capturing
 perspectives from a wide range of regions and sectors. In the United States, the survey responses came from 20 states, with
 the highest numbers reported in Mississippi (14), Alabama (13), and Florida (10) along the Gulf Coast. This concentration is
 primarily due to the survey distribution channels, which are closely connected to organizations and projects in this region.

215 The responses are subsequently grouped into two categories: academic and non-academic respondents. This
 classification is used to evaluate the differing perceptions of the proposed Angles method between these two groups.
 Academic respondents primarily included researchers, faculty, and students from various universities, while non-academic
 respondents comprised professionals from the industry, government agencies, and NGOs. This segmentation allows us to
 explore how familiarity, relevance, and clarity of the Angles method differed across these distinct sectors. The bottom panel
 220 in Figure 6 presents bar charts comparing the responses of academic ($n = 44$) and non-academic ($n = 47$) respondents to
 questions regarding their familiarity with CCF, the relevance of CCF to their work, years of experience, and familiarity with
 the concept of non-stationarity. Combining the "very well" and "extremely well" categories, non-academics show a higher
 total of 68.1% compared to 59% for academics. The relevance of CCF to respondents' work was high for both groups, with
 56.8% of academics and 70.2% of non-academics reporting it as "extremely" relevant. Similarly, regarding years of
 225 experience, non-academics showed a higher proportion (59.6%) with extensive experience compared to academics (47.7%).
 Interestingly, familiarity with non-stationarity concepts revealed a more pronounced divide, with 25% of academics
 reporting being "extremely well" familiar with non-stationarity, compared to only 14.9% of non-academics in the same
 category. This difference becomes even more pronounced when considering those who are less familiar with the concept.
 Notably, 34% of non-academics reported being "not well at all" familiar with non-stationarity, which is significantly higher
 230 than the 13.6% of academics in the same category. This disparity might reflect the theoretical and complex nature of non-
 stationarity, which may be more frequently encountered in academic research.



235 **Figure 6. Geographic distribution and knowledge assessment of the survey respondents. Top panel: World map showing the geographic distribution of survey respondents (91 total respondents) who participated in the study on compound flood risk communication. Countries are color-coded based on the number of respondents, ranging from 1 to over 10. Bottom panel: Bar charts depicting the Likert scale responses from academic (44 respondents) and non-academic (47 respondents) respondents on their familiarity with CCF, relevance of CCF to their work, years of experience, and familiarity with non-stationarity concept.**

240 Figure 7 presents a detailed comparison of academic and non-academic respondents' perceptions of various aspects of CCF risk communication, utilizing Likert scale responses. The bar chart highlights how these groups responded to various



aspects of the Angles method, including correlation understandability, non-stationarity clarity, effectiveness in CCF risk communication, and the likelihood of applying the method in their work or public communication.


First, we evaluated the understandability of correlation using strictly numerical values (i.e., correlation coefficients) versus the Angles method, which incorporates numerical values into a visual representation. For the numerical approach, academic respondents showed a higher level of agreement (50% agree or strongly agree) compared to non-academics (35.9%). When assessing the Angles method, academic respondents maintained a similar level of agreement, with 68.2% are at least slightly agree with its understandability. However, among non-academics, the level of high agreement ("agree" and "strongly agree") dropped to 19.2%. This shift can partly be related to the findings from Figure 6, where most respondents reported significant familiarity with compound flooding, and accordingly, the concept of dependence between flood drivers. This familiarity suggests that respondents are accustomed to traditional correlation coefficients, which may bias them toward these conventional methods rather than accepting a new visual approach for communicating a rather simple concept of correlation between flood drivers.

While comparing a method like the Angles approach, that incorporates familiar numerical values into a visual representation, against strictly numerical values may seem unconventional, especially among experts, it is crucial for establishing a benchmark of the new method's capabilities. Previous studies have consistently demonstrated that when communicating with the public, numbers, graphs, and technical concepts often fall short in effectively conveying the importance of hazards and risks (Morrow et al., 2015; Kuser Olsen et al., 2018). Visualization has proven to be a key tool for enhancing understanding, engagement, and decision-making (Atasoy et al., 2022; Colle et al., 2023). Thus, evaluating the Angles method against traditional numerical values was necessary to understand how well it performs relative to established approaches, even within an expert perspective. The results, although showing lower levels of familiarity among non-academics, remain promising and acceptable, underscoring the Angles method's potential as a more intuitive alternative that could bridge gaps in understanding when deployed beyond expert audiences.

Next, building upon the initial comparison, and to ensure a one-to-one comparison, we evaluated the Angles method against scatterplots for representing non-stationarity (here, in the form of evolving dependencies). For scatterplots, 40.9% of academics believe the scatterplot clearly depicts variations in dependence (agreeing or strongly agreeing), whereas only 27.7% of non-academics indicated such. However, when considering clarity for a potential audience, both groups expressed lower confidence, with only 11.3% of academics and 4.3% of non-academics agreeing or strongly agreeing. These findings clearly imply the complexity of the non-stationarity concept and the challenge of communicating it for non-academics and audience of both groups. Interestingly, both the academics and non-academics reported that the Angles method offered improved clarity for them and their potential audiences. For respondent clarity, 77.3% of academics at least slightly agreed, compared to 68.1% of non-academics. For presumed clarity to a hypothetical prospective audience, the Angles method enhanced the level of agreeing or strongly agreeing from 11.3% to 34.1% for academics, and from 4.3% to 23.4% for non-academics, compared to scatterplots.



275 Finally, Figure 7 further illustrates the effectiveness of the Angles method in CCF risk communication and the
likelihood of applying it in professional settings. Regarding the effectiveness in CCF risk communication, academic
respondents appeared more positive, with 31.8% agreeing or strongly agreeing, versus 23.4% of non-academics. Academic
respondents showed a strong consensus on the method’s practical application, with 75% expressing a likelihood of applying
it in their work or research. In contrast, non-academic respondents were more divided, with 53.2% expressing some
280 likelihood of applying it, but with a notable 21.8% disagreeing or remaining neutral, suggesting a hesitancy to adopt the
method without further familiarization. For public communication, both groups turned into higher strongly agreement and
lower agreement.



285 **Figure 7. Bar chart of Likert scale responses comparing academic ($n = 44$) and non-academic ($n = 47$) perceptions of CCF risk communication. The figure shows the percentage distribution of responses on correlation understandability, non-stationarity clarity, effectiveness in risk communication, and likelihood of applying in work/research and public communication.**

Figure 8 illustrates the relationships between various aspects of CCF understanding, risk communication, and application likelihood. For academic respondents, the figure shows that those with greater familiarity with CCF and those who find CCF highly relevant to their work tend to believe that scatterplots are not effective tools for communicating non-
290 stationarity to audiences, as indicated by the negative correlations. This pattern is similarly observed among respondents



with higher degrees, more experience, and familiarity with non-stationarity concepts, suggesting a general skepticism toward traditional scatterplot use in conveying complex, evolving relationships.

Conversely, when the Angles method is used to represent non-stationarity, there is a notable positive shift in correlations. This significant positive relationship suggests that academic respondents who were initially critical of scatterplots found the Angles method to be a more effective visual tool for communicating non-stationarity. This shift underscores the potential of the Angles approach to address perceived gaps in traditional risk communication methods among those with advanced familiarity and expertise, highlighting its value in enhancing the understanding of dynamic flood risk dependencies.

Among non-academic respondents, varying correlations are observed. This divergence might reflect differences in how these factors influence openness to the CCF communication methods in academic versus practical settings. Notably, the degree of non-academic respondents shows moderate positive correlations with years of experience in hydrologic or hydrodynamic fields (0.45) and familiarity with CCF (0.31), but weak or negative correlations with most other factors. This could suggest that while higher degrees are associated with more experience and familiarity, they don't necessarily translate to increased clarity or likelihood of applying new communication methods. The figure reveals that for non-academic respondents, the relevance of CCF to their work shows positive correlations with most factors. It exhibits stronger positive correlations with the Angles method compared to traditional methods like numerical values or scatterplots. This suggests that non-academics who find CCF relevant to their work are more likely to perceive the Angles method as an effective tool for understanding and communicating complex dependencies, compared to more traditional approaches.

It is important to note that using language like “new,” “groundbreaking,” or “different” can sometimes bias people against trying or adopting unfamiliar methods, as they tend to prefer what is familiar. In future discussions with audiences that may be hesitant to adopt the Angles method, emphasizing that it builds upon familiar concepts like correlation coefficients by adding a visual element, rather than contrasting with them, may increase the likelihood of its adoption.

Another pattern observed in the figure is that respondents who find each of the methods clear for themselves also believe it would be clear for their audience. This relationship is particularly pronounced among non-academic respondents, where there are significantly stronger positive correlations between the clarity of the methods for the respondent and its perceived clarity for the audience. This suggests that non-academics who understand these methods well are more confident in their effectiveness as a communication tool for broader audiences.




Figure 8. Correlation matrices comparing academic ($n = 44$) and non-academic respondents ($n = 47$) on familiarity, understanding, effectiveness, and likelihood of practical application of the Angles method compared to the traditional approaches (correlation values and scatterplots) for compound flood risk communication. The heatmaps display correlation coefficients, with color gradients indicating the strength and direction of correlations (blue for positive, red for negative). Circle sizes in the lower triangle represent correlation magnitudes.

4. Conclusions

This study evaluated the Angles method and the subject space as an alternative approach for visualizing and communicating the dependence structures of CF drivers, particularly in non-stationary environments. The Angles method leverages Euclidean geometry to transform numerical dependencies into visual angles, where each angle represents the relationship between flood drivers. This geometric representation allows for a more intuitive understanding of the complex dependencies compared to traditional numerical correlations (Section 2). By augmenting statistical relationships with visual



patterns, the Angles method provides an accessible way to identify changes in dependencies over time, making it a powerful tool for non-stationary risk communication (Section 3.1). Our findings reveal that the Angles method offers sensible advantages over traditional scatterplots, especially in enhancing the understanding and communication of evolving dependencies among CF drivers (Figure 7).

The survey results demonstrated that the method was primarily evaluated among a group of experienced respondents from the academic, industry, and government sectors (Figure 6). Academic respondents generally reported higher familiarity with CF dependencies and perceived the Angles method as more effective in enhancing communication of dependencies between compound flood drivers compared to traditional approaches (Figure 7). In contrast, non-academic respondents exhibited varying levels of familiarity and clarity, indicating a need for tailored communication strategies when presenting new methods like the Angles approach to diverse stakeholder groups.

Comparisons between the Angles method and scatterplots revealed that the Angles method provided a clearer and more intuitive representation of non-stationarity, particularly for academic respondents (Figure 7). This suggests that the Angles method can effectively fill existing gaps in traditional risk communication by offering a visual alternative that captures the dynamic nature of CF dependencies. Non-academic respondents also showed more positive correlations between the relevance of the Angles method to their work compared to traditional methods (Figure 8), indicating its potential alignment with practical needs in real-world flood management contexts.

The survey also highlighted a pattern where those who found the Angles method clear for themselves believed it would also be clear for others, with this effect being particularly pronounced among non-academic respondents. This underscores the Angles method's potential to facilitate effective communication beyond expert audiences, bridging gaps between scientific insights and practical applications in flood risk communication.

These findings highlight the opportunity to further develop the Angles method for communication with a non-technical audience. Given that the current evaluation focused on experienced respondents (Figure 6), future studies should explore the effectiveness of the Angles method with broader audiences, including the public and students. Engaging educational initiatives, such as those supported by the Scientific Research and Education Network (SciREN; <https://sciren.ua.edu/>), would provide valuable insights into how well this method communicates complex flood risk information to non-expert audiences. Such evaluations would not only validate the Angles method's utility across different groups but also enhance its role in scientific education and public understanding of environmental risks.



360 **Code availability**

The code used for the analysis and visualization in this study, including the implementation of the Angles method for visualizing compound flood drivers and the analysis of survey results, is available in the GitHub repository: <https://github.com/sradfar/CFnonStatViz>.

Data availability

365 The data supporting the findings of this study, including the survey results and relevant datasets used for visualizing the dependence structure of compound flood drivers, are available in the GitHub repository. Additional data, such as river discharge and sea level data, can be accessed from publicly available sources like the Global Runoff Data Center (GRDC) and the GESLA3 database. The GitHub repository can be accessed at: <https://github.com/sradfar/CFnonStatViz>.

Author contribution

370 GB, SR, and HMR conceptualized the study. SR and GB contributed to the methodology, with GB focusing on the statistical model and SR handling the survey analysis. WS, LL, and ARN assisted with preparing, reviewing, and distributing the survey. Data curation and analysis were conducted by SR and GB. SR and GB wrote the original draft, and all co-authors reviewed and provided feedback on the manuscript.

Competing interests

375 The authors declare that they have no conflict of interest.

Ethical statement

This research received ethical approval from The University of Alabama's Institutional Review Board (IRB). Informed consent was obtained from all participants prior to participation, and the survey was conducted anonymously to ensure confidentiality.

380 **Acknowledgment**

We would like to thank all the individuals who participated in our survey for their valuable input.



Financial support

This study is funded by National Science Foundation awards #2223893 and #2238000. Partial support is also provided by National Oceanic and Atmospheric Administration (NOAA), awarded to the Cooperative Institute for Research on
 385 Hydrology (CIROH) through the NOAA Cooperative Agreement with The University of Alabama, NA22NWS4320003.

References

- Atasoy, G., Ertaymaz, U., Dikmen, I., and Talat Birgonul, M.: Empowering risk communication: Use of visualizations to describe project risks, *Journal of Construction Engineering and Management*, 148, 04022015, 2022.
- Auermuller, L.: Coastal Hazards Risk Communication: A Technical Assistance Transfer Project within the National
 390 Estuarine Research Reserve System, NOAA Office for Coastal Management, 2019.
- Bevacqua, E., Vousdoukas, M. I., Zappa, G., Hodges, K., Shepherd, T. G., Maraun, D., Mentaschi, L., and Feyen, L.: More meteorological events that drive compound coastal flooding are projected under climate change, *Communications earth & environment*, 1, 47, 2020.
- Chan, F., Lu, X., Li, J., Lai, Y., Luo, M., Chen, Y. D., Wang, D., Li, N., Chen, W.-Q., and Zhu, Y.-G.: Compound flood
 395 effects, challenges and solutions: Lessons toward climate-resilient Chinese coastal cities, *Ocean & Coastal Management*, 249, 107015, 2024.
- Colle, B. A., Hathaway, J. R., Bojsza, E. J., Moses, J. M., Sanders, S. J., Rowan, K. E., Hils, A. L., Duesterhoeft, E. C., Boorboor, S., and Kaufman, A. E.: Risk Perception and Preparation for Storm Surge Flooding: A Virtual Workshop with Visualization and Stakeholder Interaction, *Bulletin of the American Meteorological Society*, 104, E1232-E1240, 2023.
 400
- Corbella, S. and Stretch, D. D.: Predicting coastal erosion trends using non-stationary statistics and process-based models, *Coastal engineering*, 70, 40-49, 2012.
- Couasnon, A., Eilander, D., Muis, S., Veldkamp, T. I., Haigh, I. D., Wahl, T., Winsemius, H. C., and Ward, P. J.: Measuring compound flood potential from river discharge and storm surge extremes at the global scale, *Natural Hazards and Earth System Sciences*, 20, 489-504, 2020.
 405
- Fakhruddin, B., Clark, H., Robinson, L., and Hieber-Girardet, L.: Should I stay or should I go now? Why risk communication is the critical component in disaster risk reduction, *Progress in disaster science*, 8, 100139, 2020.
- Farnsworth, D. L.: The Geometry of Statistics, *The College Mathematics Journal*, 31, 200-204, 2000.
- Friendly, M., Monette, G., and Fox, J.: Elliptical insights: understanding statistical methods through elliptical geometry, *Statistical Science*, 1-39, 2013.
 410
- Galiatsatou, P. and Prinos, P.: Modeling non-stationary extreme waves using a point process approach and wavelets, *Stochastic Environmental Research and Risk Assessment*, 25, 165-183, 2011.
- Ghanbari, M., Arabi, M., Kao, S. C., Obeysekera, J., and Sweet, W.: Climate change and changes in compound coastal-riverine flooding hazard along the US coasts, *Earth's Future*, 9, e2021EF002055, 2021.
- Green, J., Haigh, I., Quinn, N., Neal, J., Wahl, T., Wood, M., Eilander, D., de Ruiter, M., Ward, P., and Camus, P.: A
 415 Comprehensive Review of Compound Flooding Literature with a Focus on Coastal and Estuarine Regions, *EGUsphere*, 2024, 1-108, 2024.
- Haigh, I. D., Marcos, M., Talke, S. A., Woodworth, P. L., Hunter, J. R., Hague, B. S., Arns, A., Bradshaw, E., and Thompson, P.: GESLA version 3: A major update to the global higher-frequency sea-level dataset, *Geoscience Data Journal*, 10, 293-314, 2023.
 420
- Jane, R., Wahl, T., Santos, V. M., Misra, S. K., and White, K. D.: Assessing the potential for compound storm surge and extreme river discharge events at the catchment scale with statistical models: sensitivity analysis and recommendations for best practice, *Journal of Hydrologic Engineering*, 27, 04022001, 2022.
- Khan, S. and Mishra, J.: Critical gaps and implications of risk communication in the global agreements—SFDRR, SDGs, and UNFCCC: 3 select case studies from urban areas of tropics in South Asia, *Natural hazards*, 111, 2559-2577, 2022.
 425



- Kim, H., Villarini, G., Jane, R., Wahl, T., Misra, S., and Michalek, A.: On the generation of high-resolution probabilistic design events capturing the joint occurrence of rainfall and storm surge in coastal basins, *International Journal of Climatology*, 43, 761-771, 2023.
- 430 Kim, S. U., Son, M., Chung, E.-S., and Yu, X.: Effects of Non-Stationarity on Flood Frequency Analysis: Case Study of the Cheongmicheon Watershed in South Korea, *Sustainability*, 10, 1329, 2018.
- Kruczkiewicz, A., Klopp, J., Fisher, J., Mason, S., McClain, S., Sheekh, N., Moss, R., Parks, R., and Braneon, C.: Compound risks and complex emergencies require new approaches to preparedness, *Proceedings of the National Academy of Sciences*, 118, e2106795118, 2021.
- 435 Kuser Olsen, V., Momen, B., Langsdale, S., Galloway, G., Link, E., Brubaker, K., Ruth, M., and Hill, R.: An approach for improving flood risk communication using realistic interactive visualisation, *Journal of Flood Risk Management*, 11, S783-S793, 2018.
- Lai, Y., Li, J., Gu, X., Liu, C., and Chen, Y. D.: Global compound floods from precipitation and storm surge: Hazards and the roles of cyclones, *Journal of Climate*, 34, 8319-8339, 2021.
- 440 Milly, P. C., Betancourt, J., Falkenmark, M., Hirsch, R. M., Kundzewicz, Z. W., Lettenmaier, D. P., and Stouffer, R. J.: Stationarity is dead: Whither water management?, *Science*, 319, 573-574, 2008.
- Morrow, B. H., Lazo, J. K., Rhome, J., and Feyen, J.: Improving storm surge risk communication: Stakeholder perspectives, *Bulletin of the American Meteorological Society*, 96, 35-48, 2015.
- Naseri, K. and Hummel, M. A.: A Bayesian copula-based nonstationary framework for compound flood risk assessment along US coastlines, *Journal of Hydrology*, 610, 128005, 2022.
- 445 Nasr, A. A., Wahl, T., Rashid, M. M., Camus, P., and Haigh, I. D.: Assessing the dependence structure between oceanographic, fluvial, and pluvial flooding drivers along the United States coastline, *Hydrology and Earth System Sciences*, 25, 6203-6222, 2021.
- Pile, J., Gouramanis, C., Switzer, A. D., Rush, B., Reynolds, I., and Soria, J. L. A.: Can the risk of coastal hazards be better communicated?, *International journal of disaster risk reduction*, 27, 439-450, 2018.
- 450 Pirani, F. J. and Najafi, M. R.: Nonstationary frequency analysis of compound flooding in Canada's coastal zones, *Coastal Engineering*, 182, 104292, 2023.
- Radfar, S. and Galiatsatou, P.: Influence of nonstationarity and dependence of extreme wave parameters on the reliability assessment of coastal structures-A case study, *Ocean Engineering*, 273, 113862, 2023.
- 455 Radfar, S., Galiatsatou, P., and Wahl, T.: Application of nonstationary extreme value analysis in the coastal environment—A systematic literature review, *Weather and Climate Extremes*, 100575, 2023.
- Radfar, S., Mahmoudi, S., Moftakhari, H., Meckley, T., Bilskie, M. V., Collini, R., Alizad, K., Cherry, J. A., and Moradkhani, H.: Nature-based solutions as buffers against coastal compound flooding: Exploring potential framework for process-based modeling of hazard mitigation, *Science of The Total Environment*, 173529, 2024.
- 460 Recknagel, T., Färber, C., Plessow, H., and Looser, U.: The Global Runoff Data Centre: A building block in the chain of reproducible hydrology, *Copernicus Meetings*, 2023.
- Robins, P. E., Lewis, M. J., Elnahrawi, M., Lyddon, C., Dickson, N., and Coulthard, T. J.: Compound flooding: Dependence at sub-daily scales between extreme storm surge and fluvial flow, *Frontiers in Built Environment*, 7, 727294, 2021.
- Wahl, T., Jain, S., Bender, J., Meyers, S. D., and Luther, M. E.: Increasing risk of compound flooding from storm surge and rainfall for major US cities, *Nature Climate Change*, 5, 1093-1097, 2015.
- 465 Wickens, T. D.: *The geometry of multivariate statistics*, Psychology Press 2014.
- Zscheischler, J., Westra, S., Van Den Hurk, B. J., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch, D. N., Leonard, M., and Wahl, T.: Future climate risk from compound events, *Nature Climate Change*, 8, 469-477, 2018.