

Carbon flow and microbial activity in the hydrothermally altered sediment of Guaymas Basin

GUANG-CHAO ZHUANG¹, ANDREW MONTGOMERY²
AND SAMANTHA JOYE³

¹Ocean University of China

²University of Montana

³University of Georgia

Presenting Author: zgc@ouc.edu.cn

The Guaymas Basin, in the Gulf of California, is a deep-sea hydrothermal system characterized by rapid accumulation of organic-rich sediments and active oceanic crust production [1]. Magmatic intrusions from the spreading center alters the overlying sediments, generating a wide range of hydrocarbons such as methane, short/long chain alkanes, polycyclic aromatic hydrocarbons and other low-molecular-weight metabolites (e.g., CO₂, H₂, ammonium, organic acids and alcohols [2, 3], which constitute energetic substrates for a diverse array of microorganisms that live and thrive under extreme conditions. We integrated geochemical analysis and experimental incubations with radiotracers to constrain the carbon flow and microbial activity in hydrothermally influenced sediments from the Guaymas Basin. Abundant methane, acetate, methanol and hydrocarbons were detected in the porewaters. Rate measurements suggested methane could be produced from bicarbonate, acetate, methanol and hydrocarbons. High oxidation rates versus low methanogenesis rates indicated that these substrates were predominantly consumed by non-methanogenic processes. Methane was anaerobically oxidized and AOM was generally coupled to sulfate reduction. Microbial activity was influenced by environmental factors, which also shaped the distribution of microbial communities. These results reveal a dynamic carbon flow mediated by various microorganisms and provide new insight into the carbon metabolism and cycling in seafloor hydrothermal systems.

[1] Teske, A., de Beer, D., McKay, L.J., Tivey, M.K., Biddle, J.F., Hoer, D. et al. (2016). The Guaymas Basin Hiking Guide to Hydrothermal Mounds, Chimneys, and Microbial Mats: Complex Seafloor Expressions of Subsurface Hydrothermal Circulation. *Frontiers in Microbiology*, 7(75).

[2] Bazylinski, D. A., Farrington, J. W., & Jannasch, H. W. (1988). Hydrocarbons in surface sediments from a Guaymas Basin hydrothermal vent site. *Organic Geochemistry*, 12(6), 547-558.

[3] Zhuang, G.-C., Montgomery, A., Samarkin, V.A., Song, M., Liu, J., Schubotz, F. et al. (2019) Generation and Utilization of Volatile Fatty Acids and Alcohols in Hydrothermally Altered Sediments in the Guaymas Basin, Gulf of California. *Geophysical Research Letters* 46: 2637-2646.