

11. Goupil, N., Rayson, H., Serraille, É., Massera, A., Ferrari, P.F., Hochmann, J.-R., and Papeo, L. (2024). Visual preference for socially relevant spatial relations in humans and monkeys. *Psychol. Sci.* 35, 681–693.
12. Krupenye, C., and Hare, B. (2018). Bonobos prefer individuals that hinder others over those that help. *Curr. Biol.* 28, 280–286.e5.
13. Hochmann, J.-R. (2023). Incomplete language-of-thought in infancy. *Behav. Brain Sci.* 46, e278.
14. Hafri, A., Papafragou, A., and Trueswell, J.C. (2013). Getting the gist of events: Recognition of two-participant actions from brief displays. *J. Exp. Psychol. Gen.* 142, 880–905.
15. Hafri, A., Trueswell, J.C., and Strickland, B. (2018). Encoding of event roles from visual scenes is rapid, spontaneous, and interacts with higher-level visual processing. *Cognition* 175, 36–52.
16. Vettori, S., Odin, C., Hochmann, J.-R., and Papeo, L. (2023). A perceptual cue-based mechanism for automatic assignment of thematic agent and patient roles. Preprint at PsyArXiv, <https://doi.org/10.31234/osf.io/ydgca>.
17. Hafri, A., and Firestone, C. (2021). The perception of relations. *Trends Cogn. Sci.* 25, 475–492.
18. Vong, W.K., Wang, W., Orhan, A.E., and Lake, B.M. (2024). Grounded language acquisition through the eyes and ears of a single child. *Science* 383, 504–511.
19. Fodor, J.A., and Pylyshyn, Z.W. (1988). Connectionism and cognitive architecture: a critical analysis. *Cognition* 28, 3–71.
20. Gomes, V. (2024). Whither developmental psycholinguistics? Preprint at PsyArXiv, <https://doi.org/10.31234/osf.io/zm57t>.

Neuroscience: A big step forward for motor control in *Drosophila*

Mert Erginkaya and Jan M. Ache*

Neurobiology and Genetics, Theodor-Boveri-Institute, Biocenter, Julius-Maximilians-University of Würzburg, Am Hubland, 97074 Würzburg, Germany

*Correspondence: jan.ache@uni-wuerzburg.de
<https://doi.org/10.1016/j.cub.2024.07.097>

Connectomics approaches are fundamentally changing the way scientists investigate the brain. Recently published connectomes have enabled dissection of the intricate motor circuits in the fly's version of the spinal cord on a synaptic level. This has allowed reconstruction of complete sensorimotor pathways in *Drosophila*.

In the supermarket, you've likely never consciously controlled the precise placement of your feet as you approached the fruit section, or pondered the muscle forces needed to pick up a mango. And yet, you have accomplished such demanding motor control tasks all your life. This is possible because your spinal cord does this job for you, while your brain is busy strategizing and asking higher-level questions, like: do I really want mango? Motor control is achieved through the exquisite interplay of descending input from the brain, rhythmically active premotor networks in the spinal cord, and local feedback from sensory organs in and on the limbs¹ — all tailored to coordinating movements required to complete the task at hand. The nervous system of fruit flies, which, as the name suggests, spend a considerable amount of their lives hunting for fruit, needs to solve similar motor control tasks (Figure 1A). In fact, the fly's nervous system is able to orchestrate delicate

movements of the body during flight — controlled by four power and 13 steering muscles per wing — and walking — driven by 108 muscles controlling 30 joints across six legs. While the field has made great progress at identifying core motor circuits for specific behaviors², it remains difficult to understand how these circuits interact and coordinate with each other.

Two recent papers^{3,4} from the Tuthill and Lee labs constitute a major step towards understanding how motor circuits in the fly's version of the spinal cord, the ventral nerve cord (VNC), control leg and wing movements to turn thoughts into action. The first, by Azevedo *et al.*³, presents the synaptic wiring diagram — of a female adult nerve cord (FANC). The second, by Lesser, Azevedo *et al.*⁴, uses the FANC connectome to investigate principles of motor circuit organization in the leg and wing control systems. Together with previously published connectomes of the

Drosophila brain^{5–7} and parallel efforts in a male VNC⁸, these groundbreaking papers make it possible to trace sensorimotor pathways throughout the entire nervous system.

Azevedo *et al.*³ applied machine learning tools to analyse over 20 million electron microscopy (EM) images⁹, each depicting a thin slice of the VNC: they were able to segment 15,000 neurons, and predict 45 million synaptic connections between them to generate the FANC dataset. Ultimately, however, behavioral output is based on muscle contractions. To understand how motor circuits control movement, it is necessary to map each motor neuron to the muscle it innervates, and each muscle to a specific joint and movement direction — for example, leg extension or flexion. For this purpose, Azevedo *et al.*³ combined the FANC connectome with an X-ray dataset of a fly leg¹⁰ and high-resolution fluorescence images of motor neurons. The overall result is a comprehensive atlas of the wing






Figure 1. New connectomes link brain circuits to muscles in *Drosophila*.

(A) Schematic of a fruit fly with its central nervous system. The VNC is highlighted. (B) Anatomy of cell types within the giant fiber pathway, assembled using FANC³ and FlyWire^{7,20}. Arrows indicate leg and wing movements during take-off. (C) Connectivity diagram of the giant fiber pathway. Colors as in (B). (D) Section of FANC showing two input neurons (magenta and red) to the tibia flexor (blue). Arrowheads, presynaptic sites. Scale bar, 500 nm. FANC data from Azevedo *et al.*³, with permission from Springer Nature; FlyWire data from Dorkenwald *et al.*⁷ (CC BY-ND 4.0) and Schlegel *et al.*²⁰ (CC BY 4.0).

and front leg motor neurons, along with the motor circuits controlling them. The morphology and function of individual neurons is highly conserved within insect species, so that this atlas can serve as a blueprint for motor circuits across individuals and datasets.

Using the giant fiber escape pathway as an example (Figure 1B), Azevedo *et al.*³ illustrate how VNC circuits can control body movements. The giant fiber is the largest descending neuron in the fly nervous system. Once activated, it induces a specific sequence of leg and wing movements controlling escape jumps triggered by an approaching predator¹¹. The visual pathway that detects approaching objects and connects to the giant fiber has been characterized and described at synaptic resolution¹² — an effort aided by a full brain EM dataset⁵. In the VNC, the giant fiber connects to motor networks driving midleg extension, foreleg flexion, and wing downstroke. Together, these movements rapidly propel the body into the air¹¹ — and thus ensure flies aren't touched by humans unwilling to share their mangos. Combining FANC with the brain connectome now allows tracing this entire sensorimotor pathway at the synaptic level — from photoreceptors in

the retina to the leg and wing muscles (Figure 1B–D).

In a companion paper, Lesser, Azevedo *et al.*⁴ investigate the wiring logic of motor circuits by analysing the inputs of identified leg and wing motor neurons in FANC. They find that

premotor circuits are structured into 'motor modules'. These motor modules consist of premotor neurons targeting motor neurons controlling synergistic muscle groups, while avoiding their antagonists. This forms a synaptic basis for muscle synergies (Figure 2A). Some leg motor modules are shaped by premotor neurons providing input to all motor neurons within the module. In these cases, larger motor neurons receive more synaptic input from shared premotor neurons than smaller motor neurons (Figure 2B).

This finding is surprising: the original 'size principle' of motor neuron recruitment, established in vertebrate systems and thought to be a fundamental principle in motor control, assumes equal numbers of inputs to smaller and larger motor neurons¹³. Smaller motor neurons generate less force than larger motor neurons, but are more easily excitable; hence, smaller motor neurons are recruited earlier when receiving the same input. As Lesser, Azevedo *et al.*⁴ themselves point out, based on the large number of synaptic inputs they receive, one might assume that the largest motor neurons are the most active, and recruited earlier than small motor neurons in *Drosophila* (Figure 2B). However, the biophysical properties of individual motor neurons




Figure 2. Synergistic organization of leg motor circuits.

(A) Left: schematic of motor modules (M1–3): groups of premotor and motor neurons and the leg muscles they target. Arrows indicate tibia movement direction. Right: schematic connectivity between modules. Gray line: expected relationship based on the size principle. Bottom: smaller motor neurons are recruited before larger motor neurons during leg flexion.

counteract their input organization, so that this is not the case — as evidenced by prior electrophysiological recordings^{14,15}. Thus, leg motor neurons are recruited from small to large, as expected from the original size principle. The proportionality of motor neuron input to size is not observed within wing motor modules, further highlighting differences between the control of legs and wings. As these examples illustrate, connectomes are powerful tools to distill and test hypotheses about the fundamental organization of motor control networks — and they unfold their full potential when accompanied by physiological approaches.

Of course, there is still a lot of uncharted territory in the VNC. The biggest, perhaps, is that anatomical connectomes can only provide a roadmap for activity flow through the nervous system — which road is taken needs to be measured using physiological approaches. For example, neuromodulation can drastically shape functional connectivity in a state and context-dependent way¹⁶. Moreover, the spatial resolution of FANC is not fine enough to resolve electrical synapses, which play a key role in sensorimotor circuits^{2,17}. FANC is accompanied by a similar effort presenting the connectome of a Male Adult Nerve Cord (MANC)⁸. A complete connectome of the fly Brain And VNC (BANC) is also on the horizon, which will facilitate the identification of descending and ascending sensorimotor pathways already underway¹⁸. Hence, the motor control community will be able to contrast and compare three *Drosophila* VNCs, including male and female versions. This will allow asking questions about sex differences and individuality, for example. In FANC, Azevedo *et al.*³ managed to connect the neurons of the central nervous system to its body on the output side, by providing a motor neuron atlas and identifying their target muscles. A future upgrade to the connectome would be a similar accomplishment on the sensory side, as has recently become available in the *Drosophila* larva¹⁹. Thus, it would be possible to build even more powerful connectome-based models for the multitude of sensory contributions to behavioral control.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

- Grillner, S., and El Manira, A. (2020). Current principles of motor control, with special reference to vertebrate locomotion. *Physiol. Rev.* 100, 271–320. <https://doi.org/10.1152/physrev.00015.2019>.
- Bidaye, S.S., Bockemühl, T., and Büschges, A. (2018). Six-legged walking in insects: how CPGs, peripheral feedback, and descending signals generate coordinated and adaptive motor rhythms. *J. Neurophysiol.* 119, 459–475. <https://doi.org/10.1152/jn.00658.2017>.
- Azevedo, A., Lesser, E., Phelps, J.S., Mark, B., Elabbady, L., Kuroda, S., Sustar, A., Moussa, A., Khandelwal, A., Dallmann, C.J., *et al.* (2024). Connectomic reconstruction of a female *Drosophila* ventral nerve cord. *Nature* 631, 360–368. <https://doi.org/10.1038/s41586-024-07389-x>.
- Lesser, E., Azevedo, A.W., Phelps, J.S., Elabbady, L., Cook, A., Syed, D.S., Mark, B., Kuroda, S., Sustar, A., Moussa, A., *et al.* (2024). Synaptic architecture of leg and wing premotor control networks in *Drosophila*. *Nature* 631, 369–377. <https://doi.org/10.1038/s41586-024-07600-z>.
- Zheng, Z., Lauritzen, J.S., Perlman, E., Robinson, C.G., Nichols, M., Milkie, D., Torrens, O., Price, J., Fisher, C.B., Sharifi, N., *et al.* (2018). A complete electron microscopy volume of the brain of adult *Drosophila melanogaster*. *Cell* 174, 730–743.e22. <https://doi.org/10.1016/j.cell.2018.06.019>.
- Scheffer, L.K., Xu, C.S., Januszewski, M., Lu, Z., Takemura, S.Y., Hayworth, K.J., Huang, G.B., Shinomiya, K., Maitlin-Shepard, J., Berg, S., *et al.* (2020). A connectome and analysis of the adult *Drosophila* central brain. *eLife* 9, e57443. <https://doi.org/10.7554/eLife.57443>.
- Dorkenwald, S., Matsliah, A., Sterling, A.R., Schlegel, P., Yu, S.C., McKellar, C.E., Lin, A., Costa, M., Eichler, K., Yin, Y., *et al.* (2023). Neuronal wiring diagram of an adult brain. Preprint at bioRxiv, <https://doi.org/10.1101/2023.06.27.546656>.
- Takemura, S.-y., Hayworth, K.J., Huang, G.B., Januszewski, M., Lu, Z., Marin, E.C., Preibisch, S., Xu, C.S., Bogovic, J., Champion, A.S., *et al.* (2024). A connectome of the male *Drosophila* ventral nerve cord. *eLife* 13, RP97769. <https://doi.org/10.7554/eLife.97769>.
- Phelps, J.S., Hildebrand, D.G.C., Graham, B.J., Kuan, A.T., Thomas, L.A., Nguyen, T.M., Buhmann, J., Azevedo, A.W., Sustar, A., Agrawal, S., *et al.* (2021). Reconstruction of motor control circuits in adult *Drosophila* using automated transmission electron microscopy. *Cell* 184, 759–774.e18. <https://doi.org/10.1016/j.cell.2020.12.013>.
- Kuan, A.T., Phelps, J.S., Thomas, L.A., Nguyen, T.M., Han, J., Chen, C.L., Azevedo, A.W., Tuthill, J.C., Funke, J., Cloetens, P., *et al.* (2020). Dense neuronal reconstruction through X-ray holographic nano-tomography. *Nat. Neurosci.* 23, 1637–1643. <https://doi.org/10.1038/s41593-020-0704-9>.
- von Reyn, C.R., Breads, P., Peek, M.Y., Zheng, G.Z., Williamson, W.R., Yee, A.L., Leonardo, A., and Card, G.M. (2014). A spike-timing mechanism for action selection. *Nat. Neurosci.* 17, 962–970. <https://doi.org/10.1038/nn.3741>.
- Ache, J.M., Polksy, J., Alghailani, S., Parekh, R., Breads, P., Peek, M.Y., Bock, D.D., von Reyn, C.R., and Card, G.M. (2019). Neural basis for looming size and velocity encoding in the *Drosophila* giant fiber escape pathway. *Curr. Biol.* 29, 1073–1081.e4. <https://doi.org/10.1016/j.cub.2019.01.079>.
- Mendell, L.M. (2005). The size principle: a rule describing the recruitment of motoneurons. *J. Neurophysiol.* 93, 3024–3026. <https://doi.org/10.1152/classicsessays.0025.2005>.
- Azevedo, A.W., Dickinson, E.S., Gurung, P., Venkatasubramanian, L., Mann, R.S., and Tuthill, J.C. (2020). A size principle for recruitment of *Drosophila* leg motor neurons. *eLife* 9, e56754. <https://doi.org/10.7554/eLife.56754>.
- Gabriel, J.P., Scharstein, H., Schmidt, J., and Büschges, A. (2003). Control of flexor motoneuron activity during single leg walking of the stick insect on an electronically controlled treadmill. *J. Neurobiol.* 56, 237–251. <https://doi.org/10.1002/neu.10237>.
- Bargmann, C.I. (2012). Beyond the connectome: how neuromodulators shape neural circuits. *Bioessays* 34, 458–465. <https://doi.org/10.1002/bies.201100185>.
- Hürkey, S., Niemeyer, N., Schleimer, J.H., Rylewski, S., Schreiber, S., and Duch, C. (2023). Gap junctions desynchronize a neural circuit to stabilize insect flight. *Nature* 618, 118–125. <https://doi.org/10.1038/s41586-023-06099-0>.
- Cheong, H.S.J., Eichler, K., Stürner, T., Asinof, S.K., Champion, A.S., Marin, E.C., Oram, T.B., Sumathipala, M., Venkatasubramanian, L., Namiki, S., *et al.* (2024). Transforming descending input into behavior: The organization of premotor circuits in the *Drosophila* male adult nerve cord connectome. *eLife* 13, RP96084. <https://doi.org/10.7554/eLife.96084>.
- Richter, V., Rist, A., Kislinger, G., Laumann, M., Schoofs, A., Mirochnikow, A., Pankratz, M., Cardona, A., and Thum, A.S. (2023). Morphology and ultrastructure of external sense organs of *Drosophila* larvae. *eLife* 12, RP91155. <https://doi.org/10.7554/eLife.91155>.
- Schlegel, P., Yin, Y., Bates, A.S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D.S., Gkantia, M., dos Santos, M., Munnelly, E.J., *et al.* (2023). Whole-brain annotation and multi-connectome cell typing quantifies circuit stereotypy in *Drosophila*. Preprint at bioRxiv, <https://doi.org/10.1101/2023.06.27.546055>.