

Measurement of the inclusive WZ production cross section in pp collisions at $\sqrt{s} = 13.6$ TeV



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: The inclusive WZ production cross section is measured in proton-proton collisions at a centre-of-mass energy of 13.6 TeV, using data collected during 2022 with the CMS detector, corresponding to an integrated luminosity of 34.7 fb^{-1} . The measurement uses multileptonic final states and a simultaneous likelihood fit to the number of events in four different lepton flavour categories: eee , $ee\mu$, $\mu\mu e$, and $\mu\mu\mu$. The selection is optimized to minimize the number of background events, and relies on an efficient prompt lepton discrimination strategy. The WZ production cross section is measured in a phase space defined within a 30 GeV window around the Z boson mass, as $\sigma_{\text{total}}(\text{pp} \rightarrow \text{WZ}) = 55.2 \pm 1.2 (\text{stat}) \pm 1.2 (\text{syst}) \pm 0.8 (\text{lumi}) \pm 0.3 (\text{theo}) \text{ pb}$. In addition, the cross section is measured in a fiducial phase space closer to the detector-level requirements. All the measurements presented in this paper are in agreement with standard model predictions.

KEYWORDS: Electroweak Interaction, Hadron-Hadron Scattering, Vector Boson Production

ARXIV EPRINT: [2412.02477](https://arxiv.org/abs/2412.02477)

Contents

1	Introduction	1
2	The CMS detector	2
3	Simulated samples	2
4	Event reconstruction	3
5	Event selection	5
6	Background estimation	6
6.1	Reducible backgrounds	6
6.2	Irreducible backgrounds	7
7	Systematic uncertainties	9
8	Fit strategy and results	13
9	Summary	19
	The CMS collaboration	26

1 Introduction

Diboson production cross sections are interesting physical observables at the high energy proton-proton (pp) collisions at the CERN LHC. With relatively large event yields, high-purity multileptonic final states, and sensitivity to deviations of the trilinear gauge couplings (TGCs) [1] from their standard-model (SM) predictions, diboson production is one of the best experimental channels to study the properties of the electroweak (EW) sector of the SM.

The simultaneous production of a W and a Z boson (WZ) in pp collisions is dominated by quark-antiquark annihilation at tree level, as can be seen in figure 1. As a result, the process is sensitive to TGCs that could modify the WZ production cross section and thus provide evidence of beyond-the-SM (BSM) physics. Additionally, this process is one of the main backgrounds for many SM measurements and BSM searches in multileptonic final states.

The WZ production cross section was measured at the LHC in pp collisions at centre-of-mass energies of 5, 7, 8, and 13 TeV [2–8] by the CMS and ATLAS collaborations. All these measurements are in good agreement with the most accurate SM predictions within uncertainties. In this paper, a new measurement of the inclusive WZ production cross section in final states with three leptons that can be either electrons (e) or muons (μ) is presented. The analysed data were collected by the CMS experiment in pp collisions at a centre-of-mass energy of 13.6 TeV during 2022, the first data-taking year of the Run 3 of the LHC, and correspond to a total integrated luminosity of 34.7 fb^{-1} . The measurement is performed by classifying events according to the number of leptons of a given flavour. The cross section is determined via a simultaneous maximum likelihood fit to the event counts in all categories, as well as via fits to each category individually.




Figure 1. Leading-order Feynman diagrams for WZ production in pp collisions from the s channel (left), t channel (centre), and u channel (right).

This paper is organized as follows. Section 2 describes the CMS detector. Section 3 enumerates the simulated samples used in the measurement. Section 4 details the event reconstruction and the data sample used in this analysis. Section 5 presents the event selection and signal region definition. Section 6 explains the main SM background processes and the techniques used for their estimation. Section 7 includes a summary of the sources of uncertainty and their impact on the measurement. Section 8 presents the measurement strategy for signal extraction and the results. Finally, a summary is presented in section 9.

Tabulated results are provided in the HEPData record for this analysis [9].

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in refs. [10, 11].

Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about $4\,\mu\text{s}$ [12]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 5 kHz before data storage [13, 14].

3 Simulated samples

Signal WZ events are generated using POWHEG BOX (v2.0) [15–19] at next-to-leading order (NLO) in quantum chromodynamics (QCD), considering only the charged lepton decays of the WZ system. Renormalization (μ_R) and factorization (μ_F) scales in the signal prediction are computed dynamically based on the invariant mass of the WZ system in the generated event, according to the POWHEG method [16]. The SM background processes are simulated at NLO in QCD using POWHEG for several diboson processes (WW, ZZ) and top quark production

processes ($t\bar{t}$ and single top), at LO in QCD using MCFM (v7.0) [20] for the gluon-gluon loop induced ZZ, and with MADGRAPH5_aMC@NLO [21] (v2.9.13) for the remaining diboson processes ($Z\gamma$, $W\gamma$, and VH ; where V stands for W and Z), triboson production (WWW, WWZ, WZZ, ZZZ, $WZ\gamma$), and associated production of top quarks with other SM particles ($t\bar{t}W$, $t\bar{t}Z$, $t\bar{t}H$, $t\bar{t}\gamma$, tZq). All the MADGRAPH5_aMC@NLO processes have been generated at NLO in QCD, considering additional partons in the matrix element wherever possible. In all cases the NNPDF3.1_nnlo_hessian_pdfas [22] parton distribution function (PDF) set is used for the matrix element generation, and the generator is interfaced with PYTHIA 8 (v3.0.6) [23] for the showering, hadronization, and underlying-event modelling with the CP5 tune [24].

The propagation of the generated particles through the CMS detector and the modelling of the detector response are performed using GEANT4 [25] using the alignment and calibration from the collision data. The simulated events are processed with the same reconstruction algorithms used for the collision data.

4 Event reconstruction

The particle-flow (PF) algorithm [26] reconstructs and identifies each individual particle in an event with an optimized combination of information from the various elements of the CMS detector. The energy of photons is obtained from the ECAL measurement. The energy of electrons is determined from a combination of the electron momentum at the primary interaction vertex as determined by the tracker, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track. The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in section 9.4.1 of ref. [27]. The energy of muons is obtained from the curvature of the corresponding track, both inside the solenoid and in the return yoke. The energy of charged hadrons is determined from a combination of their momentum measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies.

All electrons in the analysis must have a transverse momentum (p_T) greater than 10 GeV and $|\eta| < 2.5$. The electron identification is based on a multivariate analysis (MVA) discriminant that combines observables describing the matching of the measurements in the tracker and the ECAL, the description of energy clusters in the ECAL, and the amount of bremsstrahlung radiation emitted during the propagation through the detector. The presence of electrons resulting from asymmetric photon conversions is reduced by requiring that their associated track have no missing hits in the innermost layers of the silicon tracker. An additional criterion that requires consistency among three measurements of the electron charge [28, 29] is applied to ensure a reliable charge determination. Muons are reconstructed by combining the information from the tracking system and the muon spectrometers in a global fit [30] that identifies the quality of the matching between the tracker and muon systems, and imposes minimal requirements in the track-related quantities. Muons are required to pass the kinematic selection requirements of $p_T > 5$ GeV and $|\eta| < 2.4$. As in the case of electrons, an additional identification requirement is imposed to reduce the

probability of charge misassignment in muons by requiring the measured curvature to have an uncertainty smaller than 20%.

An additional set of identification and isolation criteria are applied to define the restricted set of light leptons (e, μ) used in the analysis. These additional requirements target a high selection efficiency for leptons from W or Z bosons or from τ lepton decays, while rejecting those coming from other sources, such as leptons from hadron decays (non-prompt leptons). Lepton isolation requirements are imposed following the same approach as in ref. [8]. The isolation of each lepton is defined as the scalar p_T sum of all photons and charged and neutral hadrons in a cone around the lepton direction. The contributions from neutral particles originating from additional pp interactions, within the same or nearby bunch crossings (pileup) are estimated based on their average spatial energy density and the effective area covered by an isolation cone that is placed around the position of a given neutral particle in the detector. Charged particles identified to be originating from pileup vertices are discarded for this computation. A requirement on the lepton isolation relative to its p_T to be smaller than 0.4 is applied. Leptons that pass the above identification and isolation criteria are referred to as “loose” leptons.

An MVA discriminant that aims to separate the prompt and non-prompt contributions is further used to select high-quality prompt leptons [31]. Its inputs are the charged and neutral components of the lepton isolation as previously described, the properties of particles reconstructed in close proximity to the lepton, and the impact parameter with respect to the PV of the reconstructed lepton track in two and three dimensions. These selection criteria are more relevant for electrons and hence the requirements on the MVA discriminant are tighter than for muons.

For each event, hadronic jets are clustered from the PF candidates using the anti- k_T algorithm [32, 33] with a distance parameter of 0.4. Jet momentum is determined as the vectorial sum of all particle momenta in the jet, and is found from simulation to be, on average, within 5 to 10% of the true momentum over the whole p_T spectrum and detector acceptance. Pileup interactions can contribute additional tracks and calorimetric energy deposits, increasing the apparent jet momentum. The pileup-per-particle identification algorithm [34, 35] is used to mitigate the effect of pileup, making use of local shape information, event pileup properties, and tracking information. For each neutral particle, a local shape variable is computed using the surrounding charged particles compatible with the PV within $|\eta| < 2.5$, and using both charged and neutral particles in the region outside of the tracker coverage. This variable is used to distinguish between collinear and soft diffuse distributions of other objects surrounding the particle under consideration. Charged particles identified to be originating from pileup vertices are discarded. The momenta of the neutral particles are then rescaled according to their probability to originate from the PV deduced from the local shape variable, superseding the need for jet-based pileup corrections. Jets produced by the hadronization of b quarks with $p_T > 25$ GeV and $|\eta| < 2.5$ are identified using the DEEPJET algorithm [36, 37], which is based on a deep neural network that combines information from the reconstructed tracks and secondary vertices associated with the jet. The working point used in this analysis corresponds to a b tagging efficiency of 70% and a mistagging rate of 1% for light-flavoured jets and gluon jets.

The missing transverse momentum is computed as the negative sum of the p_T of all the PF candidates weighted by their probability to originate from the PV [38], and its magnitude is denoted as p_T^{miss} . The p_T^{miss} is modified to account for corrections to the energy scale of the reconstructed jets in the event. Anomalous high- p_T^{miss} events can be due to a variety of reconstruction failures, detector malfunctions, or noncollision backgrounds. Such events are rejected by event filters designed to identify more than 85–90% of the spurious high- p_T^{miss} events with a mistagging rate smaller than 0.1% [39].

5 Event selection

The data are recorded with a combination of single- and double-lepton triggers requiring the presence of one or two isolated electrons or muons with loose identification and isolation criteria. The p_T requirement of the single-lepton trigger is 32 GeV for electrons and 24 GeV for muons. The double-lepton triggers require the leading lepton to have a p_T greater than 23 (18) GeV, and the subleading lepton to have a p_T greater than 12 (8) GeV for electrons (muons).

Events are required to fulfil strict requirements designed to achieve a pure WZ signal region (SR) while keeping a high signal efficiency. The selection of events entering the SR is based on the presence of exactly three isolated leptons, with at least two leptons with electric charges of opposite sign and same flavour (OSSF). In order to exploit the kinematic properties of the WZ production, a simple assignment algorithm is used to tag each of the leptons originating from the W and Z boson decays. The algorithm works as follows: if only one OSSF lepton pair is found, the leptons are assumed to arise from the Z boson decay and, therefore, are labelled as ℓ_Z^1 and ℓ_Z^2 . In case two OSSF lepton pairs are found, the leptons of the pair with dilepton invariant mass closest to the nominal Z boson mass are selected as ℓ_Z^1 and ℓ_Z^2 . In either case, the hardest lepton not associated with the best Z boson candidate is referred to as ℓ_W . After assigning the origin of the leptons, the following additional p_T requirements are imposed in a consistent manner with the trigger requirements: $p_T(\ell_Z^1) > 25$ GeV, $p_T(\ell_Z^2) > 15$ GeV, and $p_T(\ell_W) > 25$ GeV. After this minimal preselection, ℓ_Z^1 and ℓ_Z^2 are properly tagged to the corresponding Z boson with 97% efficiency, and ℓ_W is found to be properly tagged to the W boson (when present) with a 94% efficiency. On top of this baseline selection, four channels based on the possible flavour combinations of the tagged leptons ($\ell_Z^1, \ell_Z^2, \ell_W$): eee , $ee\mu$, $\mu\mu e$, and $\mu\mu\mu$ are considered for the classification of events.

To increase the purity of WZ events, we require that the invariant mass of the ℓ_Z^1 and ℓ_Z^2 to be consistent with the Z boson mass $|m(\ell_Z^1, \ell_Z^2) - m_Z| < 15$ GeV, which removes contributions from nonresonant processes. The value of m_Z is taken from the Z mass world average given by the PDG [40]. A minimal requirement of $p_T^{\text{miss}} > 35$ GeV is included to reduce the presence of dileptonic Z production with an associated jet that can be misidentified as a lepton. Events with at least one b-tagged jets are vetoed from the SR in order to reduce contributions from top quark processes. Infrared safety in the same-flavour final states is ensured by rejecting events containing an OSSF lepton pair with invariant mass $m(\ell, \ell') < 4$ GeV. The invariant mass of the three lepton system is required to be $m(\ell_Z^1, \ell_Z^2, \ell_W) > 100$ GeV, effectively rejecting most of the contribution from $Z\gamma$ production in which the photon undergoes asymmetric conversion producing an additional final-state electron. All these requirements define the SR.

Region	N_ℓ	$p_T\{\ell_Z^1, \ell_Z^2, \ell_W(\ell_3), (\ell_4)\}$ (GeV)	N_{OSSF}	$ m(\ell_Z^1, \ell_Z^2) - m_Z $ (GeV)	p_T^{miss} (GeV)	$N_{\text{b tag}}$	$\min(m(\ell, \ell'))$ (GeV)	$m(\ell_Z^1, \ell_Z^2, \ell_W(\ell_3))$ (GeV)
SR	= 3	$>\{25, 15, 25\}$	≥ 1	< 15	> 35	= 0	> 4	> 100
ZZ CR	= 4	$>\{25, 15, 25, 15\}$	≥ 1	< 15	—	= 0	> 4	> 100
$t\bar{t}Z$ CR	= 3	$>\{25, 15, 25\}$	≥ 1	< 15	> 35	> 0	> 4	> 100
$X\gamma$ CR	= 3	$>\{25, 15, 25\}$	≥ 1	—	≤ 35	= 0	> 4	< 100

Table 1. Requirements for the definition of the signal and control regions of the analysis. Objects in parentheses relate to the ZZ CR.

Several control regions (CRs) are defined to estimate the contribution of the main background processes. Each CR is designed to be dominated by one background process and is used to estimate its normalization including the CR into the analysis likelihood fit. The CR definitions are obtained by inverting SR requirements, effectively increasing the purity of a given background without selecting events that pass the SR selection. In particular, the ZZ CR is defined by inverting the fourth-lepton veto and dropping the $p_T^{\text{miss}} > 35$ GeV requirement from the SR definition. The hardest lepton not belonging to the best OSSF pair is then referred to as ℓ_3 instead of ℓ_W in this CR, yet the requirements that are imposed are the same as the ones considered in the case with three leptons. The fourth lepton is also required to pass tight identification criteria and to have $p_T > 15$ GeV. The $t\bar{t}Z$ CR is obtained by inverting the b-tagged jet veto. Finally, the conversions CR ($X\gamma$ CR) is defined by removing the $|m(\ell_Z^1, \ell_Z^2) - m_Z| < 15$ GeV requirement and inverting the trilepton invariant mass requirement to $m(\ell_Z^1, \ell_Z^2, \ell_W) < 100$ GeV. Both modifications increase the presence of the process in which a photon is radiated by one of the final-state leptons in dileptonic Z boson decays, producing a typical trilepton resonance around the Z boson peak. To further increase the purity of $Z\gamma$ events in this phase space, p_T^{miss} is required to be smaller than 35 GeV. A summary of the definition of the SR and CRs can be found in table 1.

6 Background estimation

The background processes in the SR can be grouped into two different categories. The first category includes the reducible backgrounds that originate mainly due to the presence of non-prompt leptons, as defined in section 4, typically arising from Z +jets or $t\bar{t}$ production in which either a jet is misidentified as a lepton or a lepton is produced in the decay of a heavy-flavour hadron. The second category groups all the irreducible backgrounds including SM processes that naturally produce three or more leptons, such as boson pair production, triboson production, and associated production of top quarks with a W, Z, or H boson.

6.1 Reducible backgrounds

The SM processes with non-prompt leptons can have a contribution in the SR through the misidentification of a lepton that does not follow the prompt definition detailed in section 4, yet satisfies the tight identification requirements applied on the leptons that are used in the analysis. In final states with three leptons, these contributions are mainly attributed to Z +jets events in which a jet from ISR is misidentified as a lepton, and dileptonic $t\bar{t}$ or single top events where an additional lepton is produced in the in-flight decay of a

hadron. The lepton identification criteria have been optimized to reduce the non-prompt rate to roughly 3% of the total yield in the SR. Due to the relatively low contribution in the final selection, the non-prompt background is modelled considering the shapes and normalizations directly from MC simulation.

This MC-based non-prompt estimation is validated in a sideband region enriched in non-prompt background. This region is characterized by the presence of three final state leptons with kinematic requirements similar to the ones used in the SR, but considering events in which the tight identification criteria defined in section 4 are dropped for ℓ_W . The majority of non-prompt leptons contaminating the SR are tagged as ℓ_W according to the tagging algorithm described in section 5. Based on this premise, dropping the tight requirement on this lepton enriches the presence of the type of non-prompt backgrounds that need to be controlled in the analysis. The non-prompt purity, expressed as a percentage over the total number of selected events, is 72% (98% of it coming from Z +jets), and the remaining 28% can be broken down into a 15% of WZ events and 13% of other prompt processes. The ratio of data and MC was observed to be 0.93 in this region. We further tested the data to MC agreement for several lepton kinematic observables of ℓ_Z^1 , ℓ_Z^2 , and ℓ_W . Overall agreement was observed for both ℓ_Z^1 , ℓ_Z^2 , and a 30% normalization uncertainty was derived based on these studies to cover normalization effects that were observed particularly at low p_T (ℓ_W) and high $|\eta(\ell_W)|$. Other combinations of the non-prompt candidate was studied by dropping the tight criteria on either ℓ_Z^1 and ℓ_Z^2 , instead of ℓ_W , and a similar conclusion was reached.

6.2 Irreducible backgrounds

The main irreducible backgrounds are ZZ production, amounting to $\sim 5\%$ of the total yield in the SR, $t\bar{t}Z$ production, that contributes $\sim 2\%$ of the yield in the SR, and the associated production of a photon and other SM particles, $X\gamma$, that represents 1% of the selected events.

The purity of the ZZ contribution in the ZZ CR is about 98% of the total yield, with 90% of this contribution coming from $q\bar{q}$ annihilation ($q\bar{q} \rightarrow ZZ$), and the remaining 10% coming from gluon fusion mechanisms ($gg \rightarrow ZZ$). The distributions of several observables in this CR are shown in figure 2. Distributions of the reconstructed Z boson peak considering the two leptons tagged as ℓ_Z^1 and ℓ_Z^2 are also shown separately for electrons and muons in figure 3. All distributions show relatively good modelling of the ZZ background.

The $t\bar{t}Z$ CR is especially sensitive to both $t\bar{t}Z$ and tZq production. Residual contributions from $t\bar{t}W$ and $t\bar{t}H$ production are a factor of 20 smaller than the $t\bar{t}Z$ contribution and are grouped with $t\bar{t}Z$ into a group labelled $t\bar{t}X$. Good agreement between the observed data and predictions from simulation in this region is demonstrated in figure 4 for several variables.

The $X\gamma$ background includes processes with a prompt photon that proceeds through an asymmetric conversion due to the interaction with the detector to produce a single detected final-state lepton, or with a virtual photon that converts into a lepton pair. This background is dominated by associated $Z\gamma$ production (around 99% of the expected events), with residual contributions from $W\gamma$, $t\bar{t}\gamma$, and $WZ\gamma$ production. The distributions of several relevant observables in the $X\gamma$ CR are shown in figure 5. Good agreement is also observed in this CR.

The normalization of each major background is considered as an unconstrained parameter in the fit to the data, and the shape is obtained from the simulation. To constrain the




Figure 2. Distribution of observables in the ZZ CR accounting for the fit to data, described in section 8. From upper left (flavour composition), upper right (p_T of ℓ_Z^1), lower left (p_T of ℓ_Z^2), and lower right (p_T of ℓ_3). The hatched bands show the total uncertainty in the MC prediction. The vertical bars of the data account for the statistical uncertainty. When present, overflow events are included in the last bin of the observables. The ratio panels show the ratio between data (black markers) with respect to the total prediction after the fit to data. Processes with a small contribution to this region are grouped in the “Other” category.




Figure 3. Distribution of the invariant mass of ℓ_Z^1 and ℓ_Z^2 in the ZZ CR accounting for the fit to data. The left (right) distribution shows the case in which the leptons are electrons (muons). The hatched bands show the total uncertainty in the MC prediction. The vertical bars of the data account for the statistical uncertainty. The ratio panels show the ratio between data (black markers) with respect to the total prediction after the fit to data. Processes with a small contribution to this region are grouped in the “Other” category.

major backgrounds, the number of events in flavour categories from each of the CRs is included in the likelihood fit. All the uncertainties included in the total inclusive cross section measurement described in section 7 also apply to these CRs.

Other irreducible backgrounds contribute less than 1% to the yields in the SR and are directly estimated from simulation. These include the associated production of a vector boson and a Higgs boson (VH) and the triboson production (VVV). These backgrounds are estimated using MC simulation. Contributions from WZ produced through vector boson scattering have not been considered. This background is expected to contribute as 1% of the total yield in the SR.

7 Systematic uncertainties

The sources of systematic uncertainties can be grouped into four categories:

- Uncertainties associated with the normalization of the background sources. These are treated as uncorrelated among backgrounds.
- Uncertainties related to the modelling of instrumental effects in the simulation such as those arising from the object identification, reconstruction, and energy scales. These are typically correlated across all processes.
- Uncertainties in the measurement of the integrated luminosity collected by CMS that are also correlated among all processes estimated from simulation.
- Theoretical and modelling uncertainties that affect only the WZ process.




Figure 4. Distribution of observables in the $t\bar{t}Z$ CR accounting for the fit to data, described in section 8. From upper left (flavour composition), upper right (p_T of ℓ_Z^1), lower left (p_T of ℓ_Z^2), and lower right (p_T of ℓ_W). The hatched bands show the total uncertainty in the MC prediction. The vertical bars of the data account for the statistical uncertainty. When present, overflow events are included in the last bin of the observables. The ratio panels show the ratio between data (black markers) with respect to the total prediction after the fit to data. Processes with a small contribution to this region are grouped in the “Other” category.




Figure 5. Distribution of observables in the $X\gamma$ CR accounting for the fit to data, described in section 8. From upper left (flavour composition), upper right (p_T of ℓ_Z^1), lower left (p_T of ℓ_Z^2), and lower right (p_T of ℓ_W). The hatched bands show the total uncertainty in the MC prediction. The vertical bars of the data account for the statistical uncertainty. When present, overflow events are included in the last bin of the observables. The ratio panels show the ratio between data (black markers) with respect to the total prediction after the fit to data. Processes with a small contribution to this region are grouped in the “Other” category.

For those backgrounds that have a significant contribution to the SR (ZZ , $t\bar{t}X$, and $X\gamma$), no additional normalization uncertainty is included, as it is treated as an unconstrained parameter in the fit. As the remaining irreducible backgrounds are small and the uncertainty in their estimation contributes negligibly to the total uncertainty of the measurement, we use the experimentally measured uncertainties on each to reduce theoretical contributions to the final uncertainty: 15% for tZq [41], 50% for VVV [42], and 25% for VH [43]. Finally, a normalization uncertainty of 30% is assigned for the non-prompt lepton background, as detailed in section 5.

The uncertainties related to the mismodelling of instrumental effects in the simulation are estimated by varying the relevant parameters within their uncertainties and recomputing the event yields. The following sources of systematic uncertainties are considered: lepton identification and isolation efficiencies, jet energy scale and resolution, b tagging efficiency, p_T^{miss} scale and resolution, and pileup modelling.

The lepton identification and isolation efficiencies are estimated using the tag-and-probe technique [44] and are affected by the size of the data set in $Z \rightarrow \ell^+\ell^-$ enriched regions, where the efficiencies are derived. Both efficiencies and their uncertainties are estimated separately for electrons and muons, with the average uncertainties of $\sim 1\%$ per electron and $\sim 0.5\%$ per muon. The statistical component of these uncertainties, corresponding to the size of the data set in which the tag-and-probe technique has been applied, is also included. Uncertainties in the energy scale of electrons introduce a smaller effect that induces variations in the lepton p_T . This effect is negligible for muons.

The efficiency of the trigger selection is measured in data selected using an independent set of p_T^{miss} triggers and only the requirement of three tight leptons passing the p_T requirements of the SR described in section 5. The total uncertainty in the trigger efficiency measurement is split into a systematic component for the estimated measurement bias in the selection of the data set, and an additional statistical component originating from the limited size of this data set. Each component is applied in a decorrelated manner to different parts of the entire data set, and the uncertainties amount range between 0.5–1.0% and from 0.5–1.6% for the systematic and statistical sources, respectively. Since the central estimation of the trigger efficiency is high (over 99%), the Clopper-Pearson estimation [45] of the associated confidence interval leads to slightly asymmetric effects that slightly increase the uncertainty on the downward variations.

The energy scale and resolution of the reconstructed jets has an associated uncertainty, which is propagated to the p_T^{miss} , causing a small effect in the total cross section. The b tag veto requirement is affected by the uncertainties in the efficiency of the b tagging procedure and the mistagging rate. The effect of these uncertainties is minimized by choosing a tight working point of the b tagging algorithm. They are split into the uncertainties in the tagging of heavy-flavour, and light-flavour and gluon jets as two decorrelated sources.

The uncertainty assigned to the number of pileup events in simulation is obtained by modifying the inelastic pp cross section up and down by 4.6% [46] around its nominal value and propagating the effect to all simulated samples. The measurement of the integrated luminosity recorded by CMS introduces an additional uncertainty of 1.4% [47, 48] into the analysis. Uncertainties due to the variations of the PDFs and the missing higher order

Source	Uncertainty (%)	Processes
Integrated luminosity	1.4	All MC
Trigger efficiencies	0.5–1.6	All MC
b tagging (heavy)	0.1	All MC
b tagging (light/gluon)	< 0.1	All MC
Pileup	0.4–1.7	All MC
Jet energy scale	1.1	All MC
Electron reconstruction	3.8	All MC
Electron ident. efficiency	3.5	All MC
Electron energy scale	0.3	All MC
Muon efficiencies	1.1	All MC
Non-prompt bkg. normalization	30	Non-prompt
VVV normalization	50	VVV
tZq normalization	15	tZq
VH normalization	25	VH
ISR	0.1	WZ
FSR	0.8	WZ
WZ theory (μ_R , μ_F , PDF)	0.6	WZ

Table 2. Summary of the input relative uncertainties. Numbers are presented in percentages over the predicted signal yields of the associated process on which they have an effect. All uncertainties are treated as shape variations on the templates used for the fit, with the exception of the normalization uncertainties in the backgrounds that are treated as flat variations of the corresponding yield.

corrections in QCD, are also included in the fits. Effects due to the choice of μ_F and μ_R are evaluated by varying the parameters up and down independently by a factor of two around the nominal POWHEG μ_0 value under the constraint $0.5 \leq \mu_F/\mu_R \leq 2$. The envelope of these six variations is taken as the estimated uncertainty for the missing higher order QCD corrections. Parametric (PDF+ α_S) uncertainties are estimated using the NNPDF3.1 set. Uncertainties related to the modelling of the parton shower for the WZ signal have been estimated separately for the initial- and final-state radiation (ISR and FSR) contributions by varying the associated energy scales in simulation. Finally the uncertainty coming from the finite size of the MC samples is estimated using the Barlow-Beeston method [49]. A summary of all the input uncertainties considered in the analysis is shown in table 2.

8 Fit strategy and results

The WZ production cross section is measured in the detector-level phase space defined as SR in section 5 in each of the four flavour-exclusive channels (eee, $ee\mu$, $\mu\mu e$, and $\mu\mu\mu$), and in the flavour-inclusive channel. This is motivated by the fact that final states with three leptons of the same-flavour (eee and $\mu\mu\mu$) include additional contributions to the production modes due to interference contributions from diagrams exchanging same flavour final-state leptons. The additional combinatorics of the two same-charged leptons result in slightly

Region	Fiducial	Total
Lepton definition	Dressed (e, μ)	Dressed (e, μ , τ)
$N_\ell = 3$	✓	✓
$p_T\{\ell_Z^1, \ell_Z^2, \ell_W\} > \{25, 15, 25\} \text{ GeV}$	✓	—
$ \eta \{\ell_Z^1, \ell_Z^2, \ell_W\} < \{2.5, 2.5, 2.5\}$	✓	—
$N_{\text{OSSF}} = 1$	✓	✓
$60 < m(\ell_Z^1, \ell_Z^2) < 120 \text{ GeV}$	✓	✓
$\min(m(\ell, \ell')) > 4 \text{ GeV}$	✓	✓
$m(\ell_Z^1, \ell_Z^2, \ell_W) > 100 \text{ GeV}$	✓	—

Table 3. Definition of the fiducial and total regions. The lepton related quantities refer to “dressed” leptons. A dash symbol is used in cases the selection is not considered for the corresponding phase space.

larger cross sections. At the same time, further splitting the final states into different lepton flavour compositions leads to both the possibility of analysing deviations in the lepton flavour universality in the W and Z boson decays, and of studying possible experimental biases in the determination of electron and muon efficiencies.

The observed WZ cross section at the detector-level is extrapolated to two different particle-level phase spaces, referred to as “fiducial” (FR) and “total” (TR) regions, respectively. The TR is designed to provide a more inclusive measurement of the WZ cross section, with fewer requirements on the properties of the process. In contrast, the FR is motivated by the aim to reduce as comprehensively as possible the detector acceptance effects on the theoretical uncertainties that affect the measurement. As a result, the FR closely mirrors the SR selection criteria outlined in section 5, including lepton kinematic requirements and the exclusion of $WZ \rightarrow \tau\tau$. Both these regions are defined after hadronization and parton shower using generator-level quantities. Leptons at this level are “dressed” by adding the transverse momenta of generator-level photons within a cone of $\Delta R(\ell, \gamma) < 0.1$ to their own momenta. In addition, dressed leptons are associated to the W and Z bosons using the same tagging algorithm described in section 5. The proportion of non-FR events in the WZ sample that pass detector-level requirements is approximately 5% in all the different decay channels as well as in the inclusive channel, and it is mainly attributed to leptonic decays of at least one of the τ lepton in any of the $WZ \rightarrow \tau + X$ decay modes, where X accounts for any lepton flavour. The proportion of FR events inside the TR, known as the acceptance, is about 17% in the inclusive channel, or equivalently 4% per decay channel. These numbers have been computed considering the WZ POWHEG signal sample, and cross-checked with an alternative WZ MADGRAPH5_aMC@NLO sample. Nevertheless, they are not directly used as an input of the analysis. Instead, the non-FR events are estimated by treating them as background contributions in the extraction of the fiducial results, and as part of the signal in the TR results. The normalization of the WZ signal contributions in both cases is then measured directly in the maximum likelihood fit detailed below. The requirements of the FR and TR selections are summarised in table 3.

Process	eee	ee μ	$\mu\mu e$	$\mu\mu\mu$	Inclusive
Non-prompt	25 ± 7	13 ± 5	24 ± 7	30 ± 10	93 ± 15
ZZ	25 ± 2	37 ± 1	49 ± 3	75 ± 3	186 ± 5
X γ	12 ± 2	2.5 ± 0.3	24 ± 2	3.2 ± 0.5	41 ± 3
t $\bar{t}$ X	8.0 ± 0.8	11 ± 1	14 ± 1	21 ± 2	54 ± 3
VVV	4 ± 1	5 ± 2	7 ± 3	10 ± 4	27 ± 5
VH	3.0 ± 0.5	3.8 ± 0.7	5 ± 1	9 ± 2	20 ± 2
tZq	4.2 ± 0.5	5.3 ± 0.6	7.5 ± 0.9	11 ± 1	28 ± 2
Background	82 ± 8	78 ± 5	130 ± 9	160 ± 11	450 ± 17
WZ	410 ± 10	556 ± 12	768 ± 14	1096 ± 22	2830 ± 31
Prediction	491 ± 13	634 ± 13	898 ± 16	1256 ± 24	3280 ± 34
Data	491	643	869	1276	3279

Table 4. Number of selected events and their total uncertainty (by flavour channel) for the relevant processes in the signal region of the analysis accounting for the fit to data. The “Background” yield is the sum of all processes that are not WZ signal, and “Prediction” is the sum of all the processes, including WZ.

Results at the fiducial- and total-level are obtained from two separate maximum likelihood fits to the distributions of the number of events in different flavour channels considering the SR and all the CRs described in section 6, using the COMBINE tool [50]. In the fits, the normalization of the signal contribution to the WZ process and the ZZ, t $\bar{t}$ Z, and X γ backgrounds are considered unconstrained parameters in the fit. The systematic uncertainties described in section 7 are included as nuisance parameters with gaussian constraints correlated across SR and CRs for all shape variations, and as log-normal functions for those sources that only have a normalization effect. In the following, all presented results consider the fit results using the TR definitions unless stated otherwise. In both fits the shape for the variations originating from theoretical sources of uncertainty are normalized in a way that ensures that the sample cross section of the WZ process is the same in both the nominal templates and all the variations. This ensures that normalization effects on the theoretical uncertainties of the WZ variations do not have an effect in the measurement, keeping only the contribution that changes the overall acceptance of the FR and TR selections. The same procedure for the extraction of the cross section results is used as well for the flavour-exclusive measurements.

Figures 6 and 7 show several relevant distributions in the SR accounting for the fit to data, including the distribution used in the fit as well as other kinematic properties of the selected leptons. Observables that do not enter the fit are obtained by propagating the fit results to the observables before the fit. The event yields per process and flavour channel after the fit are shown in table 4. Good agreement after the fit is observed in all the observables.

The measured fiducial cross sections are reported in table 5 for the different flavour channels, including a comparison with theoretical predictions computed from the signal POWHEG sample, as well as the next-to-NLO (NNLO) QCD and NNLO QCD $\times$ NLO EW prediction from MATRIX [51]. In all MATRIX predictions we have adopted the main setup for the SM parameters, scale and PDF choices, and have only modified the requirements of the




Figure 6. Distributions of several observables in the SR accounting for the fit to data. From upper left (flavour composition), upper right (p_T of ℓ_Z^1), lower left (p_T of ℓ_Z^2), and lower right (p_T of ℓ_W). The hatched band includes all systematic uncertainties in the MC prediction. The vertical bars of the data account for the statistical uncertainty. When present, overflow events are included in the last bin of the observables. The ratio panels show the ratio between data (black markers) with respect to the total prediction after the fit to data. Processes with a small contribution to this region are grouped in the “Other” category.




Figure 7. Distributions of several observables in the SR accounting for the fit to data. From upper left (sum of charge of the final-state leptons), upper right (missing transverse momentum), lower left (invariant mass of the two leptons assigned to the Z boson decay), and lower right (invariant mass of the triplepton system). The hatched band includes all systematic uncertainties. When present, overflow events are included in the last bin of the observables. The ratio panels show the ratio between data (black markers) with respect to the total prediction after the fit to data. Processes with a small contribution to this region are grouped in the “Other” category.

Category	Accuracy	Fiducial cross section (fb)
eee	POWHEG, NLO QCD	$68.0^{+2.3}_{-2.1} \text{ (scale)} \pm 1.0 \text{ (PDF)}$
	MATRIX, NLO QCD	$69.9^{+3.9}_{-3.1} \text{ (scale)}$
	MATRIX, NNLO QCD	$77.0^{+1.8}_{-1.7} \text{ (scale)}$
	MATRIX, NNLO QCD $\times$ NLO EW	$75.4^{+1.7}_{-1.6} \text{ (scale)}$
	Measured	$72.0 \pm 4.0 \text{ (stat)} \pm 4.5 \text{ (syst)} \pm 1.0 \text{ (lumi)} \pm 0.4 \text{ (theo)}$
ee μ	POWHEG, NLO QCD	$68.0^{+2.3}_{-2.1} \text{ (scale)} \pm 1.0 \text{ (PDF)}$
	MATRIX, NLO QCD	$68.7^{+3.8}_{-3.0} \text{ (scale)}$
	MATRIX, NNLO QCD	$75.0^{+1.8}_{-1.6} \text{ (scale)}$
	MATRIX, NNLO QCD $\times$ NLO EW	$73.4^{+1.7}_{-1.5} \text{ (scale)}$
	Measured	$73.9 \pm 3.5 \text{ (stat)} \pm 3.1 \text{ (syst)} \pm 1.1 \text{ (lumi)} \pm 0.4 \text{ (theo)}$
$\mu\mu e$	POWHEG, NLO QCD	$68.0^{+2.3}_{-2.1} \text{ (scale)} \pm 1.0 \text{ (PDF)}$
	MATRIX, NLO QCD	$68.7^{+3.8}_{-3.0} \text{ (scale)}$
	MATRIX, NNLO QCD	$75.0^{+1.8}_{-1.6} \text{ (scale)}$
	MATRIX, NNLO QCD $\times$ NLO EW	$73.4^{+1.7}_{-1.5} \text{ (scale)}$
	Measured	$71.2 \pm 2.9 \text{ (stat)} \pm 2.0 \text{ (syst)} \pm 1.0 \text{ (lumi)} \pm 0.4 \text{ (theo)}$
$\mu\mu\mu$	POWHEG, NLO QCD	$68.0^{+2.3}_{-2.1} \text{ (scale)} \pm 1.0 \text{ (PDF)}$
	MATRIX, NLO QCD	$69.9^{+3.9}_{-3.1} \text{ (scale)}$
	MATRIX, NNLO QCD	$77.0^{+1.8}_{-1.7} \text{ (scale)}$
	MATRIX, NNLO QCD $\times$ NLO EW	$75.4^{+1.7}_{-1.6} \text{ (scale)}$
	Measured	$75.3 \pm 2.5 \text{ (stat)} \pm 1.5 \text{ (syst)} \pm 1.1 \text{ (lumi)} \pm 0.4 \text{ (theo)}$
Inclusive	POWHEG, NLO QCD	$271.9^{+9.0}_{-8.5} \text{ (scale)} \pm 3.8 \text{ (PDF)}$
	MATRIX, NLO QCD	$277.1^{+15.3}_{-12.3} \text{ (scale)}$
	MATRIX, NNLO QCD	$304.0^{+7.1}_{-6.6} \text{ (scale)}$
	MATRIX, NNLO QCD $\times$ NLO EW	$297.7^{+6.8}_{-6.3} \text{ (scale)}$
	Measured	$297.6 \pm 6.4 \text{ (stat)} \pm 6.4 \text{ (syst)} \pm 4.2 \text{ (lumi)} \pm 1.5 \text{ (theo)}$

Table 5. Measured fiducial cross sections and their corresponding uncertainties for the flavour-exclusive and flavour-inclusive categories. The predictions from both POWHEG at NLO in QCD and LO EW as well as several ones obtained from MATRIX (NLO QCD, NNLO QCD, NNLO QCD $\times$ NLO EW) are also included.

leptons in order to mimic either the fiducial or total definitions. Similarly, results for the total cross section measurement and their comparison to different predictions at different orders of the QCD and EW expansion can be found in table 6 and figure 8. The relative uncertainty in the measurement in the inclusive category of about 3.4% is compatible between the fiducial-level and total-level results. A breakdown of the different sources of systematic uncertainty and their impact in the result can be found in table 7. The values of the normalization parameters in the combined fit corresponding to each of the individual backgrounds are: $r_{ZZ} = 0.96 \pm 0.06$, $r_{t\bar{t}Z} = 1.18 \pm 0.15$, and $r_{X\gamma} = 1.10 \pm 0.13$. The measurements presented in this paper, as well as all previous WZ measurements by CMS at different centre-of-mass




Figure 8. Total WZ production cross section for each of the flavour-exclusive and for the flavour-inclusive categories. The vertical bands show different theoretical predictions for the WZ cross section at NLO in QCD (red dashed line) and NNLO QCD × NLO EW (blue solid line), as well as their corresponding scale uncertainties. For each measurement, the best fit value is denoted with a purple point, with two delimiters on the error bars that account for the statistical and total uncertainties.

energies, support the existence of large corrections beyond the NLO QCD and the good compatibility with the latest NNLO QCD × NLO EW predictions, as summarised in figure 9.

9 Summary

The inclusive WZ production cross section is measured in proton-proton collisions for the first time at a centre-of-mass energy of 13.6 TeV, using data collected during 2022 with the CMS detector, corresponding to an integrated luminosity of 34.7 fb⁻¹. The production cross sections in the total and fiducial phase spaces are measured in the inclusive case as well as in four

Accuracy	Total cross section (pb)
POWHEG, NLO QCD	$50.5^{+2.6}_{-2.1} (\text{scale}) \pm 1.1 (\text{PDF})$
MATRIX, NLO QCD	$50.4^{+2.3}_{-2.0} (\text{scale})$
MATRIX, NNLO QCD	$55.0^{+1.2}_{-1.1} (\text{scale})$
MATRIX, NNLO QCD $\times$ NLO EW	$54.7^{+1.2}_{-1.1} (\text{scale})$
Inclusive (Measured)	$55.2 \pm 1.2 (\text{stat}) \pm 1.2 (\text{syst}) \pm 0.8 (\text{lumi}) \pm 0.3 (\text{theo})$

Table 6. Flavour-inclusive total cross section result. The predictions from both POWHEG at NLO in QCD and LO EW as well as several ones obtained from MATRIX (NLO QCD, NNLO QCD, NNLO QCD $\times$ NLO EW) are also included.

Source	Inclusive (%)	eee (%)	ee μ (%)	$\mu\mu e$ (%)	$\mu\mu\mu$ (%)
Integrated luminosity	1.5	1.5	1.4	1.4	1.5
Trigger efficiencies	0.5	1.0	1.0	1.0	0.7
b tagging	0.1	0.1	0.1	0.1	0.1
Pileup	0.4	0.6	0.8	0.2	0.4
Jet energy scales	0.9	1.3	0.7	1.1	0.7
Electron reconstruction	1.2	4.0	2.9	1.1	—
Electron ident. efficiencies	0.7	3.6	2.4	1.1	—
Electron energy scale	0.1	0.1	0.1	0.0	—
Muon efficiencies	0.7	—	0.3	0.8	1.2
Non-prompt bkg. normalization	0.7	1.6	0.5	0.7	0.7
VVV normalization	0.4	0.4	0.4	0.4	0.4
tZq normalization	0.1	0.1	0.1	0.1	0.1
ZZ normalization	0.3	0.8	0.7	0.5	0.5
t $\bar{t}$ Z normalization	0.3	0.7	0.6	0.4	0.5
X γ normalization	0.2	0.7	0.3	0.4	0.2
VH normalization	0.2	0.2	0.2	0.1	0.2
ISR/FSR	0.3	0.5	0.2	0.4	0.3
WZ theory (μ_R , μ_F , PDF)	0.6	0.6	0.6	0.6	0.6
MC statistical	0.5	1.9	0.9	1.0	0.9
Statistical	2.0	5.3	4.6	3.8	3.3
Total	3.4	8.5	6.5	5.0	4.3

Table 7. Breakdown of different sources of systematic uncertainties and their relative impact in each channel, as well as in the inclusive measurement; as a percentage of the total uncertainty. The dash symbol indicates that the specific uncertainty does not apply.

combinations of final state flavour composition. The cross sections are measured in a fiducial phase space as $\sigma_{\text{fiducial}}(\text{pp} \rightarrow \text{WZ}) = 297.6 \pm 6.4 (\text{stat}) \pm 6.4 (\text{syst}) \pm 4.2 (\text{lumi}) \pm 1.5 (\text{theo}) \text{ fb}$, and in a less restricted phase space as $\sigma_{\text{total}}(\text{pp} \rightarrow \text{WZ}) = 55.2 \pm 1.2 (\text{stat}) \pm 1.2 (\text{syst}) \pm 0.8 (\text{lumi}) \pm 0.3 (\text{theo}) \text{ pb}$. All these measurements are shown to be in good agreement with the SM predictions at NNLO QCD $\times$ NLO EW.




Figure 9. Measurement obtained in this analysis (red filled marker) together with other WZ production cross section measurements at different centre-of-mass energies by the CMS [2, 7, 8] collaboration, compared to the NNLO QCD $\times$ NLO EW predictions, as well as the NLO QCD prediction, computed in all cases with MATRIX.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM

(Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science – EOS” – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany’s Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ_KI (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Centre for High Performance Computing, Big Data and Quantum Computing and FAIR – Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

Data Availability Statement. Release and preservation of data used by the CMS collaboration as the basis for publications is guided by the [CMS data preservation, re-use and open access policy](#).

Code Availability Statement. The CMS core software is publicly available on [GitHub](#).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] C. Degrande et al., *Effective Field Theory: A Modern Approach to Anomalous Couplings*, *Ann. Phys.* **335** (2013) 21 [[arXiv:1205.4231](https://arxiv.org/abs/1205.4231)] [[INSPIRE](#)].
- [2] CMS collaboration, *Measurement of the WZ production cross section in pp collisions at $\sqrt{s} = 7$ and 8 TeV and search for anomalous triple gauge couplings at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **77** (2017) 236 [[arXiv:1609.05721](https://arxiv.org/abs/1609.05721)] [[INSPIRE](#)].
- [3] ATLAS collaboration, *Measurements of $W^{\pm}Z$ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector and limits on anomalous gauge boson self-couplings*, *Phys. Rev. D* **93** (2016) 092004 [[arXiv:1603.02151](https://arxiv.org/abs/1603.02151)] [[INSPIRE](#)].
- [4] CMS collaboration, *Measurement of the WZ production cross section in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **766** (2017) 268 [[arXiv:1607.06943](https://arxiv.org/abs/1607.06943)] [[INSPIRE](#)].
- [5] ATLAS collaboration, *Measurement of the $W^{\pm}Z$ boson pair-production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector*, *Phys. Lett. B* **762** (2016) 1 [[arXiv:1606.04017](https://arxiv.org/abs/1606.04017)] [[INSPIRE](#)].
- [6] ATLAS collaboration, *Measurement of $W^{\pm}Z$ production cross sections and gauge boson polarisation in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **79** (2019) 535 [[arXiv:1902.05759](https://arxiv.org/abs/1902.05759)] [[INSPIRE](#)].
- [7] CMS collaboration, *Measurements of the electroweak diboson production cross sections in proton-proton collisions at $\sqrt{s} = 5.02$ TeV using leptonic decays*, *Phys. Rev. Lett.* **127** (2021) 191801 [[arXiv:2107.01137](https://arxiv.org/abs/2107.01137)] [[INSPIRE](#)].
- [8] CMS collaboration, *Measurement of the inclusive and differential WZ production cross sections, polarization angles, and triple gauge couplings in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2022) 032 [[arXiv:2110.11231](https://arxiv.org/abs/2110.11231)] [[INSPIRE](#)].
- [9] HEPData record for this analysis, (2024) [[DOI:10.17182/hepdata.155693](https://doi.org/10.17182/hepdata.155693)].
- [10] CMS collaboration, *The CMS Experiment at the CERN LHC*, 2008 *JINST* **3** S08004 [[INSPIRE](#)].
- [11] CMS collaboration, *Development of the CMS detector for the CERN LHC Run 3*, 2024 *JINST* **19** P05064 [[arXiv:2309.05466](https://arxiv.org/abs/2309.05466)] [[INSPIRE](#)].
- [12] CMS collaboration, *Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2020 *JINST* **15** P10017 [[arXiv:2006.10165](https://arxiv.org/abs/2006.10165)] [[INSPIRE](#)].
- [13] CMS collaboration, *The CMS trigger system*, 2017 *JINST* **12** P01020 [[arXiv:1609.02366](https://arxiv.org/abs/1609.02366)] [[INSPIRE](#)].
- [14] CMS collaboration, *Performance of the CMS high-level trigger during LHC Run 2*, 2024 *JINST* **19** P11021 [[arXiv:2410.17038](https://arxiv.org/abs/2410.17038)] [[INSPIRE](#)].
- [15] P. Nason, *A New method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040 [[hep-ph/0409146](https://arxiv.org/abs/hep-ph/0409146)] [[INSPIRE](#)].
- [16] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with Parton Shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[arXiv:0709.2092](https://arxiv.org/abs/0709.2092)] [[INSPIRE](#)].

- [17] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [18] T. Melia, P. Nason, R. Röntsch and G. Zanderighi, *W^+W^- , WZ and ZZ production in the POWHEG BOX*, *JHEP* **11** (2011) 078 [[arXiv:1107.5051](#)] [[INSPIRE](#)].
- [19] P. Nason and G. Zanderighi, *W^+W^- , WZ and ZZ production in the POWHEG-BOX-V2*, *Eur. Phys. J. C* **74** (2014) 2702 [[arXiv:1311.1365](#)] [[INSPIRE](#)].
- [20] J.M. Campbell and R.K. Ellis, *MCFM for the Tevatron and the LHC*, *Nucl. Phys. B Proc. Suppl.* **205–206** (2010) 10 [[arXiv:1007.3492](#)] [[INSPIRE](#)].
- [21] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [22] NNPDF collaboration, *Parton distributions from high-precision collider data*, *Eur. Phys. J. C* **77** (2017) 663 [[arXiv:1706.00428](#)] [[INSPIRE](#)].
- [23] C. Bierlich et al., *A comprehensive guide to the physics and usage of PYTHIA 8.3*, *SciPost Phys. Codebases* **2022** (2022) 8 [[arXiv:2203.11601](#)] [[INSPIRE](#)].
- [24] CMS collaboration, *Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements*, *Eur. Phys. J. C* **80** (2020) 4 [[arXiv:1903.12179](#)] [[INSPIRE](#)].
- [25] GEANT4 collaboration, *GEANT4 — A Simulation Toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [26] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, *2017 JINST* **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].
- [27] D. Contardo et al., *Technical Proposal for the Phase-II Upgrade of the CMS Detector*, CERN-LHCC-2015-010 (2015) [CMS-TDR-15-02] [[DOI:10.17181/CERN.VU8I.D59J](#)] [[INSPIRE](#)].
- [28] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, *2021 JINST* **16** P05014 [[arXiv:2012.06888](#)] [[INSPIRE](#)].
- [29] CMS collaboration, *Performance of Electron Reconstruction and Selection with the CMS Detector in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV*, *2015 JINST* **10** P06005 [[arXiv:1502.02701](#)] [[INSPIRE](#)].
- [30] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, *2018 JINST* **13** P06015 [[arXiv:1804.04528](#)] [[INSPIRE](#)].
- [31] CMS collaboration, *Muon identification using multivariate techniques in the CMS experiment in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *2024 JINST* **19** P02031 [[arXiv:2310.03844](#)] [[INSPIRE](#)].
- [32] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].
- [33] M. Cacciari, G.P. Salam and G. Soyez, *FastJet User Manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [34] CMS collaboration, *Pileup mitigation at CMS in 13 TeV data*, *2020 JINST* **15** P09018 [[arXiv:2003.00503](#)] [[INSPIRE](#)].
- [35] D. Bertolini, P. Harris, M. Low and N. Tran, *Pileup Per Particle Identification*, *JHEP* **10** (2014) 059 [[arXiv:1407.6013](#)] [[INSPIRE](#)].

- [36] CMS collaboration, *Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV*, **2018 JINST** **13** P05011 [[arXiv:1712.07158](#)] [[INSPIRE](#)].
- [37] E. Bols, J. Kieseler, M. Verzetti, M. Stoye and A. Stakia, *Jet Flavour Classification Using DeepJet*, **2020 JINST** **15** P12012 [[arXiv:2008.10519](#)] [[INSPIRE](#)].
- [38] CMS collaboration, *Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector*, **2019 JINST** **14** P07004 [[arXiv:1903.06078](#)] [[INSPIRE](#)].
- [39] CMS collaboration, *Mitigation of anomalous missing transverse momentum measurements in data collected by CMS at $\sqrt{s} = 13$ TeV during the LHC Run 2*, **CMS-DP-2020-018** (2020).
- [40] ALEPH et al. collaborations, *Precision electroweak measurements on the Z resonance*, **Phys. Rep.** **427** (2006) 257 [[hep-ex/0509008](#)] [[INSPIRE](#)].
- [41] CMS collaboration, *Observation of Single Top Quark Production in Association with a Z Boson in Proton-Proton Collisions at $\sqrt{s} = 13$ TeV*, **Phys. Rev. Lett.** **122** (2019) 132003 [[arXiv:1812.05900](#)] [[INSPIRE](#)].
- [42] CMS collaboration, *Observation of the Production of Three Massive Gauge Bosons at $\sqrt{s} = 13$ TeV*, **Phys. Rev. Lett.** **125** (2020) 151802 [[arXiv:2006.11191](#)] [[INSPIRE](#)].
- [43] CMS collaboration, *Observation of Higgs boson decay to bottom quarks*, **Phys. Rev. Lett.** **121** (2018) 121801 [[arXiv:1808.08242](#)] [[INSPIRE](#)].
- [44] CMS collaboration, *Measurements of Inclusive W and Z Cross Sections in pp Collisions at $\sqrt{s} = 7$ TeV*, **JHEP** **01** (2011) 080 [[arXiv:1012.2466](#)] [[INSPIRE](#)].
- [45] C.J. Clopper and E.S. Pearson, *The use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial*, **Biometrika** **26** (1934) 404 [[INSPIRE](#)].
- [46] CMS collaboration, *Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV*, **JHEP** **07** (2018) 161 [[arXiv:1802.02613](#)] [[INSPIRE](#)].
- [47] CMS collaboration, *Luminosity measurement in proton-proton collisions at 13.6 TeV in 2022 at CMS*, **CMS-PAS-LUM-22-001** (2024).
- [48] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS*, **Eur. Phys. J. C** **81** (2021) 800 [[arXiv:2104.01927](#)] [[INSPIRE](#)].
- [49] R.J. Barlow and C. Beeston, *Fitting using finite Monte Carlo samples*, **Comput. Phys. Commun.** **77** (1993) 219 [[INSPIRE](#)].
- [50] CMS collaboration, *The CMS Statistical Analysis and Combination Tool: Combine*, **Comput. Softw. Big Sci.** **8** (2024) 19 [[arXiv:2404.06614](#)] [[INSPIRE](#)].
- [51] M. Grazzini, S. Kallweit, J.M. Lindert, S. Pozzorini and M. Wiesemann, *NNLO QCD + NLO EW with Matrix+OpenLoops: precise predictions for vector-boson pair production*, **JHEP** **02** (2020) 087 [[arXiv:1912.00068](#)] [[INSPIRE](#)].

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan ¹

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic , L. Benato , T. Bergauer , S. Chatterjee , K. Damanakis ,
M. Dragicevic , P.S. Hussain , M. Jeitler ², N. Krammer , A. Li , D. Liko , I. Mikulec ,
J. Schieck ², R. Schöffbeck ², D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz ²

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , T. Van Laer 

Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen,
S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat ³, G.P. Van Onsem ,
S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das , I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart ,
P. Gianneios , J. Jaramillo , A. Khalilzadeh , F.A. Khan , K. Lee , A. Malara , M.A. Shahzad,
L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , Y. Hong , J. Knolle , L. Lambrecht , D. Marckx ,
K. Mota Amarilo , K. Skovpen , N. Van Den Bossche , J. van der Linden , L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere ,
I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre , J. Lidrych ,
P. Mastrapasqua , T.T. Tran , S. Turkcapar 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , G. Correia Silva , C. Hensel , T. Menezes De Oliveira ,
C. Mora Herrera ⁴, P. Rebello Teles , M. Soeiro , E.J. Tonelli Manganote ⁵, A. Vilela Pereira ⁴

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho ,
J. Chinellato ⁶, E.M. Da Costa , G.G. Da Silveira ⁷, D. De Jesus Damiao ,
S. Fonseca De Souza , R. Gomes De Souza , T. Laux Kuhn ⁷, M. Macedo , J. Martins ,
L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes ⁷, L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores ,
I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova ,
G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu

Institute of High Energy Physics, Beijing, China

G.M. Chen ⁸, H.S. Chen ⁸, M. Chen ⁸, F. Iemmi , C.H. Jiang, A. Kapoor ⁹, H. Liao ,
Z.-A. Liu ¹⁰, R. Sharma ¹¹, J.N. Song¹⁰, J. Tao , C. Wang⁸, J. Wang , Z. Wang⁸, H. Zhang ,
J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, C. Jiang ,
A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang,
L. Zhang , Y. Zhao, C. Zhou 

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer¹², B. Li¹³, K. Yi ¹⁴, J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li

Zhejiang University, Hangzhou, Zhejiang, ChinaZ. Lin , C. Lu , M. Xiao **Universidad de Los Andes, Bogota, Colombia**C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega**Universidad de Antioquia, Medellin, Colombia**C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez **University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac **University of Split, Faculty of Science, Split, Croatia**M. Kovac , A. Petkovic , T. Sculac **Institute Rudjer Boskovic, Zagreb, Croatia**P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov ¹⁵, T. Susa **University of Cyprus, Nicosia, Cyprus**A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , J. Mousa , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov **Charles University, Prague, Czech Republic**M. Finger , M. Finger Jr. , A. Kveton **Escuela Politecnica Nacional, Quito, Ecuador**E. Ayala **Universidad San Francisco de Quito, Quito, Ecuador**E. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**A.A. Abdelalim ^{16,17}, S. Elgammal¹⁸, A. Ellithi Kamel¹⁹**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**M. Abdullah Al-Mashad , M.A. Mahmoud **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken **Department of Physics, University of Helsinki, Helsinki, Finland**H. Kirschenmann , K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew ,
K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Myllymäki ,
M.M. Rantanen , H. Siikonen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour ,
P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi ,
L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro ²⁰, P. Simkina , M. Titov ,
M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi ,
T.D. Cuisset , F. Damas , O. Davignon , A. De Wit , I.T. Ehle ,
B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi ,
B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen , C. Ochando , R. Salerno ,
J.B. Sauvan , Y. Sirois , L. Urda Gómez , E. Vernazza , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram ²¹, J. Andrea , D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert ,
C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet ,
G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo ,
P. Depasse , C. Dozen ²², H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch ,
C. Greenberg , G. Grenier , B. Ille , E. Jourdhuy, I.B. Laktineh, M. Lethuillier , L. Mirabito,
S. Perries, A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

Georgian Technical University, Tbilisi, Georgia

I. Lomidze , T. Toriashvili ²³, Z. Tsamalaidze ²⁴

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser ,
A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann ,
M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung 

M.Y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , A. Stein , F. Torres Da Silva De Araujo ²⁵, S. Wiedenbeck , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman ²⁶, K. Borras ²⁷, A. Campbell , A. Cardini , C. Cheng , F. Colombina , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , E. Gallo ²⁶, A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁸, W. Lohmann ²⁹, F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , Y. Otariid, J. Park , E. Ranken , A. Raspereza , D. Rastorguev , J. Rübenach, L. Rygaard, A. Saggio , M. Scham ^{30,27}, S. Schnake ²⁷, P. Schütze , C. Schwanenberger ²⁶, D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford , F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann, L. Wiens ²⁷, C. Wissing , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , S. Bollweg, M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , D. Hundhausen, H.R. Jabusch , G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer , C.C. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, B. Wiederspan, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , T. Chwalek , A. Dierlamm , A. Droll, U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ³¹, R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , O. Lavoryk , J.M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis , A. Papadopoulos³¹, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi ,
K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos ,
E. Siamarkou, G. Tsipolitis , A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas ,
P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath ^{32,33}, K. Márton, A.J. Rádl ³⁴, F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti ³⁵, M.M.A. Gadallah ³⁶, Á. Kadlecik , P. Major ,
G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo ³⁵, F. Nemes ³⁵, T. Novak 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra ³⁷, A. Kaur ,
A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar ,
M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman ³⁸, S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana , P. Kalbhor ,
S. Kamble , J.R. Komaragiri ³⁹, D. Kumar ³⁹, T. Mishra , B. Parida ⁴⁰, P.R. Pujahari ,
N.R. Saha , A. Sharma , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, G.B. Mohanty , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar ,
G. Majumdar , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati , C. Kar , D. Maity , P. Mal , K. Naskar , A. Nayak , S. Nayak,
K. Pal , P. Sadangi, S.K. Swain , S. Varghese , D. Vats 

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya , A. Alpana , S. Dube , B. Gomber , P. Hazarika , B. Kansal , A. Laha ,
B. Sahu , S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi , A. Jafari , M. Zeinali 

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie ,
M. Mohammadi Najafabadi , S. Tizchang 

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia , A. Colaleo , D. Creanza , B. D'Anzi , N. De Filippis ,
M. De Palma , W. Elmetenawee , N. Ferrara , L. Fiore , G. Iaselli , L. Longo ,
M. Louka , G. Maggi , M. Maggi , I. Margjeka , V. Mastrapasqua , S. My ,
S. Nuzzo , A. Pellecchia , A. Pompili , G. Pugliese , R. Radogna , D. Ramos ,
A. Ranieri , L. Silvestris , F.M. Simone , Ü. Sözbilir , A. Stamerra , D. Troiano ,
R. Venditti , P. Verwilligen , A. Zaza 

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi , C. Battilana , D. Bonacorsi , P. Capiluppi , A. Castro ,
F.R. Cavallo , M. Cuffiani , G.M. Dallavalle , T. Diotallevi , F. Fabbri ,
A. Fanfani , D. Fasanella , P. Giacomelli , L. Giommi , C. Grandi , L. Guiducci ,
S. Lo Meo , M. Lorusso , L. Lunerti , S. Marcellini , G. Masetti , F.L. Navarria ,
G. Paggi , A. Perrotta , F. Primavera , A.M. Rossi , S. Rossi Tisbeni ,
T. Rovelli , G.P. Siroli 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa , A. Di Mattia , A. Lapertosa , R. Potenza , A. Tricomi , C. Tuve 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras , G. Barbagli , G. Bardelli , B. Camaiani , A. Cassese , R. Ceccarelli ,
V. Ciulli , C. Civinini , R. D'Alessandro , E. Focardi , T. Kello , G. Latino 

P. Lenzi ^{a,b}, M. Lizzo ^a, M. Meschini ^a, S. Paoletti ^a, A. Papanastassiou^{a,b}, G. Sguazzoni ^a,
L. Viliani ^a

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola ⁵¹, D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira ^a, F. Ferro ^a, E. Robutti ^a, S. Tosi ^{a,b}

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia ^a, F. Brivio ^a, F. Cetorelli ^{a,b}, F. De Guio ^{a,b}, M.E. Dinardo ^{a,b}, P. Dini ^a,
S. Gennai ^a, R. Gerosa ^{a,b}, A. Ghezzi ^{a,b}, P. Govoni ^{a,b}, L. Guzzi ^a, M.T. Lucchini ^{a,b},
M. Malberti ^a, S. Malvezzi ^a, A. Massironi ^a, D. Menasce ^a, L. Moroni ^a, M. Paganoni ^{a,b},
S. Palluotto ^{a,b}, D. Pedrini ^a, A. Perego ^{a,b}, B.S. Pinolini^a, G. Pizzati ^{a,b}, S. Ragazzi ^{a,b},
T. Tabarelli de Fatis ^{a,b}

INFN Sezione di Napoli^a, Università di Napoli ‘Federico II’^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo ^a, A. Cagnotta ^{a,b}, F. Carnevali^{a,b}, N. Cavallo ^{a,c}, F. Fabozzi ^{a,c},
A.O.M. Iorio ^{a,b}, L. Lista ^{a,b,52}, P. Paolucci ^{a,31}, B. Rossi ^a

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino ^a, P. Azzi ^a, N. Bacchetta ^{a,53}, D. Bisello ^{a,b}, P. Bortignon ^a, G. Bortolato^{a,b},
A. Bragagnolo ^{a,b}, A.C.M. Bulla ^a, R. Carlin ^{a,b}, P. Checchia ^a, T. Dorigo ^{a,54},
U. Gasparini ^{a,b}, S. Giorgetti^a, E. Lusiani ^a, M. Margoni ^{a,b}, M. Michelotto ^a,
M. Migliorini ^{a,b}, F. Montecassiano ^a, J. Pazzini ^{a,b}, P. Ronchese ^{a,b}, R. Rossin ^{a,b},
F. Simonetto ^{a,b}, M. Tosi ^{a,b}, A. Triossi ^{a,b}, S. Ventura ^a, M. Zanetti ^{a,b}, P. Zotto ^{a,b},
A. Zucchetta ^{a,b}, G. Zumerle ^{a,b}

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

A. Braghieri ^a, S. Calzaferri ^a, D. Fiorina ^a, P. Montagna ^{a,b}, V. Re ^a, C. Riccardi ^{a,b},
P. Salvini ^a, I. Vai ^{a,b}, P. Vitulo ^{a,b}

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal ^{a,b}, M.E. Ascioti^{a,b}, G.M. Bilei ^a, C. Carrivale^{a,b}, D. Ciangottini ^{a,b}, L. Fanò ^{a,b},
M. Magherini ^{a,b}, V. Mariani ^{a,b}, M. Menichelli ^a, F. Moscatelli ^{a,55}, A. Rossi ^{a,b},
A. Santocchia ^{a,b}, D. Spiga ^a, T. Tedeschi ^{a,b}

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C. Aimè ^a, C.A. Alexe ^{a,c}, P. Asenov ^{a,b}, P. Azzurri ^a, G. Bagliesi ^a, R. Bhattacharya ^a,
L. Bianchini ^{a,b}, T. Boccali ^a, E. Bossini ^a, D. Bruschini ^{a,c}, R. Castaldi ^a, M.A. Ciocci ^{a,b},
M. Cipriani ^{a,b}, V. D’Amante ^{a,d}, R. Dell’Orso ^a, S. Donato ^a, A. Giassi ^a, F. Ligabue ^{a,c},
A.C. Marini ^a, D. Matos Figueiredo ^a, A. Messineo ^{a,b}, S. Mishra ^a,

V.K. Muraleedharan Nair Bindhu ^{a,b,42}, M. Musich ^{a,b}, F. Palla ^a, A. Rizzi ^{a,b}, G. Rolandi ^{a,c}, S. Roy Chowdhury ^a, T. Sarkar ^a, A. Scribano ^a, P. Spagnolo ^a, R. Tenchini ^a, G. Tonelli ^{a,b}, N. Turini ^{a,d}, F. Vaselli ^{a,c}, A. Venturi ^a, P.G. Verdini ^a

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

C. Baldenegro Barrera ^{a,b}, P. Barria ^a, C. Basile ^{a,b}, F. Cavallari ^a, L. Cunqueiro Mendez ^{a,b}, D. Del Re ^{a,b}, E. Di Marco ^{a,b}, M. Diemoz ^a, F. Errico ^{a,b}, R. Gargiulo^{a,b}, E. Longo ^{a,b}, L. Martikainen ^{a,b}, J. Mijuskovic ^{a,b}, G. Organtini ^{a,b}, F. Pandolfi ^a, R. Paramatti ^{a,b}, C. Quaranta ^{a,b}, S. Rahatlou ^{a,b}, C. Rovelli ^a, F. Santanastasio ^{a,b}, L. Soffi ^a, V. Vladimirov^{a,b}

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane ^{a,b}, R. Arcidiacono ^{a,c}, S. Argiro ^{a,b}, M. Arneodo ^{a,c}, N. Bartosik ^a, R. Bellan ^{a,b}, A. Bellora ^{a,b}, C. Biino ^a, C. Borca ^{a,b}, N. Cartiglia ^a, M. Costa ^{a,b}, R. Covarelli ^{a,b}, N. Demaria ^a, L. Finco ^a, M. Grippo ^{a,b}, B. Kiani ^{a,b}, F. Legger ^a, F. Luongo ^{a,b}, C. Mariotti ^a, L. Markovic ^{a,b}, S. Maselli ^a, A. Mecca ^{a,b}, L. Menzio^{a,b}, P. Meridiani ^a, E. Migliore ^{a,b}, M. Monteno ^a, R. Mulargia ^a, M.M. Obertino ^{a,b}, G. Ortona ^a, L. Pacher ^{a,b}, N. Pastrone ^a, M. Pelliccioni ^a, M. Ruspa ^{a,c}, F. Siviero ^{a,b}, V. Sola ^{a,b}, A. Solano ^{a,b}, A. Staiano ^a, C. Tarricone ^{a,b}, D. Trocino ^a, G. Umoret ^{a,b}, R. White ^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar ^{a,b}, S. Belforte ^a, V. Candelise ^{a,b}, M. Casarsa ^a, F. Cossutti ^a, K. De Leo ^a, G. Della Ricca ^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra ^a, J. Hong ^a, B. Kim ^a, J. Kim, D. Lee, H. Lee, S.W. Lee ^a, C.S. Moon ^a, Y.D. Oh ^a, M.S. Ryu ^a, S. Sekmen ^a, B. Tae, Y.C. Yang ^a

Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim ^a

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak ^a, P. Gwak ^a, H. Kim ^a, D.H. Moon ^a

Hanyang University, Seoul, Korea

E. Asilar ^a, J. Choi ^{a,56}, D. Kim ^a, T.J. Kim ^a, J.A. Merlin, Y. Ryou

Korea University, Seoul, Korea

S. Choi ^a, S. Han, B. Hong ^a, K. Lee, K.S. Lee ^a, S. Lee ^a, J. Yoo ^a

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh ^a, S. Yang ^a

Sejong University, Seoul, Korea

H.S. Kim ^a, Y. Kim, S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , Y.W. Kim , S. Ko , H. Kwon ,
H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh,
I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

S. Ha , K. Hwang , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 

Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs, A. Potrebko , M. Seidel ,
D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff ⁵⁷, Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Schrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo ,
I. Heredia-De La Cruz ⁵⁸, R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera,
A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, MexicoI. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada **University of Montenegro, Podgorica, Montenegro**I. Bubanja , N. Raicevic **University of Canterbury, Christchurch, New Zealand**P.H. Butler **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan **AGH University of Krakow, Krakow, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad **Warsaw University of Technology, Warsaw, Poland**P. Fokow , K. Pozniak , W. Zabolotny **Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**M. Araujo , D. Bastos , C. Beirão Da Cruz E. Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff **Faculty of Physics, University of Belgrade, Belgrade, Serbia**P. Adzic , P. Milenovic **VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**D. Devetak, M. Dordevic , J. Milosevic , L. Nadder , V. Rekovic**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C.M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , S. Sánchez Navas , J. Sastre , J. Vazquez Escobar 

Universidad Autónoma de Madrid, Madrid, SpainJ.F. de Trocóniz **Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain**

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras ,
 I. Gonzalez Caballero , P. Leguina , M. Obeso Menendez , E. Palencia Cortezon ,
 J. Prado Pico , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote ,
 C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros ,
 M. Fernandez , G. Gomez , C. Lasasoa García , R. Lopez Ruiz , C. Martinez Rivero ,
 P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos ,
 J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri LankaB. Kailasapathy ⁵⁹, D.D.C. Wickramarathna **University of Ruhuna, Department of Physics, Matara, Sri Lanka**W.G.D. Dharmaratna ⁶⁰, K. Liyanage , N. Perera **CERN, European Organization for Nuclear Research, Geneva, Switzerland**

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney ,
 A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borgonovi , C. Botta ,
 E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d’Enterria , A. Dabrowski ,
 A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni ,
 W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , J. Hegeman , J.K. Heikkilä , B. Huber ,
 V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban ²⁹, S. Laurila ,
 P. Lecoq , E. Leutgeb , C. Lourenço , L. Malgeri , M. Mannelli , M. Matthewman,
 A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat ,
 M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer ,
 M. Pierini , H. Qu , D. Rabady , B. Ribeiro Lopes , F. Riti , M. Rovere , H. Sakulin ,
 R. Salvatico , S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma ,
 K. Shchelina , P. Silva , P. Sphicas ⁶¹, A.G. Stahl Leiton , A. Steen , S. Summers ,
 D. Treille , P. Tropea , D. Walter , J. Wanczyk ⁶², J. Wang, S. Wuchterl , P. Zehetner ,
 P. Zejdl , W.D. Zeuner

PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua ⁶³, L. Caminada ⁶³, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram ,
 H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶³, L. Noehte ⁶³, T. Rohe , A. Samalan

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , A. Calandri , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'Archiac ⁶², A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani , L. Marchese , A. Mascellani ⁶², F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , B. Ristic , R. Seidita , J. Steggemann ⁶², A. Tarabini , D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler ⁶⁴, P. Bärtshi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechi , A. Macchiolo , P. Meiring , F. Meng , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger , R. Tramontano 

National Central University, Chung-Li, Taiwan

C. Adloff⁶⁵, D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari ³⁹

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.G. Chen, A. De Iorio , W.-S. Hou , T.H. Hsu, Y.W. Kao, S. Karmakar , G. Kole , Y.Y. Li , R.-S. Lu , E. Paganis , X.F. Su , J. Thomas-Wilsker , L.S. Tsai, D. Tsionou, H.Y. Wu, E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusitanand 

Tunis El Manar University, Tunis, Tunisia

Y. Maghrbi 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu ⁶⁶, E. Eskut , Y. Guler ⁶⁷, E. Gurbinar Guler ⁶⁷, C. Isik , O. Kara, A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir ⁶⁸, A. Polatoz , B. Tali ⁶⁹, U.G. Tok , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Sokmen, M. Yalvac ⁷⁰

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya ⁷¹, O. Kaya ⁷², S. Tekten ⁷³

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak ^{66,74}, G.G. Dincer ⁶⁶, S. Sen ⁷⁵

Istanbul University, Istanbul, Turkey

O. Aydilek ⁷⁶, B. Hacisahinoglu , I. Hos ⁷⁷, B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, TurkeyS. Cerci , B. Isildak ⁷⁸, D. Sunar Cerci , T. Yetkin **Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine**A. Boyaryntsev , B. Grynyov **National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine**L. Levchuk **University of Bristol, Bristol, United Kingdom**D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw, V.J. Smith , K. Walkingshaw Pass**Rutherford Appleton Laboratory, Didcot, United Kingdom**A.H. Ball, K.W. Bell , A. Belyaev ⁷⁹, C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis, K. Harder , S. Harper , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams **Imperial College, London, United Kingdom**I. Andreou , R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, C.A. Carrillo Montoya , G.S. Chahal ⁸⁰, D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer, G. Fedi , G. Hall , A. Howard, G. Iles , C.R. Knight , P. Krueper, J. Langford , K.H. Law , J. León Holgado , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios, M. Mieskolainen , J. Nash ⁸¹, M. Pesaresi , P.B. Pradeep, B.C. Radburn-Smith , A. Richards, A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee ³¹, M. Vojinovic , N. Wardle , D. Winterbottom **Brunel University, Uxbridge, United Kingdom**J.E. Cole , A. Khan, P. Kyberd , I.D. Reid **Baylor University, Waco, Texas, USA**S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann , K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson **Catholic University of America, Washington, DC, USA**R. Bartek , A. Dominguez , A.E. Simsek , S.S. Yu **The University of Alabama, Tuscaloosa, Alabama, USA**B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson, C.U. Perez , P. Rumerio ⁸², E. Usai , R. Yi 

Boston University, Boston, Massachusetts, USA

A. Akpinar^{[ID](#)}, C. Cosby^{[ID](#)}, G. De Castro, Z. Demiragli^{[ID](#)}, C. Erice^{[ID](#)}, C. Fangmeier^{[ID](#)},
C. Fernandez Madrazo^{[ID](#)}, E. Fontanesi^{[ID](#)}, D. Gastler^{[ID](#)}, F. Golf^{[ID](#)}, S. Jeon^{[ID](#)}, J. O’Cain, I. Reed^{[ID](#)},
J. Rohlf^{[ID](#)}, K. Salyer^{[ID](#)}, D. Sperka^{[ID](#)}, D. Spitzbart^{[ID](#)}, I. Suarez^{[ID](#)}, A. Tsatsos^{[ID](#)}, A.G. Zecchinelli^{[ID](#)}

Brown University, Providence, Rhode Island, USA

G. Barone^{[ID](#)}, G. Benelli^{[ID](#)}, D. Cutts^{[ID](#)}, L. Gouskos^{[ID](#)}, M. Hadley^{[ID](#)}, U. Heintz^{[ID](#)}, K.W. Ho^{[ID](#)},
J.M. Hogan^{[ID](#)}⁸³, T. Kwon^{[ID](#)}, G. Landsberg^{[ID](#)}, K.T. Lau^{[ID](#)}, J. Luo^{[ID](#)}, S. Mondal^{[ID](#)}, T. Russell,
S. Sagir^{[ID](#)}⁸⁴, X. Shen^{[ID](#)}, F. Simpson^{[ID](#)}, M. Stamenkovic^{[ID](#)}, N. Venkatasubramanian

University of California, Davis, Davis, California, USA

S. Abbott^{[ID](#)}, B. Barton^{[ID](#)}, C. Brainerd^{[ID](#)}, R. Breedon^{[ID](#)}, H. Cai^{[ID](#)},
M. Calderon De La Barca Sanchez^{[ID](#)}, M. Chertok^{[ID](#)}, M. Citron^{[ID](#)}, J. Conway^{[ID](#)}, P.T. Cox^{[ID](#)},
R. Erbacher^{[ID](#)}, F. Jensen^{[ID](#)}, O. Kukral^{[ID](#)}, G. Mocellin^{[ID](#)}, M. Mulhearn^{[ID](#)}, S. Ostrom^{[ID](#)}, W. Wei^{[ID](#)},
S. Yoo^{[ID](#)}, F. Zhang^{[ID](#)}

University of California, Los Angeles, California, USA

K. Adamidis, M. Bachtis^{[ID](#)}, D. Campos, R. Cousins^{[ID](#)}, A. Datta^{[ID](#)}, G. Flores Avila^{[ID](#)}, J. Hauser^{[ID](#)},
M. Ignatenko^{[ID](#)}, M.A. Iqbal^{[ID](#)}, T. Lam^{[ID](#)}, Y.F. Lo, E. Manca^{[ID](#)}, A. Nunez Del Prado, D. Saltzberg^{[ID](#)},
V. Valuev^{[ID](#)}

University of California, Riverside, Riverside, California, USA

R. Clare^{[ID](#)}, J.W. Gary^{[ID](#)}, G. Hanson^{[ID](#)}

University of California, San Diego, La Jolla, California, USA

A. Aportela, A. Arora^{[ID](#)}, J.G. Branson^{[ID](#)}, S. Cittolin^{[ID](#)}, S. Cooperstein^{[ID](#)}, D. Diaz^{[ID](#)}, J. Duarte^{[ID](#)},
L. Giannini^{[ID](#)}, Y. Gu, J. Guiang^{[ID](#)}, R. Kansal^{[ID](#)}, V. Krutelyov^{[ID](#)}, R. Lee^{[ID](#)}, J. Letts^{[ID](#)},
M. Masciovecchio^{[ID](#)}, F. Mokhtar^{[ID](#)}, S. Mukherjee^{[ID](#)}, M. Pieri^{[ID](#)}, D. Primosch, M. Quinnan^{[ID](#)},
V. Sharma^{[ID](#)}, M. Tadel^{[ID](#)}, E. Vourliotis^{[ID](#)}, F. Würthwein^{[ID](#)}, Y. Xiang^{[ID](#)}, A. Yagil^{[ID](#)}

University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas^{[ID](#)}, L. Brennan^{[ID](#)}, C. Campagnari^{[ID](#)}, K. Downham^{[ID](#)}, C. Grieco^{[ID](#)}, M.M. Hussain,
J. Incandela^{[ID](#)}, J. Kim^{[ID](#)}, A.J. Li^{[ID](#)}, P. Masterson^{[ID](#)}, H. Mei^{[ID](#)}, J. Richman^{[ID](#)}, S.N. Santpur^{[ID](#)},
U. Sarica^{[ID](#)}, R. Schmitz^{[ID](#)}, F. Setti^{[ID](#)}, J. Sheplock^{[ID](#)}, D. Stuart^{[ID](#)}, T.Á. Vámi^{[ID](#)}, X. Yan^{[ID](#)}, D. Zhang

California Institute of Technology, Pasadena, California, USA

S. Bhattacharya^{[ID](#)}, A. Bornheim^{[ID](#)}, O. Cerri, A. Latorre, J. Mao^{[ID](#)}, H.B. Newman^{[ID](#)},
G. Reales Gutiérrez, M. Spiropulu^{[ID](#)}, J.R. Vlimant^{[ID](#)}, C. Wang^{[ID](#)}, S. Xie^{[ID](#)}, R.Y. Zhu^{[ID](#)}

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison^{[ID](#)}, S. An^{[ID](#)}, P. Bryant^{[ID](#)}, M. Cremonesi, V. Dutta^{[ID](#)}, T. Ferguson^{[ID](#)},
T.A. Gómez Espinosa^{[ID](#)}, A. Harilal^{[ID](#)}, A. Kallil Tharayil, C. Liu^{[ID](#)}, T. Mudholkar^{[ID](#)}, S. Murthy^{[ID](#)},
P. Palit^{[ID](#)}, K. Park, M. Paulini^{[ID](#)}, A. Roberts^{[ID](#)}, A. Sanchez^{[ID](#)}, W. Terrill^{[ID](#)}

University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , G. Karathanasis , N. Manganello ,
J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner ,
N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Dickinson , J. Fan ,
X. Fan , S. Hogan , P. Kotamnives, J. Monroy , M. Oshiro , J.R. Patterson , M. Reid ,
A. Ryd , J. Thom , P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick ,
D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati ,
H.W.K. Cheung , F. Chlebana , G. Cummings , I. Dutta , V.D. Elvira , Y. Feng ,
J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl ,
D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , B. Jayatilaka ,
S. Jindariani , M. Johnson , U. Joshi , T. Klijsma , B. Klima , K.H.M. Kwok ,
S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima ,
C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba ,
D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena ⁸⁵, F. Ravera ,
A. Reinsvold Hall ⁸⁶, L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha ,
L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger ,
E.W. Vaandering , I. Zoi 

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh ,
E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez ,
G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig ,
Y. Takahashi , J. Wang 

Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , R. Hashmi , R.S. Kim , S. Kim ,
T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay , J. Zhang

Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , T. Elkafrawy ⁸⁷, M. Hohlmann , E. Yanes

University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov ,
C.E. Gerber , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.H. Lee , D.S. Lemos ,
A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti,
T. Roy , S. Rudrabhatla , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye ,
J. Yoo 

The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend, K. Dilsiz ⁸⁸, L. Emediato , G. Karaman , O.K. Köseyan ,
J.-P. Merlo, A. Mestvirishvili ⁸⁹, O. Neogi, H. Ogul ⁹⁰, Y. Onel , A. Penzo , C. Snyder,
E. Tiras ⁹¹

Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou ,
P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean ,
Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu ,
J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu ⁹², C. Rogan , C. Royon ,
S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, USA

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli ,
D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen ,
R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili ,
C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova ⁹³, A. Shevelev , L. Wang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

J. Bendavid , I.A. Cali , P.C. Chou , M. D'Alfonso , J. Eysermans , C. Freer ,
G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa ,
L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus ,
C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang ,
B. Wyslouch , T.J. Yang 

University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi ,
M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla ,
J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.W. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava ,
M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma, M. Campana , J. Dervan , Y. Haddad ,
Y. Han , I. Israr , A. Krishna , J. Li , M. Lu , R. McCarthy , D.M. Morse , V. Nguyen ,
T. Orimoto , A. Parker , L. Skinnari , E. Tsai , D. Wood 

Northwestern University, Evanston, Illinois, USA

J. Bueghly, S. Dittmer , K.A. Hahn , D. Li , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco

University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore , Y. Musienko ²⁴, H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, D.A. Wenzl, B.L. Winer , B.R. Yates 

Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , K. Coldham, P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully , L.H. Vage

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma

Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer, J.F. Schulte , M. Stojanovic , J. Thieman , A.K. Viridi , F. Wang , A. Wildridge , W. Xie , Y. Yao 

Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, USA

D. Acosta , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar , P. Parygin ⁹³, R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslawski , S. Konstantinou , I. Laflotte , A. Lath , R. Montalvo, K. Nash, J. Reichert , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti ,
N. Karunaratna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov ⁶², O. Bouhali ⁹⁴, R. Eusebi , J. Gilmore ,
T. Huang , T. Kamon ⁹⁵, H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens ,
A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , N. Gogate , A. Hussain , Y. Kazhykarim, K. Lamichhane ,
S.W. Lee , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli ,
A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovsky , C. Neu 

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts ,
C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless ,
J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida ,
L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi ,
W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin [†], I. Gorbunov , V. Karjavine ,
V. Korenkov , A. Lanev , A. Malakhov , V. Matveev ⁹⁶, V. Palichik , V. Perelygin ,
M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin ,
B.S. Yuldashev⁹⁷, A. Zarubin , I. Zhizhin , Yu. Andreev , A. Dermenev , S. Gninenko ,
N. Golubev , A. Karneyeu , D. Kirpichnikov , M. Kirsanov , N. Krasnikov , I. Tlisova ,
A. Toropin 

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

G. Gavrillov , V. Golovtcov , Y. Ivanov , V. Kim ⁹⁸, P. Levchenko ⁹⁹, V. Murzin ,
V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev[†], T. Aushev , K. Ivanov ,
V. Gavrillov , N. Lychkovskaya , A. Nikitenko ^{100,101}, V. Popov , A. Zhokin ,
M. Chadeeva ⁹⁸, R. Chistov ⁹⁸, S. Polikarpov ⁹⁸, V. Andreev , M. Azarkin , M. Kirakosyan,
A. Terkulov , E. Boos , V. Bunichev , M. Dubinin ⁸⁵, L. Dudko , A. Gribushin 

V. Klyukhin^{}, O. Kodolova^{}¹⁰¹, O. Lukina^{}, S. Obraztsov^{}, S. Petrushanko^{}, V. Savrin^{},
A. Snigirev^{}, V. Blinov⁹⁸, T. Dimova^{}⁹⁸, A. Kozyrev^{}⁹⁸, O. Radchenko^{}⁹⁸, Y. Skovpen^{}⁹⁸,
V. Kachanov^{}, S. Slabospitskii^{}, A. Uzunian^{}, A. Babaev^{}, V. Borshch^{}, D. Druzhkin^{}

[†] Deceased

¹ Also at Yerevan State University, Yerevan, Armenia

² Also at TU Wien, Vienna, Austria

³ Also at Ghent University, Ghent, Belgium

⁴ Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵ Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

⁶ Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸ Also at University of Chinese Academy of Sciences, Beijing, China

⁹ Also at China Center of Advanced Science and Technology, Beijing, China

¹⁰ Also at University of Chinese Academy of Sciences, Beijing, China

¹¹ Also at China Spallation Neutron Source, Guangdong, China

¹² Now at Henan Normal University, Xinxiang, China

¹³ Also at University of Shanghai for Science and Technology, Shanghai, China

¹⁴ Now at The University of Iowa, Iowa City, Iowa, USA

¹⁵ Also at an institute formerly covered by a cooperation agreement with CERN

¹⁶ Also at Helwan University, Cairo, Egypt

¹⁷ Now at Zewail City of Science and Technology, Zewail, Egypt

¹⁸ Now at British University in Egypt, Cairo, Egypt

¹⁹ Now at Cairo University, Cairo, Egypt

²⁰ Also at Purdue University, West Lafayette, Indiana, USA

²¹ Also at Université de Haute Alsace, Mulhouse, France

²² Also at Istinye University, Istanbul, Turkey

²³ Also at Tbilisi State University, Tbilisi, Georgia

²⁴ Also at an international laboratory covered by a cooperation agreement with CERN

²⁵ Also at The University of the State of Amazonas, Manaus, Brazil

²⁶ Also at University of Hamburg, Hamburg, Germany

²⁷ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁸ Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

²⁹ Also at Brandenburg University of Technology, Cottbus, Germany

³⁰ Also at Forschungszentrum Jülich, Jülich, Germany

³¹ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

³² Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

³³ Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

³⁴ Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³⁵ Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

³⁶ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

³⁷ Also at Punjab Agricultural University, Ludhiana, India

³⁸ Also at University of Visva-Bharati, Santiniketan, India

³⁹ Also at Indian Institute of Science (IISc), Bangalore, India

⁴⁰ Also at Amity University Uttar Pradesh, Noida, India

⁴¹ Also at IIT Bhubaneswar, Bhubaneswar, India

⁴² Also at Institute of Physics, Bhubaneswar, India

⁴³ Also at University of Hyderabad, Hyderabad, India

⁴⁴ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany

⁴⁵ Also at Isfahan University of Technology, Isfahan, Iran

⁴⁶ Also at Sharif University of Technology, Tehran, Iran

- ⁴⁷ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁸ Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁹ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁵⁰ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁵¹ Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁵² Also at Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Napoli, Italy
- ⁵³ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵⁴ Also at Lulea University of Technology, Lulea, Sweden
- ⁵⁵ Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵⁶ Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁷ Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁸ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- ⁵⁹ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁶⁰ Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁶¹ Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶² Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶³ Also at Universität Zürich, Zurich, Switzerland
- ⁶⁴ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶⁵ Also at Laboratoire d’Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- ⁶⁶ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁷ Also at Konya Technical University, Konya, Turkey
- ⁶⁸ Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁹ Also at Adiyaman University, Adiyaman, Turkey
- ⁷⁰ Also at Bozok Universitеси Rektörlüğü, Yozgat, Turkey
- ⁷¹ Also at Marmara University, Istanbul, Turkey
- ⁷² Also at Milli Savunma University, Istanbul, Turkey
- ⁷³ Also at Kafkas University, Kars, Turkey
- ⁷⁴ Now at Istanbul Okan University, Istanbul, Turkey
- ⁷⁵ Also at Hacettepe University, Ankara, Turkey
- ⁷⁶ Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ⁷⁷ Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁸ Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁹ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁸⁰ Also at IPPP Durham University, Durham, United Kingdom
- ⁸¹ Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸² Also at Università di Torino, Torino, Italy
- ⁸³ Also at Bethel University, St. Paul, Minnesota, USA
- ⁸⁴ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸⁵ Also at California Institute of Technology, Pasadena, California, USA
- ⁸⁶ Also at United States Naval Academy, Annapolis, Maryland, USA
- ⁸⁷ Also at Ain Shams University, Cairo, Egypt
- ⁸⁸ Also at Bingol University, Bingol, Turkey
- ⁸⁹ Also at Georgian Technical University, Tbilisi, Georgia
- ⁹⁰ Also at Sinop University, Sinop, Turkey
- ⁹¹ Also at Erciyes University, Kayseri, Turkey
- ⁹² Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ⁹³ Now at another institute formerly covered by a cooperation agreement with CERN
- ⁹⁴ Also at Texas A&M University at Qatar, Doha, Qatar
- ⁹⁵ Also at Kyungpook National University, Daegu, Korea
- ⁹⁶ Also at another international laboratory covered by a cooperation agreement with CERN

⁹⁷ *Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan*

⁹⁸ *Also at another institute formerly covered by a cooperation agreement with CERN*

⁹⁹ *Also at Northeastern University, Boston, Massachusetts, USA*

¹⁰⁰ *Also at Imperial College, London, United Kingdom*

¹⁰¹ *Now at Yerevan Physics Institute, Yerevan, Armenia*