

---

# 249-2 - Booth No. 72: PROVENANCE OF THE UPPERMOST CARBONIFEROUS–LOWER TRIASSIC SANDSTONES, BOGDA MOUNTAINS, NW CHINA: IMPLICATION ON THE LATE PALEOZOIC TECTONIC HISTORY OF THE SOUTHERN CENTRAL ASIAN OROGENIC BELT



Wednesday, September 25, 2024



8:00 AM - 5:30 PM



Hall D (Anaheim Convention Center)

---

## Booth No. 72

---

### Abstract

The Permian-Triassic time is a significant stage in the Paleozoic continental amalgamation and Cenozoic orogenic reactivation of southern Central Asian Orogenic Belt (CAOB). Field, petrographic, and detrital zircon U-Pb geochronological data of the uppermost Carboniferous–Lower Triassic sandstones from 3 sections in Bogda Mountains, greater Turpan-Junggar basin, NW China, are used to decipher the tectonic history. The sections are Tarlong-Taodonggou (TT) and Zhaobishan (ZBS) in the south and Dalongkou (DLK) in the north, 100 km apart and ~7,000 m in total thickness. Four petrofacies of 229 sandstones and U-Pb dates of 3505 zircons of 35 sandstones form the basis for interpretation. During Gzhelian–Asselian, andesite and basalt are the major source lithologies in TT. Zircon ages peak at ~300 Ma. During Sakmarian–Kungurian, basalt and andesite are the main source rocks in TT and ZBS; and zircon ages of both areas peak at ~300 Ma. The Roadian–Wordian is represented by a regional unconformity. The Guadalupian source lithology and zircon date show a major change. Andesite is the common and rhyolite and basalt the minor source lithologies for TT and DLK; but rhyolite for ZBS. A unimodal peak at ~305 Ma occurs in TT; but two peaks at 305 and 455 Ma with common Precambrian dates in ZBS; and peaks of 310–295 Ma in DLK. During Wuchiapingian–mid Olenekian, andesite and rhyolite are the common source lithologies for TT and DLK; but rhyolite as the primary volcanic lithology for ZBS. In TT, Wuchiapingian–Induan samples have a major age peak at ~300 Ma, and an Olenekian sample has two peaks at ~300 and ~450 Ma. In ZBS, the age pattern is similar to that of the Guadalupian sample. In DLK, samples have a major peak at ~310 Ma and a minor peak at ~450 Ma. The comparable age clusters identified by multi-dimensional scaling indicate that North Tianshan is the source for TT and ZBS during the latest Carboniferous–early Permian. But in mid Permian, south Central Tianshan became the main source solely to ZBS. During late Permian–Early Triassic, both north and central Tianshan became the common sources to all three areas due to enhanced denudation. The source change in mid-Permian across

a regional unconformity is synchronous with Paleo-Asian Ocean closure and arc-continent and continent-continent collisions, which occurred along the southern margin of Turpan-Junggar basin no later than Guadalupian.

---

Geological Society of America Abstracts with Programs. Vol. 56, No. 5, 2024  
doi: 10.1130/abs/2024AM-401642

© Copyright 2024 The Geological Society of America (GSA), all rights reserved.

## Author

---



**Sixuan Wu**  
Missouri University of Science and Technology



**Wan Yang**  
Missouri University of Science and Technology

---

---

## View Related

---

---