



Review

Bioengineering tissue morphogenesis and function in human neural organoids



Nikolai J. Fedorchak^a, Nisha Iyer^a, Randolph S. Ashton^{a,b,*}

^a Wisconsin Institute for Discovery, University of Wisconsin–Madison, Madison, WI 53715, United States

^b Department of Biomedical Engineering, University of Wisconsin–Madison, Madison, WI 53706, United States

ARTICLE INFO

Keywords:

Tissue morphology
Signaling centers
Morphogenetic patterning
Assembloids
Vascularization
Circuit formation
Network maturation

ABSTRACT

Over the last decade, scientists have begun to model CNS development, function, and disease *in vitro* using human pluripotent stem cell (hPSC)-derived organoids. Using traditional protocols, these 3D tissues are generated by combining the innate emergent properties of differentiating hPSC aggregates with a bioreactor environment that induces interstitial transport of oxygen and nutrients and an optional supportive hydrogel extracellular matrix (ECM). During extended culture, the hPSC-derived neural organoids (hNOs) obtain millimeter scale sizes with internal microscale cytoarchitectures, cellular phenotypes, and neuronal circuit behaviors mimetic of those observed in the developing brain, eye, or spinal cord. Early studies evaluated the cytoarchitectural and phenotypical character of these organoids and provided unprecedented insight into the morphogenetic processes that govern CNS development. Comparisons to human fetal tissues revealed their significant similarities and differences. While hNOs have current disease modeling applications and significant future promise, their value as anatomical and physiological models is limited because they fail to form reproducibly and recapitulate more mature *in vivo* features. These include biomimetic macroscale tissue morphology, positioning of morphogen signaling centers to orchestrate appropriate spatial organization and intra- and inter-connectivity of discrete tissue regions, maturation of physiologically relevant neural circuits, and formation of vascular networks that can support sustained *in vitro* tissue growth. To address these inadequacies scientists have begun to integrate organoid culture with bioengineering techniques and methodologies including genome editing, biomaterials, and microfabricated and microfluidic platforms that enable spatiotemporal control of cellular differentiation or the biochemical and biophysical cues that orchestrate organoid morphogenesis. This review will examine recent advances in hNO technologies and culture strategies that promote reproducible *in vitro* morphogenesis and greater biomimicry in structure and function.

1. Introduction

In 1998, James Thomson and colleagues reported the isolation of human embryonic stem cells (hESCs) from the inner cell mass of human blastocysts [1]. This generated excitement for the potential of hESC-derived cellular therapeutics. Yet, utilization of human pluripotent stem cells (hPSCs) for research purposes such as the study of cell fate acquisition, tissue development and physiology, disease modeling, and drug screening have been their most impactful applications to date. A decade after Thomson's publication, Yoshiki Sasai demonstrated that 3D cellular aggregates of both mouse and human embryonic stem cells (hESCs) could spontaneously morph into polarized cortical tissues *in vitro*. These cortical tissues featured microscale cytoarchitectures

reminiscent of the developing neocortex [2]. With this seminal demonstration, hPSC-derived neural organoid (hNO) culture, a practice that attempts to recapitulate central nervous system (CNS) morphogenesis *in vitro*, emerged as a new experimental modality to study early signaling processes that are essential to brain, retina, and spinal cord development, function, and disease. Since Sasai's publication, organoids featuring cytoarchitectures and functions similar to a variety of other CNS tissues have been demonstrated. This includes the developing human telencephalon, optic cup, diencephalon, midbrain, cerebellum, and spinal cord [3–8].

The importance of hNOs as an experimental model cannot be understated. Classical models of CNS morphogenesis were first derived from examination of non-mammalian vertebrate organisms [9,10].

* Corresponding author at: Wisconsin Institute for Discovery, University of Wisconsin–Madison, Madison, WI 53715, United States.

E-mail address: rashton2@wisc.edu (R.S. Ashton).

These early characterizations informed the development of non-human mammalian models [11]. However, those models are not generalizable to humans due to non-conserved CNS features between species of different vertebrate clades [12]. For instance, the developing human neocortex features both an inner and outer subventricular zone (iSVZ, oSVZ), with oSVZ radial glial cells dividing rapidly to increase the number and potentially the diversity of cortical neural cell fates [13,14]. The oSVZ radial glial cell phenotype even differs between humans and other primates, and it likely has a role in the evolutionary expansion of the human neocortex [15,16]. Thus, there are inherent limitations in using other vertebrate species to draw conclusion about human CNS development and disease pathologies. The use of hNOs enables the study of some facets of human CNS development while circumventing modeling and resource limitations associated with using animal or human fetal tissue.

Human neural organoids are distinct from traditional neurospheres and 3D tissue engineered constructs consisting of cells seeded on a 3D scaffold. As described by Marti-Figueroa et al. and others, organoids are derived from hPSCs or organ progenitors and form tissues that self-organize through cell sorting and spatially restricted lineage commitment in a biomimetic manner [17–19]. They initiate as an aggregate of stem cells receptive to biophysical and biochemical inputs. Over time, extrinsic inputs and intrinsic cellular signaling propel the aggregated stem cells to morph, i.e. differentiate and self-assemble, into a more complex tissue. This morphogenic process results in a level of biomimetic cell phenotypic diversity and microscale tissues cytoarchitecture that is not achievable by neurospheres and 3D tissue engineered constructs [20,21].

Traditionally, there are two main strategies to culture neural organoids. One strategy utilizes an unguided approach wherein minimal cues coax cellular aggregates to differentiate and self-organize [22,23]. Despite its minimalistic appearance, this method often generates ‘mixed’ organoids where the majority of the cells acquire various neuroectodermal fates but a small portion adopt phenotypes of other ectodermal, mesodermal, or endodermal lineages. Alternatively, directed culture strategies utilize more extrinsic cues and growth factors to pattern neural organoids towards a specific CNS regional fate [4,6,7, 24–27]. This latter strategy reduces the semi-stochastic cell fate acquisition associated with the unguided approach but simultaneously reduces the diversity of neuroectodermal-derived tissue and cell phenotypes that emerge within the neural organoid. Despite their modeling superiority to standard cell culture methodologies, hNOs grown using the previously mentioned strategies have persistent biomimicry and standardization issues that could be rectified by enabling more precise spatiotemporal control of their *in vitro* morphogenesis.

Here, we review bioengineering approaches and methodologies that have been used to improve hNO recapitulation of CNS morphogenesis, tissue structure, and function. The review is not meant to be exhaustive, but instead, will highlight exemplars that encapsulate the various approaches found within the hNO field. In particular, the review focuses on the integration of bioengineering techniques to help improve the reproducibility of hNO cellular composition and anatomy [22,28,29], formation of signaling centers to produce stereotype morphogenetic patterning [30], generation of biomimetic circuits including inter-regional connectivity [28], and vascularization to enhance maturation [31] and reduce cell death [22]. Initially, hNO morphogenesis was thought to be an inherently variable, spontaneous process. However, bioengineering strategies are now being developed and integrated to effectively control such emergent behaviors.

2. Bioengineering hNO morphology

CNS morphogenesis is orchestrated *in vivo* by myriad, spatially and temporally discrete, biochemical and biophysical factors. Lack of control over such factors during hNO derivation is the source of organoid variability and failure, in some aspects, to recapitulate normal

developmental processes. Application of biophysical constraints during the organoid derivation process is a tractable feat. Thus, researchers have begun integrating bioengineering approaches with hNO derivation protocols to more effectively control tissue morphology (Fig. 1).

For over two decades, soft lithography techniques such as micro-contact printing of alkanethiol self-assembled monolayers have been used to pattern cell adhesion at the microscale on optically transparent substrates [32]. Recently, Knight et al. used these techniques to investigate how a hPSC aggregate's microscale morphology would influence subsequent neuroepithelial tissue emergence, i.e. the inception of hNO morphogenesis [33,34]. They hypothesized that the spontaneous neurulation-like events that generate polarized neuroepithelial tissues, a.k.a. neural rosettes, within hNOs could be reproducibly induced in neurally differentiating hPSC aggregates of controlled microscale dimensions. In their experiments, hPSCs were seeded at cell densities sufficient to form a confluent monolayer on micropatterned regions, and over 5 days in neural induction media, the cells proliferated to form 3-D hemispherical tissues. They observed that hPSCs differentiated to a forebrain neuroepithelial fate and cultured on 250 μm diameter circular micropatterns reproducibly formed tissues with a singularly polarized neural rosette, which is mimetic of the embryonic neural tube's cytoarchitecture. Interestingly, reproducible formation of spinal neuroepithelial tissues with a singularly polarized rosette required smaller 150 μm diameter circular micropatterns due to differences in the cells' biomechanical properties [34]. Their observed effect of hPSC aggregate morphology on the emergence of polarized forebrain neuroepithelium was also reproduced by Harema et al. during derivation of micro-patterned human ‘neuruloids’ [35]. In this study, reproducible and spatially stereotyped ectodermal morphogenesis, which entails emergence of neuroepithelial, neural crest, sensory placode and epidermal tissues, was achieved after forming hPSCs aggregates on 500 μm diameter circular micropatterns and with exposure to bone morphogenic protein-4 (BMP4). These examples demonstrate that even in adherent cultures, inceptive tissue morphology is a biophysical characteristic that can be engineered to effectively regulate subsequent neural morphogenesis.

When deriving suspension hNOs, the initial spherical hPSC aggregate size has also been proven to be an important determinant for protocol reproducibility. For example, Zhu and colleagues engineered a micro-pillar array chip in which seeded hPSCs aggregate within the array's inter-pillar regions to form spherical embryoid bodies (EBs) of uniform shape and size [36]. EB size can be varied by adjusting the micropillar array's dimensions, and hNOs were most consistent when the micro-pillars were 800 μm in height with a 50 μm pitch. More commonly, researchers form cell aggregates from a specified number of hPSCs using low-attachment V-bottom well or AggreWell™ plates [37,38]. In these platforms, cell aggregate size is indirectly controlled via cell seeding density and well dimensions. Subsequent dorsal forebrain patterning of such aggregates in suspension culture has been implemented at scale to generate cortical hNOs with neuronal subtype lamination similar to the inside-out patterning observed *in vivo* [24,38–40]. At ~7 weeks post aggregation, astrogenesis can be observed within these hNOs, and at ~10-weeks post aggregation, cortical hNOs possess transcriptional profiles equivalent to the second trimester human fetal cortex [39]. Single-cell RNA-seq analysis proved that hNOs derived using such controlled conditions yielded highly reproducible cellular compositions, even across multiple hPSC lines [37]. Moreover, the diversity of constituent cell phenotypes and their transcriptional profile was similar to that of fetal human cortex at ~6 months of gestation [38]. While the described cortical hNO protocols begin with forming spheroidal cell aggregates of controlled, uniform size, they also exclude embedding the organoid within a xenogenic extracellular matrix (EMC) hydrogel during long-term culture. Hence, the protocols reproducibility may not be solely attributed to its inception via uniformly sized hPSC aggregates.

As an alternative approach to focusing on the initial spheroidal cell aggregate's size, Lancaster and colleagues used poly(lactic-co-glycolic




Fig. 1. Approaches to bioengineer human neural organoid (hNO) morphology. Human pluripotent stem cell (hPSC) aggregate morphology can be regulated at the microscale using microfilaments, micropatterned substrates, micropillar arrays, microwells, molded or 3D printed hydrogels, and microfabricated chamber devices. Gene editing tools may also indirectly exert control over hNO morphology (e.g. folding) by regulating cell behavior (e.g. proliferation).

acid) microfilaments to seed the formation of elongated hPSCs aggregates with increased surface area-to-volume ratios [41]. They hypothesized that the increased surface-area-to-volume ratio would expose a larger fraction of the aggregate's cells to its exterior surface resulting in more homogenous neural induction. Indeed, their elongated hPSCs aggregates preferentially differentiated into neuroepithelium with negligible presence of mesodermal and endodermal phenotypes. The resulting microfilament-engineered cerebral organoids (encORs) displayed a significant and reproducible increase in the formation of forebrain tissues at the expense of more caudal midbrain and hindbrain tissues. The observed alteration in organoid composition due to the initial hPSC aggregate's microfilament-guided increase in surface area-to-volume ratio further exemplifies the influence of the inceptive aggregate's biophysical characteristics on organoid morphogenesis.

Regulating hNO morphology during latter stages of derivation protocols is challenging due to the near ubiquitous need for agitated suspension culture and the organoid's constantly increasing size [22]. McNulty and Marti-Figueroa et al. attempted to address this issue using sacrificially molded alginate hydrogels to encapsulate hPSC aggregates and provide sustained imposition of a cylindrical morphology with defined microscale dimensions [42]. The combined rigidity and porosity of alginate hydrogels enabled stirred-tank bioreactor culture of the entire organoid-hydrogel composite, and robust neuroepithelium formation was observed over the course of 16 days. However, longer-term culture was not feasible due to the hydrogel cavity's inability to expand with the growing neural organoid. Alternatively, Karzbrun and colleagues avoided agitated suspension culture altogether while using microfabricated culture chambers to constrain the morphology of developing hNOs [43]. This biophysical constraint induced compression forces within the organoid's expanding neuroepithelium, and mechanical instabilities resulting from local changes in cell density as well as apical cytoskeletal contraction caused 'wrinkling.' The authors argue that this is a model of folding during normal human brain development [43]. However, a more convincing model of human brain folding was developed by Li et al.'s derivation of hNOs from hPSCs with *PTEN* loss of function mutations [44]. The loss of normal *PTEN* activity

increased proliferation of the organoids' neural progenitor cells, including that of *HOPX*⁺ outer radial glial cells in the sub-ventricular zone. Stresses generated by the resulting rapid tissue expansion induced folding of the emerging cortical plate [44]. Such genetic approaches for imposing biophysical factors that regulate organoid morphology can be instituted during long-term agitated suspension culture. Further integration of both genetic and previously discussed physical approaches could eventually enable constant control of biophysical factors throughout hNO morphogenesis.

3. Bioengineering symmetry breaking and morphogenetic patterning

Although inceptive hPSC aggregates are initially quite homogenous, the mere formation of a 3-D aggregate creates heterogeneous cellular microenvironments that elicit spontaneous, anisotropic patterns of differentiation. This phenomenon is observed at the earliest stages of normal development. For example, cells on the surface of the 32-cell morula differentiate into trophectoderm while cells on the morula's interior experience a different microenvironment and differentiate into an inner cell mass fate [45]. In neural organoid culture, this phenomenon induces spontaneous symmetry breaking and body axis-like patterning as powerfully exemplified by rostro-caudal, dorso-ventral, and medio-lateral tissue organizations observed in mESC-derived cerebral [46] and spinal organoids [47] and gastruloids [48]. Clearly organoids possess the potential for extensive morphogenetic patterning to generate tissues containing biomimetic body axis-like structure. The challenge is how to reproducibly engineer such morphogenetic patterning to increase the biomimicry of hNOs while maintaining consistency in each organoid's anatomy.

Within the developing CNS, morphogens emanating from signaling/organizing centers' modulate local cellular differentiation to impart axial patterning. In 2013, Lancaster et al. demonstrated that a diverse spectrum of brain tissues could develop within hPSC-derived cerebral organoids (hCOs) [22]. This prompted the realization that such signaling centers could spontaneously arise within hNOs. In 2017,

Renner et al. verified the presence of such signaling centers as well as coordinated axial patterning through a detailed analysis of hCO forebrain tissues [30]. In vivo, morphogenesis of the rostral telencephalon is orchestrated by fibroblast growth factors (FGFs) secreted from the rostral anterior neural ridge, Wnts and BMPs secreted from the dorso-medial cortical hem, and Wnt inhibitors, EGF, and other signaling molecules secreted from the antihem (AH; Fig. 2A) [49–52]. The discrete spatial presence of these morphogenetic fields instructs dorso-ventral patterning of pallial and subpallial telencephalic domains. In their experiments, Renner et al. generated 53 hCOs with FGF-2 media supplementation during the initial embryoid body culture stage. They observed that 11 of the 53 hCOs generated both PSPB-like and cortical hem-like signaling centers. Specifically, the expression of Wnt-2b and BMP-6 overlapped with the presence of cortical hem tissues, and proper spatial abutment of choroid plexus-hem-dorsal pallium and lateral pallium-subpallial ganglionic eminence tissues was observed across serial cryosections [30,53]. This suggests that morphogen molecules are present alongside signal center-like structures within hCOs.

Discrete signaling centers also form within hPSC-derived spinal organoids (hSOs) [8,54]. In spinal cord development, rostro-caudal patterning of spinal neuroepithelial cell fate requires FGF, Wnt, and retinoic acid signaling [55]. Subsequently, dorso-ventral patterning of the spinal neuroepithelium is initiated by BMPs and Wnts secreted from the dorsal ectodermal roof plate and sonic hedgehog (Shh) secreted from the ventral mesodermal notochord (Fig. 2B) [56,57]. In an analogous manner, caudalization of hPSCs to spinal neuroepithelial cell fates also requires FGF, Wnt, and retinoic acid signaling [58,59]. In hSO culture, Ogura et al. demonstrated that roof-plate-like signaling centers can spontaneously emerge and secrete BMP4 that activates BMP and Wnt signaling within adjacent tissue regions (Fig. 2C) [8]. This induced axial patterning of the remaining dorsal progenitor domains within the hSO's neuroepithelium. In a similar hSO approach, Duval et al. observed that application of exogenous BMP4 has the strongest dorsalizing effects at early exposure times suggesting the existence of an ideal temporal window for effective morphogenetic patterning [54]. Conversely, development of a ventral signaling center within hSO's is not spontaneous, but requires media supplementation with a Shh small molecule agonists [8]. At specific levels of Shh activation, Ogura et al.

demonstrated that floor plate-like signaling centers, which develop within the embryo's spinal neuroepithelium immediately adjacent to the notochord and secrete Shh, were observed alongside other ventral progenitor domains in hSOs (Fig. 2C). Moreover, the floor plate and ventral progenitor domains were spatially organized in a manner consistent with morphogenetic patterning of the ventral spinal cord [8]. It remains to be demonstrated whether a complete dorso-ventral patterning axis can be established within a single hSO. This will likely require further integration of bioengineering methods.

In an effort to move from spontaneous to more instructed morphogenetic patterning of hCOs, Cederquist et al. used TALEN-mediated genome editing to engineer a doxycycline inducible Shh-expressing hPSC line (iShh) that could serve as a prechordal plate-like signaling center surrogate (Fig. 2A and C) [60]. Using round bottom microwells, a small iShh cell spheroid was formed first before subsequent aggregation as part of a larger hPSC spheroid. Upon further culture and doxycycline exposure, the iShh cells secreted Shh and acted as a signaling center to impart both dorso-ventral and rostro-caudal patterning of telencephalic and diencephalic forebrain tissues within the hCO and in a biomimetic distance dependent manner. This demonstration provides proof-of-concept that synthetic signaling centers can orchestrate hNO symmetry breaking and morphogenetic patterning. However, the use of cellular surrogates may not be ideal since their uncontrolled migration throughout the developing organoid confounds standardization of the morphogenetic process [60].

The previously discussed studies demonstrate the feasibility of standardizing long-range morphogenetic patterning of axial anatomic structure within hNOs by either inducing cellular signaling centers, i.e. via biochemical or optogenetic stimuli, or using artificial surrogates. However, reproducibility of signaling center formation and standardization of the spatial orientation and extent of axial pattern remains inconsistent from organoid-to-organoid. A major limitation to achieving more consistency is the requisite use of agitated suspension culture, which ensures sufficient interstitial nutrient and oxygen diffusion to sustain protracted organoid growth and maturation [8,19,39,60]. If this limitation can be overcome, then microfluidic culture platforms can be used to reproducibly instruct hNO symmetry breaking and morphogenetic patterning. For example, Manfrin et al. recently developed a




Fig. 2. Signaling centers and assembloids. (A) The developing vertebrate forebrain is patterned by multiple signaling centers including the anterior neural ridge (ANR), the cortical hem (CH), the prechordal plate (PCP), and the antihem (AH, at dorsal-ventral telencephalon boundary). (B) The developing spinal cord is patterned by multiple signaling centers including the roof plate (RP), the floor plate (FP), and paraxial mesoderm (PM). It contains both dorsal and ventral progenitors. (C) Human neural organoids (hNOs) can be bioengineered as assembloids that contain inducible signaling centers (e.g. inducible Shh (iShh) domains). Moreover, interregional phenomena like unidirectional migration of cortical interneuron (cIN)-like cells or axonal extension and preferential targeting may also be observed. In spinal cord organoids, exposure to select growth factors induces spontaneous emergence of both dorsal and ventral signaling centers that morphogenetically pattern adjacent organoid regions.

polydimethylsiloxane (PDMS)-based microfluidic device that uses simple Fickian diffusion to create stationary morphogen gradients across a micropatterned, geometrically restricted hPSC colony [61]. By applying counteracting gradients of BMP4 and Noggin—a BMP4 antagonist, they demonstrated controlled induction of differential germ layer fate acquisition across the colonies and in a direction parallel to the gradients. Integration of analogous microfluidic devices with hNO derivation protocols will be critical for scalable manufacture of tissues with reproducible and biomimetic anatomical structure.

4. Bioengineering interactions between organoid tissues of different CNS regions

Assembloids of intra- and inter-cephalic forebrain organoids have been developed as rudimentary models of inter-regional cellular migration and axonal targeting followed by synaptogenesis (Fig. 2C) [5]. The first assembloid studies characterized the fusion of dorsal and ventral telencephalic organoids [27,62,63]. Fusion was initiated by culturing prepatterned EBs side-by-side in a microcentrifuge tube/microwell [27,63] or within a globule of Matrigel [62]. Within these intra-telencephalic assembloids, biomimetic emergent properties were observed such as unidirectional migration of nascent cortical interneurons from the ventral towards the dorsal organoid region. As a disease modeling exemplar, the assembloid's ventral portion was derived from hiPSCs bearing a Timothy Syndrome (TS) gain-of-function mutation [27]. Distinct from their wild-type counterparts, cortical interneurons generated within mutant ventral tissues displayed abnormal migration patterns, which could be partially corrected using inhibitors of L-type calcium channels.

As an inter-cephalic assembloid example, Xiang et al. devised a patterning strategy to generate human thalamic-like organoids (hThOs) and fused them with human cortical organoids (hCrO) within a Matrigel droplet [5]. The thalamus emerges within the diencephalic vesicle and is an important routing and regulatory structure containing many primary and higher-order nuclei of differing functions. It is the major relay between the cortex and subcortical tissues [64]. Within hThO-CrO assembloids, Xiang et al. observed bundled axonal projections akin to reciprocal corticothalamic and thalamocortical projections. The reciprocal axonal projections preferentially targeted and synapsed with post-mitotic neuronal layers instead of progenitor regions within the assembloids, which is consistent with cortical-thalamic targeting in the developing forebrain [5,65]. Despite obtaining a biomimetic axial organization and rudimentary circuitry, it remains to be examined as to whether assembloid fusion led to maturation of the hThO as evidenced by emergence of higher order thalamic-like nuclei. More generally, a potential limitation of the assembloid approach is the possible exclusion of critical inter-regional tissue structures generated via morphogenetic patterning during normal CNS development.

5. Bioengineering models of human circuit formation and maturation

Understanding the association between neuronal circuit function and neuropathology is a major objective of neuroscience research. Mechanisms underlying neuropsychiatric disorders such as Autism Spectrum Disorder, Epilepsy, Down Syndrome, and Schizophrenia indicate irregularities in developmental processes that govern circuit formation and network activity. These irregularities may cause imbalances in the inhibitory (GABAergic) and excitatory (glutamatergic) signaling within local networks [66–73]. Despite the historical use of animal models to discern the properties of neuronal circuitry, there are significant concerns that some neuropsychiatric phenotypes are not conserved between model organisms and humans [73]. This has motivated efforts to develop hNOs as models of circuit formation and network activity.

Extensive scRNA-sequencing has documented that the repertoire of fetal neuronal and glial subtypes required for biomimetic neuronal

circuit formation exist within both cerebral [74] and cortical organoids [75] after ~6–8 months of culture. Substantial neuronal maturation occurs as evidenced by synapse and dendritic spine formation [74]. Moreover, both glutamatergic and GABAergic neurons contribute to the organoids' patterns of electrical activity, which increase consistently with culture duration as measured by microelectrode arrays (Fig. 3). Analogous to human development, the spatiotemporal patterns of electrical activity evolve from discontinuous, non-correlated events (immature organoids) to periodic, synchronized bursting events (>6 mo. matured organoids). This progression indicates spontaneous formation of circuit networks within hNOs [74–76]. After 8 months of culture, retinal-like tissues within cerebral organoids contain photoreceptor-like cells that can be stimulated to fire action potentials using exogenous 530 nm light [74]. After 10 months of culture, neural network firing patterns within cortical organoids demonstrate attributes of complex, oscillatory behaviors observed in human fetal electroencephalograms (EEGs). For example, human fetal EEGs at 7 months of gestation are characterized by intermittent burst of activity separated by periods of quiescence. Within these bursts, the superimposition of both low (e.g. 0.3–2 Hz delta waves) and rhythmic, high frequency (e.g. 8–25 Hz) activity can be detected [76]. Trujillo et al. documented an analogous superimposition within their cortical organoid's oscillatory network activity (i.e. cross-frequency phase-amplitude coupling) using microelectrode arrays (MEA). However, as discussed by the study's authors, multiple experimental and clinical variables like skull filtering properties, electrode placement, and potential neurological conditions limit direct EEG and MEA signal comparison [75]. Moreover, the non-stereotyped cytoarchitecture of hNOs makes it difficult to correlate the MEA's local field potential measurements with specific, physiologically relevant CNS tissue structures or networks.

Registration of intra-organoid circuits with a physiologically relevant network output will likely require integration of novel organoid culture and optogenetic techniques. For example, Giandomenico et al. combined the previously discussed enCOR method with air-liquid interface cerebral organoid slice culture (ALI-COs) [41,77,78]. This culture method reduced cell death and increased the number of cortical neuron populations as well as short and long-range axonal tract-like projections as compared to whole organoid controls [77]. Also, ALI-COs could be maintained in long-term culture to allow neuronal maturation and spontaneous network formation. Notably, the development of intracortical callosal-like projections that were responsive to endogenous axonal guidance cues and corticofugal-like projection tracts that extend away from the ALI-CO gross structure were observed. Using retrograde tracing, it was proven that CUX2⁺ (superficial layer) neurons accounted for the vast majority of internal callosal-like projections, whereas two-thirds of all exterior projecting neurons were CTIP2⁺ (deep layer) neurons. This distribution of CUX2⁺ and CTIP2⁺ neurons is mimetic of cortical and corticofugal projections *in vivo*, respectively. Moreover, Giandomenico et al. demonstrated that their corticofugal-like projections could synapse with explant mouse spinal cord tissue and generate contractions in mouse paraspinal muscle [77]. ALI-CO slice culture tissue morphology permits direct manipulation of constituents cells and circuits for facile integration of synthetic biology tools, e.g. optogenetics [79] and rabies virus-mediated monosynaptic retrograde tracing [80]. Such integration would help to further decipher and potentially modulate circuit network structure, dynamics, and maturation.

6. Bioengineering hNO vascularization

Poor gas and nutrient diffusion are a persistent issue with suspension hNO culture. It reduces cell survival within the organoids' medulla during long-term culture [19]. Organoid culture within spinner flasks or oscillatory bioreactors improves gas and nutrients diffusion resulting in improved cell viability compared to stationary culture [22,26]. However, agitating bioreactors still do not support sustained organoid




Fig. 3. Circuit Network Formation and Vascularization. Neuronal activity and maturation of circuits within hNOs can be measured using multielectrode arrays (MEAs) to detect the spatial and temporal patterns of neuronal activity. An increase in electrophysiological activity over culture duration and the presence of oscillatory waves indicates network formation and maturation. Human neural organoid (hNO) vascularization can be achieved upon invasion of host vasculature after implantation or using co-differentiated or supplemented endothelial cells *in vitro*.

expansion without formation of a hypoxic medulla [77]. The desire to expand hNO growth in the absence of hypoxia as well as include a critical factor of the neurogenic niche [81] and regulator of CNS physiology and disease [82] has motivated development of approaches to vascularize hNOs [79,83–85].

Bioengineering approaches to achieve hNO vascularization have varied from co-derivation of cerebral organoids and blood vessels [83] to direct organoid implantation into rodents (Fig. 3) [79]. Ham et al. demonstrated that VEGF could be included within hCO derivation protocols to induce co-differentiate of endothelial-like cells (ECs) without inhibiting neural morphogenesis [83]. The ECs formed tubules within the organoid and surrounding Matrigel hydrogel and expressed blood-brain barrier (BBB) tight junction protein markers. Also, further supplementation with Wnt7a induced co-derivation of pericyte-like cells within the hCOs [83]. As an analogous co-derivation approach, Cakir et al. engineered doxycycline-inducible *ETV2* expression into a hESC line [84]. Using an EB composed of 20 % genome edited cells, *ETV2*-induced reprogramming could be used to generate hCOs containing ECs that formed perfusable vascular structures *in vitro* and *in vivo*. As a third approach, Pham et al. re-embedded day 34 hNOs in Matrigel containing iPSC-derived endothelial cells (ECs) [85]. After 20 days of *in vitro* culture and 2 weeks of implantation in an immunodeficient NSG mouse, human EC-based tubular structures penetrated the hNO's core sustaining transplant viability *in vivo* compared to non-EC containing organoids [85]. The fourth approach entailed implantation of 40–50 day hCOs directly into the retrosplenial cortex of NOD-SCID mice [79]. In the absence of hPSC-derived ECs, host vasculature possessing BBB-like features invaded and perfused the organoids supporting long-term viability. Moreover, the formation of vascular structures enhanced neural differentiation as day 50 implanted hNOs exhibited greater numbers of NeuN⁺ cells compared to day 102, stage-matched, non-implanted controls. Interestingly, synaptic connectivity between the vascularized hCO and host brain tissues was observed via electrophysiological analysis and immunostaining [79]. Vascularization of hNOs is clearly feasible and several transplantation approaches can yield functional vascular networks with blood flow [79,84]. However, reproducibility of the vascularization time course, vascular tree structure, and homogeneity of spatial distribution within the organoid would be improved by further incorporation of sacrificial molding [17,42,86].

or 3D printing [87,88] techniques.

7. Conclusion

The hNO experimental platform holds tremendous promise for modeling CNS development, physiology, and disease. However, as in all product development pipelines, standardization of hNO derivation protocols is critical to reproducibly bioengineer CNS morphogenesis *ex vivo* to create tissues with biomimetic cellular composition, cytoarchitecture, and anatomy. As reviewed previously [17], numerous tissue engineering methodologies can be applied to hNO culture to precisely control biochemical and biophysical microenvironmental facets. It is only via such integration that we will be able to decipher the biological rules and implement the necessary spatiotemporal control to fully bioengineer CNS development *ex vivo*. Here, we have provided exemplars of bioengineering approaches to control hNO tissue morphology, axial patterning via morphogens emanating from signaling centers or assembly of regionally distinct tissues, circuit formation and maturation, and vascularization. While these studies advance the field and respond to continued pressure to create ever more complex organoid systems, it is equally important for the practical application of hNO science to synthesize their collective biological findings and use this new knowledge to develop bioreactor systems and protocols capable of standardizing comprehensive hNO manufacture. After all, “organoid biology is developmental biology” [18]. If we can understand the rules of human CNS development, then we should be able to precisely bioengineer its morphogenesis.

Acknowledgments

This work is supported by EPA-G2013-STAR-L1 grant #83573701, NSF CAREER award #1651646, and NIHR33NS082618 and UG3TR003150 grants awarded to RSA. NJF and RSA equally contributed to the manuscript's conceptualization and composition. NI aided in figure preparation and manuscript proof-reading and is supported by NIH F32NS106740.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.semcd.2020.05.025>.

References

- [1] J.A. Thomson, J. Itskovitz-Eldor, S.S. Shapiro, M.A. Waknitz, J.J. Swiergiel, V. S. Marshall, et al., Embryonic stem cell lines derived from human blastocysts, *Science* 282 (1998) 1145–1147.
- [2] M. Eiraku, K. Watanabe, M. Matsuo-Takasaki, M. Kawada, S. Yonemura, M. Matsumura, et al., Self-organized formation of polarized cortical tissues from ESCs and its active manipulation by extrinsic signals, *Cell Stem Cell* 3 (2008) 519–532.
- [3] J. Mariani, M.V. Simonini, D. Palejev, L. Tomasini, G. Coppola, A.M. Szekely, et al., Modeling human cortical development in vitro using induced pluripotent stem cells, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 12770–12775.
- [4] T. Nakano, S. Ando, N. Takata, M. Kawada, K. Muguruma, K. Sekiguchi, et al., Self-formation of optic cups and storable stratified neural retina from human ESCs, *Cell Stem Cell* 10 (2012) 771–785.
- [5] Y. Xiang, Y. Tanaka, B. Cakir, B. Patterson, K.-Y. Kim, P. Sun, et al., hESC-derived thalamic organoids form reciprocal projections when fused with cortical organoids, *Cell Stem Cell* 24 (2019) 487–497.
- [6] J. Jo, Y. Xiao, A.X. Sun, E. Cukuroglu, H.-D. Tran, J. Göke, et al., Midbrain-like organoids from human pluripotent stem cells contain functional dopaminergic and neuromelanin-producing neurons, *Cell Stem Cell* 19 (2016) 248–257.
- [7] K. Muguruma, A. Nishiyama, H. Kawakami, K. Hashimoto, Y. Sasai, Self-organization of polarized cerebellar tissue in 3D culture of human pluripotent stem cells, *Cell Rep.* 10 (2015) 537–550.
- [8] T. Ogura, H. Sakaguchi, S. Miyamoto, J. Takahashi, Three-dimensional induction of dorsal, intermediate and ventral spinal cord tissues from human pluripotent stem cells, *Development* 145 (2018) dev162214.
- [9] C.J. Herrick, The morphology of the forebrain in amphibia and reptilia, *J. Comp. Neurol.* 20 (1910) 413–547.
- [10] H.J. Orr, Contribution to the embryology of the lizard, *J. Morphol.* 1 (1887) 331–372.
- [11] L. Puelles, M. Harrison, G. Paxinos, C. Watson, A developmental ontology for the mammalian brain based on the prosomeric model, *Trends Neurosci.* 36 (2013) 570–578.
- [12] S. Herculano-Houzel, Not all brains are made the same: new views on brain scaling in evolution, *Brain Behav. Evol.* 78 (2011) 22–36.
- [13] D.V. Hansen, J.H. Lui, P.R.L. Parker, A.R. Kriegstein, Neurogenic radial glia in the outer subventricular zone of human neocortex, *Nature* 464 (2010) 554–561.
- [14] J.H. Lui, D.V. Hansen, A.R. Kriegstein, Development and evolution of the human neocortex, *Cell* 146 (2011) 18–36.
- [15] A. Kriegstein, S. Noctor, V. Martínez-Cerdeño, Patterns of neural stem and progenitor cell division may underlie evolutionary cortical expansion, *Nat. Rev. Neurosci.* 7 (2006) 883–890.
- [16] A.A. Pollen, A. Bhaduri, M.G. Andrews, T.J. Nowakowski, O.S. Meyerson, M. A. Mostajo-Radji, et al., Establishing cerebral organoids as models of human-specific brain evolution, *Cell* 176 (2019) 743–756.
- [17] C.R. Martí-Figueroa, R.S. Ashton, The case for applying tissue engineering methodologies to instruct human organoid morphogenesis, *Acta Biomater.* 54 (2017) 35–44.
- [18] M. Huch, J.A. Knoblich, M.P. Lutolf, A. Martínez Arias, The hope and the hype of organoid research, *Development* 144 (2017) 938–941.
- [19] M.A. Lancaster, J.A. Knoblich, Generation of cerebral organoids from human pluripotent stem cells, *Nat. Protoc.* 9 (2014) 2329–2340.
- [20] E.L. Elson, G.M. Genin, Tissue constructs: platforms for basic research and drug discovery, *Interface Focus* 6 (2016) 20150095.
- [21] D. Pamies, A human brain microphysiological system derived from induced pluripotent stem cells to study neurological diseases and toxicity, *Altex-Altern. Anim. Ex.* 34 (2017) 362–376.
- [22] M.A. Lancaster, M. Renner, C.-A. Martin, D. Wenzel, L.S. Bicknell, M.E. Hurles, et al., Cerebral organoids model human brain development and microcephaly, *Nature* 501 (2013) 373–379.
- [23] B.A. Lindborg, J.H. Brekke, A.L. Vegoe, C.B. Ulrich, K.T. Haider, S. Subramaniam, et al., Rapid induction of cerebral organoids from human induced pluripotent stem cells using a chemically defined hydrogel and defined cell culture medium: cerebral organoid induction using defined reagents, *Stem Cell Transl. Med.* 5 (2016) 970–979.
- [24] T. Kadoshima, H. Sakaguchi, T. Nakano, M. Soen, S. Ando, M. Eiraku, et al., Self-organization of axial polarity, inside-out layer pattern, and species-specific progenitor dynamics in human ES cell-derived neocortex, *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 20284–20289.
- [25] H. Sakaguchi, T. Kadoshima, M. Soen, N. Narii, Y. Ishida, M. Ohgushi, et al., Generation of functional hippocampal neurons from self-organizing human embryonic stem cell-derived dorsomedial telencephalic tissue, *Nat. Commun.* 6 (2015) 8896.
- [26] X. Qian, H.N. Nguyen, M.M. Song, C. Hadiono, S.C. Ogden, C. Hammack, et al., Brain-region-specific organoids using mini-bioreactors for modeling ZIKV exposure, *Cell* 165 (2016) 1238–1254.
- [27] F. Birey, J. Andersen, C.D. Makinson, S. Islam, W. Wei, N. Huber, et al., Assembly of functionally integrated human forebrain spheroids, *Nature* 545 (2017) 54–59.
- [28] I. Kelava, M.A. Lancaster, Dishing out mini-brains: current progress and future prospects in brain organoid research, *Dev. Biol.* 420 (2016) 199–209.
- [29] J.A. Brassard, M.P. Lutolf, Engineering stem cell self-organization to build better organoids, *Cell Stem Cell* 24 (2019) 860–876.
- [30] M. Renner, M.A. Lancaster, S. Bian, H. Choi, T. Ku, A. Peer, et al., Self-organized developmental patterning and differentiation in cerebral organoids, *Embo. J.* 36 (2017) 1316–1329.
- [31] S.P. Paşa, The rise of three-dimensional human brain cultures, *Nature* 553 (2018) 437–445.
- [32] M. Mrksich, L.E. Dike, J. Tien, D.E. Ingber, G.M. Whitesides, Using microcontact printing to pattern the attachment of mammalian cells to self-assembled monolayers of alkanethiolates on transparent films of gold and silver, *Exp. Cell Res.* 235 (1997) 305–313.
- [33] G.T. Knight, J. Sha, R.S. Ashton, Micropatterned, clickable culture substrates enable *in situ* spatiotemporal control of human PSC-derived neural tissue morphology, *Chem. Commun.* 51 (2015) 5238–5241.
- [34] G.T. Knight, B.F. Lundin, N. Iyer, L.M. Ashton, W.A. Sethares, R.M. Willett, et al., Engineering induction of singular neural rosette emergence within hPSC-derived tissues, *ELife* 7 (2018) e37549.
- [35] T. Haremakki, J.J. Metzger, T. Rito, M.Z. Ozair, F. Etoc, A.H. Brivanlou, Self-organizing neuruloids model developmental aspects of Huntington's disease in the ectodermal compartment, *Nat. Biotechnol.* 37 (2019) 1198–1208.
- [36] Y. Zhu, L. Wang, H. Yu, F. Yin, Y. Wang, H. Liu, et al., *In situ* generation of human brain organoids on a micropillar array, *Lab Chip* 17 (2017) 2941–2950.
- [37] S.-J. Yoon, L.S. Elahi, A.M. Paşa, R.M. Marton, A. Gordon, O. Revah, et al., Reliability of human cortical organoid generation, *Nat. Methods* 16 (2019) 75–78.
- [38] S. Velasco, A.J. Kedaigle, S.K. Simmons, A. Nash, M. Rocha, G. Quadrato, et al., Individual brain organoids reproducibly form cell diversity of the human cerebral cortex, *Nature* 570 (2019) 523–527.
- [39] A.M. Paşa, S.A. Sloan, L.E. Clarke, Y. Tian, C.D. Makinson, N. Huber, et al., Functional cortical neurons and astrocytes from human pluripotent stem cells in 3D culture, *Nat. Methods* 12 (2015) 671–678.
- [40] P. Rakic, Evolution of the neocortex: a perspective from developmental biology, *Nat. Rev. Neurosci.* 10 (2009) 724–735.
- [41] M.A. Lancaster, N.S. Corsini, S. Wolfinger, E.H. Gustafson, A.W. Phillips, T. R. Burkard, et al., Guided self-organization and cortical plate formation in human brain organoids, *Nat. Biotechnol.* 35 (2017) 659–666.
- [42] J.D. McNulty, C. Martí-Figueroa, F. Seipel, J.Z. Plantz, T. Ellingham, L.J. L. Duddleston, et al., Micro-injection molded, poly(vinyl alcohol)-calcium salt templates for precise customization of 3D hydrogel internal architecture, *Acta Biomater.* 95 (2019) 258–268.
- [43] E. Karzbrun, A. Kshirsagar, S.R. Cohen, J.H. Hanna, O. Reiner, Human brain organoids on a chip reveal the physics of folding, *Nat. Phys.* 14 (2018) 515–522.
- [44] Y. Li, J. Muffat, A. Omer, I. Bosch, M.A. Lancaster, M. Sur, et al., Induction of expansion and folding in human cerebral organoids, *Cell Stem Cell.* 20 (2016) 385–396.
- [45] G. Guo, M. Huss, G.Q. Tong, C. Wang, L. Sun, Li, N.D. Clarke, et al., Resolution of cell fate decisions revealed by single-cell gene expression analysis from zygote to blastocyst, *Dev. Cell* 18 (2010) 675–685.
- [46] N. Takata, E. Sakakura, M. Eiraku, T. Kasukawa, Y. Sasai, Self-patterning of rostral-caudal neuroectoderm requires dual role of fgf signaling for localized wnt antagonism, *Nat. Commun.* 8 (2017) 1339.
- [47] A. Meinhardt, D. Eberle, A. Tazaki, A. Ranga, M. Niesche, M. Wilsch-Bräuninger, et al., 3D reconstitution of the patterned neural tube from embryonic stem cells, *Stem Cell Rep.* 3 (2014) 987–999.
- [48] L. Beccari, N. Moris, M. Girgin, D.A. Turner, P. Baillie-Johnson, A.C. Cossy, et al., Multi-axial self-organization properties of mouse embryonic stem cells into gastruloids, *Nature* 562 (2018) 272–276.
- [49] E.A. Grove, S. Tole, J. Limon, L. Yip, C.W. Ragsdale, The hem of the embryonic cerebral cortex is defined by the expression of multiple wnt genes and is compromised in Gli3-deficient mice, *Development* 125 (1998) 2315–2325.
- [50] T. Fukuchi-Shimogori, E.A. Grove, Neocortex patterning by the secreted signaling molecule FGF8, *Science* 294 (2001) 1071–1074.
- [51] J.M. Hébert, Y. Mishina, S.K. McConnell, BMP signaling is required locally to pattern the dorsal telencephalic midline, *Neuron* 35 (2002) 1029–1041.
- [52] S. Assimacopoulos, E.A. Grove, C.W. Ragsdale, Identification of a Pax6-dependent epidermal growth factor family signaling source at the lateral edge of the embryonic cerebral cortex, *J. Neurosci.* 23 (2003) 6399–6403.
- [53] V.V. Chizhikov, A.G. Lindgren, Y. Mishima, R.W. Roberts, K.A. Aldinger, G. R. Miesegaes, et al., Lmx1a regulates fate and location of cells originating from the cerebellar rhombic lip and telencephalic cortical hem, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 10725–10730.
- [54] N. Duval, C. Vaslin, T. Barata, Y. Frarma, V. Contremoulin, X. Baudin, et al., BMP4 patterns smad activity and generates stereotyped cell fate organisation in spinal organoids, *Development* (2019) dev.175430.
- [55] I. Olivera-Martínez, K.G. Storey, Wnt signals provide a timing mechanism for the FGF-retinoid differentiation switch during vertebrate body axis extension, *Development* 134 (2007) 2125–2135.
- [56] J. Briscoe, J. Ericson, The specification of neuronal identity by graded sonic hedgehog signalling, *Semin. Cell Dev. Biol.* 10 (1999) 353–362.
- [57] D. Zechner, T. Müller, H. Wende, I. Walther, M.M. Taketo, E.B. Crenshaw 3rd, et al., Bmp and Wnt/beta-catenin signals control expression of the transcription factor Olig3 and the specification of spinal cord neurons, *Dev. Biol.* 303 (2007) 181–190.
- [58] M. Gouti, A. Tsakiridis, F.J. Wymeersch, Y. Huang, J. Kleinjung, V. Wilson, et al., *In vitro* generation of neuromesodermal progenitors reveals distinct roles for wnt

signalling in the specification of spinal cord and paraxial mesoderm identity, *PLoS Biol.* 12 (2014) e1001937.

[59] E.S. Lippmann, C.E. Williams, D.A. Ruhl, M.C. Estevez-Silva, E.R. Chapman, J. J. Coon, et al., Deterministic HOX patterning in human pluripotent stem cell-derived neuroectoderm, *Stem Cell Rep.* 4 (2015) 632–644.

[60] G.Y. Cederquist, J.J. Asciola, J. Tchieu, R.M. Walsh, D. Cornacchia, M.D. Resh, et al., Specification of positional identity in forebrain organoids, *Nat. Biotechnol.* 37 (2019) 436–444.

[61] A. Manfrin, Y. Tabata, E.R. Paquet, A.R. Vuaridel, F.R. Rivest, F. Naeff, et al., Engineered signaling centers for the spatially controlled patterning of human pluripotent stem cells, *Nat. Methods* 16 (2019) 640–648.

[62] J.A. Bagley, D. Reumann, S. Bian, J. Levi-Strauss, J.A. Knoblich, Fused cerebral organoids model interactions between brain regions, *Nat. Methods* 14 (2017) 743–751.

[63] Y. Xiang, Y. Tanaka, B. Patterson, Y.J. Kang, G. Govindaiah, N. Roselaar, et al., Fusion of regionally specified hPSC-derived organoids models human brain development and interneuron migration, *Cell Stem Cell* 21 (3) (2017) 383–398.

[64] R.W. Guillery, S.M. Sherman, Thalamic relay functions and their role in corticocortical communication: generalizations from the visual system, *Neuron* 33 (2002) 163–175.

[65] G. López-Bendito, Z. Molnár, Thalamocortical development: how are we going to get there? *Nat. Rev. Neurosci.* 4 (2003) 276–289.

[66] K.M. Spencer, P.G. Nestor, M.A. Niznikiewicz, D.F. Salisbury, M.E. Shenton, R. W. McCarley, Abnormal neural synchrony in schizophrenia, *J. Neurosci.* 23 (2003) 7407–7411.

[67] D.A. Lewis, T. Hashimoto, D.W. Volk, Cortical inhibitory neurons and schizophrenia, *Nat. Rev. Neurosci.* 6 (2005) 312–324.

[68] M. Bartos, I. Vida, P. Jonas, Synaptic mechanisms of synchronized gamma oscillations in inhibitory interneuron networks, *Nat. Rev. Neurosci.* 8 (2007) 45–56.

[69] H.-Q. Huo, Z.-Y. Qu, F. Yuan, L. Ma, L. Yao, M. Xu, et al., Modeling down syndrome with patient iPSCs reveals cellular and migration deficits of GABAergic neurons, *Stem Cell Rep.* 10 (2018) 1251–1266.

[70] A. Brooks-Kayal, Epilepsy and autism spectrum disorders: are there common developmental mechanisms? *Brain Dev-JPN* 32 (2010) 731–738.

[71] B. Zikopoulos, H. Barbas, Altered neural connectivity in excitatory and inhibitory cortical circuits in autism, *Front. Hum. Neurosci.* 7 (2013) 609.

[72] J. DeFelipe, Chandelier cells and epilepsy, *Brain* 122 (1999) 1807–1822.

[73] E.J. Nestler, S.E. Hyman, Animal models of neuropsychiatric disorders, *Nat. Neurosci.* 13 (2010) 1161–1169.

[74] G. Quadrato, T. Nguyen, E.Z. Macosko, J.L. Sherwood, S. Min Yang, D.R. Berger, et al., Cell diversity and network dynamics in photosensitive human brain organoids, *Nature* 545 (2017) 48–53.

[75] C.A. Trujillo, R. Gao, P.D. Negräes, J. Gu, J. Buchanan, S. Preissl, et al., Complex oscillatory waves emerging from cortical organoids model early human brain network development, *Cell Stem Cell* 25 (2019) 558–569.

[76] R. Khazipov, H.J. Luhmann, Early patterns of electrical activity in the developing cerebral cortex of humans and rodents, *Trends Neurosci.* 29 (2006) 414–418.

[77] S.L. Giandomenico, S.B. Mierau, G.M. Gibbons, L.M.D. Wenger, L. Masullo, T. Sit, et al., Cerebral organoids at the air–liquid interface generate diverse nerve tracts with functional output, *Nat. Neurosci.* 22 (2019) 669–679.

[78] R.A.M. Daza, C. Englund, R.F. Hevner, Organotypic slice culture of embryonic brain tissue, *CSH Protoc.* (2007) i4914, <https://doi.org/10.1101/pdb.prot4914>.

[79] A.A. Mansour, J.T. Gonçalves, C.W. Bloyd, H. Li, S. Fernandes, D. Quang, et al., An *in vivo* model of functional and vascularized human brain organoids, *Nat. Biotechnol.* 36 (2018) 432–441.

[80] J.H. Marshel, T. Mori, K.J. Nielsen, E.M. Callaway, Targeting single neuronal networks for gene expression and cell labeling *in vivo*, *Neuron* 67 (2010) 562–574.

[81] T.D. Palmer, A.R. Willhoite, F.H. Gage, Vascular niche for adult hippocampal neurogenesis, *J. Comp. Neurol.* 425 (2000) 479–494.

[82] M.D. Sweeney, K. Kisler, A. Montagne, A.W. Toga, B.V. Zlokovic, The role of brain vasculature in neurodegenerative disorders, *Nat. Neurosci.* 21 (2018) 1318–1331.

[83] O. Ham, Y.B. Jin, J. Kim, M.-O. Lee, Blood vessel formation in cerebral organoids formed from human embryonic stem cells, *Biochem. Biophys. Res. Co. Epub.* (2019), <https://doi.org/10.1016/j.bbrc.2019.10.079>.

[84] B. Cakir, Y. Xiang, Y. Tanaka, M.H. Kural, P. Maxime, Y.J. Kang, et al., Engineering of human brain organoids with a functional vascular-like system, *Nat. Methods* 16 (2019) 1169–1175.

[85] M.T. Pham, K.M. Pollock, M.D. Rose, W.A. Cary, H.R. Stewart, P. Zhou, et al., Generation of human vascularized brain organoids, *Neuroreport* 29 (2018) 588–593.

[86] J.S. Miller, K.R. Stevens, M.T. Yang, B.M. Baker, D.H. Nguyen, D.M. Cohen, et al., Rapid casting of patterned vascular networks for perfusable engineered three-dimensional tissues, *Nat. Mater.* 11 (2012) 768–774.

[87] T.J. Hinton, Q. Jallerat, R.N. Palchesko, J.H. Park, M.S. Grodzicki, H.J. Shue, et al., Three-dimensional printing of complex biological structures by freeform reversible embedding of suspended hydrogels, *Sci. Adv.* 1 (2015) e1500758.

[88] B. Grigoryan, S.J. Paulsen, D.C. Corbett, D.W. Sazer, C.L. Fortin, A.J. Zaita, et al., Multivascular networks and functional intravascular topologies within biocompatible hydrogels, *Science* 364 (2019) 458–464.