

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch

Journal:	<i>Ecology and Evolution</i>
Manuscript ID	ECE-2025-09-02337.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Robertson, Erica C.N.; Colorado State University Department of Biology Brown, Timothy ; University of California Santa Cruz Department of Ecology and Evolutionary Biology Deitch, Sophie; Colorado State University Department of Biology Bossu, Christen; Colorado State University, Biology Zavaleta, Erika ; University of California Santa Cruz Department of Ecology and Evolutionary Biology Hooten, Mevin B.; The University of Texas at Austin College of Natural Sciences Ruegg, Kristen; University of Colorado, Biology
Category:	Molecular Evolution
Habitat:	Terrestrial
Organism:	Vertebrate
Approach:	Molecular Genetics
Abstract:	Understanding patterns and mechanisms underlying local adaptation is becoming increasingly important for species conservation amid anthropogenically driven environmental change. Alpine systems are experiencing particularly intense pressure from environmental change resulting from increased rates of warming and corresponding loss of snow and ice. We integrate morphological and genetic analyses to identify traits important for local adaptation in one of the highest elevation breeding birds in North America, the Sierra Nevada Gray-crowned Rosy Finch. We performed an in-depth analysis of how traits with known links to thermoregulation in birds such as wing length, bill size, and feather microstructure vary between two populations at sites with contrasting climate and environmental conditions. We identified loci underlying these traits using a genome-wide association study and further examined regions of the genome related to altitude adaptation and cold tolerance using F_{ST} outlier tests. Together, these results indicate that temperature, food availability, and alpine landscape features may impose multifaceted and potentially conflicting selective pressures on morphological traits important to adaptation in alpine birds. Overall, this work represents one of the first in-depth analyses of the genetic basis of adaptation in an alpine specialist songbird.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



SCHOLARONE™
Manuscripts

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch

Running Title: Local adaptation in an alpine finch

Authors: Robertson, Erica C.N.^{1*}, Brown, Timothy M.², Deitch, Sophie¹, Bossu, Christine M.¹, Zavaleta, Erika S.², Hooten, Mevin B.³, Ruegg, Kristen C.¹

Affiliations:

1. Department of Biology, Colorado State University, Fort Collins, CO, USA

*Corresponding author: ericacnrobertson@gmail.com

2. Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, California, USA

3. Department of Statistics and Data Sciences, The University of Texas at Austin, Austin, Texas

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Understanding patterns and mechanisms underlying local adaptation is becoming increasingly important for species conservation amid anthropogenically driven environmental change. Alpine systems are experiencing particularly intense pressure from environmental change resulting from increased rates of warming and corresponding loss of snow and ice. We integrate morphological and genetic analyses to identify traits important for local adaptation in one of the highest elevation breeding birds in North America, the Sierra Nevada Gray-crowned Rosy-Finch. We performed an in-depth analysis of how traits with known links to thermoregulation in birds such as wing length, bill size, and feather microstructure vary between two populations at sites with contrasting climate and environmental conditions. We identified loci underlying these traits using a genome-wide association study and further examined regions of the genome related to altitude adaptation and cold tolerance using F_{ST} outlier tests. Together, these results indicate that temperature, food availability, and alpine landscape features may impose multifaceted and potentially conflicting selective pressures on morphological traits important to adaptation in alpine birds. Overall, this work represents one of the first in-depth analyses of the genetic basis of adaptation in an alpine specialist songbird.

Keywords

local adaptation, conservation genomics, genome-wide association study, F_{ST} outlier test, alpine

22 Introduction

23 Local adaptation occurs when natural selection favors traits that confer fitness
24 advantages in specific habitats (Endler, 1977; Kawecki & Ebert, 2004; Savolainen et al.,
25 2013). A species' adaptive capacity determines the extent to which it will be able to
26 respond to rapid environmental change (Forester et al., 2022, 2023; Lande & Shannon,
27 1996; Meek et al., 2023). As a result, identifying genetic and morphological traits
28 involved in local adaptation is a key component of effective species conservation
29 planning (Hoban et al., 2016; Meek et al., 2023). Historically, assessments of the capacity
30 for local adaptation relied on reciprocal transplant experiments that allowed researchers
31 to link variation in putatively ecologically important traits to differences in fitness across
32 environmental gradients (Blanquart et al., 2013; Clausen et al., 1941; Kawecki & Ebert,
33 2004; Savolainen et al., 2013). However, recent advances in genomic methods have
34 made it possible to detect signatures of selection across the genome and to associate
35 these with environmental variation and ecologically important morphological variation
36 (Capblancq et al., 2020; Hoban et al., 2016; Lotterhos & Whitlock, 2015). We leveraged
37 recent advances in genomic methods to identify morphological and genetic traits
38 important to local adaptation in an understudied alpine bird system.

39 Many montane species undergo upward range shifts with climate change
40 (Mamantov et al., 2021). However, for those species already living near the top of a
41 mountain system, there are constraints that may limit the ability to track their niche, a
42 phenomenon termed the "escalator to extinction" (Freeman et al., 2018; Urban, 2018).
43 Alternatively, local adaptation can impact adaptive responses to the environment and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

allow species to persist despite a restricted capacity for range shifts (Aitken et al., 2008; Capblancq et al., 2020). A first step toward improving our understanding of how high-altitude species will respond to climate change is to identify morphological traits and corresponding regions of the genome involved in adaptation to extreme alpine environments (Blanquart et al., 2013; Pritchard & Di Rienzo, 2010).

Birds living in alpine regions are exposed to extreme environments throughout the year, including variable snow cover, hypoxic conditions, and dramatic temperature changes (Grabherr et al., 2010; Körner, 2003). As a result, traits that contribute directly to thermoregulation are expected to be under strong selective pressure in alpine species. Previous work has shown climate-linked traits often follow ecogeographical rules such as Bergmann’s and Allen’s rule. Allen’s rule posits that animals adapted to colder environments will have shorter appendages than animals adapted to warm climates (Allen, 1877). In birds, this rule predicts that beaks and wings will be longer in warmer environments to facilitate greater heat dissipation across a larger surface area, and shorter in colder environments to facilitate heat conservation (Greenberg et al., 2012; Lewden et al., 2023; Symonds & Tattersall, 2010; Tattersall et al., 2017; Ward et al., 1999; Weeks et al., 2020, 2025). Bergmann’s rule, which states that body size tends to decrease as temperature increases, has also been supported in birds (Ashton, 2002; He et al., 2023). Following these rules, we predicted that birds will have shorter beaks and smaller wings as well as larger overall body sizes in higher, colder environments.

An additional aspect of beak morphology that should be investigated in the context of thermoregulation is nares length. Nares are the nostril of the bird and are located at the top of the beak. Grinnell (1913) documented subspecific divergence in

1
2
3 67 *Leucosticte tephrocotis dawsoni*, including differences in beak length and bill depth from
4
5 68 the nostril. While nare size per se was not the focus, these measurements provide a
6
7 69 morphological basis for interpreting variation in nasal exposure and potential heat
8
9 70 exchange, particularly in the context of ecological differences between alpine sites.
10
11 71 While there has not, to our knowledge, been work directly linking nare size to
12
13 72 thermoregulatory function, this may be an important and understudied aspect of beak
14
15 73 morphology as it relates to thermoregulation in birds. Following the logic for broader
16
17 74 beak morphology, we can expect to see larger nares in the warmer environments to
18
19 75 facilitate the dumping of excess heat.
20
21
22
23

24 76 Although less studied in the context of local adaptation, feather microstructure
25
26 77 may also be important to avian species' adaptation to environmental variation in
27
28 78 temperature. While substantial evidence suggests feather microstructure plays a key
29
30 79 role in determining the thermoregulatory capacity of a bird's plumage (Barve et al., 2021;
31
32 80 Stettenheim, 2000; Stoutjesdijk, 2003; Wolf & Walsberg, 2000), there is some confusion
33
34 81 over how environmental variation may drive feather microstructure differences (D'alba
35
36 82 et al., 2017; Koskenpato et al., 2016; Lei et al., 2002; Pap et al., 2017, 2020). In one of
37
38 83 the few studies that has been conducted at the population level, Koskenpato et al.
39
40 84 (2016) found that tawny owls living in colder environments have significantly denser
41
42 85 plumulaceous contour feathers than those living in warmer environments. This suggests
43
44 86 that feather density may facilitate heat retention. In contrast, however, Pap et al. (2017)
45
46 87 found that European bird species wintering in colder areas had less dense feathers than
47
48 88 those wintering in warmer places. Thus, there is a clear need for studies specifically
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

89 linking environmental variation with population-level differences in feather
90 microstructure.

91 Recent advances in population genomics have enabled the detection of loci
92 involved in local adaptation, even in non-model and wild species (Faria et al., 2014).
93 Among these, F_{ST} outlier tests are commonly used to identify genomic regions that
94 exhibit elevated population differentiation, which may signal divergent selection (Hoban
95 et al., 2016). However, many adaptive traits are likely polygenic, involving small effects
96 at numerous loci (Pritchard & Di Rienzo, 2010; Yeaman, 2015), and such loci will not be
97 picked up by F_{ST} outlier tests. To address this, genome-wide association studies
98 (GWAS) can be used to link more subtle genetic variation to phenotypic variation in
99 ecologically important traits suspected to be under selection (Stinchcombe & Hoekstra,
100 2008). Taken together, combining approaches that detect both population-level
101 divergence and polygenic trait architecture offers a powerful framework for investigating
102 local adaptation.

103 We investigated the potential for local adaptation in the Sierra Nevada Gray-
104 crowned Rosy-Finch (*Leucosticte tephrocotis dawsoni*), one of the highest-breeding
105 songbirds in North America. Of the roughly four groups of rosy-finch found in North
106 America, this subspecies is more localized, found only in California's alpine, and is a
107 short-distance migrant. In comparison, the broader Gray-crowned Rosy-Finch species
108 range extends along the entire western edge of North America from California to
109 Alaska. Despite long-standing interest (Grinnell, 1913, 1917; Twining, 1940), research
110 on this subspecies has been limited due to the difficulty of accessing its extreme nesting
111 sites located on the rocky cliff faces of the Sierra Nevada and White Mountains of

California, more than 3,000 meters above sea level. Additionally, this subspecies may act as a strong model of local adaptation due to the variation in habitat experienced within a restricted range. To help fill this knowledge gap we conducted morphological and genomic analysis of two populations breeding at distinct places in the elevational and thermal range. Piute Pass, located in the Sierra Nevada, is lower in elevation and warmer than the White Mountains (Fig 1A). To identify traits potentially involved in local adaptation, we quantified variation in beak and feather morphology between the two populations. Based on Allen's rule, we predicted that birds in the warmer Piute Pass population would have longer wings and larger beaks. Following Bergmann's rule, we predicted that birds in Piute Pass would also have smaller overall body size. Further, we predicted that birds in the colder White Mountains location would exhibit feather traits associated with increased insulation, specifically, longer barbules and higher node density. Lastly, we used genome-wide association studies (GWAS) to identify genes linked to feather and bill morphology and F_{ST} outlier tests to identify other potentially important sources of genetic variation (Fig 1B-C). Overall, our findings provide new insights into the process of local adaptation in an extreme alpine specialist bird and provide key baseline information on the capacity for adaptation in the face of global environmental change.

Methods

Site Selection

Our study compares Rosy-Finch populations from two sites: one in the Sierra Nevada and one in the White Mountains. Sierra Gray-crowned Rosy-Finches occupy alpine habitats from approximately 2,750-4,000 m across California (Brown et al., in

1
2
3 135 review), and our study sites span a substantial portion of this range. The Piute Pass
4
5 136 location is in the Sierra Nevada at 3,470-3,625m in elevation and has a warmer average
6
7 137 temperature during the breeding season of 8°C (with a range of 3.8-13.0°C). The White
8
9 138 Mountain location, near White Mountain Peak, is higher in elevation at 4,285-4,344m
10
11 139 and colder with a mean average temperature during the breeding season of 6°C (with a
12
13 140 range of 1.5-10.7°C) (extracted from AdaptWest Project, 2022, see supplemental
14
15 141 methods and SupFig1). Because it lies in the Sierra Nevada rain shadow, the White
16
17 142 Mountain site receives roughly one-third the precipitation of areas at comparable
18
19 143 elevations in the Sierras (Rundel et al., 2008). Between the two sites, Piute Pass
20
21 144 receives more precipitation as snow and has a more persistent snowpack than the
22
23 145 White Mountain location. Additionally, aquatic environments such as streams, lakes and
24
25 146 ephemeral pools are more abundant in the Sierra Nevada than the White Mountains
26
27 147 (Rundel et al., 2008). The alpine ecosystems of these mountains can be delineated as
28
29 148 communities occurring above tree line (Rundel & Millar, 2016). Environmental stressors
30
31 149 such as extreme winter temperatures, short growing seasons, high winds, low partial
32
33 150 pressures of O₂, and limited water availability are characteristic of these alpine
34
35 151 environments (Grabherr et al., 2010; Rundel & Millar, 2016). Separating the western
36
37 152 Sierra Nevada Mountains and the eastern White Mountains is the Owens Valley, a
38
39 153 subregion of the Great Basin Desert (Belnap et al., 2016).

46
47
48 154 **Sample Collection**

49 155 A total of 171 Sierra Nevada Gray-crowned Rosy-Finches were captured using
50
51 156 potter traps at White Mountain (n=98) and Piute Pass (n=73) in June and July, the peak
52
53 157 of the species' breeding season (MacDougall-Shackleton et al., 2000). Captures within
54
55
56
57
58
59
60

1
2
3 158 each mountain range occurred across multiple locations within approximately 1 km² at
4
5
6 159 each site (White Mountain or Piute Pass) to maximize sample sizes while maintaining
7
8 160 site-specific environmental characteristics. The sampling thus, in total, represents only
9
10 161 these two sites. Not all morphology was able to be collected for every bird, resulting in
11
12 162 slight variation in the types of data collected for each individual (see details in
13
14 163 supplemental methods, see sample sizes for each trait in Supplementary Table 1).
15
16
17 164 Blood samples were collected from 150 of the captured birds using the brachial wing
18
19 165 vein and stored in Queen's lysis buffer at room temperature (Owen, 2011; Seutin et al.,
20
21 166 1991). At the same time, morphological measurements - tarsus length, wing chord,
22
23 167 beak width, beak length, beak depth, nare length, and mass - were also collected from
24
25 168 154 of the birds using an electronic caliper and 5-10 body feathers were collected from
26
27 169 the breast of all individuals. Additionally, birds were aged and sexed following the
28
29 170 guidelines established in Pyle (2022) (see supplemental methodology). Birds were
30
31 171 banded to allow for identification of the individuals and recognize recaptures, then
32
33 172 released. All handling and banding of birds was done following the guidelines and
34
35 173 protocols of the U.S Geological Survey (USGS) Bird Banding Laboratory (BBL) and
36
37 174 complied with permits and permissions from federal and state agencies. All
38
39
40 175 measurements were taken by two researchers (TB and EZ), with one (TB) collecting
41
42 176 approximately 80% of the data. Because the majority of measurements were taken by a
43
44 177 single observer, and all followed the same standardized protocol, observer identity was
45
46
47
48
49 178 not included as a covariate in subsequent analyses.
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Feather Microstructure Measurements

We measured the microstructure of three feathers for each individual. For each feather, three photos were taken using an Olympus BX51 Microscope: one photo each of an unbroken pennaceous and plumulaceous barb from the center of each region was taken at 4x objective, and one photo of a plumulaceous barbule was taken at 10x objective (Fig 3A). Structures were then measured from the photos using the segmented line tool in ImageJ. For the pennaceous and plumulaceous barbule length, we measured one barbule from each region starting from the distal tip to where the barbule meets the barb. For the pennaceous and plumulaceous barbule density, we drew a 0.5mm line along the middle of a barb, counting the number of barbules along that line. And finally, for plumulaceous node density, we drew a 0.2mm line along the center of a barbule, counting the number of nodes along that line. Each replicate measure for a feather trait was averaged per individual prior to downstream analysis. Approximately 95% of feathers were measured by one researcher (SD) and observer was not included as a covariate in subsequent analyses.

Statistical analysis for local adaptation

Body size and sex can confound morphological comparisons among individual birds (D. C. Adams et al., 2020). To account for these effects in our investigation of Allen's rule, we included sex and tarsus length as predictor variables in our models. We used tarsus length as an index of body size because it correlates strongly with mass but is less affected by short-term fluctuations in condition; this approach has been used in recent large-scale morphological studies (Demery et al., 2021; Weeks et al., 2020; Zimova et al., 2023). As additional validation of tarsus as a proxy size for body size, we generated a PC axis that incorporated multiple allometric traits and found that this was

highly correlated (0.7) with tarsus (see supplement for additional details). We proceeded with tarsus as our proxy for body size in all further analyses. To assess trait variation between sites, we used both univariate and multivariate approaches (see SupTable 2-3 for a summary of trait values). We began by constructing linear models with tarsus length, sex, age, and site code as predictors and used AIC weights to identify the combination of variables that best explained trait variation. Traits for which site code was a significant predictor were considered significantly different between the two populations. We opted not to use a Bonferroni correction for the univariate models as each model had a specific hypothesis associated with it. We used *emmeans* (Lenth, 2025) to calculate average trait values while controlling for predictors other than site code, and these adjusted means were overlaid on the raw data in a violin plot (Fig 2 & 3B). To test our hypotheses derived from Bergmann's rule, we compared both tarsus length and a principal component axis representing overall body size using linear models. Each model included age and sex as covariates to account for potential confounding effects on trait variation. We then assessed whether body size differed significantly between the two populations by including site as a predictor.

We also conducted a multivariate analysis to account for potential correlations among traits in our investigation of Allen's rule. For our multivariate approach we incorporated all body morphology traits (wing chord, nare length, beak depth, beak width, beak length) into a MANOVA model with site, age, sex, and tarsus as predictors. The same analysis was performed with all of the feather traits (pennaceous barbule density and length, plumulaceous barbule density, length, and node density) together.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Sequencing and Genotyping

DNA was extracted from blood samples of 150 individuals using Qiagen’s DNeasy Blood and Tissue Extraction Kit protocol and quantified with the Qubit dsDNA HS Assay Kit. We used a modified version of Illumina’s Nextera Library Preparation protocol to prepare WGS libraries and pooled the libraries by equal mass before sequencing. The resulting libraries were sequenced on 2 lanes of an Illumina NovaSeq 6000 at Novogene Corporation.

To process the raw sequence reads and detect variants, we utilized Snakemake (Molder et al. 2021), a workflow management system that provides efficiency, adaptability, and reproducibility. The pipeline we adapted to our species can be found on Github (<https://github.com/eriqande/mega-non-model-wgs-snakeflow/>). To summarize the workflow, sequence data were trimmed using fastp (S. Chen et al., 2018) to remove adaptor sequences and polyG tails using a sliding window, and then aligned to a Brown-capped Rosy-Finch (*Leucosticte australis*) reference genome (GenBank: GCA_025504685.1) using Burrows-Wheeler Aligner software (H. Li & Durbin, 2009). We marked PCR duplicates using SAMtools (Danecek et al., 2021) and read groups (sample, lane, library) were added using Picard (<http://broadinstitute.github.io/picard>). Individual coverage was estimated using SAMtools, and, given the range of coverage (7.4-19X), we downsampled bam files to 10x using the SAMtools subsample function (Danecek et al., 2021). Individual gvcf files were created using the GATK HaplotypeCaller (Poplin et al., 2018) with high base quality score filters (--min-base-quality-score 33 --minimum-mapping-quality 20) to remove batch effects (Lou & Therikildsen, 2022). We then parallelized the calling of genotypes across 3 million bp

regions across the genome using the GenomicsDBImport and GenotypeGVCFs functions (Auwera & O'Connor, 2020; McKenna et al., 2010). GATK versions above 4.0 call missing data as homozygous reference, which can bias downstream analysis. To avoid this, all loci with a depth of zero were manually marked as missing. To remove systematic errors according to GATK variant quality score recalibration (VQSR) best practices, we hard filtered variants with the following parameters: StrandOddsRatio (>3.0), FisherStrand (>60.0), MappingQuality (>40.0), and Quality by Depth (>2.0). We further filtered for Single Nucleotide Polymorphisms (SNPs), removing indels, filtering for missingness (<80%), allele frequencies (>0.05 and <0.95), depth (>4x), quality (>30.0) using BCFtools (Danecek et al., 2021). After filtering, the resulting high-quality SNPs were passed through BEAGLE 4.1 (Browning & Browning, 2016) to impute missing genotypes.

Population genetic structure analysis

Population structure was assessed using a Principal Component Analysis (PCA) followed by ADMIXTURE (Alexander et al., 2009). SNPs identified from sequence data were pruned for linkage disequilibrium in PLINK (Purcell et al., 2007) using a 50kb window, 10 SNP window step size, and a R^2 threshold of 0.2. Eigenvalues for the PCA were then generated based on the pruned SNPs using PLINK and the first two principal components (PC) were plotted in R. To further confirm a lack of population structure ADMIXTURE was used to test $k=1-3$. The results were plotted using R and the cv values were evaluated.

Genome-Wide Association Study

Two GWAS models were implemented on all traits using GEMMA (Zhou et al., 2013): a Bayesian sparse linear model, BSLMM, and a univariate linear model, ULMM. These GWAS approaches allow us to detect SNPs associated with a given trait while explicitly accounting for population structure and relatedness through the incorporation of random effects whose correlation structure is characterized by a relatedness matrix. Input files were generated with PLINK (Purcell et al., 2007). We implemented a Bayesian sparse linear mixed model (BSLMM) using 500,000 MCMC burn-in iterations that were discarded and 5 million MCMC iterations that were saved. We then filtered for SNPs that had a mean posterior inclusion probability (PIP) above 0.01. A univariate linear mixed model (ULMM) employing the Wald test was also conducted (see SupTable 7 for λ GC values for each model and SupFig6 for PP plots for each trait). The Wald test assesses the significance of the SNP genotypes in a linear regression model and is used to test the null hypothesis that the effect size of a genetic variant on the phenotypic trait is zero (i.e., there is no association between the variant and the trait). SNPs were then filtered based on a p-value threshold of 5×10^{-8} , a threshold widely used in GWAS studies (e.g. Chen et al., 2021).

Genome Wide F_{ST}

To estimate per site F_{ST} , we used OutFLANK (Whitlock & Lotterhos, 2015). OutFLANK identifies outlier F_{ST} SNPs by modeling the distribution of F_{ST} values across loci using a trimmed likelihood approach to exclude loci under strong selection. It then fits a chi-square distribution to the neutral F_{ST} values and identifies SNPs with unusually high or low F_{ST} as potential outliers, suggesting loci under selection. We used a false discovery rate threshold of 0.005 to determine outlier SNPs. We took the 90th quantile

of these F_{ST} values to narrow down the top SNPs for literature review and plotted these on a Manhattan plot. This allowed for isolation of the most divergent loci as well as easier visualization. We then determined the F_{ST} values for the SNPs previously identified in the GWAS analyses based on the values calculated by OutFLANK.

Identification of associated genes, enrichment analysis and literature review

To identify genes associated with significant SNPs from our GWAS and F_{ST} analysis, we implemented BEDTools (Quinlan & Hall, 2010) closest function with a previously created Brown-capped Rosy-Finch gene annotation (Funk et al., 2023) to identify genes close to the SNP positions. The output from this tool was filtered to find only protein coding genes within 25kb of the given SNP. For plotting purposes, we used the NUCmer function in MUMmer4.x (Marçais et al., 2018) to align the Brown-capped Rosy-Finch genome assembly to the Zebra Finch (*Taeniopygia guttata*) genome assembly (GenBank: GCF_003957565.2). We then filtered the delta alignment output of NUCmer to keep only matches that were greater than 400 bps, and used show-coords to display the coordinates. We then used custom R scripts to convert our scaffold and positions to Zebra Finch chromosome positions for plotting SNPs in Manhattan plots.

Enrichment in gene ontology (GO) terms was performed with Panther 19.0 (Thomas et al., 2022). For this analysis, *Gallus gallus* was used as the background gene set. Enrichment thresholds were set to $p < 0.05$ after Bonferroni correction. Gene network analysis was conducted using STRING (Szklarczyk et al., 2023) with zebra finch as the organismal reference for background genes. STRING (Search Tool for the Retrieval of Interacting Genes/Proteins) is a database that predicts and visualizes protein-protein interactions based on known and predicted associations from

1
2
3 316 experiments, literature, and computational methods (Szklaarczyk et al., 2023). For F_{ST}
4
5 317 genes, all genes associated with the OutFLANK outliers were input into enrichment
6
7 318 analyses. Genes resulting from the GWAS analysis were grouped with all beak
8
9
10 319 morphology trait genes run as a single analysis, all feather trait genes run together, and
11
12 320 wing chord genes run in another analysis.
13

14
15 321 We performed a literature search for all of the candidate genes using google
16
17 322 scholar and the search terms gene name followed by, bird, trait specific term (when
18
19 323 applicable), environmental adaptation, local adaptation (e.g., “‘XIAP’ gene beak”, or
20
21 324 “‘DSCAM’ gene environmental adaptation”). For F_{ST} SNPs, only those genes associated
22
23
24 325 with the 90th quantile of SNPs were evaluated in the literature. Genes were grouped by
25
26 326 their key functions or associations as described in the literature.
27
28
29

30 327 **Results**

31
32 328 **Local adaptation of body and feather morphology**

33
34 329 For the univariate approach, AIC model selection indicated five traits for which
35
36 330 site was a significant predictor. For wing chord, the best model included tarsus length,
37
38 331 age, sex, and site. The model showed high explanatory power (adjusted $R^2=0.673$,
39
40 332 $p<0.001$) and showed that birds at the White Mountain location had shorter wing chords
41
42
43 333 compared to Piute Pass (PIPA) ($p=0.003$). The best model for nare length included
44
45 334 tarsus, sex, and site and had lower explanatory power (adjusted $R^2=0.197$) but was still
46
47 335 highly significant ($p<0.001$). This model found that nare length was shorter in the White
48
49
50 336 Mountain population ($p<0.001$). Similarly, beak width had tarsus, age and site as
51
52 337 significant predictors (adjusted $R^2=0.135$, $p<0.001$) with White Mountain having beaks
53
54 338 that are wider than those in Piute Pass ($p<0.001$). Beak depth had tarsus, age and site
55
56
57
58
59
60

339 as the best predictors (adjusted $R^2=0.250$) and found that White Mountain has
 340 significantly deeper beaks than Piute Pass ($p=0.003$). Beak length was not significantly
 341 different between the two locations and was best predicted by sex (adjusted $R^2=0.516$,
 342 $p<0.001$). For feather microstructure, the only trait that had site as a significant predictor
 343 was plumulaceous node density. This model included tarsus and age as well as site but
 344 had low explanatory power (adjusted $R^2=0.076$, $p<0.001$). It showed that plumulaceous
 345 node density was higher in the White Mountain populations ($p<0.001$). The best model
 346 for plumulaceous barbule length based on AIC weights included site as a predictor;
 347 however, the effect size was uncertain, and the effect of site was not statistically
 348 significant (adjusted $R^2=0.142$, $p=0.078$). Pennaceous barbule density had tarsus and
 349 age as the best predictors (adjusted $R^2=0.365$, $p<0.001$), pennaceous barbule length
 350 had sex as the best predictor (adjusted $R^2=0.307$, $p<0.001$) and plumulaceous barbule
 351 density had tarsus and age as the best predictors (adjusted $R^2=0.076$, $p=0.004$). We
 352 found that body size did not vary between sites. In our model using tarsus (adjusted
 353 $R^2=0.165$, $p<0.001$) both age and sex were significant predictors ($p<0.001$ and $p<0.05$),
 354 however site was not significant ($p=0.417$). Our comparison using the PC axis
 355 representing size (adjusted $R^2=0.4639$, $p<0.001$) had similar results with age and sex
 356 as significant and site as not ($p=0.53$).

357 When looking at the results of the multivariate analysis, we see that for body
 358 morphology traits site was a highly significant explanatory variable ($p=1.327e-11$).
 359 Tarsus, age and sex were also highly significant ($p<2.2e-16$). The MANOVA on feather
 360 traits also had site as a significant predictor although the p-value was less significant
 361 ($p=0.036$). Age was also significant ($p=1.5e-14$) but tarsus and sex were not ($p>0.05$).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For details on the models tested, models chosen, and full model results, see
Supplementary Table 4-6.

Evaluation of population structure

Whole-genome sequencing was performed on 150 individuals and 146 were
included in downstream analysis. Four individuals were removed because their
collection location could not be validated. Variant filtering resulted in 7,739,836 SNPs
for subsequent genetic analysis. Linkage pruning removed 5,526,104 SNPs. Visualizing
PLINK PCA results revealed little population structure with most individuals clustering
together (SupFig 2). ADMIXTURE results confirmed that there is no population structure
in these samples because k=1 had the lowest cv error (0.599) compared to k=2 or 3
(0.612 and 0.627).

Genes identified with genome-wide association approach

The univariate linear mixed model identified a total of 64 SNPs for wing chord,
beak length, beak width, pennaceous barbule length, and pennaceous barbule density.
Using a Brown-capped Rosy-Finch gene annotation we identified a total of 26 unique
genes corresponding to these SNPs (Table 1). We filtered out genes that were not fully
characterized (e.g., “LOC100221041”) leaving 14 genes. The Bayesian sparse linear
mixed model identified 266 SNPs for wing chord, nare length, beak length, depth and
width, pennaceous barbule length and density, and plumulaceous barbule length and
node density. We identified 146 genes associated with these SNPs (Table 1), 110 of
which had been fully characterized. For each trait the global view of p-values for all
SNPs and associated genes across both models are represented by Manhattan plots

(Fig 4A-C, SupFig 3-5). Given the sample size ($n = 146$), this GWAS is best viewed as conservative, detecting only loci with relatively strong effects on morphology.

GO term analysis using Panther produced significant results for beak depth and a combined set of all feather traits. For beak depth, the top terms were presynaptic membrane assembly ($p=1.44 \times 10^{-5}$), synaptic membrane adhesion (specifically cell adhesion) ($p=1.01 \times 10^{-5}$) and heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules ($p=1.13 \times 10^{-5}$). For the feather traits (plumulaceous barbule and node density combined with pennaceous barbule length), fructosamine catabolic process was a significant term ($p=1.12 \times 10^{-6}$). Wing chord had no significant GO terms. STRING analysis for beak and feather traits as well as wing chord did not find significantly more interactions than expected.

Genes identified through high Fixation index (F_{ST}) between the two populations

Using OutFLANK 831 loci were identified as having significantly high F_{ST} when comparing the White Mountain and Piute Pass populations, corresponding to 265 genes (Fig 4D). No GWAS SNPs were flagged as having significant F_{ST} values. The minimum F_{ST} to be considered an outlier was 0.17513. The mean F_{ST} of SNPs not considered outliers was 0.0045. Using a 90th quantile we determined $F_{ST}=0.235$ to be the threshold, narrowing down the SNPs to 105 loci. For the smaller set of loci, bedtools identified 61 related genes (Table 2). We ran STRING and GO term enrichments on both the larger and smaller set of genes. The smaller set did not have a significant number of connections over all or any significantly enriched GO terms. The larger set of 264 genes, however, had an overall significant number of connections ($p<0.05$) and had multiple significant annotated keywords (sourced from UniProt). The most significant

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

was the keyword repeat (FDR<0.001), the next was ubl conjugation pathways (FDR<0.05).

Discussion

Understanding the genetic and morphological basis of local adaptation is critical for predicting how species respond to environmental change. Alpine species face unique conservation challenges due to their extreme environments and thus are a high priority for studies of adaptive capacity. In this study, we investigated morphological and genetic differences between two populations of an alpine specialist songbird located in distinct parts of its environmental range. Consistent with our predictions, we found wings were shorter and feathers were denser in the colder environment. However, in contrast to our prediction, beak depth and width were larger in the colder environment, suggesting other factors like diet may place contrasting selective pressures on this ecologically important trait. We also identified numerous genes with known links to limb, facial, and feather development that potentially underlie variation in wing, bill and feather microstructure. Further F_{ST} outlier analyses revealed numerous significant differences between cold and warm populations at SNPs located within genes with clear links to altitudinal adaptation. Together, our results provide important insights into the morphological traits and associated genomic regions associated with microgeographic variation in a high-altitude songbird species.

Trait differences between warm and cold habitats

Ecogeographic rules, such as Allen’s and Bergmann’s rules, are useful for predicting broad geographic trends in morphology in response to environmental conditions, but observed patterns do not always align with expectations. In our study,

we found that wing chord was smaller in the colder environment, suggesting that this morphological trait may play a role in thermoregulation, consistent with Allen's rule. Heat dissipation from the wings occurs across the vascularized brachial regions, primarily during flight when these areas are exposed to air flow (Lewden et al., 2023; Ward et al., 1999). Wing bone length reflects the extent of vascularized areas on the wings and has been shown to be longer in warmer environments across a broad sample of passerine species (Weeks et al., 2025). However, studies of other alpine bird species have found the opposite pattern: that birds from high-elevation sites have longer wings than those from low-elevation sites (Bears et al., 2008; Ceresa et al., 2024). In these cases, researchers have suggested that the demand for more energetically efficient flight at high elevations may outweigh thermoregulatory selective pressures, demonstrating the often contrasting selective pressures on ecologically important traits. Although outside of the scope of the current work, additional wing measurements could be collected that would allow for a more thorough investigation into the components of flight morphology. It is possible that migratory behavior is influencing wing length as previous work has shown that both within (Egbert & Belthoff, 2003; Grilli et al., 2017) and between (Lockwood et al., 1998) species those who migrate further have longer wings. To date, no formal study has been conducted on the migratory patterns of the Sierra Nevada Gray-crowned subspecies and so future work could investigate the links between wing morphology and movement patterns. Overall, our morphological analyses support the idea that thermoregulatory demands may play an important role in shaping wing morphology in the Sierra Nevada Gray-crowned Rosy-Finch, potentially

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

outweighing the benefits of flight efficiency in this high-elevation, cold-adapted species but not excluding the potentially parallel pressure of migratory distance.

Contrary to our predictions based on Allen’s rule, beak depth and width were larger in the colder environment. Although temperature is often a key environmental variable exerting selective pressure on alpine taxa, variation in beak morphology between our populations may instead reflect divergent selective pressures associated with foraging ecology, a well-documented driver of beak divergence in birds (Grant et al., 1976; Lamichhaney et al., 2018). In our study system, the Piute Pass population likely has access to more aquatic insects due to the higher availability of aquatic habitats proximate to Rosy-Finch breeding sites in the Sierra Nevada (Epanchin et al., 2010; Rundel & Millar, 2016). For example, Epanchin et al. (2010) found that, when abundant, mayflies can comprise up to 38% of the Sierra Nevada Gray-Crowned Rosy-Finch diet. In contrast, the White Mountain population may rely more heavily on insects deposited in snowfields, a behavior that could favor a more robust, conical bill for extracting frozen prey from the snow surface. Work is currently underway to investigate the dietary differences between these two populations of Rosy-Finch (Tim Brown, personal correspondence), and future work should explore the selective pressures experienced by this species. The other aspect of beak morphology assessed, nares length, was significantly larger in the warmer environment. If nares is playing a role in thermoregulation, these results would be consistent with larger nares allowing for more heat to be lost. Additionally, smaller nares in the colder environment would potentially prevent snow from entering the nasal passages while foraging on snow. Overall, our results on beak morphology emphasize the complexity of identifying drivers of local

adaptation, especially in alpine systems where extreme environments, food scarcity, and variation in migration strategies can create conflicting selection pressures.

Our comparison of body size did not have significant results, suggesting that Bergmann's rule is not contributing to morphological differences in this subspecies. He et al (2023) found that there was significant variation in both strength and direction of correlation between latitude and body size within and between taxa. With our results, this could imply that Sierra Nevada Gray-crowned Rosy-Finch are an exception to the rule. In another meta-analysis, it was found that sedentary bird species were more likely to conform to Bergmann's rule than migratory ones and suggest that winter temperatures may exert a higher pressure than breeding temperatures (Meiri & Dayan, 2003). The Sierra Nevada Gray-crowned Rosy-Finch is a short-distance, altitudinal migrant. It is possible that both White Mountain and Piute Pass populations move to similar areas or climates during the winter season, thus decreasing the divergent selection pressure that would drive differences in Bergmann's rule.

Of our feather comparisons, we identified significant differences between cold and warm sites for only one trait: plumulaceous node density. In line with our predictions, plumulaceous node density was higher in the colder, high elevation White Mountain population. Higher node density is known to improve air trapping ability, thus trapping warmth against the body surface (King & McLelland, 1984). Consistent with our results, a study on sparrows found that higher elevation alpine forms had higher node density relative to lowland forms (Lei et al., 2002). A further multi-species comparison found that species wintering in cold and windy conditions had downy feathers with higher node densities (D'alba et al., 2017, 2017). These interspecific trends are mirrored

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

in our results, suggesting that the White Mountain population has feather adaptations consistent with an increased need for heat retention. The lack of significant results for other feather traits may be explained by the amount of variation in the measurements as well as the sample sizes being small, resulting in an underpowered comparison. An alternative cause for the non-significant results is that alternative selection pressures not considered in the present work are driving morphology of these traits. This is somewhat supported by the overall low explanatory power of our models used in this study. Additional work on this aspect of local adaptation would benefit from investigating the influences of alternative environmental traits and utilizing a more controlled setting to evaluate the functionality of intraspecific differences in feather morphology.

Genome-wide analysis

We used GWAS to identify SNPs within genes involved in traits that differ between warm- and cold-adapted populations described previously: beak width and depth, nare length, wing cord length, and pennaceous barbule density. Because ours is one of the first studies to investigate the genetic basis of feather microstructure variation, we also report on the genes associated with other measured feather traits, even though high variance in these traits prevented our ability to robustly investigate differences between populations occupying cold and warm environments.

Our literature review of GWAS results identified genes with diverse associations. For wing chord, we identified limb and bone development, heat response, and vascularization. Notably, the genes *GUCA1C* and *NIPBL* have been linked to limb development in ducks and zebrafish (G. Li et al., 2020; Mohammadi, 2024; Muto et al.,

2014) and may play a similar developmental role in songbirds. We also detected the genes *PGLYRP2* and *PACRG*, which have been shown to be upregulated in chickens experiencing heat stress (Kim et al., 2021; Tian et al., 2020). Together, the limb development and heat response genes we identified as important further support the potential connection between wing length and thermoregulation suggested by our morphological finding of shorter wings in the colder environment. Although GWAS results do not establish a direct functional link to specific traits (Tam et al., 2019), our findings suggest that future research should investigate the potential role of thermoregulatory selective pressures on genes involved in wing development.

When focusing on genes associated with beak morphology, BSLMM found more SNPs than LMM and identified several genes potentially associated with facial structure development and climate adaptation. One gene associated with beak depth was *XIAP*. Duplications of this gene have been found to result in facial dysmorphism in humans, including the underdevelopment of cheekbones and protruding jaws (Di Benedetto et al., 2014). Although a link with beak development in birds has not yet been formally established, this could be an area for further research. Another gene identified for beak width was *MINDY2*, which belongs to a group of deubiquitinating enzymes. These genes are related to the regulation of the Wnt pathway (Park et al., 2020), a pathway that is well established as important for beak development (Nierop 2002). Additional genes such as *RASL11A*, *TINAG*, and *PLPP3* have been directly linked to beak development in avian species (Abernathy, 2021; Bai et al., 2018). Two genes associated with beak depth in our study (*LRP1B* and *UTRN*) were shown to be associated with breeding temperature in a study that used a GWAS approach to identify

1
2
3 542 loci associated with climate variables related to morphology in hermit thrush (N. Adams
4
5 543 et al., 2025). Although the beak morphology of our species did not entirely fall in line
6
7 544 with predictions based around their thermoregulatory role, the elucidation of genes
8
9 545 indicating a relationship between thermoregulation and beak morphology suggests that
10
11 546 beaks still play a thermoregulatory role in this species.
12
13
14

15 547 Our GWAS on feather traits was, to the best of our knowledge, the first genome-
16
17 548 wide association study of feather microstructural traits. Identifying the genetic basis of
18
19 549 these fine-scale traits opens new avenues for studying how feather structure evolves in
20
21 550 response to environmental pressures and functional demands. One of the genes
22
23 551 associated with pennaceous barbule length in our study, *HBE1*, has also been identified
24
25 552 in previous research on feather development. Using RNA sequencing, Limber et al.
26
27 553 (2024) found that *HBE1*, a hemoglobin gene (Mao et al., 2023), was highly expressed in
28
29 554 feather pulp, the central tissue of the developing feather germ. Additional genes
30
31 555 associated with feather traits in our study have been identified as being related to
32
33 556 plumage coloration in birds, including *EFNA2*, *NRXN1*, and *HERC4* (Shakya, 2020; L.
34
35 557 Sun et al., 2020; X. Zhang et al., 2023). Many other genes identified as being
36
37 558 associated with feather traits in our work have not been shown to be associated with
38
39 559 feather development or morphology in the literature. Feather morphology remains
40
41 560 understudied in the context of local adaptation, and the genetic architecture of these
42
43 561 microstructural traits is still poorly understood. Our study offers a foundation for deeper
44
45 562 investigation. Expanding this work to include comparative genomic studies across
46
47 563 populations or species with more strongly differentiated feather morphology could also
48
49
50
51
52
53
54
55
56
57
58
59
60

help link genotype to phenotype and provide insight into the adaptive significance of feather microstructure.

It is important to note that, in this system, we are limited in our ability to assess the role of phenotypic plasticity in driving trait differences. Bringing individuals together from both populations into a common aviary, as was done in Bears et al. (2008), would allow for better examination of these effects. It could also be argued that showing relatively high F_{ST} values for GWAS loci might suggest that the morphological differentiation is due to divergence rather than phenotypic plasticity. Our original intent with the F_{ST} outlier tests was to determine if any of the genes identified in the GWAS had variants that diverged significantly between the cold and warm environment populations. However, when comparing our F_{ST} results with SNPs identified in the GWAS, we found no overlap between the two sets. Although this could suggest that phenotypic plasticity is primarily driving the trait divergence, one must also consider that this result is consistent with known limitations of F_{ST} -based outlier scans, which are biased toward detecting loci under strong divergent selection and with large effect sizes (Hoban et al., 2016). F_{ST} measures allele frequency differences between populations and loci with large phenotypic effects are more likely to experience strong selection and exhibit pronounced divergence. In contrast, highly polygenic traits are shaped by small-effect variants that shift subtly in frequency—patterns that often fall below the detection threshold of F_{ST} scans. Our findings underscore the value of combining F_{ST} outlier analyses with GWAS approaches: while the former is designed to pinpoint loci under strong selection, the latter is better suited for uncovering the complex genetic

1
2
3 586 architecture of polygenic traits. Together, these methods can generate a more complete
4
5 587 understanding of the genetic basis for local adaptation.
6

7
8 588 While we did not find overlap between the F_{ST} and the GWAS results, our F_{ST}
9
10 589 analysis highlighted numerous genes involved in hypoxia response, altitude adaptation,
11
12 590 and cold adaptation. These results are consistent with our hypothesis that elevation and
13
14 591 associated climatic variables are applying divergent selection pressures in this species.
15
16 592 Of these genes, *NOX4* was found to be involved in mediating hypoxia-inducible
17
18 593 transcription factors, a key component of the cellular response to hypoxia (Diebold et
19
20 594 al., 2010). Similarly, *MAML2* and *SUPT20H* were found to be associated with altitude
21
22 595 adaptation in deer mice (Schweizer et al., 2021a) and *ADGRD1* was associated with
23
24 596 altitude adaptation in Tibetan chickens (Nan et al., 2023). Multiple studies identified
25
26 597 *PPP1R1C* as being related to cold adaptation in chickens (Fedorova et al., 2022a;
27
28 598 Romanov et al., 2023). These are all traits that would lend themselves to survival in
29
30 599 alpine environments, and divergent selection is likely a product of varying selection
31
32 600 pressures between the higher and colder White Mountain location and the warmer and
33
34 601 lower Piute Pass location.
35
36
37
38
39

40 602 Our F_{ST} analysis also identified several genes that might be involved with
41
42 603 plumage color differences in Rosy-Finches. Another gene that showed divergence
43
44 604 between these two populations of finch was *PHIP* which has been shown to be related
45
46 605 to differentiation in plumage coloration in House Finch (Balenger et al., 2015).
47
48 606 Interestingly, when comparing candidate outlier genes from this work with studies
49
50 607 identifying genes involved with the notable plumage coloration differences across the
51
52 608 broader rosy-finch species complex, there were some overlapping genes. In a GWAS
53
54
55
56
57
58
59
60

on crown coloration, Funk et al (2023) identified *NDUFAF2*, a gene that was also significant in this work. For body plumage coloration, our results overlap with those of Funk et al (2023) for the genes *KCNV2*, *PUM3*, and *VLDLR*. Although it was not a component of the current work, other studies have shown that plumage coloration is involved in avian thermoregulation, with darker plumage putatively facilitating heat absorption (Medina et al., 2018; Rogalla et al., 2022), and can show adaptation to local environment (Romano et al., 2019; Sandoval & Barrantes, 2019; Sirkiä et al., 2010). This provides an important avenue for future work with this species investigating the differences in plumage coloration between populations.

An additional result of our F_{ST} analysis was the identification of a gene, *CTIF*, that has been previously identified as being a climate-adapted gene in a landscape genomic study on the closely related Brown-capped Rosy-Finch (DeSaix et al., 2022). Interestingly, this gene was also identified through a contrasting approach using a mixed linear model on beak shape for two species of Darwin's finches, *Geospiza fortis* and *G. scandens* (Lawson & Petren, 2017). The authors of this paper summarize their results as indicating that inter and intraspecies beak shape variation is a result of a small suite of traits evolving in concert, corresponding to many genes. It is possible that we did not identify *CTIF* genes in our GWAS method because of low power due to smaller variation in the beak traits measured, a result of using continuous measurements between two populations versus categorical comparisons between different species as in Lawson & Petren (2017). Because of the large number of SNPs obtained for our two populations, our F_{ST} analysis likely had higher power and thus detected this gene, in addition to the fact that it had a high F_{ST} of 0.263 (averaged

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

across the 5 SNPs associated with this trait). The overlap of these genes in the two studies described as well as ours makes it a strong candidate as a genetic component of adaptation to environment, potentially through the mechanism of beak morphology.

An important outcome of local adaptation is the maintenance of ecologically important genetic variation (Savolainen et al., 2013). Conserving genetic variation prevents loss of adaptive potential and can underlie rapid adaptive responses to environmental change (Forester et al., 2023). We integrated phenotypic measurements, genome-wide association studies, and F_{ST} analyses, to understand the morphological and genetic basis of local adaptation of an alpine bird species, the Sierra Gray-crowned Rosy-Finch. When comparing traits suspected to be involved in thermoregulation for this species, we found that morphological variation across populations only partially aligned with ecogeographic expectations. Wing chord was shorter in the colder environment, consistent with Allen’s rule and likely reflecting thermoregulatory adaptation. However, beak depth and width were larger in colder environments, contrary to predictions, suggesting additional selective pressures such as diet specialization may be playing a role. Comparisons of feather microstructure further showed thermoregulatory adaptation, with higher plumulaceous node density in the high-elevation population likely enhancing insulation. When coupled with the results of our GWAS and F_{ST} outlier scans we gained a clearer picture of traits and genes important to adaptation in alpine birds. Importantly, we found that many of the genes underlying wing length, beak depth and feather microstructure had links to both development of similar traits in other organisms as well as thermoregulation. Our

654 genomic and morphological results provide evidence that this alpine species is locally
655 adapted to temperature variation.

656 For those traits that are involved in thermoregulation, we can hypothesize about
657 the implication of the intraspecific differences in light of future climate change. Recent
658 work has shown that, for populations whose morphology and genetics misalign with the
659 environment as it changes due to anthropogenic pressures there is a decrease in
660 fitness (Pelletier & Coltman, 2018; Rodriguez et al., 2025). As climate change drives
661 increasing variability in temperature, understanding both simple and complex genetic
662 architectures underlying local adaptation is essential for predicting species' responses
663 to global change. Although we cannot directly assess fitness consequences in this
664 species, this work enhances our understanding of the morphological traits and genetic
665 mechanisms underlying local adaptation in an alpine species and lays a ground work for
666 future investigation The resulting insights will be used to inform the delineation of
667 conservation units and support management strategies that seek to preserve the
668 evolutionary processes essential for long-term persistence of alpine organisms in a
669 changing world (Allendorf et al., 2010; Flanagan et al., 2018; Shafer et al., 2015).

670 **Acknowledgements**

671 We thank Sarah Albright, Alena Arnold, Emma Brown, Carl Brown, Alison Brunschwiler,
672 Vincent Chevreuil, Ryan Cheung, Daisy Cortes, Jessica Gallardo, Karim Hanna, Raquel
673 Lozano, Juan Rebellon, Leticia Santillana, Kevin Silberberg, and Juyung Yoo for their
674 participating in catching, handling, banding, measuring, and sampling birds. We thank
675 Bailey Lubash for contributing to the methodology for measuring feather microstructure
676 characteristics. We thank NovoGene for their assistance with next generation
677 sequencing. We thank Christine Rayne and Amanda Carpenter for library preparation,

and the Ruegg and Funk labs at Colorado State University for detailed feedback on early iterations of the project and manuscript. This work utilized the RMACC Summit supercomputer, which is supported by the National Science Foundation (awards ACI-1532235 and ACI-1532236), the University of Colorado Boulder and Colorado State University. The RMACC Summit supercomputer is a joint effort of the University of Colorado Boulder and Colorado State University. Figures were created with [BioRender.com](https://www.biorender.com) and illustrations were created by Erica Robertson. Funding for this project was made possible by the National Science Foundation (NSF 2222524, NSF 2222525, and NSF 2222526). Handling, banding, and sampling of Grey-crowned Rosy-Finch was performed under Federal Banding Permit 24401 and CFWScientific Collecting Permit (SCP) S-190610001-22081-001-02.

References

Abernathy, E. (2021). *The Genomic Basis of Population and Adaptive Divergence in Buteo Sister Species Across Multiple Evolutionary and Geographic Scales* [Ph.D., University of California, Davis].
<https://www.proquest.com/docview/2638107596/abstract/BB50D6B5F64067PQ/1>

Adams, D. C., Glynne, E., & Kaliontzopoulou, A. (2020). Interspecific allometry for sexual shape dimorphism: Macroevolution of multivariate sexual phenotypes with application to Rensch’s rule. *Evolution; International Journal of Organic Evolution*, 74(9), 1908–1922. <https://doi.org/10.1111/evo.14049>

Adams, N., Dias, T., Skeen, H. R., Pegan, T., Willard, D. E., Winger, B., Ruegg, K., Weeks, B. C., & Bay, R. (2025). Genetic and morphological shifts associated with climate change in a migratory bird. *BMC Biology*, 23(1), 3.
<https://doi.org/10.1186/s12915-024-02107-5>

AdaptWest Project. (2022). *Gridded current and projected climate data for North America at 1km resolution, generated using the ClimateNA v7.30 software (T. Wang et al., 2022)* [Dataset].

Ahmad, H. I., Asif, A. R., Ahmad, M. J., Jabbar, F., Adnan, M., Ahmed, S., Afzal,

- G., Saleem, A. H., Li, L., Jiang, H., Simirgiotis, M. J., Rauf, M., & Chen, J. (2020). Adaptive evolution of peptidoglycan recognition protein family regulates the innate signaling against microbial pathogens in vertebrates. *Microbial Pathogenesis*, 147, 104361. <https://doi.org/10.1016/j.micpath.2020.104361>
- Aitken, S. N., Yeaman, S., Holliday, J. A., Wang, T., & Curtis-McLane, S. (2008). Adaptation, migration or extirpation: Climate change outcomes for tree populations. *Evolutionary Applications*, 1(1), 95–111. <https://doi.org/10.1111/j.1752-4571.2007.00013.x>
- Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, 19(9), 1655–1664. <https://doi.org/10.1101/gr.094052.109>
- Allen, J. A. (1877). *The Influence of Physical Conditions in the Genesis of Species*.
- Allendorf, F. W., Hohenlohe, P. A., & Luikart, G. (2010). Genomics and the future of conservation genetics. *Nature Reviews Genetics*, 11(10), 697–709. <https://doi.org/10.1038/nrg2844>
- Alsoufi, M. A., Liu, Y., Cao, C., Zhao, J., Kang, J., Li, M., Wang, K., He, Y., & Ge, C. (2023). Integrated Transcriptomics Profiling in Chahua and Digao Chickens' Breast for Assessment Molecular Mechanism of Meat Quality Traits. *Genes*, 14(1), Article 1. <https://doi.org/10.3390/genes14010095>
- Ashton, K. G. (2002). Patterns of within-species body size variation of birds: Strong evidence for Bergmann's rule. *Global Ecology and Biogeography*, 11(6), 505–523. <https://doi.org/10.1046/j.1466-822X.2002.00313.x>

1
2
3 729 Auwera, G. V. der, & O'Connor, B. D. (2020). *Genomics in the cloud: Using*
4
5 730 *Docker, GATK, and WDL in Terra* (First edition). O'Reilly.
6
7
8 731 Bai, H., Sun, Y., Liu, N., Xue, F., Li, Y., Xu, S., Ye, J., Zhang, L., Chen, Y., &
9
10 732 Chen, J. (2018). Single SNP- and pathway-based genome-wide association
11
12 733 studies for beak deformity in chickens using high-density 600K SNP arrays. *BMC*
13
14 734 *Genomics*, 19(1), 501. <https://doi.org/10.1186/s12864-018-4882-8>
15
16
17 735 Bai, H., Zhao, N., Li, X., Ding, Y., Guo, Q., Chen, G., & Chang, G. (2024). Whole-
18
19 736 genome resequencing identifies candidate genes associated with heat
20
21 737 adaptation in chickens. *Poultry Science*, 103(10), 104139.
22
23
24 738 <https://doi.org/10.1016/j.psj.2024.104139>
25
26 739 Bai, H., Zhu, J., Sun, Y., Liu, R., Liu, N., Li, D., Wen, J., & Chen, J. (2014).
27
28 740 Identification of Genes Related to Beak Deformity of Chickens Using Digital
29
30 741 Gene Expression Profiling. *PLOS ONE*, 9(9), e107050.
31
32 742 <https://doi.org/10.1371/journal.pone.0107050>
33
34
35 743 Balenger, S. L., Bonneaud, C., Sefick, S. A., Edwards, S. V., & Hill, G. E. (2015).
36
37 744 Plumage color and pathogen-induced gene expression in a wild bird. *Behavioral*
38
39 745 *Ecology*, 26(4), 1100–1110. <https://doi.org/10.1093/beheco/arv055>
40
41
42 746 Barnes, N. E., Mendoza, K. M., Strasburg, G. M., Velleman, S. G., & Reed, K. M.
43
44 747 (2019). Thermal challenge alters the transcriptional profile of the breast muscle in
45
46 748 turkey poults. *Poultry Science*, 98(1), 74–91. <https://doi.org/10.3382/ps/pey401>
47
48
49 749 Barve, S., Ramesh, V., Dotterer, T. M., & Dove, C. J. (2021). Elevation and body
50
51 750 size drive convergent variation in thermo-insulative feather structure of
52
53 751 Himalayan birds. *Ecography*, 44(5), 680–689. <https://doi.org/10.1111/ecog.05376>
54
55
56
57
58
59
60

- 1
2
3 752 Bears, H., C. Drever, M., & Martin, K. (2008). Comparative morphology of dark-
4
5 753 eyed juncos *Junco hyemalis* breeding at two elevations: A common aviary
6
7 754 experiment. *Journal of Avian Biology*, 39(2), 152–162.
8
9
10 755 <https://doi.org/10.1111/j.2008.0908-8857.04191.x>
11
12 756 Belnap, J., Webb, R. H., Esque, T. C., Brooks, M. L., Defalco, L. A., &
13
14 757 Macmahon, J. A. (2016). Deserts. In *Ecosystems of California* (1st ed., pp. 635–
15
16 758 668). University of California Press.
17
18
19 759 <http://www.jstor.org/stable/10.1525/j.ctv1xxzp6.35>
20
21 760 Bhavana, K., Foote, D. J., Srikanth, K., Balakrishnan, C. N., Prabhu, V. R.,
22
23 761 Sankaralingam, S., Singha, H. S., Gopalakrishnan, A., & Nagarajan, M. (2022).
24
25 762 Comparative transcriptome analysis of Indian domestic duck reveals candidate
26
27 763 genes associated with egg production. *Scientific Reports*, 12(1), 10943.
28
29
30 764 <https://doi.org/10.1038/s41598-022-15099-5>
31
32 765 Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to
33
34 766 measuring local adaptation. *Ecology Letters*, 16(9), 1195–1205.
35
36
37 767 <https://doi.org/10.1111/ele.12150>
38
39 768 Boer, E. F., Maclary, E. T., & Shapiro, M. D. (2021). Complex genetic
40
41 769 architecture of three-dimensional craniofacial shape variation in domestic
42
43 770 pigeons. *Evolution & Development*, 23(6), 477–495.
44
45
46 771 <https://doi.org/10.1111/ede.12395>
47
48 772 Borges, R., Fonseca, J., Gomes, C., Johnson, W. E., O'Brien, S. J., Zhang, G.,
49
50 773 Gilbert, M. T. P., Jarvis, E. D., & Antunes, A. (2019). Avian Binocularity and
51
52 774 Adaptation to Nocturnal Environments: Genomic Insights from a Highly Derived
53
54
55
56
57
58
59
60

1
2
3 775 Visual Phenotype. *Genome Biology and Evolution*, 11(8), 2244–2255.
4
5 776 <https://doi.org/10.1093/gbe/evz111>
6
7
8 777 Browning, B. L., & Browning, S. R. (2016). Genotype Imputation with Millions of
9
10 778 Reference Samples. *American Journal of Human Genetics*, 98(1), 116–126.
11
12 779 <https://doi.org/10.1016/j.ajhg.2015.11.020>
13
14
15 780 Buroker, N. E., Ning, X.-H., Zhou, Z.-N., Li, K., Cen, W.-J., Wu, X.-F., Zhu, W.-Z.,
16
17 781 Scott, C. R., & Chen, S.-H. (2012). AKT3, ANGPTL4, eNOS3, and VEGFA
18
19 782 associations with high altitude sickness in Han and Tibetan Chinese at the
20
21 783 Qinghai-Tibetan Plateau. *International Journal of Hematology*, 96(2), 200–213.
22
23 784 <https://doi.org/10.1007/s12185-012-1117-7>
24
25
26 785 Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Exposito-Alonso, M., & Keller, S. R.
27
28 786 (2020). Genomic Prediction of (Mal)Adaptation Across Current and Future
29
30 787 Climatic Landscapes. *Annual Review of Ecology, Evolution, and Systematics*,
31
32 788 51(Volume 51, 2020), 245–269. [https://doi.org/10.1146/annurev-ecolsys-020720-](https://doi.org/10.1146/annurev-ecolsys-020720-042553)
33
34 789 042553
35
36
37 790 Ceresa, F., Brambilla, M., Kvist, L., Vitulano, S., Pes, M., Tomasi, L., Pedrini, P.,
38
39 791 Bettega, C., Anderle, M., Hilpold, A., & Kranebitter, P. (2024). Wing morphology
40
41 792 changes with habitat availability and elevation in an alpine-specialist bird. *Global*
42
43 793 *Ecology and Conservation*, 56, e03276.
44
45 794 <https://doi.org/10.1016/j.gecco.2024.e03276>
46
47
48 795 Chen, C.-K., Chang, Y.-M., Jiang, T.-X., Yue, Z., Liu, T.-Y., Lu, J., Yu, Z., Lin, J.-
49
50 796 J., Vu, T.-D., Huang, T.-Y., Harn, H. I.-C., Ng, C. S., Wu, P., Chuong, C.-M., & Li,
51
52 797 W.-H. (2024). Conserved regulatory switches for the transition from natal down to
53
54
55
56
57
58
59
60

juvenile feather in birds. *Nature Communications*, 15(1), 4174.

<https://doi.org/10.1038/s41467-024-48303-3>

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, 34(17), i884–i890.

<https://doi.org/10.1093/bioinformatics/bty560>

Chen, X., Bai, X., Liu, H., Zhao, B., Yan, Z., Hou, Y., & Chu, Q. (2022).

Population Genomic Sequencing Delineates Global Landscape of Copy Number Variations that Drive Domestication and Breed Formation of in Chicken. *Frontiers in Genetics*, 13. <https://doi.org/10.3389/fgene.2022.830393>

Chen, Z., Boehnke, M., Wen, X., & Mukherjee, B. (2021). Revisiting the genome-wide significance threshold for common variant GWAS. *G3*

Genes|Genomes|Genetics, 11(2), jkaa056.

<https://doi.org/10.1093/g3journal/jkaa056>

Cheng, Y., Miller, M. J., Zhang, D., Song, G., Jia, C., Qu, Y., & Lei, F. (2020).

Comparative Genomics Reveals Evolution of a Beak Morphology Locus in a High-Altitude Songbird. *Molecular Biology and Evolution*, 37(10), 2983–2988.

<https://doi.org/10.1093/molbev/msaa157>

Cho, E., Kim, M., Cho, S., So, H.-J., Lee, K.-T., Cha, J., Jin, D., & Lee, J. H.

(2023). A genome-wide association study for the fatty acid composition of breast

meat in an F2 crossbred chicken population. *Journal of Animal Science and*

Technology, 65(4), 735–747. <https://doi.org/10.5187/jast.2023.e1>

Cho, Y. S., Jun, J. H., Kim, J. A., Kim, H.-M., Chung, O., Kang, S.-G., Park, J.-Y.,

Kim, H.-J., Kim, S., Kim, H.-J., Jang, J., Na, K.-J., Kim, J., Park, S. G., Lee, H.-

1
2
3 821 Y., Manica, A., Mindell, D. P., Fuchs, J., Edwards, J. S., ... Bhak, J. (2019).
4
5 822 Raptor genomes reveal evolutionary signatures of predatory and nocturnal
6
7 823 lifestyles. *Genome Biology*, 20(1), 181. [https://doi.org/10.1186/s13059-019-1793-](https://doi.org/10.1186/s13059-019-1793-1)
8
9 824 1
10
11
12 825 Chung, O., Jin, S., Cho, Y. S., Lim, J., Kim, H., Jho, S., Kim, H.-M., Jun, J., Lee,
13
14 826 H., Chon, A., Ko, J., Edwards, J., Weber, J. A., Han, K., O'Brien, S. J., Manica,
15
16 827 A., Bhak, J., & Paek, W. K. (2015). The first whole genome and transcriptome of
17
18 828 the cinereous vulture reveals adaptation in the gastric and immune defense
19
20 829 systems and possible convergent evolution between the Old and New World
21
22 830 vultures. *Genome Biology*, 16(1), 215. [https://doi.org/10.1186/s13059-015-0780-](https://doi.org/10.1186/s13059-015-0780-4)
23
24 831 4
25
26
27 832 Clark, J., Lim, H. C., Olsen, B., & Kovach, A. (2024). *Whole-genome data*
28
29 833 *uncover the basis of local adaptation with gene flow in a threatened coastal*
30
31 834 *songbird*. <https://doi.org/10.22541/au.172114945.59127531/v1>
32
33
34 835 Clausen, J., Keck, D. D., & Hiesey, W. M. (1941). Regional Differentiation in
35
36 836 Plant Species. *The American Naturalist*, 75(758), 231–250.
37
38 837 <https://doi.org/10.1086/280955>
39
40
41 838 Crates, R., von Takach, B., Young, C. M., Stojanovic, D., Neaves, L. E., Murphy,
42
43 839 L., Gautschi, D., Hogg, C. J., Heinsohn, R., Bell, P., & Farquharson, K. A. (2024).
44
45 840 Genomic insights into the critically endangered King Island scrubtit. *Journal of*
46
47 841 *Heredity*, 115(5), 552–564. <https://doi.org/10.1093/jhered/esae029>
48
49
50 842 da Costa Moreno, H. (2022). *Molecular Evolution of Vertebrate Genes Involved*
51
52 843 *in Mitochondrial Biogenesis: Unravelling the Origins of Birds' Endothermy*
53
54
55
56
57
58
59
60

- 844 [Masters Thesis, Universidade do Porto]. ProQuest Dissertations & Theses.
- 845 D'alba, L., Carlsen, T. H., Ásgeirsson, Á., Shawkey, M. D., & Jónsson, J. E.
- 846 (2017). Contributions of feather microstructure to eider down insulation
- 847 properties. *Journal of Avian Biology*, 48(8), 1150–1157.
- 848 <https://doi.org/10.1111/jav.01294>
- 849 Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,
- 850 Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve
- 851 years of SAMtools and BCFtools. *GigaScience*, 10(2), giab008.
- 852 <https://doi.org/10.1093/gigascience/giab008>
- 853 Delmore, K., Illera, J. C., Pérez-Tris, J., Segelbacher, G., Lugo Ramos, J. S.,
- 854 Durieux, G., Ishigohoka, J., & Liedvogel, M. (2020). The evolutionary history and
- 855 genomics of European blackcap migration. *eLife*, 9, e54462.
- 856 <https://doi.org/10.7554/eLife.54462>
- 857 Demery, A.-J. C., Burns, K. J., & Mason, N. A. (2021). Bill size, bill shape, and
- 858 body size constrain bird song evolution on a macroevolutionary scale.
- 859 *Ornithology*, 138(2), ukab011. <https://doi.org/10.1093/ornithology/ukab011>
- 860 DeSaix, M., George, T., Seglund, A., Spellman, G., Zavaleta, E., & Ruegg, K.
- 861 (2022). Forecasting climate change response in an alpine specialist songbird
- 862 reveals the importance of considering novel climate. *Diversity and Distributions*,
- 863 28. <https://doi.org/10.1111/ddi.13628>
- 864 Di Benedetto, D., Musumeci, S. A., Avola, E., Alberti, A., Buono, S., Scuderi, C.,
- 865 Grillo, L., Galesi, O., Spalletta, A., Giudice, M. L., Luciano, D., Vinci, M., Bianca,
- 866 S., Romano, C., & Fichera, M. (2014). Definition of minimal duplicated region

1
2
3 867 encompassing the XIAP and STAG2 genes in the Xq25 microduplication
4
5 868 syndrome. *American Journal of Medical Genetics. Part A*, 164A(8), 1923–1930.
6
7 869 <https://doi.org/10.1002/ajmg.a.36570>
8
9
10 870 Diebold, I., Petry, A., Hess, J., & Görlach, A. (2010). The NADPH Oxidase
11
12 871 Subunit NOX4 Is a New Target Gene of the Hypoxia-inducible Factor-1.
13
14 872 *Molecular Biology of the Cell*, 21(12), 2087–2096.
15
16 873 <https://doi.org/10.1091/mbc.e09-12-1003>
17
18
19 874 Ducrest, A.-L., San-Jose, L. M., Neuenschwander, S., Schmid-Siegert, E.,
20
21 875 Simon, C., Pagni, M., Iseli, C., Richter, H., Guex, N., Cumer, T., Beaudoin, E.,
22
23 876 Dupasquier, M., Charruau, P., Ducouret, P., Xenarios, I., Goudet, J., & Roulin, A.
24
25 877 (2025). Melanin and Neurotransmitter Signalling Genes Are Differentially Co-
26
27 878 Expressed in Growing Feathers of White and Rufous Barn Owls. *Pigment Cell &*
28
29 879 *Melanoma Research*, 38(2), e70001. <https://doi.org/10.1111/pcmr.70001>
30
31
32 880 Egbert, J. R., & Belthoff, J. R. (2003). Wing Shape in House Finches Differs
33
34 881 Relative to Migratory Habit in Eastern and Western North America. *The Condor*,
35
36 882 105(4), 825–829. <https://doi.org/10.1093/condor/105.4.825>
37
38
39 883 Endler, J. A. (1977). *Geographic Variation, Speciation and Clines. (MPB-10)*,
40
41 884 *Volume 10*. Princeton University Press. <https://doi.org/10.2307/j.ctvx5wbdg>
42
43
44 885 Epanchin, P. N., Knapp, R. A., & Lawler, S. P. (2010). Nonnative trout impact an
45
46 886 alpine-nesting bird by altering aquatic-insect subsidies. *Ecology*, 91(8), 2406–
47
48 887 2415. <https://doi.org/10.1890/09-1974.1>
49
50
51 888 Faria, R., Renaut, S., Galindo, J., Pinho, C., Melo-Ferreira, J., Melo, M., Jones,
52
53 889 F., Salzburger, W., Schluter, D., & Butlin, R. (2014). Advances in Ecological
54
55
56
57
58
59
60

- 890 Speciation: An integrative approach. *Molecular Ecology*, 23(3), 513–521.
- 891 <https://doi.org/10.1111/mec.12616>
- 892 Farias-Virgens, M., Peede, D., Deacon, T., Okanoya, K., White, S. A., & Huerta-
893 Sanchez, E. (2025). The genomics of the domestication syndrome in a songbird
894 model species. *Communications Biology*, 8(1), 853.
- 895 <https://doi.org/10.1038/s42003-025-08235-0>
- 896 Fedorova, E. S., Dementieva, N. V., Shcherbakov, Y. S., & Stanishevskaya, O. I.
897 (2022a). Identification of Key Candidate Genes in Runs of Homozygosity of the
898 Genome of Two Chicken Breeds, Associated with Cold Adaptation. *Biology*,
899 11(4), Article 4. <https://doi.org/10.3390/biology11040547>
- 900 Feng, J., Zhu, W., Shi, H., Peng, D., Zang, L., Wang, Y., ZhaXi, L., BaiMa, J.,
901 Amevor, F. K., Wang, X., Ma, X., & Zhao, X. (2023). Analysis of the Selection
902 Signal of the Tibetan Black Chicken Genome Based on Whole-Genome
903 Sequencing. *Genes*, 14(9), Article 9. <https://doi.org/10.3390/genes14091672>
- 904 Flanagan, S. P., Forester, B. R., Latch, E. K., Aitken, S. N., & Hoban, S. (2018).
905 Guidelines for planning genomic assessment and monitoring of locally adaptive
906 variation to inform species conservation. *Evolutionary Applications*, 11(7), 1035–
907 1052. <https://doi.org/10.1111/eva.12569>
- 908 Forester, B. R., Beever, E. A., Darst, C., Szymanski, J., & Funk, W. C. (2022).
909 Linking evolutionary potential to extinction risk: Applications and future directions.
910 *Frontiers in Ecology and the Environment*, 20(9), 507–515.
- 911 <https://doi.org/10.1002/fee.2552>
- 912 Forester, B. R., Day, C. C., Ruegg, K., & Landguth, E. L. (2023). Evolutionary

1
2
3 913 potential mitigates extinction risk under climate change in the endangered
4
5 914 southwestern willow flycatcher. *Journal of Heredity*, 114(4), 341–353.
6
7
8 915 <https://doi.org/10.1093/jhered/esac067>
9
10 916 Franchini, P., Irisarri, I., Fudickar, A., Schmidt, A., Meyer, A., Wikelski, M., &
11
12 917 Partecke, J. (2017). Animal tracking meets migration genomics: Transcriptomic
13
14 918 analysis of a partially migratory bird species. *Molecular Ecology*, 26(12), 3204–
15
16
17 919 3216. <https://doi.org/10.1111/mec.14108>
18
19 920 Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018).
20
21 921 Climate change causes upslope shifts and mountaintop extirpations in a tropical
22
23 922 bird community. *Proceedings of the National Academy of Sciences*, 115(47),
24
25
26 923 11982–11987. <https://doi.org/10.1073/pnas.1804224115>
27
28 924 Fridolfsson, A.-K., & Ellegren, H. (2000). Molecular Evolution of the Avian CHD1
29
30 925 Genes on the Z and W Sex Chromosomes. *Genetics*, 155(4), 1903–1912.
31
32 926 <https://doi.org/10.1093/genetics/155.4.1903>
33
34 927 Funk, E. R., Spellman, G. M., Winker, K., Withrow, J. J., Ruegg, K. C., & Taylor,
35
36 928 S. A. (2023). The genetic basis of plumage coloration and elevation adaptation in
37
38 929 a clade of recently diverged alpine and arctic songbirds. *Evolution; International*
39
40 930 *Journal of Organic Evolution*, 77(3), 705–717.
41
42 931 <https://doi.org/10.1093/evolut/qpac064>
43
44 932 Gao, G., Zhang, K., Zhao, X., Xu, G., Xie, Y., Zhou, L., Zhang, C., & Wang, Q.
45
46 933 (2023). Identification of Molecular Markers Associated with Goose Egg Quality
47
48 934 Through Genome-Wide Association Analysis. *Scientia Agricultura Sinica*, 56(19),
49
50
51 935 3894–3904. <https://doi.org/10.3864/j.issn.0578-1752.2023.19.015>
52
53
54
55
56
57
58
59
60

- 936 García, N. C., Campagna, L., Rush, A. C., Bowie, R. C. K., & Lovette, I. J.
937 (2023). Comparative genomics of two Empidonax flycatchers reveal candidate
938 genes for bird song production. *Evolution*, 77(8), 1818–1828.
939 <https://doi.org/10.1093/evolut/qpad096>
- 940 Gesemann, M., & Neuhauss, S. C. F. (2023). Evolution of visual guanylyl
941 cyclases and their activating proteins with respect to clade and species-specific
942 visual system adaptation. *Frontiers in Molecular Neuroscience*, 16.
943 <https://doi.org/10.3389/fnmol.2023.1131093>
- 944 Gheyas, A. A., Vallejo-Trujillo, A., Kebede, A., Lozano-Jaramillo, M., Dessie, T.,
945 Smith, J., & Hanotte, O. (2021). Integrated Environmental and Genomic Analysis
946 Reveals the Drivers of Local Adaptation in African Indigenous Chickens.
947 *Molecular Biology and Evolution*, 38(10), 4268–4285.
948 <https://doi.org/10.1093/molbev/msab156>
- 949 Gienapp, P., Laine, V. N., Mateman, A. C., van Oers, K., & Visser, M. E. (2017).
950 Environment-Dependent Genotype-Phenotype Associations in Avian Breeding
951 Time. *Frontiers in Genetics*, 8. <https://doi.org/10.3389/fgene.2017.00102>
- 952 Grabherr, G., Gottfried, M., & Pauli, H. (2010). Climate Change Impacts in Alpine
953 Environments. *Geography Compass*, 4(8), 1133–1153.
954 <https://doi.org/10.1111/j.1749-8198.2010.00356.x>
- 955 Grant, P. R., Grant, B. R., Smith, J. N., Abbott, I. J., & Abbott, L. K. (1976).
956 Darwin's finches: Population variation and natural selection. *Proceedings of the*
957 *National Academy of Sciences*, 73(1), 257–261.
958 <https://doi.org/10.1073/pnas.73.1.257>

1
2
3 959 Greenberg, R., Cadena, V., Danner, R. M., & Tattersall, G. (2012). Heat Loss
4
5 960 May Explain Bill Size Differences between Birds Occupying Different Habitats.
6
7 961 *PLOS ONE*, 7(7), e40933. <https://doi.org/10.1371/journal.pone.0040933>
8
9
10 962 Grilli, M. G., Lambertucci, S. A., Therrien, J.-F., & Bildstein, K. L. (2017). Wing
11
12 963 size but not wing shape is related to migratory behavior in a soaring bird. *Journal*
13
14 964 *of Avian Biology*, 48(5), 669–678. <https://doi.org/10.1111/jav.01220>
15
16
17 965 Grinnell, J. (1913). *Leucosticte tephrocotis dawsoni*: A New Race of Rosy Finch
18
19 966 from the Sierra Nevada. *The Condor*, 15(2), 76–79.
20
21 967 <https://doi.org/10.2307/1362056>
22
23
24 968 Grinnell, J. (1917). Field Tests of Theories Concerning Distributional Control. *The*
25
26 969 *American Naturalist*, 51(602), 115–128. <https://doi.org/10.1086/279591>
27
28 970 Guo, J., Sun, C., Qu, L., Shen, M., Dou, T., Ma, M., Wang, K., & Yang, N. (2017).
29
30 971 Genetic architecture of bone quality variation in layer chickens revealed by a
31
32 972 genome-wide association study. *Scientific Reports*, 7(1), 45317.
33
34 973 <https://doi.org/10.1038/srep45317>
35
36
37 974 He, J., Tu, J., Yu, J., & Jiang, H. (2023). A global assessment of Bergmann’s rule
38
39 975 in mammals and birds. *Global Change Biology*, 29(18), 5199–5210.
40
41 976 <https://doi.org/10.1111/gcb.16860>
42
43
44 977 Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D.
45
46 978 B., Poss, M. L., Reed, L. K., Storfer, A., & Whitlock, M. C. (2016). Finding the
47
48 979 Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future
49
50 980 Directions. *The American Naturalist*, 188(4), 379–397.
51
52 981 <https://doi.org/10.1086/688018>
53
54
55
56
57
58
59
60

- 982 Hu, X., Zhang, X., Liu, Z., Li, S., Zheng, X., Nie, Y., Tao, Y., Zhou, X., Wu, W.,
983 Yang, G., Zhao, Q., Zhang, Y., Xu, Q., & Mou, C. (2020). Exploration of key
984 regulators driving primary feather follicle induction in goose skin. *Gene*, 731,
985 144338. <https://doi.org/10.1016/j.gene.2020.144338>
- 986 Johnsson, M., Jonsson, K. B., Andersson, L., Jensen, P., & Wright, D. (2015).
987 Genetic Regulation of Bone Metabolism in the Chicken: Similarities and
988 Differences to Mammalian Systems. *PLOS Genetics*, 11(5), e1005250.
989 <https://doi.org/10.1371/journal.pgen.1005250>
- 990 Johnston, R. A., Paxton, K. L., Moore, F. R., Wayne, R. K., & Smith, T. B. (2016).
991 Seasonal gene expression in a migratory songbird. *Molecular Ecology*, 25(22),
992 5680–5691. <https://doi.org/10.1111/mec.13879>
- 993 Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation.
994 *Ecology Letters*, 7(12), 1225–1241. [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2004.00684.x)
995 0248.2004.00684.x
- 996 Kim, H., Kim, H., Seong, P., Arora, D., Shin, D., Park, W., & Park, J.-E. (2021).
997 Transcriptomic Response under Heat Stress in Chickens Revealed the
998 Regulation of Genes and Alteration of Metabolism to Maintain Homeostasis.
999 *Animals: An Open Access Journal from MDPI*, 11(8), 2241.
1000 <https://doi.org/10.3390/ani11082241>
- 1001 King, A. S., & McLelland, J. (1984). *Birds: Their structure and function* (2nd ed).
1002 Baillière Tindall.
- 1003 Kinsella, C. M., Ruiz-Ruano, F. J., Dion-Côté, A.-M., Charles, A. J., Gossmann,
1004 T. I., Cabrero, J., Kappei, D., Hemmings, N., Simons, M. J. P., Camacho, J. P.

1
2
3 1005 M., Forstmeier, W., & Suh, A. (2019). Programmed DNA elimination of germline
4
5 1006 development genes in songbirds. *Nature Communications*, 10(1), 5468.
6
7
8 1007 <https://doi.org/10.1038/s41467-019-13427-4>
9
10 1008 Körner, C. (2003). Alpine climate. In C. Körner (Ed.), *Alpine Plant Life: Functional*
11
12 1009 *Plant Ecology of High Mountain Ecosystems* (pp. 21–30). Springer.
13
14
15 1010 https://doi.org/10.1007/978-3-642-18970-8_3
16
17 1011 Koskenpato, K., Ahola, K., Karstinen, T., & Karell, P. (2016). Is the denser
18
19 1012 contour feather structure in pale grey than in pheomelanic brown tawny owls
20
21 1013 *Strix aluco* an adaptation to cold environments? *Journal of Avian Biology*, 47(1),
22
23 1014 1–6. <https://doi.org/10.1111/jav.00746>
24
25
26 1015 Kuttiyarthu Veetil, N., Henschen, A. E., Hawley, D. M., Melepat, B., Dalloul, R. A.,
27
28 1016 Beneš, V., Adelman, J. S., & Vinkler, M. (2024). Varying conjunctival immune
29
30 1017 response adaptations of house finch populations to a rapidly evolving bacterial
31
32 1018 pathogen. *Frontiers in Immunology*, 15.
33
34
35 1019 <https://doi.org/10.3389/fimmu.2024.1250818>
36
37
38 1020 Lamichhaney, S., Han, F., Webster, M. T., Andersson, L., Grant, B. R., & Grant,
39
40 1021 P. R. (2018). Rapid hybrid speciation in Darwin’s finches. *Science*, 359(6372),
41
42 1022 224–228. <https://doi.org/10.1126/science.aao4593>
43
44
45 1023 Lande, R., & Shannon, S. (1996). The role of genetic variation in adaptation and
46
47 1024 population persistence in a changing environment. *Evolution*, 50(1), 434–437.
48
49 1025 <https://doi.org/10.1111/j.1558-5646.1996.tb04504.x>
50
51
52 1026 Lawson, L. P., & Petren, K. (2017). The adaptive genomic landscape of beak
53
54 1027 morphology in Darwin’s finches. *Molecular Ecology*, 26(19), 4978–4989.
55
56
57
58
59
60

- 1028 <https://doi.org/10.1111/mec.14166>
- 1029 Lei, F.-M., Qu, Y.-H., Gan, Y.-L., Gebauer, A., & Kaiser, M. (2002). The feather
1030 microstructure of Passerine sparrows in China. *Journal Für Ornithologie*, 143(2),
1031 205–212. <https://doi.org/10.1046/j.1439-0361.2002.02001.x>
- 1032 Lennon, O. (2022). *Do Candidate Genes Distinguish Migratory and Nonmigratory*
1033 *Avian Species?* [Undergraduate Research Scholars Thesis, Texas A&M
1034 University]. <https://hdl.handle.net/1969.1/196498>
- 1035 Lenth, R. (2025). *emmeans: Estimated Marginal Means, aka Least-Squares*
1036 *Means* (Version 1.10.7-100002) [R package].
- 1037 Lewden, A., Bishop, C. M., & Askew, G. N. (2023). How birds dissipate heat
1038 before, during and after flight. *Journal of the Royal Society Interface*, 20(209),
1039 20230442. <https://doi.org/10.1098/rsif.2023.0442>
- 1040 Li, G.-S., Liu, W.-W., Zhang, F., Zhu, F., Yang, F.-X., Hao, J.-P., & Hou, Z.-C.
1041 (2020). Genome-wide association study of bone quality and feed efficiency-
1042 related traits in Pekin ducks. *Genomics*, 112(6), 5021–5028.
1043 <https://doi.org/10.1016/j.ygeno.2020.09.023>
- 1044 Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with
1045 Burrows–Wheeler transform. *Bioinformatics*, 25(14), 1754–1760.
1046 <https://doi.org/10.1093/bioinformatics/btp324>
- 1047 Limber, C., Wagner, G. P., & Prum, R. O. (2024). *Genetic Characterization of the*
1048 *Cell Types of in Developing Feathers, and the Evolution of Feather Complexity*
1049 (p. 2024.11.21.624532). bioRxiv. <https://doi.org/10.1101/2024.11.21.624532>
- 1050 Lockwood, R., Swaddle, J. P., & Rayner, J. M. V. (1998). Avian Wingtip Shape

1
2
3 1051 Reconsidered: Wingtip Shape Indices and Morphological Adaptations to
4
5 1052 Migration. *Journal of Avian Biology*, 29(3), 273–292.
6
7
8 1053 <https://doi.org/10.2307/3677110>
9
10 1054 Loma, J. D., Vicente, M., Tirado, N., Ascui, F., Parada, L. A., Gardon, J.,
11
12 1055 Schlebusch, C., & Broberg, K. (2025). Human adaptation in the Andes
13
14 1056 Mountains. *Human Population Genetics and Genomics*, 5(1).
15
16
17 1057 <https://doi.org/10.47248/hpgg2505010002>
18
19 1058 Lotterhos, K. E., & Whitlock, M. C. (2015). The relative power of genome scans
20
21 1059 to detect local adaptation depends on sampling design and statistical method.
22
23 1060 *Molecular Ecology*, 24(5), 1031–1046. <https://doi.org/10.1111/mec.13100>
24
25
26 1061 Lou, R. N., & Therkildsen, N. O. (2022). Batch effects in population genomic
27
28 1062 studies with low-coverage whole genome sequencing data: Causes, detection
29
30 1063 and mitigation. *Molecular Ecology Resources*, 22(5), 1678–1692.
31
32
33 1064 <https://doi.org/10.1111/1755-0998.13559>
34
35 1065 Lovell, P. V., Carleton, J. B., & Mello, C. V. (2013). Genomics analysis of
36
37 1066 potassium channel genes in songbirds reveals molecular specializations of brain
38
39 1067 circuits for the maintenance and production of learned vocalizations. *BMC*
40
41 1068 *Genomics*, 14(1), 470. <https://doi.org/10.1186/1471-2164-14-470>
42
43
44 1069 Luo, X., Guo, J., Zhang, J., Ma, Z., & Li, H. (2024). Overview of chicken embryo
45
46 1070 genes related to sex differentiation. *PeerJ*, 12, e17072.
47
48
49 1071 <https://doi.org/10.7717/peerj.17072>
50
51 1072 Ma, S., Liu, H., Wang, J., Wang, L., Xi, Y., Liu, Y., Xu, Q., Hu, J., Han, C., Bai, L.,
52
53 1073 Li, L., & Wang, J. (2021). Transcriptome Analysis Reveals Genes Associated
54
55
56
57
58
59
60

- 1074 With Sexual Dichromatism of Head Feather Color in Mallard. *Frontiers in*
1075 *Genetics*, 12. <https://doi.org/10.3389/fgene.2021.627974>
- 1076 MacDougall-Shackleton, S. A., Johnson, R. E., & Hahn, T. P. (2000). Gray-
1077 crowned Rosy-Finch (*Leucosticte tephrocotis*). In *The Birds of North America*.
1078 Cornell Lab of Ornithology.
- 1079 Mamantov, M. A., Gibson-Reinemer, D. K., Linck, E. B., & Sheldon, K. S. (2021).
1080 Climate-driven range shifts of montane species vary with elevation. *Global*
1081 *Ecology and Biogeography*, 30(4), 784–794. <https://doi.org/10.1111/geb.13246>
- 1082 Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., & Zimin,
1083 A. (2018). MUMmer4: A fast and versatile genome alignment system. *PLOS*
1084 *Computational Biology*, 14(1), e1005944.
1085 <https://doi.org/10.1371/journal.pcbi.1005944>
- 1086 Mascarenhas, R., Meirelles, P. M., & Batalha-Filho, H. (2023). Urbanization
1087 drives adaptive evolution in a Neotropical bird. *Current Zoology*, 69(5), 607–619.
1088 <https://doi.org/10.1093/cz/zoac066>
- 1089 Matsuda, Y., & Makino, T. (2024). Comparative genomics reveals convergent
1090 signals associated with the high metabolism and longevity in birds and bats.
1091 *Proceedings of the Royal Society B: Biological Sciences*, 291(2029), 20241068.
1092 <https://doi.org/10.1098/rspb.2024.1068>
- 1093 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,
1094 Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The
1095 Genome Analysis Toolkit: A MapReduce framework for analyzing next-
1096 generation DNA sequencing data. *Genome Research*, 20(9), 1297–1303.

1
2
3 1097 <https://doi.org/10.1101/gr.107524.110>
4
5 1098 Medina, I., Newton, E., Kearney, M. R., Mulder, R. A., Porter, W. P., & Stuart-
6
7 1099 Fox, D. (2018). Reflection of near-infrared light confers thermal protection in
8
9 birds. *Nature Communications*, 9(1), 3610. [https://doi.org/10.1038/s41467-018-](https://doi.org/10.1038/s41467-018-05898-8)
10 1100
11 05898-8
12 1101
13
14 1102 Meek, M. H., Beever, E. A., Barbosa, S., Fitzpatrick, S. W., Fletcher, N. K.,
15
16 Mittan-Moreau, C. S., Reid, B. N., Campbell-Staton, S. C., Green, N. F., &
17 1103
18 Hellmann, J. J. (2023). Understanding Local Adaptation to Prepare Populations
19 1104
20 for Climate Change. *BioScience*, 73(1), 36–47.
21 1105
22 <https://doi.org/10.1093/biosci/biac101>
23 1106
24
25 Meiri, S., & Dayan, T. (2003). On the validity of Bergmann’s rule. *Journal of*
26 1107
27 *Biogeography*, 30(3), 331–351. <https://doi.org/10.1046/j.1365-2699.2003.00837.x>
28 1108
29
30 Mesner, L. D., Ray, B., Hsu, Y.-H., Manichaikul, A., Lum, E., Bryda, E. C., Rich,
31 1109
32 S. S., Rosen, C. J., Criqui, M. H., Allison, M., Budoff, M. J., Clemens, T. L., &
33 1110
34 Farber, C. R. (2014). *Bicc1* is a genetic determinant of osteoblastogenesis and
35 1111
36 bone mineral density. *The Journal of Clinical Investigation*, 124(6), 2736–2749.
37 1112
38 <https://doi.org/10.1172/JCI73072>
39 1113
40
41 Mohammadi, H. (2024a). A Genome-wide Association Study based on Pathway
42 1114
43 Analysis Related to the Angle Wing in Pekin Ducks using Whole-Genome
44 1115
45 Sequencing Data. *Research On Animal Production*, 15(2), 32–43.
46 1116
47 <https://doi.org/10.61186/rap.15.2.32>
48 1117
49
50 Muto, A., Ikeda, S., Lopez-Burks, M. E., Kikuchi, Y., Calof, A. L., Lander, A. D., &
51 1118
52 Schilling, T. F. (2014). Nipbl and Mediator Cooperatively Regulate Gene
53 1119
54
55
56
57
58
59
60

- 1
2
3 1120 Expression to Control Limb Development. *PLOS Genetics*, 10(9), e1004671.
4
5 1121 <https://doi.org/10.1371/journal.pgen.1004671>
6
7
8 1122 Nan, J., Yang, S., Zhang, X., Leng, T., Zhuoma, J., Zhuoma, R., Yuan, J., Pi, J.,
9
10 1123 Sheng, Z., & Li, S. (2023). Identification of candidate genes related to highland
11
12 1124 adaptation from multiple Chinese local chicken breeds by whole genome
13
14 1125 sequencing analysis. *Animal Genetics*, 54(1), 55–67.
15
16 1126 <https://doi.org/10.1111/age.13268>
17
18
19 1127 Owen, J. C. (2011). Collecting, processing, and storing avian blood: A review.
20
21 1128 *Journal of Field Ornithology*, 82(4), 339–354. <https://doi.org/10.1111/j.1557->
22
23 1129 9263.2011.00338.x
24
25
26 1130 Pap, P. L., Osváth, G., Daubner, T., Nord, A., & Vincze, O. (2020). Down feather
27
28 1131 morphology reflects adaptation to habitat and thermal conditions across the avian
29
30 1132 phylogeny. *Evolution*, 74(10), 2365–2376. <https://doi.org/10.1111/evo.14075>
31
32
33 1133 Pap, P. L., Vincze, O., Wekerle, B., Daubner, T., Vágási, C. I., Nudds, R. L.,
34
35 1134 Dyke, G. J., & Osváth, G. (2017). A phylogenetic comparative analysis reveals
36
37 1135 correlations between body feather structure and habitat. *Functional Ecology*,
38
39 1136 31(6), 1241–1251. <https://doi.org/10.1111/1365-2435.12820>
40
41
42 1137 Park, H.-B., Kim, J.-W., & Baek, K.-H. (2020). Regulation of Wnt Signaling
43
44 1138 through Ubiquitination and Deubiquitination in Cancers. *International Journal of*
45
46 1139 *Molecular Sciences*, 21(11), Article 11. <https://doi.org/10.3390/ijms21113904>
47
48
49 1140 Pelletier, F., & Coltman, D. W. (2018). Will human influences on evolutionary
50
51 1141 dynamics in the wild pervade the Anthropocene? *BMC Biology*, 16(1), 7.
52
53 1142 <https://doi.org/10.1186/s12915-017-0476-1>
54
55
56
57
58
59
60

1
2
3 1143 Poplin, R., Ruano-Rubio, V., DePristo, M. A., Fennell, T. J., Carneiro, M. O.,
4
5 1144 Auwera, G. A. V. der, Kling, D. E., Gauthier, L. D., Levy-Moonshine, A., Roazen,
6
7 1145 D., Shakir, K., Thibault, J., Chandran, S., Whelan, C., Lek, M., Gabriel, S., Daly,
8
9 1146 M. J., Neale, B., MacArthur, D. G., & Banks, E. (2018). *Scaling accurate genetic*
11
12 1147 *variant discovery to tens of thousands of samples* (p. 201178). bioRxiv.
13
14 1148 <https://doi.org/10.1101/201178>
15
16
17 1149 Pritchard, J. K., & Di Rienzo, A. (2010). Adaptation – not by sweeps alone.
18
19 1150 *Nature Reviews Genetics*, 11(10), 665–667. <https://doi.org/10.1038/nrg2880>
20
21 1151 Prost, S., Armstrong, E. E., Nylander, J., Thomas, G. W. C., Suh, A., Petersen,
22
23 1152 B., Dalen, L., Benz, B. W., Blom, M. P. K., Palkopoulou, E., Ericson, P. G. P., &
24
25 1153 Irestedt, M. (2018). *Genome analyses show strong selection on coloration,*
26
27 1154 *morphological and behavioral phenotypes in birds-of-paradise* (p. 287086).
28
29 1155 bioRxiv. <https://doi.org/10.1101/287086>
30
31
32
33 1156 Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender,
34
35 1157 D., Maller, J., Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007).
36
37 1158 PLINK: A Tool Set for Whole-Genome Association and Population-Based
38
39 1159 Linkage Analyses. *American Journal of Human Genetics*, 81(3), 559–575.
40
41
42 1160 Pyle, P. (2022). *Identification Guide to North American Birds, Part I* (Second
43
44 1161 Edition). Slate Creek Press.
45
46
47 1162 Qi, X., Zhang, Q., He, Y., Yang, L., Zhang, X., Shi, P., Yang, L., Liu, Z., Zhang,
48
49 1163 F., Liu, F., Liu, S., Wu, T., Cui, C., Ouzhuluobu, Bai, C., Baimakangzhuo, Han, J.,
50
51 1164 Zhao, S., Liang, C., & Su, B. (2019). The Transcriptomic Landscape of Yaks
52
53 1165 Reveals Molecular Pathways for High Altitude Adaptation. *Genome Biology and*
54
55
56
57
58
59
60

- 1
2
3 1166 *Evolution*, 11(1), 72–85. <https://doi.org/10.1093/gbe/evy264>
4
5 1167 Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for
6
7 1168 comparing genomic features. *Bioinformatics*, 26(6), 841–842.
8
9
10 1169 <https://doi.org/10.1093/bioinformatics/btq033>
11
12 1170 Ricchetti, I., Taucher, T., Loebick, R., Sin, S. Y. W., Palacios, C., &
13
14 1171 Lamichhaney, S. (2024). Gene expression variation in geographically diverse
15
16 1172 populations of two North American songbird species. *Frontiers in Bird Science*, 3.
17
18 1173 <https://doi.org/10.3389/fbirs.2024.1382657>
19
20
21 1174 Rodriguez, M. D., Bay, R. A., & Ruegg, K. C. (2025). Telomere Length
22
23 1175 Differences Indicate Climate Change-Induced Stress and Population Decline in a
24
25 1176 Migratory Bird. *Molecular Ecology*, 34(21), e17642.
26
27 1177 <https://doi.org/10.1111/mec.17642>
28
29
30 1178 Rogalla, S., Shawkey, M. D., & D'Alba, L. (2022). Thermal effects of plumage
31
32 1179 coloration. *Ibis*, 164(4), 933–948. <https://doi.org/10.1111/ibi.13100>
33
34
35 1180 Romano, A., Séchaud, R., Hirzel, A. H., & Roulin, A. (2019). Climate-driven
36
37 1181 convergent evolution of plumage colour in a cosmopolitan bird. *Global Ecology*
38
39 1182 *and Biogeography*, 28(4), 496–507. <https://doi.org/10.1111/geb.12870>
40
41
42 1183 Romanov, M. N., Abdelmanova, A. S., Fisinin, V. I., Gladyr, E. A., Volkova, N. A.,
43
44 1184 Koshkina, O. A., Rodionov, A. N., Vetokh, A. N., Gusev, I. V., Anshakov, D. V.,
45
46 1185 Stanishevskaya, O. I., Dotsev, A. V., Griffin, D. K., & Zinovieva, N. A. (2023).
47
48 1186 Selective footprints and genes relevant to cold adaptation and other phenotypic
49
50 1187 traits are unscrambled in the genomes of divergently selected chicken breeds.
51
52 1188 *Journal of Animal Science and Biotechnology*, 14(1), 35.
53
54
55
56
57
58
59
60

1
2
3 1189 <https://doi.org/10.1186/s40104-022-00813-0>
4
5 1190 Rundel, P. W., Gibson, A. C., & Sharifi, M. R. (2008). The Alpine Flora of the
6
7 1191 White Mountains, California. *Madroño*, 55(3), 202–215.
8
9
10 1192 <https://doi.org/10.3120/0024-9637-55.3.202>
11
12 1193 Rundel, P. W., & Millar, C. I. (2016). Alpine Ecosystems. In H. Mooney & E.
13
14 1194 Zavaleta (Eds.), *Ecosystems of California* (pp. 613–634). University of California
15
16 Press. [https://www.degruyterbrill.com/document/doi/10.1525/9780520962170-](https://www.degruyterbrill.com/document/doi/10.1525/9780520962170-033/pdf?licenseType=restricted)
17 1195
18 033/pdf?licenseType=restricted
19 1196
20
21 1197 Sadeesh, E. M., Lahamge, M. S., Malik, A., & Ampadi, A. N. (2025). Nuclear
22
23 1198 Genome-Encoded Mitochondrial OXPHOS Complex I Genes in Female Buffalo
24
25 Show Tissue-Specific Differences. *Molecular Biotechnology*, 67(6), 2411–2427.
26 1199
27
28 1200 <https://doi.org/10.1007/s12033-024-01206-6>
29
30
31 1201 Sandoval, L., & Barrantes, G. (2019). Is black plumage an adaptation to high
32
33 1202 elevations in a cosmopolitan bird genus? *Journal of Avian Biology*, 50(7).
34
35 1203 <https://doi.org/10.1111/jav.02041>
36
37
38 1204 Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local
39
40 1205 adaptation. *Nature Reviews Genetics*, 14(11), 807–820.
41
42 1206 <https://doi.org/10.1038/nrg3522>
43
44
45 1207 Schweizer, R. M., Jones, M. R., Bradburd, G. S., Storz, J. F., Senner, N. R.,
46
47 1208 Wolf, C., & Cheviron, Z. A. (2021a). Broad Concordance in the Spatial
48
49 1209 Distribution of Adaptive and Neutral Genetic Variation across an Elevational
50
51 1210 Gradient in Deer Mice. *Molecular Biology and Evolution*, 38(10), 4286–4300.
52
53 1211 <https://doi.org/10.1093/molbev/msab161>
54
55
56
57
58
59
60

- 1212 Sebastianelli, M., Lukhele, S. M., Secomandi, S., de Souza, S. G., Haase, B.,
1213 Moysi, M., Nikiforou, C., Hutfluss, A., Mountcastle, J., Balacco, J., Pelan, S.,
1214 Chow, W., Fedrigo, O., Downs, C. T., Monadjem, A., Dingemanse, N. J., Jarvis,
1215 E. D., Brelsford, A., vonHoldt, B. M., & Kirschel, A. N. G. (2024). A genomic basis
1216 of vocal rhythm in birds. *Nature Communications*, 15(1), 3095.
1217 <https://doi.org/10.1038/s41467-024-47305-5>
- 1218 Seki, R., Li, C., Fang, Q., Hayashi, S., Egawa, S., Hu, J., Xu, L., Pan, H., Kondo,
1219 M., Sato, T., Matsubara, H., Kamiyama, N., Kitajima, K., Saito, D., Liu, Y.,
1220 Gilbert, M. T. P., Zhou, Q., Xu, X., Shiroishi, T., ... Zhang, G. (2017). Functional
1221 roles of Aves class-specific cis-regulatory elements on macroevolution of bird-
1222 specific features. *Nature Communications*, 8(1), 14229.
1223 <https://doi.org/10.1038/ncomms14229>
- 1224 Semenov, G. A., Sonnenberg, B. R., Branch, C. L., Heinen, V. K., Welklin, J. F.,
1225 Padula, S. R., Patel, A. M., Bridge, E. S., Pravosudov, V. V., & Taylor, S. A.
1226 (2024). Genes and gene networks underlying spatial cognition in food-caching
1227 chickadees. *Current Biology*, 34(9), 1930-1939.e4.
1228 <https://doi.org/10.1016/j.cub.2024.03.058>
- 1229 Seutin, G., White, B. N., & Boag, P. T. (1991). Preservation of avian blood and
1230 tissue samples for DNA analyses. *Canadian Journal of Zoology*, 69(1), 82–90.
1231 <https://doi.org/10.1139/z91-013>
- 1232 Shafer, A. B. A., Wolf, J. B. W., Alves, P. C., Bergström, L., Bruford, M. W.,
1233 Brännström, I., Colling, G., Dalén, L., De Meester, L., Ekblom, R., Fawcett, K. D.,
1234 Fior, S., Hajibabaei, M., Hill, J. A., Hoezel, A. R., Höglund, J., Jensen, E. L.,

1
2
3 1235 Krause, J., Kristensen, T. N., ... Zieliński, P. (2015). Genomics and the
4
5 1236 challenging translation into conservation practice. *Trends in Ecology & Evolution*,
6
7 1237 30(2), 78–87. <https://doi.org/10.1016/j.tree.2014.11.009>
8
9
10 1238 Shakya, S. B. (2020). *Genomics and Population History of Black-Headed Bulbul*
11
12 1239 (*Brachypodius Atriceps*) Color Morphs [Ph.D., Louisiana State University and
13
14 1240 Agricultural & Mechanical College].
15
16
17 1241 [https://www.proquest.com/docview/2665131551/abstract/2200EDB181A4591PQ/](https://www.proquest.com/docview/2665131551/abstract/2200EDB181A4591PQ/1)
18
19 1242 1
20
21 1243 Sirkiä, P. M., Virolainen, M., & Laaksonen, T. (2010). Melanin coloration has
22
23 1244 temperature-dependent effects on breeding performance that may maintain
24
25 1245 phenotypic variation in a passerine bird. *Journal of Evolutionary Biology*, 23(11),
26
27 1246 2385–2396. <https://doi.org/10.1111/j.1420-9101.2010.02100.x>
28
29
30 1247 Stettenheim, P. R. (2000). The Integumentary Morphology of Modern Birds—An
31
32 1248 Overview. *American Zoologist*, 40(4), 461–477.
33
34 1249 <https://doi.org/10.1093/icb/40.4.461>
35
36
37 1250 Steven, S. (2021). *Genome-Wide Analyses of Genomic Variation Regulating*
38
39 1251 *Differential Stress Responses in Japanese Quail* [Doctoral dissertation,
40
41 1252 University of Arkansas]. ProQuest Dissertations & Theses.
42
43
44 1253 Stinchcombe, J. R., & Hoekstra, H. E. (2008). Combining population genomics
45
46 1254 and quantitative genetics: Finding the genes underlying ecologically important
47
48 1255 traits. *Heredity*, 100(2), 158–170. <https://doi.org/10.1038/sj.hdy.6800937>
49
50
51 1256 Stoutjesdijk, F. (2003). Birds in the cold: The effects of plumage structure and
52
53 1257 environment on operative temperature, shown by spherical models. *Ornis*
54
55
56
57
58
59
60

- 1258 *Svecica*, 13(4), Article 4. <https://doi.org/10.34080/os.v13.22804>
- 1259 Struk, A. A., Mugon, J., Huston, A., Scholer, A. A., Stadler, G., Higgins, E. T.,
1260 Sokolowski, M. B., & Danckert, J. (2019). Self-regulation and the foraging gene
1261 (PRKG1) in humans. *Proceedings of the National Academy of Sciences*, 116(10),
1262 4434–4439. <https://doi.org/10.1073/pnas.1809924116>
- 1263 Sun, L., Zhou, T., Wan, Q.-H., & Fang, S.-G. (2020). Transcriptome Comparison
1264 Reveals Key Components of Nuptial Plumage Coloration in Crested Ibis.
1265 *Biomolecules*, 10(6), Article 6. <https://doi.org/10.3390/biom10060905>
- 1266 Sun, Y., Li, Y., Jiang, X., Wu, Q., Lin, R., Chen, H., Zhang, M., Zeng, T., Tian, Y.,
1267 Xu, E., Zhang, Y., & Lu, L. (2024). Genome-wide association study identified
1268 candidate genes for egg production traits in the Longyan Shan-ma duck. *Poultry*
1269 *Science*, 103(9), 104032. <https://doi.org/10.1016/j.psj.2024.104032>
- 1270 Symonds, M. R. E., & Tattersall, G. J. (2010). Geographical variation in bill size
1271 across bird species provides evidence for Allen's rule. *The American Naturalist*,
1272 176(2), 188–197. <https://doi.org/10.1086/653666>
- 1273 Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R.,
1274 Gable, A. L., Fang, T., Doncheva, N. T., Pyysalo, S., Bork, P., Jensen, L. J., &
1275 von Mering, C. (2023). The STRING database in 2023: Protein-protein
1276 association networks and functional enrichment analyses for any sequenced
1277 genome of interest. *Nucleic Acids Research*, 51(D1), D638–D646.
1278 <https://doi.org/10.1093/nar/gkac1000>
- 1279 Tam, V., Patel, N., Turcotte, M., Bossé, Y., Paré, G., & Meyre, D. (2019).
1280 Benefits and limitations of genome-wide association studies. *Nature Reviews*

1
2
3 1281 *Genetics*, 20(8), 467–484. <https://doi.org/10.1038/s41576-019-0127-1>
4
5 1282 Tattersall, G. J., Arnaout, B., & Symonds, M. R. E. (2017). The evolution of the
6
7 1283 avian bill as a thermoregulatory organ. *Biological Reviews*, 92(3), 1630–1656.
8
9
10 1284 <https://doi.org/10.1111/brv.12299>
11
12 1285 Thomas, P. D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P., & Mi,
13
14 1286 H. (2022). PANTHER: Making genome-scale phylogenetics accessible to all.
15
16
17 1287 *Protein Science*, 31(1), 8–22. <https://doi.org/10.1002/pro.4218>
18
19 1288 Tian, S., Zhou, X., Phuntsok, T., Zhao, N., Zhang, D., Ning, C., Li, D., & Zhao, H.
20
21 1289 (2020). Genomic Analyses Reveal Genetic Adaptations to Tropical Climates in
22
23 1290 Chickens. *iScience*, 23(11). <https://doi.org/10.1016/j.isci.2020.101644>
24
25
26 1291 Twining, H. (1940). Foraging Behavior and Survival in the Sierra Nevada Rosy
27
28 1292 Finch. *The Condor*, 42(1), 64–72. <https://doi.org/10.2307/1364320>
29
30
31 1293 Twumasi, G., Wang, H., Xi, Y., Qi, J., Li, L., Bai, L., & Liu, H. (2024). Genome-
32
33 1294 Wide Association Studies Reveal Candidate Genes Associated with
34
35 1295 Pigmentation Patterns of Single Feathers of Tianfu Nonghua Ducks. *Animals*,
36
37 1296 14(1), Article 1. <https://doi.org/10.3390/ani14010085>
38
39
40 1297 Urban, M. C. (2018). Escalator to extinction. *Proceedings of the National*
41
42 1298 *Academy of Sciences*, 115(47), 11871–11873.
43
44
45 1299 <https://doi.org/10.1073/pnas.1817416115>
46
47 1300 Videla Rodriguez, E. A., Mitchell, J. B. O., & Smith, V. A. (2022). A Bayesian
48
49 1301 network structure learning approach to identify genes associated with stress in
50
51 1302 spleens of chickens. *Scientific Reports*, 12(1), 7482.
52
53
54 1303 <https://doi.org/10.1038/s41598-022-11633-7>
55
56
57
58
59
60

- 1
2
3 1304 Wang, C., Li, S., Yu, W., Xin, Q., Li, C., Feng, Y., Peng, X., & Gong, Y. (2011).
4
5 1305 Cloning and expression profiling of the VLDLR gene associated with egg
6
7 1306 performance in duck (*Anas platyrhynchos*). *Genetics Selection Evolution*, 43(1),
8
9 1307 29. <https://doi.org/10.1186/1297-9686-43-29>
10
11
12 1308 Ward, S., Rayner, J. M., Möller, U., Jackson, D. M., Nachtigall, W., & Speakman,
13
14 1309 J. R. (1999). Heat transfer from starlings *sturnus vulgaris* during flight. *Journal of*
15
16 1310 *Experimental Biology*, 202 (Pt 12), 1589–1602.
17
18 1311 <https://doi.org/10.1242/jeb.202.12.1589>
19
20
21 1312 Weeks, B. C., Harvey, C., Tobias, J. A., Sheard, C., Zhou, Z., & Fouhey, D. F.
22
23 1313 (2025). Longer Wing Bones in Warmer Climates Suggest a Role of
24
25 1314 Thermoregulation in Bird Wing Evolution. *Global Ecology and Biogeography*,
26
27 1315 34(4), e70033. <https://doi.org/10.1111/geb.70033>
28
29
30 1316 Weeks, B. C., Willard, D. E., Zimova, M., Ellis, A. A., Witynski, M. L., Hennen, M.,
31
32 1317 & Winger, B. M. (2020). Shared morphological consequences of global warming
33
34 1318 in North American migratory birds. *Ecology Letters*, 23(2), 316–325.
35
36 1319 <https://doi.org/10.1111/ele.13434>
37
38
39 1320 Whitlock, M. C., & Lotterhos, K. E. (2015). Reliable Detection of Loci Responsible
40
41 1321 for Local Adaptation: Inference of a Null Model through Trimming the Distribution
42
43 1322 of F(ST). *The American Naturalist*, 186 Suppl 1, S24-36.
44
45 1323 <https://doi.org/10.1086/682949>
46
47
48 1324 Wolf, B. O., & Walsberg, G. E. (2000). The Role of the Plumage in Heat Transfer
49
50 1325 Processes of Birds1. *American Zoologist*, 40(4), 575–584.
51
52 1326 <https://doi.org/10.1093/icb/40.4.575>
53
54
55
56
57
58
59
60

1
2
3 1327 Wu, Y., Yan, Y., Zhao, Y., Gu, L., Wang, S., & Johnson, D. H. (2021). Genomic
4
5 1328 bases underlying the adaptive radiation of core landbirds. *BMC Ecology and*
6
7 1329 *Evolution*, 21(1), 162. <https://doi.org/10.1186/s12862-021-01888-5>
8
9
10 1330 Xiao, J., Wu, Y., Chen, R., Lin, Y., Wu, L., Tian, W., & Liu, L. (2008). Expression
11
12 1331 of Pcp4 gene during osteogenic differentiation of bone marrow mesenchymal
13
14 1332 stem cells in vitro. *Molecular and Cellular Biochemistry*, 309(1), 143–150.
15
16
17 1333 <https://doi.org/10.1007/s11010-007-9652-x>
18
19 1334 Xie, J., Tang, L., Lu, L., Zhang, L., Xi, L., Liu, H.-C., Odle, J., & Luo, X. (2014).
20
21 1335 Differential Expression of Heat Shock Transcription Factors and Heat Shock
22
23 1336 Proteins after Acute and Chronic Heat Stress in Laying Chickens (*Gallus gallus*).
24
25
26 1337 *PLOS ONE*, 9(7), e102204. <https://doi.org/10.1371/journal.pone.0102204>
27
28 1338 Xiong, D.-H., Liu, X.-G., Guo, Y.-F., Tan, L.-J., Wang, L., Sha, B.-Y., Tang, Z.-H.,
29
30 1339 Pan, F., Yang, T.-L., Chen, X.-D., Lei, S.-F., Yerges, L. M., Zhu, X.-Z., Wheeler,
31
32 1340 V. W., Patrick, A. L., Bunker, C. H., Guo, Y., Yan, H., Pei, Y.-F., ... Deng, H.-W.
33
34 1341 (2009). Genome-wide Association and Follow-Up Replication Studies Identified
35
36 1342 ADAMTS18 and TGFBR3 as Bone Mass Candidate Genes in Different Ethnic
37
38 1343 Groups. *The American Journal of Human Genetics*, 84(3), 388–398.
39
40 1344 <https://doi.org/10.1016/j.ajhg.2009.01.025>
41
42 1345 Yan, C., Zhang, Z.-Y., Lv, Y., Wang, Z., Jiang, K., & Li, J.-T. (2022). Genome of
43
44 1346 *Laudakia sacra* Provides New Insights into High-Altitude Adaptation of
45
46 1347 Ectotherms. *International Journal of Molecular Sciences*, 23(17), Article 17.
47
48 1348 <https://doi.org/10.3390/ijms231710081>
49
50 1349 Yeaman, S. (2015). Local Adaptation by Alleles of Small Effect. *The American*
51
52
53
54
55
56
57
58
59
60

- 1
2
3 1350 *Naturalist*, 186 Suppl 1, S74-89. <https://doi.org/10.1086/682405>
4
5 1351 Zhang, C., Liu, C., Zhang, J., Zheng, L., Chang, Q., Cui, Z., & Liu, S. (2022).
6
7 1352 Analysis on the desert adaptability of indigenous sheep in the southern edge of
8
9 Taklimakan Desert. *Scientific Reports*, 12(1), 12264.
10 1353
11
12 1354 <https://doi.org/10.1038/s41598-022-15986-x>
13
14 1355 Zhang, W., Fan, Z., Han, E., Hou, R., Zhang, L., Galaverni, M., Huang, J., Liu,
15
16 H., Silva, P., Li, P., Pollinger, J. P., Du, L., Zhang, X., Yue, B., Wayne, R. K., &
17 1356
18 Zhang, Z. (2014). Hypoxia Adaptations in the Grey Wolf (*Canis lupus chanco*)
19 1357
20 from Qinghai-Tibet Plateau. *PLOS Genetics*, 10(7), e1004466.
21 1358
22 <https://doi.org/10.1371/journal.pgen.1004466>
23 1359
24
25 Zhang, X., Yang, F., Zhu, T., Zhao, X., Zhang, J., Wen, J., Zhang, Y., Wang, G.,
26 1360
27 Ren, X., Chen, A., Wang, X., Wang, L., Lv, X., Yang, W., Qu, C., Wang, H., Ning,
28 1361
29 Z., & Qu, L. (2024). Whole genome resequencing reveals genomic regions
30 1362
31 related to red plumage in ducks. *Poultry Science*, 103(6), 103694.
32 1363
33 <https://doi.org/10.1016/j.psj.2024.103694>
34 1364
35
36 Zhang, X., Zhu, T., Wang, L., Lv, X., Yang, W., Qu, C., Li, H., Wang, H., Ning, Z.,
37 1365
38 & Qu, L. (2023). Genome-Wide Association Study Reveals the Genetic Basis of
39 1366
40 Duck Plumage Colors. *Genes*, 14(4), Article 4.
41 1367
42 <https://doi.org/10.3390/genes14040856>
43 1368
44
45 Zheng, S., Ouyang, J., Liu, S., Tang, H., Xiong, Y., Yan, X., & Chen, H. (2023).
46 1369
47 Genomic signatures reveal selection in Lingxian white goose. *Poultry Science*,
48 1370
49 102(1), 102269. <https://doi.org/10.1016/j.psj.2022.102269>
50 1371
51
52 Zhou, X., Carbonetto, P., & Stephens, M. (2013). Polygenic Modeling with
53 1372
54
55
56
57
58
59
60

1
2
3 1373 Bayesian Sparse Linear Mixed Models. *PLOS Genetics*, 9(2), e1003264.
4
5 1374 <https://doi.org/10.1371/journal.pgen.1003264>
6
7
8 1375 Zhuang, Z.-X., Chen, S.-E., Chen, C.-F., Lin, E.-C., & Huang, S.-Y. (2019).
9
10 1376 Genome-wide association study on the body temperature changes of a broiler-
11
12 1377 type strain Taiwan country chickens under acute heat stress. *Journal of Thermal*
13
14
15 1378 *Biology*, 82, 33–42. <https://doi.org/10.1016/j.jtherbio.2019.03.007>
16
17 1379 Zimova, M., Weeks, B. C., Willard, D. E., Giery, S. T., Jirinec, V., Burner, R. C., &
18
19 1380 Winger, B. M. (2023). Body size predicts the rate of contemporary morphological
20
21 1381 change in birds. *Proceedings of the National Academy of Sciences of the United*
22
23
24 1382 *States of America*, 120(20), e2206971120.
25
26 1383 <https://doi.org/10.1073/pnas.2206971120>
27
28
29

30 1384 **Data Accessibility**
31
32 1385 Raw sequence data have been deposited in the NCBI Sequence Read Archive under
33 1386 BioProject accession PRJNA1297057.
34
35 1387 Statistical analysis, GWAS and F_{ST} results are available at
36 1388 <http://datadryad.org/share/DqvYM4bZd7lZHEXy5O9FGkrQ86Q90YoGz3ZOAJoLHmY>
37
38 1389 Code and scripts used for processing and analysis of sequence data are available at
39 1390 <https://github.com/erigande/mega-non-model-wgs-snakeflow/>. All other scripts as well
40
41 1391 as sample metadata are available on GitHub at
42 1392 https://github.com/ecnrobertson/GCRF_Pub.
43
44


45 1393 **Benefit-Sharing Statement**
46
47 1394 Benefits Generated: Benefits from this research accrue from the sharing of our data and
48 1395 results on public databases as described above.
49
50

51 1396 **Statement of Contributions**
52
53 1397 E.C.N.R, T.B., E.Z., K.C.R conceived of the study and E.C.N.R. wrote the majority of
54 1398 the manuscript and oversaw the majority of the analysis. C.M.B. contributed to the
55 1399 bioinformatic analysis. S.D. performed and conceived of the methodology for all feather
56
57
58
59
60


morphology measurements. T.B. and E.Z. contributed ideas and performed all field sampling. K.C.R., E.Z. and M.B.H. acquired funding. All authors discussed the results and contributed to the final manuscript.

Figures

A Sampling Scheme



B GWAS Methods



C F_{ST} Methods




Figure 1. A) A map of California highlighting the sampling location and design for two populations of Grey-crowned Rosy-Finch, Piute Pass and White Mountain. Elevation is noted for each site along with an elevation transect. B) Overview of methods for GWAS analysis. Morphological measurements were corrected for the effects of sex, age, and body size before being input into BSLMM and LMM models. C) Overview of the methods for the F_{ST} analysis. The output of both analyses were analyzed to identify significant SNPs and related genes.




Figure 2. Violin plots of beak morphology raw measurement: A) beak width, B) nares length, and C) beak depth. Overlaid are the corrected mean trait measurements shown with a crossbar. Significance of the effect of site on the morphological trait is shown for each comparison. Piute Pass is the warm site and White Mountain is the cold site. We found that nares were longer while beak widths and depths were smaller in the warmer environment.

Figure 3. A) Photograph showing a body feather from a Sierra Nevada Gray-crowned Rosy-Finch captured at White Mountain. Highlighted is a microscopic image of the pennaceous and plumulaceous barbule structure. B) Violin plots of node density raw measurements, with illustrations above of the structure of plumulaceous barbule nodes. Overlaid on the plots are the corrected mean trait measurements shown with a crossbar. Significance of the effect of site on the morphological trait is shown for each comparison.





Figure 4. Manhattan plot depicting the results of the LMM analysis on beak depth (A), wing chord (B), and pennaceous barbule length (C) with $-\log_{10}(\text{p-value})$ on the y-axis and the mapped Zebra Finch chromosome location on the x-axis. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted. Also plotted are the results of the per-site F_{ST} results (D), using the 90th quantile threshold of 0.235. Corresponding genes are also shown and functional categories are highlighted by color category. See Table 1 for more

1505 functional details.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch

Running Title: Local adaptation in an alpine finch

Authors: Robertson, Erica C.N.^{1*}, Brown, Timothy M.², Deitch, Sophie¹, Bossu, Christine M.¹, Zavaleta, Erika S.², Hooten, Mevin B.³, Ruegg, Kristen C.¹

Affiliations:

^{1.} Department of Biology, Colorado State University, Fort Collins, CO, USA

^{1.} ^{*}Corresponding author: ericacnrobertson@gmail.com

^{2.} ^{2.} Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, California, USA

^{3.} ^{3.} Department of Statistics and Data Sciences, The University of Texas at Austin, Austin, Texas

Corresponding Author: ~~Erica C.N. Robertson~~

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Understanding patterns and mechanisms underlying local adaptation is becoming increasingly important for species conservation amid anthropogenically driven environmental change. Alpine systems are experiencing particularly intense pressure from environmental change resulting from increased rates of warming and corresponding loss of snow and ice. We integrate morphological and genetic analyses to identify traits important for local adaptation in one of the highest elevation breeding birds in North America, the Sierra Nevada Gray-crowned Rosy-Finch. We performed an in-depth analysis of how traits with known links to thermoregulation in birds such as wing length, bill size, and feather microstructure vary between two populations at sites with contrasting climate and environmental conditions. We identified loci underlying these traits using a genome-wide association study and further examined regions of the genome related to altitude adaptation and cold tolerance using F_{ST} outlier tests. Together, these results indicate that temperature, food availability, and alpine landscape features may impose multifaceted and potentially conflicting selective pressures on morphological traits important to adaptation in alpine birds. Overall, this work represents one of the first in-depth analyses of the genetic basis of adaptation in an alpine specialist songbird.

Keywords

local adaptation, conservation genomics, genome-wide association study, F_{ST} outlier test, alpine

Introduction

Local adaptation occurs when natural selection favors traits that confer fitness advantages in specific habitats (Endler, 1977; Kawecki & Ebert, 2004; Savolainen et al., 2013). A species' adaptive capacity determines the extent to which it will be able to respond to rapid environmental change (Forester et al., 2022, 2023; Lande & Shannon, 1996; Meek et al., 2023). As a result, identifying genetic and morphological traits involved in local adaptation is a key component of effective species conservation planning (Hoban et al., 2016; Meek et al., 2023). Historically, assessments of the capacity for local adaptation relied on reciprocal transplant experiments that allowed researchers to link variation in putatively ecologically important traits to differences in fitness across environmental gradients (Blanquart et al., 2013; Clausen et al., 1941; Kawecki & Ebert, 2004; Savolainen et al., 2013). However, recent advances in genomic methods have made it possible to detect signatures of selection across the genome and to associate these with environmental variation and ecologically important morphological variation (Capblancq et al., 2020; Hoban et al., 2016; Lotterhos & Whitlock, 2015). We leveraged recent advances in genomic methods to identify morphological and genetic traits important to local adaptation in an understudied alpine bird system.

Many montane species undergo upward range shifts with climate change (Mamantov et al., 2021). However, for those species already living near the top of a mountain system, there are constraints that may limit the ability to track their niche, a phenomenon termed the "escalator to extinction" (Freeman et al., 2018; Urban, 2018). Alternatively, local adaptation can impact adaptive responses to the environment and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

allow species to persist despite a restricted capacity for range shifts (Aitken et al., 2008; Capblancq et al., 2020). A first step toward improving our understanding of how high-altitude species will respond to climate change is to identify morphological traits and corresponding regions of the genome involved in adaptation to extreme alpine environments (Blanquart et al., 2013; Pritchard & Di Rienzo, 2010).

Birds living in alpine regions are exposed to extreme environments throughout the year, including variable snow cover, hypoxic conditions, and dramatic temperature changes (Grabherr et al., 2010; Körner, 2003). As a result, traits that contribute directly to thermoregulation are expected to be under strong selective pressure in alpine species. Previous work has shown climate-linked traits often follow [ecogeographical rules such as Bergmann's and Allen's rule](#). ~~Allen's rule, a hypothesis that~~ posits [that animals adapted to colder environments will have shorter appendages than animals adapted to warm climates \(Allen, 1877\)](#). In birds, this rule predicts that beaks and wings will be longer in warmer environments to facilitate greater heat dissipation across a larger surface area, and shorter in colder environments to facilitate heat conservation (Greenberg et al., 2012; Lewden et al., 2023; Symonds & Tattersall, 2010; Tattersall et al., 2017; Ward et al., 1999; Weeks et al., 2020, 2025). ~~For this reason, we predicted that birds in high-altitude alpine systems will have shorter beaks and smaller wings.~~ [Bergmann's rule, which states that body size tends to decrease as temperature increases, has also been supported in birds \(Ashton, 2002; He et al., 2023\)](#). [Following these rules, we predicted that birds will have shorter beaks and smaller wings as well as larger overall body sizes in higher, colder environments.](#)

An additional aspect of beak morphology that should be investigated in the context of thermoregulation is nares length. Nares are the nostril of the bird and are located at the top of the beak. Grinnell (1913) documented subspecific divergence in *Leucosticte tephrocotis dawsoni*, including differences in beak length and bill depth from the nostril. While nares size per se was not the focus, these measurements provide a morphological basis for interpreting variation in nasal exposure and potential heat exchange, particularly in the context of ecological differences between alpine sites. While there has not, to our knowledge, been work directly linking nares size to thermoregulatory function, this may be an important and understudied aspect of beak morphology as it relates to thermoregulation in birds. Following the logic for broader beak morphology, we can expect to see larger nares in the warmer environments to facilitate the dumping of excess heat.

Although less studied in the context of local adaptation, feather microstructure may also be important to avian species' adaptation to environmental variation in temperature. While substantial evidence suggests feather microstructure plays a key role in determining the thermoregulatory capacity of a bird's plumage (Barve et al., 2021; Stettenheim, 2000; Stoutjesdijk, 2003; Wolf & Walsberg, 2000), there is some confusion over how environmental variation may drive feather microstructure differences (D'alba et al., 2017; Koskenpato et al., 2016; Lei et al., 2002; Pap et al., 2017, 2020). In one of the few studies that has been conducted at the population level, Koskenpato et al. (2016) found that tawny owls living in colder environments have significantly denser plumulaceous contour feathers than those living in warmer environments. This suggests that feather density may facilitate heat retention. In contrast, however, Pap et al. (2017)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

found that European bird species wintering in colder areas had less dense feathers than those wintering in warmer places. Thus, there is a clear need for studies specifically linking environmental variation with population-level differences in feather microstructure.

Recent advances in population genomics have enabled the detection of loci involved in local adaptation, even in non-model and wild species (Faria et al., 2014). Among these, F_{ST} outlier tests are commonly used to identify genomic regions that exhibit elevated population differentiation, which may signal divergent selection (Hoban et al., 2016). However, many adaptive traits are likely polygenic, involving small effects at numerous loci (Pritchard & Di Rienzo, 2010; Yeaman, 2015), and such loci will not be picked up by F_{ST} outlier tests. To address this, genome-wide association studies (GWAS) can be used to link more subtle genetic variation to phenotypic variation in ecologically important traits suspected to be under selection (Stinchcombe & Hoekstra, 2008). Taken together, combining approaches that detect both population-level divergence and polygenic trait architecture offers a powerful framework for investigating local adaptation.

We investigated the potential for local adaptation in the [Sierra Nevada](#) Gray-crowned Rosy-Finch (*Leucosticte tephrocotis dawsoni*), one of the highest-breeding songbirds in North America. [Of the roughly four groups of rosy-finch found in North America, this subspecies is more localized, found only in California's alpine, and is a short-distance migrant. In comparison, the broader Gray-crowned Rosy-Finch species range extends along the entire western edge of North America from California to Alaska.](#) Despite long-standing interest (Grinnell, 1913, 1917; Twining, 1940), research

on this subspecies has been limited due to the difficulty of accessing its extreme nesting sites located in the rocky cliff faces of the Sierra Nevada and White Mountains of California, more than 3,000 meters above sea level. Additionally, this subspecies may act as a strong model of local adaptation due to the variation in habitat experienced within a restricted range. To help fill this knowledge gap we conducted morphological and genomic analysis of two populations breeding at distinct contrasting places in the elevational and thermal range. Piute Pass, located in the Sierra Nevada, is lower in elevation and warmer than the White Mountains (Fig. 1A). To identify traits potentially involved in local adaptation, we quantified variation in beak and feather morphology between the two populations. Based on Allen's rule, we predicted that birds in the warmer Piute Pass population would have longer wings and larger beaks. Following Bergmann's rule, we predicted that birds in Piute Pass would also have smaller overall body size. Further, we predicted that birds in the colder White Mountains location would exhibit feather traits associated with increased insulation, specifically, longer barbules and higher node density. Lastly, we used genome-wide association studies (GWAS) to identify genes linked to feather and bill morphology and F_{ST} outlier tests to identify other potentially important sources of genetic variation (Fig 1B-C). Overall, our findings provide new insights into the process of local adaptation in an extreme alpine specialist bird and provide key baseline information on the capacity for adaptation in the face of global environmental change.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

132 **Methods**

133 **Site Selection**

134 Our study compares Rosy-Finch populations from two sites: [one](#) in the Sierra
135 Nevada and [one in the](#) White Mountains. [Sierra Gray-crowned Rosy-Finches occupy](#)
136 [alpine habitats from approximately 2,750–4,000 m across California \(Brown et al., in](#)
137 [review\), and our study sites span a substantial portion of this range.](#) The Piute Pass
138 location is in the Sierra Nevada at [3,470–3,625m](#) in elevation and has a warmer average
139 temperature during the [breeding season warmest month](#) of [8°C \(with a range of 3.8–](#)
140 [13.0°C\)](#). The White Mountain location, near White Mountain Peak, is higher in elevation
141 at [4,285–4,344m](#) and colder with a mean average temperature during the [Rosy-Finch](#)
142 [breeding season of 6°C \(with a range of 1.5–10.7°C\) \(extracted from AdaptWest Project,](#)
143 [2022, see supplemental methods and SupFig1\).](#) Because it lies in the Sierra Nevada
144 rain shadow, the White Mountain site receives roughly one-third the precipitation of
145 areas at comparable elevations in the Sierras (Rundel et al., 2008). Between the two
146 sites, Piute Pass receives more precipitation as snow and has a more persistent
147 snowpack than the White Mountain location. Additionally, aquatic environments such as
148 streams, lakes and ephemeral pools are more abundant in the Sierra Nevada than the
149 White Mountains (Rundel et al., 2008). The alpine ecosystems of these mountains can
150 be delineated as communities occurring above tree line (Rundel & Millar, 2016).
151 Environmental stressors such as extreme winter temperatures, short growing seasons,
152 high winds, low partial pressures of O₂, and limited water availability are characteristic of
153 these alpine environments (Grabherr et al., 2010; Rundel & Millar, 2016). Separating
154 the western Sierra Nevada Mountains and the eastern White Mountains is the Owens
155 Valley, a subregion of the Great Basin Desert (Belnap et al., 2016).

Sample Collection

A total of 171 [Sierra Nevada](#) Gray-crowned Rosy-Finches were ~~captured~~collected using potter traps ~~at~~ White Mountains (n=98) and Piute Pass (n=73) in June and July, the peak of the species' breeding season (MacDougall-Shackleton et al., 2000). [Captures within each mountain range occurred across multiple locations within approximately 1 km² at each site \(White Mountain or Piute Pass\) to maximize sample sizes while maintaining site-specific environmental characteristics. The sampling thus, in total, represents only these two sites. Not all morphology was able to be collected for every bird,](#) resulting in slight variation in the types of data collected for each individual [\(see details in supplemental methods, see sample sizes for each trait in SupTable 1\)](#). Blood samples were collected from 150 of the captured birds using the brachial wing vein and stored in Queen's lysis buffer at room temperature (Owen, 2011; Seutin et al., 1991). At the same time, morphological measurements - tarsus length, wing chord, beak width, beak length, beak depth, ~~and~~nare length, ~~and~~mass - were also collected from 154 of the birds using an electronic caliper and 5-10 body feathers were collected from the breast of all individuals. [Additionally, birds were aged and sexed following the guidelines established in Pyle \(2022\) \(see supplemental methodology\)](#). Birds were banded to allow for identification of the individuals and ~~recognize~~to avoid recaptures, then released. All handling and banding of birds was done following the guidelines and protocols of the U.S Geological Survey (USGS) Bird Banding Laboratory (BBL) and ~~complied~~comply with permits and permissions from federal and state agencies. [All measurements were taken by two researchers \(TB and EZ\), with one \(TB\) collecting approximately 80% of the data. Because the majority of measurements were](#)

Commented [1]: Make sure this citation is included

1
2
3
4
5
6
7
8
9
10 179 [taken by a single observer, and all followed the same standardized protocol, observer](#)
11 180 [identity was not included as a covariate in subsequent analyses.](#)
12

13
14 181 **Feather Microstructure Measurements**

15 182 We measured the microstructure of three feathers for each individual. For each
16
17 183 feather, three photos were taken using an Olympus BX51 Microscope: one photo each
18
19 184 of an unbroken pennaceous and plumulaceous barb from the center of each region was
20
21 185 taken at 4x objective, and one photo of a plumulaceous barbule was taken at 10x
22
23 186 objective [\(Fig 3A\)](#). Structures were then measured from the photos using the
24
25 187 segmented line tool in ImageJ. For the pennaceous and plumulaceous barbule length,
26
27 188 we measured one barbule from each region starting from the distal tip to where the
28
29 189 barbule meets the barb. For the pennaceous and plumulaceous barbule density, we
30
31 190 drew a 0.5mm line along the middle of a barb, counting the number of barbules along
32
33 191 that line. And finally, for plumulaceous node density, we drew a 0.2mm line along the
34
35 192 center of a barbule, counting the number of nodes along that line. ~~[\(Fig 2A\)](#)~~. Each
36
37 193 replicate measure for a feather trait was averaged per individual prior to downstream
38
39 194 analysis. [Approximately 95% of feathers were measured by one researcher \(SD\) and](#)
40
41 195 [observer was not included as a covariate in subsequent analyses.](#)

40 196 **Statistical analysis for local adaptation**

41 197 Body size and sex can confound morphological comparisons among individual
42
43 198 birds (D. C. Adams et al., 2020). To account for these effects [in our investigation of](#)
44
45 199 [Allen's rule](#), we included sex and tarsus length ~~[\(a proxy for body size\)](#)~~ as predictor
46
47 200 variables in our models. [We used tarsus length as an index of body size because it](#)
48
49 201 [correlates strongly with mass but is less affected by short-term fluctuations in condition;](#)
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10 202 [this approach has been used in recent large-scale morphological studies \(Demery et al.,](#)
11 203 [2021; Weeks et al., 2020; Zimova et al., 2023\).](#) As additional validation of tarsus as a
12 204 [proxy size for body size, we generated a PC axis that incorporated multiple allometric](#)
13 205 [traits and found that this was highly correlated \(0.7\) with tarsus \(see supplement for](#)
14 206 [additional details\). We proceeded with tarsus as our proxy for body size in all further](#)
15 207 [analyses. To assess trait variation between sites, we used both univariate and](#)
16 208 [multivariate approaches \(see SupTable 2-3 for a summary of trait values\).](#) We began by
17 209 [constructing](#) For each trait, we constructed linear models with tarsus length, sex, age,
18 210 and site code as predictors and used AIC [weightstests](#) to identify the combination of
19 211 variables that best explained trait variation. Traits for which site code was a significant
20 212 predictor were considered significantly different between the two populations. [We opted](#)
21 213 [not to use a Bonferroni correction for the univariate models as each model had a](#)
22 214 [specific hypothesis associated with it. All p-values for site code predictors were](#)
23 215 [corrected for multiple testing using a Bonferroni correction, and corrected p-values are](#)
24 216 [reported.](#) We used *emmeans* (Lenth, 2025) to calculate average trait values while
25 217 controlling for predictors other than site code, and these adjusted means were overlaid
26 218 on the raw data in a violin plot (Fig 2 & 3B). [To test our hypotheses derived from](#)
27 219 [Bergmann's rule, we compared both tarsus length and a principal component axis](#)
28 220 [representing overall body size using linear models. Each model included age and sex](#)
29 221 [as covariates to account for potential confounding effects on trait variation. We then](#)
30 222 [assessed whether body size differed significantly between the two populations by](#)
31 223 [including site as a predictor.](#)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

224 We also ~~conducted~~include a multivariate analysis to account for potential
225 correlations ~~among~~between traits in our investigation of Allen's rule. For our multivariate
226 approach we~~re~~ incorporated all body morphology traits (wing chord, nare length, beak
227 depth, beak width, beak length) into a MANOVA model with site, age, sex, and tarsus
228 as predictors. The same analysis was ~~performed~~run with all of the feather traits
229 (pennaceous barbule density and length, plumulaceous barbule density, length, and
230 node density) together.

22
23
24
25
26
27
28
29
30
31
32
33

231 **Sequencing and Genotyping**

23
24
25
26
27
28
29
30
31
32
33

232 DNA was extracted from blood samples of 150 individuals using Qiagen's
233 DNeasy Blood and Tissue Extraction Kit protocol and quantified with the Qubit dsDNA
234 HS Assay Kit. We used a modified version of Illumina's Nextera Library Preparation
235 protocol to prepare WGS libraries and pooled the libraries by equal mass before
236 sequencing. The resulting libraries were sequenced on 2 lanes of an Illumina NovaSeq
237 6000 at Novogene Corporation.

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

238 To process the raw sequence reads and detect variants, we utilized Snakemake
239 (Molder et al. 2021), a workflow management system that provides efficiency,
240 adaptability, and reproducibility. The pipeline we adapted to our species can be found
241 on Github (<https://github.com/eriqande/mega-non-model-wgs-snakeflow/>). [To](#)
242 [summarize the workflow, s](#)Sequence data were trimmed using fastp (S. Chen et al.,
243 2018) to remove adaptor sequences and polyG tails using a sliding window, and then
244 aligned to a Brown-capped Rosy-Finch (*Leucosticte australis*) reference genome
245 (GenBank: GCA_025504685.1) using Burrows-Wheeler Aligner software (H. Li &
246 Durbin, 2009). We marked PCR duplicates using ~~SAMtools~~SAMtools (Danecek et al.,

2021) and read groups (sample, lane, library) were added using [Picard](#) (<http://broadinstitute.github.io/picard>). Individual coverage was estimated using [SAMtools](#), and, given the range of coverage (7.4-19X), we downsampled bam files to 10x using the SAMtools subsample function (Danecek et al., 2021). Individual gvcf files were created using [the](#) GATK's HaplotypeCaller (Poplin et al., 2018) with high base quality score filters (--min-base-quality-score 33 --minimum-mapping-quality 20) to remove batch effects (Lou & Therikildsen, 2022). We then parallelized the calling of genotypes across 3 million bp regions across the genome using the GenomicsDBImport and GenotypeGVCFs functions (Auwerda & O'Connor, 2020; McKenna et al., 2010). GATK versions above 4.0 call missing data as homozygous reference, which can bias downstream analysis. To avoid this, all loci with a depth of zero were manually marked as missing. To remove systematic errors according to GATK variant quality score recalibration (VQSR) best practices, we hard filtered variants with the following parameters: [StrandOddsRatio](#) (>3.0), [FisherStrand](#) (>60.0), [MappingQuality](#) (>40.0), and [Quality by Depth](#) (>2.0). We further filtered [for](#) Single Nucleotide Polymorphisms (SNPs), removing indels, filtering for missingness (<80.50%), allele frequencies (>0.05 and <0.95), depth (>4x), quality (>30.0) using BCFtools (Danecek et al., 2021). After filtering, the resulting high-quality SNPs were passed through BEAGLE 4.1 (Browning & Browning, 2016) to impute missing genotypes.

Population genetic structure analysis

Population structure was assessed using a Principal Component Analysis (PCA) followed by ADMIXTURE (Alexander et al., 2009). SNPs identified from sequence data were pruned for linkage disequilibrium in PLINK (Purcell et al., 2007) using a 50kb

1
2
3
4
5
6
7
8
9
10 270 window, 10 [SNPbp](#) window step size, and a R^2 threshold of 0.[24](#). Eigenvalues for the
11 271 PCA were then generated based on the pruned SNPs using PLINK and the first two
12 272 principal components (PC) were plotted in R. To further confirm a lack of population
13 273 structure ADMIXTURE was used to test $k=1$ -[32](#). The results were plotted using R and
14 274 the cv values were evaluated.

18
19 275 **Genome-Wide Association Study**

20 276 Two GWAS models were implemented on all traits using GEMMA (Zhou et al.,
21 277 2013); a Bayesian sparse linear model, BSLMM, and a univariate linear model, ULMM.
22 278 These GWAS approaches allow us to detect SNPs associated with a given trait while
23 279 explicitly accounting for population structure and relatedness through the incorporation
24 280 of random effects whose correlation structure is characterized by a relatedness matrix.
25 281 Input files were generated with PLINK (Purcell et al., 2007). We implemented a
26 282 Bayesian sparse linear mixed model (BSLMM) using 500,000 MCMC burn-in iterations
27 283 that were discarded and 5 million MCMC iterations that were saved. We then filtered for
28 284 SNPs that had a mean posterior inclusion probability (PIP) above 0.01. [\(Fig 4 A & B\)](#). A
29 285 univariate linear mixed model (ULMM) employing the Wald test was also conducted
30 286 [\(see SupTable 7 for \$\lambda\$ GC values for each model and SupFig6 for PP plots for each trait\)](#).
31 287 The Wald test assesses the significance of the SNP genotypes in a linear regression
32 288 model and is used to test the null hypothesis that the effect size of a genetic variant on
33 289 the phenotypic trait is zero (i.e., there is no association between the variant and the
34 290 trait). SNPs were then filtered based on a p-value threshold of 5×10^{-8} , a threshold widely
35 291 used in GWAS studies (e.g. Chen et al., 2021). [\(Fig 4 A & B\)](#).

36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Genome Wide F_{ST}

To estimate per site F_{ST} , we used OutFLANK (Whitlock & Lotterhos, 2015). OutFLANK identifies outlier F_{ST} SNPs by modeling the distribution of F_{ST} values across loci using a trimmed likelihood approach to exclude loci under strong selection. It then fits a chi-square distribution to the neutral F_{ST} values and identifies SNPs with unusually high or low F_{ST} as potential outliers, suggesting loci under selection. We used a false discovery rate threshold of 0.005 to determine outlier SNPs. We took the 90th quantile of these F_{ST} values to narrow down the top SNPs for [literature](#) review and plotted these on a Manhattan plot ([Fig 4 C](#)). [This allowed for isolation of the most divergent loci as well as easier visualization.](#) We then determined the F_{ST} values for the SNPs previously identified in the GWAS analyses based on the values calculated by OutFLANK.

Identification of associated genes, enrichment analysis and literature review

To identify genes associated with significant SNPs from our GWAS and F_{ST} analysis, we implemented BEDTools (Quinlan & Hall, 2010) closest function with a previously created Brown-capped Rosy-Finch gene annotation (Funk et al., 2023) to identify genes close to the SNP positions. The output from this tool was filtered to find only protein coding genes within 25kb of the given SNP. For plotting purposes, we used the [NUCmer](#) function in MUMmer4.x (Marçais et al., 2018) to align the Brown-capped Rosy-Finch genome assembly to the Zebra Finch (*Taeniopygia guttata*) genome assembly (GenBank: GCF_003957565.2). We then filtered the delta alignment output of [NUCmer](#) to keep only matches that were greater than 400 bps, and used [the show-coords](#) to display the coordinates. We then used custom R scripts to convert

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

our scaffold and positions to Zebra Finch chromosome positions for plotting SNPs in
Manhattan plots.

Enrichment in gene ontology (GO) terms was performed with Panther 19.0
(Thomas et al., 2022). [For this analysis, Gallus gallus was used as the background
gene set.](#) Enrichment thresholds were set to $p < 0.05$ after Bonferroni correction. Gene
network analysis was conducted using STRING (Szklarczyk et al., 2023) with zebra
finch as the organismal reference [for background genes](#). STRING (Search Tool for the
Retrieval of Interacting Genes/Proteins) is a database that predicts and visualizes
protein-protein interactions based on known and predicted associations from
experiments, literature, and computational methods (Szklarczyk et al., 2023). For F_{ST}
genes, all genes associated with the OutFLANK outliers were input into enrichment
analyses. Genes [resulting from the GWAS analysis](#) were grouped with all beak
morphology trait genes run as a single analysis, all feather trait genes run together, and
wing chord genes run in another analysis.

We performed a literature search for all of the candidate genes using google
scholar and the search terms gene name followed by, bird, trait specific term (when
applicable), environmental adaptation, local adaptation (e.g., [“XIA” gene beak](#), or
[“DSCAM” gene environmental adaptation](#)). For F_{ST} SNPs, only those genes
associated with the 90th quantile of SNPs were evaluated in the literature. Genes were
grouped by their key functions or associations as described in the literature. ~~(Table 1).~~

Results

Local adaptation of body and feather morphology

For the univariate approach, AIC model selection indicated five traits for which where site was a significant predictor. For wing chord, the best model included tarsus length, age, sex, and site. The model showed high explanatory power (adjusted $R^2=0.673$, $p<0.001$) and showed that birds at the White Mountain location (~~Whites~~WMTN) had shorter wings chords compared to Piute Pass (PIPA) ($p=0.003$). The best model for tare length included tarsus, sex, and site and had lower explanatory power (adjusted $R^2=0.197$) but was still highly significant ($p<0.001$). This model found that tare length was shorter in the White MountainWMTN population ($p<0.001$). Similarly, beak width had tarsus, age and site as significant predictors (adjusted $R^2=0.135$, $p<0.001$) with White MountainWMTN having beaks that are wider than those in Piute PassPIPA ($p<0.001$ ~~$p=0.004$~~). Beak depth had tarsus, age and site as the best predictors (adjusted $R^2=0.250$) and found that White MountainWMTN has significantly deeper beaks than Piute PassPIPA ($p=0.003$ ~~$p=0.004$~~). Beak length was not significantly different between the two locations and was best predicted by sex (adjusted $R^2=0.516$, $p<0.001$). For feather microstructure, the only trait that had site as a significant predictor was plumulaceous node density. This model included tarsus and age as well as site but had low explanatory power (adjusted $R^2=0.076$ ~~$R^2=0.076$~~ , $p<0.001$). It and showed that plumulaceous node density was higher in the White MountainWMTN populations ($p<0.001$ ~~$p=0.037$~~). The best model for plumulaceous barbule length based on weighted AIC weights included site as a predictor; however, the effect size was uncertain, and the effect of site was not statistically significant (adjusted $R^2=0.142$, $p=0.078$). Pennaceous barbule density had tarsus and age as the best predictors (adjusted $R^2=0.365$,

1
2
3
4
5
6
7
8
9
10 359 p<0.001), pennaceous barbule length had sex as the best predictor (adjusted R²=0.307,
11 360 p<0.001) and plumulaceous barbule density had tarsus and age as the best predictors
12
13 361 (adjusted R²=0.076, p=0.004). [We found that body size did not vary between sites. In](#)
14
15 362 [our model using tarsus \(adjusted R²=0.165, p<0.001\) both age and sex were significant](#)
16
17 363 [predictors \(p<0.001 and p<0.05\), however site was not significant \(p=0.417\). Our](#)
18 364 [comparison using the PC axis representing size \(adjusted R²=0.4639, p<0.001\) had](#)
19
20 365 [similar results with age and sex as significant and site as not \(p=0.53\).](#)
21
22 366 [When looking at the results of the multivariate analysis, we see that for body](#)
23 367 [morphology traits site was a highly significant explanatory variable \(p=1.327e-11\).](#)
24
25 368 [Tarsus, age and sex were also highly significant \(p<2.2e-16\). The MANOVA on feather](#)
26
27 369 [traits also had site as a significant predictor although the p-value was less significant](#)
28
29 370 [\(p=0.036\). Age was also significant \(p=1.5e-14\) but tarsus and sex were not \(p>0.05\).](#)
30
31 371 [For details on the models tested, models chosen, and full model results, see](#)
32 372 [Supplementary Table 4-6.](#)

Commented [2]: Which table?
Commented [3]: making sure you catch this

34
35 373 **Evaluation of population structure**
36 374 Whole-genome sequencing was performed on 150 individuals [and 146 were](#)
37
38 375 [included in downstream analysis. Four](#)Two individuals were removed [because](#)as their
39 376 collection location [could not be validated](#)~~was not available~~. Variant filtering resulted in
40
41 377 7,739,836 SNPs for subsequent genetic analysis. Linkage pruning removed 5,526,104
42
43 378 SNPs. Visualizing PLINK PCA results revealed little population structure with most
44
45 379 individuals clustering together (SupFig [24](#)). ADMIXTURE results confirmed that there is
46
47 380 no population structure in these samples [because](#)as k=1 had the lowest cv error (0.599)
48 381 compared to k=2 or 3 (0.612 and 0.627).
49
50
51
52
53
54
55
56
57
58
59
60

Genes identified with genome-wide association approach

The univariate linear mixed model identified a total of 64 SNPs for ~~traits~~ wing chord, beak length, beak width, pennaceous barbule length, and pennaceous barbule density. Using a Brown-capped Rosy-Finch gene annotation we identified a total of 26 unique genes corresponding to these SNPs (Table 1). We filtered out genes that ~~were~~ have not ~~been~~ fully characterized (e.g., "LOC100221041") leaving 14 genes. The Bayesian sparse linear mixed model identified 266 SNPs for wing chord, ~~nare~~ length, beak length, depth and width, pennaceous barbule length and density, and plumulaceous barbule length and node density. We identified 146 genes associated with these SNPs (Table 1), 110 of which had been fully characterized. For each trait the global view of p-values for all SNPs and associated genes across both models are represented by Manhattan plots (Fig 4-A-C, SupFig 3-52-5). Given the sample size ($n = 146$), this GWAS is best viewed as conservative, detecting only loci with relatively strong effects on morphology.

GO term analysis using Panther produced significant results for beak depth and a combined set of all feather traits ~~combined~~. For beak depth, the top terms were presynaptic membrane assembly ($p=1.44 \times 10^{-5}$), synaptic membrane adhesion (specifically cell adhesion) ($p=1.01 \times 10^{-5}$) and heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules ($p=1.13 \times 10^{-5}$). For the feather traits (plumulaceous barbule and node density combined with pennaceous barbule length), fructosamine catabolic process was a significant term ($p=1.12 \times 10^{-6}$). Wing chord had no significant GO terms. STRINGString analysis for beak and feather traits as well as wing chord did not find significantly more interactions than expected.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Genes identified through high Fixation index (F_{ST}) between the two populations
Using OutFLANK 831 loci were identified as having significantly high F_{ST} when comparing the White Mountain and Piute Pass populations, corresponding to 265 genes (Fig 4D, [Sup Table 2](#)). No GWAS SNPs were flagged as having significant F_{ST} values. The minimum F_{ST} to be considered an outlier was 0.17513. The mean F_{ST} of SNPs not considered outliers was 0.0045. Using a 90th quantile we determined $F_{ST}=0.235$ to be the threshold, narrowing down the SNPs to 105 loci. For the smaller set of loci, bedtools identified 61 related genes (Table 2). We ran STRING and GO term enrichments on both the larger and smaller set of genes. The smaller set did not have a significant number of connections over all or any significantly enriched GO terms. The larger set of 264 genes, however, had an overall significant number of connections ($p<0.05$) and had multiple significant annotated keywords (sourced from UniProt). The most significant was the keyword repeat ($FDR_p<0.001$), the next was ubl conjugation pathways ($FDR_p<0.05$).

Discussion

Understanding the genetic and morphological basis of local adaptation is critical for predicting how species respond to environmental change. Alpine species face unique conservation challenges due to their extreme environments and thus are a high priority for studies of adaptive capacity. In this study, we investigated morphological and genetic differences between two populations of an alpine specialist songbird located [in](#) [at distinctthe extreme partsends](#) of its environmental range. Consistent with our predictions, we found wings were shorter and feathers were denser in the colder environment. However, in contrast to our prediction, beak depth and width were larger

in the colder environment, suggesting other factors like diet may place contrasting selective pressures on this ecologically important trait. We also identified numerous genes with known links to limb, facial, and feather development that potentially underlie variation in wing, bill and feather microstructure. Further F_{ST} outlier analyses revealed numerous significant differences between cold and warm populations at SNPs located within genes with clear links to altitudinal adaptation. Together, our results provide important insights into the morphological traits and associated genomic regions associated with microgeographic variation in a high-altitude songbird species.

Trait differences between warm and cold habitats

Ecogeographic rules, such as Allen's [and Bergmann's rules](#), are useful for predicting broad geographic trends in morphology in response to environmental conditions, but observed patterns do not always align with expectations. In our study, we found that wing chord was smaller in the colder environment, suggesting that this morphological trait may play a role in thermoregulation, consistent with Allen's rule. Heat dissipation from the wings occurs across the vascularized brachial regions, primarily during flight when these areas are exposed to air flow (Lewden et al., 2023; Ward et al., 1999). Wing bone length reflects the extent of vascularized areas on the wings and has been shown to be longer in warmer environments across a broad sample of passerine species (Weeks et al., 2025). However, studies of other alpine bird species have found the opposite pattern: that birds from high-elevation sites have longer wings than those from low-elevation sites (Bears et al., 2008; Ceresa et al., 2024). In these cases, researchers have suggested that the demand for more energetically efficient flight at high elevations may outweigh thermoregulatory selective pressures,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

demonstrating the often contrasting selective pressures on ecologically important traits. Although outside of the scope of the current work, additional wing measurements could be collected that would allow for a more thorough investigation into the components of flight morphology. It is possible that migratory behavior is influencing wing length as previous work has shown that both within (Egbert & Belthoff, 2003; Grilli et al., 2017) and between (Lockwood et al., 1998) species those who migrate further have longer wings. To date, no formal study has been conducted on the migratory patterns of the Sierra Nevada Gray-crowned subspecies and so future work could investigate the links between wing morphology and movement patterns. Overall, our morphological analyses support the idea that thermoregulatory demands may play an important role important dominant role in shaping wing morphology in the Sierra Nevada Gray-crowned Rosy-Finch, potentially outweighing the benefits of flight efficiency in this high-elevation, cold-adapted species but not excluding the potentially parallel pressure of migratory distance.

Contrary to our predictions based on Allen’s rule, beak depth and width were larger in the colder environment. Although temperature is often a key environmental variable exerting selective pressure on alpine taxa, variation in beak morphology between our populations may instead reflect divergent selective pressures associated with foraging ecology, a well-documented driver of beak divergence in birds (Grant et al., 1976; Lamichhaney et al., 2018). In our study system, the Piute Pass population likely has access to more aquatic insects due to the higher availability of aquatic habitats proximate to Rosy-Finch breeding sites in the Sierra Nevada (Epanchin et al., 2010; Rundel & Millar, 2016). For example, Epanchin et al. (2010) found that, when abundant, mayflies can comprise up to

38% of the Sierra Nevada Gray-Crowned Rosy-Finch diet. In contrast, the White Mountain population may rely more heavily on insects deposited in snowfields, a behavior that could favor a more robust, conical bill for extracting frozen prey from the snow surface. Work is currently underway to investigate the dietary differences between these two populations of Rosy-Finch (Tim Brown, personal correspondence), and future work should explore the selective pressures experienced by this species. The other aspect of beak morphology assessed, nares length, was significantly larger in the warmer environment. If nares is playing a role in thermoregulation, these results would be consistent with larger nares allowing for more heat to be lost. Additionally, smaller nares in the colder environment would potentially prevent snow from entering the nasal passages while foraging on snow. Overall, our results on beak morphology These results emphasize the complexity of identifying drivers of local adaptation, especially in alpine systems where extreme environments, food scarcity, and variation in migration strategies can create conflicting selection pressures.

Our comparison of body size did not have significant results, suggesting that Bergmann's rule is not contributing to morphological differences in this subspecies. He et al (2023) found that there was significant variation in both strength and direction of correlation between latitude and body size within and between taxa. With our results, this could imply that Sierra Nevada Gray-crowned Rosy-Finch are an exception to the rule. In another meta-analysis, it was found that sedentary bird species were more likely to conform to Bergmann's rule than migratory ones and suggest that winter temperatures may exert a higher pressure than breeding temperatures (Meiri & Dayan,

Commented [4]: Including thoughts on nares differences here.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2003). The Sierra Nevada Gray-crowned Rosy-Finch is a short-distance, altitudinal migrant. It is possible that both White Mountain and Piute Pass populations move to similar areas or climates during the winter season, thus decreasing the divergent selection pressure that would drive differences in Bermann's rule.

Of our feather comparisons, we identified significant differences between cold and warm sites for only one trait— plumulaceous node density. In line with our predictions, plumulaceous node density was higher in the colder, high elevation White Mountain population. Higher node density is known to improve air trapping ability, thus trapping warmth against the body surface (King & McLelland, 1984). Consistent with our results, a study on sparrows found that higher elevation alpine forms had higher node density relative to lowland forms (Lei et al., 2002). A further multi-species comparison found that species wintering in cold and windy conditions had downy feathers with higher node densities (D'alba et al., 2017, 2017). These interspecific trends are mirrored in our results, suggesting that the White Mountain population has feather adaptations consistent with an increased need for heat retention. The lack of significant results for other feather traits may be explained by the amount of variation in the measurements as well as the sample sizes being small, resulting in an underpowered comparison. An alternative cause for the non-significant results is that alternative selection pressures not considered in the present work are driving morphology of these traits. This is somewhat supported by the overall low explanatory power of our models used in this study. Additional work on this aspect of local adaptation would benefit from investigating the influences of alternative environmental traits and utilizing a more controlled setting to evaluate the functionality of intraspecific differences in feather morphology.

Genome-wide analysis

We used GWAS to identify SNPs within genes involved in traits that differ between warm- and cold-adapted populations described previously: beak width and depth, nares length, wing cord length, and pennaceous barbule density. Because ours is one of the first studies to investigate the genetic basis of feather microstructure variation, we also report on the genes associated with other measured feather traits, even though high variance in these traits prevented our ability to robustly investigate differences between populations occupying cold and warm environments.

Our literature review of GWAS results identified genes with diverse associations. For wing chord, we identified limb and bone development, heat response, and vascularization. Notably, the genes *GUCA1C* and *NIPBL* have been linked to limb development in ducks and zebrafish (G. Li et al., 2020; Mohammadi, 2024; Muto et al., 2014) and may play a similar developmental role in songbirds. We also detected the genes *PGLYRP2* and *PACRG*, which have been shown to be upregulated in chickens experiencing heat stress (Kim et al., 2021; Tian et al., 2020). Together, the limb development and heat response genes we identified as important further support the potential connection between wing length and thermoregulation suggested by our morphological finding of shorter wings in the colder environment. Although GWAS results do not establish a direct functional link to specific traits (Tam et al., 2019), our findings suggest that future research should investigate the potential role of thermoregulatory selective pressures on genes involved in wing development.

1
2
3
4
5
6
7
8
9
10 540 When focusing on looking at genes associated with beak morphology, BSLMM
11 541 found more SNPs than LMM and identified several genes potentially associated with
12 542 facial structure development and climate adaptation. One gene associated with beak
13 543 depth was *XIAP*. Duplications of this gene have been found to result in facial
14 544 dysmorphism in humans, including the underdevelopment of cheekbones and
15 545 protruding jaws (Di Benedetto et al., 2014). Although a link with beak development in
16 546 birds has not yet been formally established, this could be an area for further research.
17 547 Another gene identified for beak width was *MINDY2*, which belongs to a group of
18 548 deubiquitinating enzymes. These genes are related to the regulation of the Wnt pathway
19 549 (Park et al., 2020), a pathway that is well established as important for beak
20 550 development (Nierop 2002). Additional genes such as *RASL11A*, *TINAG*, and *PLPP3*
21 551 have been directly linked to beak development in avian species (Abernathy, 2021; Bai
22 552 et al., 2018). Two genes associated with beak depth in our study (*LRP1B* and *UTRN*)
23 553 were shown to be associated with breeding temperature in a study that used a GWAS
24 554 approach to identify loci associated with climate variables related to morphology in
25 555 hermit thrush (N. Adams et al., 2025). Although the beak morphology of our species did
26 556 not entirely fall in line with predictions based around their thermoregulatory role, the
27 557 elucidation of genes indicating a relationship between thermoregulation and beak
28 558 morphology suggests that beaks still play a thermoregulatory role in this species.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43 559 Our GWAS on feather traits was, to the best of our knowledge, the first genome-
44 560 wide association study of feather microstructural traits. Identifying the genetic basis of
45 561 these fine-scale traits opens new avenues for studying how feather structure evolves in
46 562 response to environmental pressures and functional demands. One of the genes
47
48
49
50
51
52
53
54
55
56
57
58
59
60

associated with pennaceous barbule length in our study, *HBE1*, has also been identified in previous research on feather development. Using RNA sequencing, Limber et al. (2024) found that *HBE1*, a hemoglobin gene (Mao et al., 2023), was highly expressed in feather pulp, the central tissue of the developing feather germ. Additional genes associated with feather traits in our study have been identified as being related to plumage coloration in birds, including *EFNA2*, *NRXN1*, and *HERC4* (Shakya, 2020; L. Sun et al., 2020; X. Zhang et al., 2023). Many other genes identified as being associated with feather traits in our work have not been shown to be associated with feather development or morphology in the literature. Feather morphology remains understudied in the context of local adaptation, and the genetic architecture of these microstructural traits is still poorly understood. Our study offers a foundation for deeper investigation. Expanding this work to include comparative genomic studies across populations or species with more strongly differentiated feather morphology could also help link genotype to phenotype and provide insight into the adaptive significance of feather microstructure.

~~Our original intent with the F_{ST} -outlier tests was to determine if any of the genes identified in the GWAS had variants that diverged significantly between the cold and warm environment populations. However, when comparing our F_{ST} results with SNPs identified in the GWAS, we found no overlap between the two sets. It is important to note that, in this system, we are limited in our ability to assess the role of phenotypic plasticity in driving trait differences. Bringing individuals together from both populations into a common aviary, as was done in Bears et al. (2008), would allow for better examination of these effects. It could also be argued that showing relatively high F_{ST}~~

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

values for GWAS loci might suggest that the morphological differentiation is due to divergence rather than phenotypic plasticity. Our original intent with the F_{ST} outlier tests was to determine if any of the genes identified in the GWAS had variants that diverged significantly between the cold and warm environment populations. However, when comparing our F_{ST} results with SNPs identified in the GWAS, we found no overlap between the two sets. Although this could suggest that phenotypic plasticity is primarily driving the trait divergence, one must also consider that this result is consistent with known limitations of F_{ST} -based outlier scans, which are biased toward detecting loci under strong divergent selection and with large effect sizes (Hoban et al., 2016). F_{ST} measures allele frequency differences between populations and loci with large phenotypic effects are more likely to experience strong selection and exhibit pronounced divergence. In contrast, highly polygenic traits are shaped by small-effect variants that shift subtly in frequency—patterns that often fall below the detection threshold of F_{ST} scans. Our findings underscore the value of combining F_{ST} outlier analyses with GWAS approaches: while the former is designed to pinpoint loci under strong selection, the latter is better suited for uncovering the complex genetic architecture of polygenic traits. Together, these methods can generate a more complete understanding of the genetic basis for local adaptation.

While we did not find overlap between the F_{ST} and the GWAS results, our F_{ST} analysis highlighted numerous genes involved in hypoxia response, altitude adaptation, and cold adaptation. These results are consistent with our hypothesis that elevation and associated climatic variables are applying divergent selection pressures in this species. Of these genes, *NOX4* was found to be involved in mediating hypoxia-inducible

transcription factors, a key component of the cellular response to hypoxia (Diebold et al., 2010). Similarly, *MAML2* and *SUPT20H* were found to be associated with altitude adaptation in deer mice (Schweizer et al., 2021a) and *ADGRD1* was associated with altitude adaptation in Tibetan chickens (Nan et al., 2023). Multiple studies identified *PPP1R1C* as being related to cold adaptation in chickens (Fedorova et al., 2022a; Romanov et al., 2023). These are all traits that would lend themselves to survival in alpine environments, and divergent selection is likely a product of varying selection pressures between the higher and colder White Mountain location and the warmer and lower Piute Pass location.

Our F_{ST} analysis also identified several genes that might be involved with plumage color differences in Rosy-Finches. Another gene that showed divergence between these two populations of finch was *PHIP* which has been shown to be related to differentiation in plumage coloration in House Finch (Balenger et al., 2015). Interestingly, when comparing candidate outlier genes from this work with [studieswork](#) identifying genes involved with the notable plumage coloration differences across the broader *rRosy-fFinch* species complex, there were some overlapping genes. In a GWAS on crown coloration, Funk et al (2023) identified *NDUFAF2*, a gene that was also significant in this work. For body plumage coloration, our results overlap with those of Funk et al (2023) for the genes *KCNV2*, *PUM3*, and *VLDLR*. Although it was not a component of the current work, other studies have shown that plumage coloration is involved in avian thermoregulation, with darker plumage putatively facilitating heat absorption (Medina et al., 2018; Rogalla et al., 2022), and can show adaptation to local environment (Romano et al., 2019; Sandoval & Barrantes, 2019; Sirkiä et al., 2010). This

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

632 provides an important avenue for future work with this species investigating the
633 differences in plumage coloration between populations.

634 An additional result of our F_{ST} analysis was the identification of a gene, *CTIF*, that
635 has been previously identified as being a climate-adapted gene in a landscape genomic
636 study on the closely related Brown-capped Rosy-Finch (DeSaix et al., 2022).
637 Interestingly, this gene was also identified through a contrasting approach using a
638 mixed linear model on beak shape for two species of Darwin’s finches, *Geospiza fortis*
639 and *G. scandens* (Lawson & Petren, 2017). The authors of this paper summarize their
640 results as indicating that inter and intraspecies beak shape variation is a result of a
641 small suite of traits evolving in concert, corresponding to many genes. It is possible that
642 we did not identify *CTIF* genes in our GWAS method because of low power due to
643 smaller variation in the beak traits measured, a result of using continuous
644 measurements between two populations versus categorical comparisons between
645 different species as in Lawson & Petren (2017). Because of the large number of SNPs
646 obtained for our two populations, our F_{ST} analysis likely had higher power and thus
647 detected this gene, in addition to the fact that it had a high F_{ST} of 0.263 (averaged
648 across the 5 SNPs associated with this trait). The overlap of these genes in the two
649 studies described as well as ours makes it a strong candidate as a genetic component
650 of adaptation to environment, potentially through the mechanism of beak morphology.

651 An important outcome of local adaptation is the maintenance of ecologically
652 important genetic variation (Savolainen et al., 2013). Conserving genetic variation
653 prevents loss of adaptive potential and can underlie rapid adaptive responses to
654 environmental change (Forester et al., 2023). We integrated phenotypic measurements,

genome-wide association studies, and F_{ST} analyses, to understand the morphological and genetic basis of local adaptation of an alpine bird species, the Sierra Gray-crowned Rosy-Finch. When comparing traits suspected to be involved in thermoregulation for this species, we found that morphological variation across populations only partially aligned with ecogeographic expectations. Wing chord was shorter in the colder environment, consistent with Allen's rule and likely reflecting thermoregulatory adaptation. However, beak depth and width were larger in colder environments, contrary to predictions, suggesting additional selective pressures such as diet specialization may be playing a role. Comparisons of feather microstructure further showed thermoregulatory adaptation, with higher plumulaceous node density in the high-elevation population likely enhancing insulation. When coupled with the results of our GWAS and F_{ST} outlier scans we gained a clearer picture of traits and genes important to adaptation in alpine birds. Importantly, we found that many of the genes underlying wing length, beak depth and feather microstructure had links to both development of similar traits in other organisms as well as thermoregulation. Our genomic and morphological results provide strong evidence that this alpine species is locally adapted to temperature variation.

For those traits that are involved in thermoregulation, we can hypothesize about the implication of the intraspecific differences in light of future climate change. Recent work has shown that, for populations whose morphology and genetics misalign with the environment as it changes due to anthropogenic pressures there is a decrease in

1
2
3
4
5
6
7
8
9
10 676 [fitness \(Pelletier & Coltman, 2018; Rodriguez et al., 2025\). As climate change drives](#)
11 677 [increasing variability in temperature, understanding both simple and complex genetic](#)
12 678 [architectures underlying local adaptation is essential for predicting species' responses](#)
13 679 [to global change. Although we cannot directly assess fitness consequences in this](#)
14 680 [species. Overall,](#) this work enhances our understanding of the morphological traits and
15 681 genetic mechanisms underlying local adaptation in an alpine species [and lays a ground](#)
16 682 [work for future investigation.](#) The resulting insights will be used to inform the delineation
17 683 of conservation units and support management strategies that seek to preserve the
18 684 evolutionary processes essential for long-term persistence of alpine organisms in a
19 685 changing world (Allendorf et al., 2010; Flanagan et al., 2018; Shafer et al., 2015).

20
21
22
23
24
25
26
27
28
29
30 686 **Acknowledgements**

31 687 We thank Sarah Albright, Alena Arnold, Emma Brown, Carl Brown, Alison Brunschwiler,
32 688 Vincent Chevreuil, Ryan Cheung, Daisy Cortes, Jessica Gallardo, Karim Hanna, Raquel
33 689 Lozano, Juan Rebellon, Leticia Santillana, Kevin Silberberg, and Juyung Yoo for their
34 690 participating in catching, handling, banding, measuring, and sampling birds. We thank
35 691 Bailey Lubash for contributing to the methodology for measuring feather microstructure
36 692 characteristics. We thank NovoGene for their assistance with next generation
37 693 sequencing. We thank Christine Rayne and Amanda Carpenter for library preparation,
38 694 and the Ruegg and Funk labs at Colorado State University for detailed feedback on
39 695 early ~~iterations interactions~~ of the project and manuscript. This work utilized the RMACC
40 696 Summit supercomputer, which is supported by the National Science Foundation
41 697 (awards ACI-1532235 and ACI-1532236), the University of Colorado Boulder and
42 698 Colorado State University. The RMACC Summit supercomputer is a joint effort of the
43 699 University of Colorado Boulder and Colorado State University. Figures were created
44 700 with [BioRender.com](#) ~~and illustrations were created by Erica Robertson.~~ Funding for this
45 701 project was made possible by the National Science Foundation (NSF 2222524, NSF
46 702 2222525, and NSF 2222526). Handling, banding, and sampling of Grey-crowned Rosy-
47 703 Finch was performed under Federal Banding Permit 24401 and CFWSscientific
48 704 Collecting Permit (SCP) S-190610001-22081-001-02.

References

- Abernathy, E. (2021). *The Genomic Basis of Population and Adaptive Divergence in Buteo Sister Species Across Multiple Evolutionary and Geographic Scales* [Ph.D., University of California, Davis].
<https://www.proquest.com/docview/2638107596/abstract/BB50D6B5F64067PQ/1>
- Adams, D. C., Glynne, E., & Kaliontzopoulou, A. (2020). Interspecific allometry for sexual shape dimorphism: Macroevolution of multivariate sexual phenotypes with application to Rensch's rule. *Evolution; International Journal of Organic Evolution*, 74(9), 1908–1922. <https://doi.org/10.1111/evo.14049>
- Adams, N., Dias, T., Skeen, H. R., Pegan, T., Willard, D. E., Winger, B., Ruegg, K., Weeks, B. C., & Bay, R. (2025). Genetic and morphological shifts associated with climate change in a migratory bird. *BMC Biology*, 23(1), 3.
<https://doi.org/10.1186/s12915-024-02107-5>
- AdaptWest Project. (2022). *Gridded current and projected climate data for North America at 1km resolution, generated using the ClimateNA v7.30 software (T. Wang et al., 2022)* [Dataset].
- Ahmad, H. I., Asif, A. R., Ahmad, M. J., Jabbar, F., Adnan, M., Ahmed, S., Afzal, G., Saleem, A. H., Li, L., Jiang, H., Simirgiotis, M. J., Rauf, M., & Chen, J. (2020). Adaptive evolution of peptidoglycan recognition protein family regulates the innate signaling against microbial pathogens in vertebrates. *Microbial Pathogenesis*, 147, 104361.
<https://doi.org/10.1016/j.micpath.2020.104361>
- Aitken, S. N., Yeaman, S., Holliday, J. A., Wang, T., & Curtis-McLane, S. (2008).

1
2
3
4
5
6
7
8
9
10 727 Adaptation, migration or extirpation: Climate change outcomes for tree
11 728 populations. *Evolutionary Applications*, 1(1), 95–111.
12
13 729 <https://doi.org/10.1111/j.1752-4571.2007.00013.x>
14
15 730 Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of
16
17 731 ancestry in unrelated individuals. *Genome Research*, 19(9), 1655–1664.
18
19 732 <https://doi.org/10.1101/gr.094052.109>
20
21 733 Allen, J. A. (1877). *The Influence of Physical Conditions in the Genesis of Species*.
22
23 734 Allendorf, F. W., Hohenlohe, P. A., & Luikart, G. (2010). Genomics and the future of
24
25 735 conservation genetics. *Nature Reviews Genetics*, 11(10), 697–709.
26
27 736 <https://doi.org/10.1038/nrg2844>
28
29 737 Alsoufi, M. A., Liu, Y., Cao, C., Zhao, J., Kang, J., Li, M., Wang, K., He, Y., & Ge, C.
30 738 (2023). Integrated Transcriptomics Profiling in Chahua and Digao Chickens’ Breast
31
32 739 for Assessment Molecular Mechanism of Meat Quality Traits. *Genes*, 14(1), Article
33
34 740 1. <https://doi.org/10.3390/genes14010095>
35
36 741 Ashton, K. G. (2002). Patterns of within-species body size variation of birds: Strong
37
38 742 evidence for Bergmann’s rule. *Global Ecology and Biogeography*, 11(6), 505–523.
39
40 743 <https://doi.org/10.1046/j.1466-822X.2002.00313.x>
41
42 744 Auwera, G. V. der, & O’Connor, B. D. (2020). *Genomics in the cloud: Using Docker,*
43 745 *GATK, and WDL in Terra* (First edition). O’Reilly.
44
45 746 Bai, H., Sun, Y., Liu, N., Xue, F., Li, Y., Xu, S., Ye, J., Zhang, L., Chen, Y., & Chen, J.
46
47 747 (2018). Single SNP- and pathway-based genome-wide association studies for beak
48
49 748 deformity in chickens using high-density 600K SNP arrays. *BMC Genomics*, 19(1),
50
51
52
53
54
55
56
57
58
59
60

501. <https://doi.org/10.1186/s12864-018-4882-8>
- Bai, H., Zhao, N., Li, X., Ding, Y., Guo, Q., Chen, G., & Chang, G. (2024). Whole-genome resequencing identifies candidate genes associated with heat adaptation in chickens. *Poultry Science*, 103(10), 104139. <https://doi.org/10.1016/j.psj.2024.104139>
- Bai, H., Zhu, J., Sun, Y., Liu, R., Liu, N., Li, D., Wen, J., & Chen, J. (2014). Identification of Genes Related to Beak Deformity of Chickens Using Digital Gene Expression Profiling. *PLOS ONE*, 9(9), e107050. <https://doi.org/10.1371/journal.pone.0107050>
- Balenger, S. L., Bonneaud, C., Sefick, S. A., Edwards, S. V., & Hill, G. E. (2015). Plumage color and pathogen-induced gene expression in a wild bird. *Behavioral Ecology*, 26(4), 1100–1110. <https://doi.org/10.1093/beheco/arv055>
- Barnes, N. E., Mendoza, K. M., Strasburg, G. M., Velleman, S. G., & Reed, K. M. (2019). Thermal challenge alters the transcriptional profile of the breast muscle in turkey poults. *Poultry Science*, 98(1), 74–91. <https://doi.org/10.3382/ps/pey401>
- Barve, S., Ramesh, V., Dotterer, T. M., & Dove, C. J. (2021). Elevation and body size drive convergent variation in thermo-insulative feather structure of Himalayan birds. *Ecography*, 44(5), 680–689. <https://doi.org/10.1111/ecog.05376>
- Bears, H., C. Drever, M., & Martin, K. (2008). Comparative morphology of dark-eyed juncos *Junco hyemalis* breeding at two elevations: A common aviary experiment. *Journal of Avian Biology*, 39(2), 152–162. <https://doi.org/10.1111/j.2008.0908-8857.04191.x>

1
2
3
4
5
6
7
8
9
10 771 Belnap, J., Webb, R. H., Esque, T. C., Brooks, M. L., Defalco, L. A., & Macmahon, J.
11 772 A. (2016). Deserts. In *Ecosystems of California* (1st ed., pp. 635–668). University of
12 773 California Press. <http://www.jstor.org/stable/10.1525/j.ctv1xxzp6.35>
13 774 Bhavana, K., Foote, D. J., Srikanth, K., Balakrishnan, C. N., Prabhu, V. R.,
14 775 Sankaralingam, S., Singha, H. S., Gopalakrishnan, A., & Nagarajan, M. (2022).
15 776 Comparative transcriptome analysis of Indian domestic duck reveals candidate
16 777 genes associated with egg production. *Scientific Reports*, 12(1), 10943.
17 778 <https://doi.org/10.1038/s41598-022-15099-5>
18 779 Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to
19 780 measuring local adaptation. *Ecology Letters*, 16(9), 1195–1205.
20 781 <https://doi.org/10.1111/ele.12150>
21 782 Boer, E. F., Maclary, E. T., & Shapiro, M. D. (2021). Complex genetic architecture of
22 783 three-dimensional craniofacial shape variation in domestic pigeons. *Evolution &*
23 784 *Development*, 23(6), 477–495. <https://doi.org/10.1111/ede.12395>
24 785 Borges, R., Fonseca, J., Gomes, C., Johnson, W. E., O'Brien, S. J., Zhang, G., Gilbert,
25 786 M. T. P., Jarvis, E. D., & Antunes, A. (2019). Avian Binocularity and Adaptation to
26 787 Nocturnal Environments: Genomic Insights from a Highly Derived Visual Phenotype.
27 788 *Genome Biology and Evolution*, 11(8), 2244–2255.
28 789 <https://doi.org/10.1093/gbe/evz111>
29 790 Browning, B. L., & Browning, S. R. (2016). Genotype Imputation with Millions of
30 791 Reference Samples. *American Journal of Human Genetics*, 98(1), 116–126.
31 792 <https://doi.org/10.1016/j.ajhg.2015.11.020>
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- Buroker, N. E., Ning, X.-H., Zhou, Z.-N., Li, K., Cen, W.-J., Wu, X.-F., Zhu, W.-Z.,
Scott, C. R., & Chen, S.-H. (2012). AKT3, ANGPTL4, eNOS3, and VEGFA associations
with high altitude sickness in Han and Tibetan Chinese at the Qinghai-Tibetan
Plateau. *International Journal of Hematology*, 96(2), 200–213.
<https://doi.org/10.1007/s12185-012-1117-7>
- Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Exposito-Alonso, M., & Keller, S. R.
(2020). Genomic Prediction of (Mal)Adaptation Across Current and Future Climatic
Landscapes. *Annual Review of Ecology, Evolution, and Systematics*, 51(Volume 51,
2020), 245–269. <https://doi.org/10.1146/annurev-ecolsys-020720-042553>
- Ceresa, F., Brambilla, M., Kvist, L., Vitulano, S., Pes, M., Tomasi, L., Pedrini, P.,
Bettega, C., Anderle, M., Hilpold, A., & Kranebitter, P. (2024). Wing morphology
changes with habitat availability and elevation in an alpine-specialist bird. *Global
Ecology and Conservation*, 56, e03276.
<https://doi.org/10.1016/j.gecco.2024.e03276>
- Chen, C.-K., Chang, Y.-M., Jiang, T.-X., Yue, Z., Liu, T.-Y., Lu, J., Yu, Z., Lin, J.-J., Vu,
T.-D., Huang, T.-Y., Harn, H. I.-C., Ng, C. S., Wu, P., Chuong, C.-M., & Li, W.-H.
(2024). Conserved regulatory switches for the transition from natal down to juvenile
feather in birds. *Nature Communications*, 15(1), 4174.
<https://doi.org/10.1038/s41467-024-48303-3>
- Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics*, 34(17), i884–i890.
<https://doi.org/10.1093/bioinformatics/bty560>

1
2
3
4
5
6
7
8
9
10 815 Chen, X., Bai, X., Liu, H., Zhao, B., Yan, Z., Hou, Y., & Chu, Q. (2022). Population
11 816 Genomic Sequencing Delineates Global Landscape of Copy Number Variations that
12 Drive Domestication and Breed Formation of in Chicken. *Frontiers in Genetics*, 13.
13 817 <https://doi.org/10.3389/fgene.2022.830393>
14 818
15 819 Chen, Z., Boehnke, M., Wen, X., & Mukherjee, B. (2021). Revisiting the genome-wide
16 significance threshold for common variant GWAS. *G3 Genes[Genomes]Genetics*,
17 820 11(2), jkaa056. <https://doi.org/10.1093/g3journal/jkaa056>
18 821
19 822 Cheng, Y., Miller, M. J., Zhang, D., Song, G., Jia, C., Qu, Y., & Lei, F. (2020).
20 Comparative Genomics Reveals Evolution of a Beak Morphology Locus in a High-
21 Altitude Songbird. *Molecular Biology and Evolution*, 37(10), 2983–2988.
22 823 <https://doi.org/10.1093/molbev/msaa157>
23 824
24 825 Cho, E., Kim, M., Cho, S., So, H.-J., Lee, K.-T., Cha, J., Jin, D., & Lee, J. H. (2023). A
25 genome-wide association study for the fatty acid composition of breast meat in an
26 F2 crossbred chicken population. *Journal of Animal Science and Technology*, 65(4),
27 828 735–747. <https://doi.org/10.5187/jast.2023.e1>
28 829
29 830 Cho, Y. S., Jun, J. H., Kim, J. A., Kim, H.-M., Chung, O., Kang, S.-G., Park, J.-Y., Kim,
30 H.-J., Kim, S., Kim, H.-J., Jang, J., Na, K.-J., Kim, J., Park, S. G., Lee, H.-Y., Manica, A.,
31 Mindell, D. P., Fuchs, J., Edwards, J. S., ... Bhak, J. (2019). Raptor genomes reveal
32 evolutionary signatures of predatory and nocturnal lifestyles. *Genome Biology*,
33 833 20(1), 181. <https://doi.org/10.1186/s13059-019-1793-1>
34 834
35 835 Chung, O., Jin, S., Cho, Y. S., Lim, J., Kim, H., Jho, S., Kim, H.-M., Jun, J., Lee, H.,
36 Chon, A., Ko, J., Edwards, J., Weber, J. A., Han, K., O'Brien, S. J., Manica, A., Bhak, J.,

- 837 & Paek, W. K. (2015). The first whole genome and transcriptome of the cinereous
838 vulture reveals adaptation in the gastric and immune defense systems and possible
839 convergent evolution between the Old and New World vultures. *Genome Biology*,
840 16(1), 215. <https://doi.org/10.1186/s13059-015-0780-4>
- 841 Clark, J., Lim, H. C., Olsen, B., & Kovach, A. (2024). *Whole-genome data uncover the*
842 *basis of local adaptation with gene flow in a threatened coastal songbird.*
843 <https://doi.org/10.22541/au.172114945.59127531/v1>
- 844 Clausen, J., Keck, D. D., & Hiesey, W. M. (1941). Regional Differentiation in Plant
845 Species. *The American Naturalist*, 75(758), 231–250.
846 <https://doi.org/10.1086/280955>
- 847 Crates, R., von Takach, B., Young, C. M., Stojanovic, D., Neaves, L. E., Murphy, L.,
848 Gautschi, D., Hogg, C. J., Heinsohn, R., Bell, P., & Farquharson, K. A. (2024).
849 Genomic insights into the critically endangered King Island scrubtit. *Journal of*
850 *Heredity*, 115(5), 552–564. <https://doi.org/10.1093/jhered/esae029>
- 851 da Costa Moreno, H. (2022). *Molecular Evolution of Vertebrate Genes Involved in*
852 *Mitochondrial Biogenesis: Unravelling the Origins of Birds' Endothermy* [Masters
853 Thesis, Universidade do Porto]. ProQuest Dissertations & Theses.
- 854 D'alba, L., Carlsen, T. H., Ásgeirsson, Á., Shawkey, M. D., & Jónsson, J. E. (2017).
855 Contributions of feather microstructure to eider down insulation properties. *Journal*
856 *of Avian Biology*, 48(8), 1150–1157. <https://doi.org/10.1111/jav.01294>
- 857 Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,
858 Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve

1
2
3
4
5
6
7
8
9
10 859 years of SAMtools and BCFtools. *GigaScience*, 10(2), giab008.
11 860 <https://doi.org/10.1093/gigascience/giab008>
12 861 Delmore, K., Illera, J. C., Pérez-Tris, J., Segelbacher, G., Lugo Ramos, J. S., Durieux,
13 862 G., Ishigohoka, J., & Liedvogel, M. (2020). The evolutionary history and genomics of
14 863 European blackcap migration. *eLife*, 9, e54462. <https://doi.org/10.7554/eLife.54462>
15 864 Demery, A.-J. C., Burns, K. J., & Mason, N. A. (2021). Bill size, bill shape, and body
16 865 size constrain bird song evolution on a macroevolutionary scale. *Ornithology*,
17 866 138(2), ukab011. <https://doi.org/10.1093/ornithology/ukab011>
18 867 DeSaix, M., George, T., Seglund, A., Spellman, G., Zavaleta, E., & Ruegg, K. (2022).
19 868 Forecasting climate change response in an alpine specialist songbird reveals the
20 869 importance of considering novel climate. *Diversity and Distributions*, 28.
21 870 <https://doi.org/10.1111/ddi.13628>
22 871 Di Benedetto, D., Musumeci, S. A., Avola, E., Alberti, A., Buono, S., Scuderi, C.,
23 872 Grillo, L., Galesi, O., Spalletta, A., Giudice, M. L., Luciano, D., Vinci, M., Bianca, S.,
24 873 Romano, C., & Fichera, M. (2014). Definition of minimal duplicated region
25 874 encompassing the XIAP and STAG2 genes in the Xq25 microduplication syndrome.
26 875 *American Journal of Medical Genetics. Part A*, 164A(8), 1923–1930.
27 876 <https://doi.org/10.1002/ajmg.a.36570>
28 877 Diebold, I., Petry, A., Hess, J., & Görlach, A. (2010). The NADPH Oxidase Subunit
29 878 NOX4 Is a New Target Gene of the Hypoxia-inducible Factor-1. *Molecular Biology of*
30 879 *the Cell*, 21(12), 2087–2096. <https://doi.org/10.1091/mbc.e09-12-1003>
31 880 Ducrest, A.-L., San-Jose, L. M., Neuenschwander, S., Schmid-Siegert, E., Simon, C.,
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- Pagni, M., Iseli, C., Richter, H., Guex, N., Cumer, T., Beaudoin, E., Dupasquier, M.,
Charruau, P., Ducouret, P., Xenarios, I., Goudet, J., & Roulin, A. (2025). Melanin and
Neurotransmitter Signalling Genes Are Differentially Co-Expressed in Growing
Feathers of White and Rufous Barn Owls. *Pigment Cell & Melanoma Research*,
38(2), e70001. <https://doi.org/10.1111/pcmr.70001>
- Egbert, J. R., & Belthoff, J. R. (2003). Wing Shape in House Finches Differs Relative to
Migratory Habit in Eastern and Western North America. *The Condor*, 105(4), 825–
829. <https://doi.org/10.1093/condor/105.4.825>
- Endler, J. A. (1977). *Geographic Variation, Speciation and Clines. (MPB-10), Volume*
10. Princeton University Press. <https://doi.org/10.2307/j.ctvx5wbdg>
- Epanchin, P. N., Knapp, R. A., & Lawler, S. P. (2010). Nonnative trout impact an
alpine-nesting bird by altering aquatic-insect subsidies. *Ecology*, 91(8), 2406–2415.
<https://doi.org/10.1890/09-1974.1>
- Faria, R., Renaut, S., Galindo, J., Pinho, C., Melo-Ferreira, J., Melo, M., Jones, F.,
Salzburger, W., Schluter, D., & Butlin, R. (2014). Advances in Ecological Speciation:
An integrative approach. *Molecular Ecology*, 23(3), 513–521.
<https://doi.org/10.1111/mec.12616>
- Farias-Virgens, M., Peede, D., Deacon, T., Okanoya, K., White, S. A., & Huerta-
Sanchez, E. (2025). The genomics of the domestication syndrome in a songbird
model species. *Communications Biology*, 8(1), 853.
<https://doi.org/10.1038/s42003-025-08235-0>
- Fedorova, E. S., Dementieva, N. V., Shcherbakov, Y. S., & Stanishevskaya, O. I.

1
2
3
4
5
6
7
8
9
10 903 (2022a). Identification of Key Candidate Genes in Runs of Homozygosity of the
11 904 Genome of Two Chicken Breeds, Associated with Cold Adaptation. *Biology*, 11(4),
12 Article 4. <https://doi.org/10.3390/biology11040547>
13 905
14
15 906 Feng, J., Zhu, W., Shi, H., Peng, D., Zang, L., Wang, Y., ZhaXi, L., BaiMa, J., Amevor,
16
17 907 F. K., Wang, X., Ma, X., & Zhao, X. (2023). Analysis of the Selection Signal of the
18
19 908 Tibetan Black Chicken Genome Based on Whole-Genome Sequencing. *Genes*,
20
21 909 14(9), Article 9. <https://doi.org/10.3390/genes14091672>
22
23 910 Flanagan, S. P., Forester, B. R., Latch, E. K., Aitken, S. N., & Hoban, S. (2018).
24
25 911 Guidelines for planning genomic assessment and monitoring of locally adaptive
26
27 912 variation to inform species conservation. *Evolutionary Applications*, 11(7), 1035–
28 913 1052. <https://doi.org/10.1111/eva.12569>
29
30 914 Forester, B. R., Beever, E. A., Darst, C., Szymanski, J., & Funk, W. C. (2022). Linking
31
32 915 evolutionary potential to extinction risk: Applications and future directions.
33
34 916 *Frontiers in Ecology and the Environment*, 20(9), 507–515.
35
36 917 <https://doi.org/10.1002/fee.2552>
37
38 918 Forester, B. R., Day, C. C., Ruegg, K., & Landguth, E. L. (2023). Evolutionary potential
39
40 919 mitigates extinction risk under climate change in the endangered southwestern
41 920 willow flycatcher. *Journal of Heredity*, 114(4), 341–353.
42
43 921 <https://doi.org/10.1093/jhered/esac067>
44
45 922 Franchini, P., Irisarri, I., Fudickar, A., Schmidt, A., Meyer, A., Wikelski, M., &
46
47 923 Partecke, J. (2017). Animal tracking meets migration genomics: Transcriptomic
48
49 924 analysis of a partially migratory bird species. *Molecular Ecology*, 26(12), 3204–3216.

- <https://doi.org/10.1111/mec.14108>
- Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018). Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the National Academy of Sciences*, 115(47), 11982–11987. <https://doi.org/10.1073/pnas.1804224115>
- Fridolfsson, A.-K., & Ellegren, H. (2000). Molecular Evolution of the Avian CHD1 Genes on the Z and W Sex Chromosomes. *Genetics*, 155(4), 1903–1912. <https://doi.org/10.1093/genetics/155.4.1903>
- Funk, E. R., Spellman, G. M., Winker, K., Withrow, J. J., Ruegg, K. C., & Taylor, S. A. (2023). The genetic basis of plumage coloration and elevation adaptation in a clade of recently diverged alpine and arctic songbirds. *Evolution; International Journal of Organic Evolution*, 77(3), 705–717. <https://doi.org/10.1093/evolut/qpac064>
- Gao, G., Zhang, K., Zhao, X., Xu, G., Xie, Y., Zhou, L., Zhang, C., & Wang, Q. (2023). Identification of Molecular Markers Associated with Goose Egg Quality Through Genome-Wide Association Analysis. *Scientia Agricultura Sinica*, 56(19), 3894–3904. <https://doi.org/10.3864/j.issn.0578-1752.2023.19.015>
- García, N. C., Campagna, L., Rush, A. C., Bowie, R. C. K., & Lovette, I. J. (2023). Comparative genomics of two Empidonax flycatchers reveal candidate genes for bird song production. *Evolution*, 77(8), 1818–1828. <https://doi.org/10.1093/evolut/qpac096>
- Gesemann, M., & Neuhauss, S. C. F. (2023). Evolution of visual guanylyl cyclases and their activating proteins with respect to clade and species-specific visual

1
2
3
4
5
6
7
8
9
10 947 system adaptation. *Frontiers in Molecular Neuroscience*, 16.
11 948 <https://doi.org/10.3389/fnmol.2023.1131093>
12
13 949 Gheyas, A. A., Vallejo-Trujillo, A., Kebede, A., Lozano-Jaramillo, M., Dessie, T.,
14
15 950 Smith, J., & Hanotte, O. (2021). Integrated Environmental and Genomic Analysis
16
17 951 Reveals the Drivers of Local Adaptation in African Indigenous Chickens. *Molecular*
18
19 952 *Biology and Evolution*, 38(10), 4268–4285.
20
21 953 <https://doi.org/10.1093/molbev/msab156>
22
23 954 Gienapp, P., Laine, V. N., Mateman, A. C., van Oers, K., & Visser, M. E. (2017).
24
25 955 Environment-Dependent Genotype-Phenotype Associations in Avian Breeding Time.
26
27 956 *Frontiers in Genetics*, 8. <https://doi.org/10.3389/fgene.2017.00102>
28
29 957 Grabherr, G., Gottfried, M., & Pauli, H. (2010). Climate Change Impacts in Alpine
30
31 958 Environments. *Geography Compass*, 4(8), 1133–1153.
32
33 959 <https://doi.org/10.1111/j.1749-8198.2010.00356.x>
34
35 960 Grant, P. R., Grant, B. R., Smith, J. N., Abbott, I. J., & Abbott, L. K. (1976). Darwin's
36
37 961 finches: Population variation and natural selection. *Proceedings of the National*
38
39 962 *Academy of Sciences*, 73(1), 257–261. <https://doi.org/10.1073/pnas.73.1.257>
40
41 963 Greenberg, R., Cadena, V., Danner, R. M., & Tattersall, G. (2012). Heat Loss May
42
43 964 Explain Bill Size Differences between Birds Occupying Different Habitats. *PLOS*
44
45 965 *ONE*, 7(7), e40933. <https://doi.org/10.1371/journal.pone.0040933>
46
47 966 Grilli, M. G., Lambertucci, S. A., Therrien, J.-F., & Bildstein, K. L. (2017). Wing size
48
49 967 but not wing shape is related to migratory behavior in a soaring bird. *Journal of Avian*
50
51
52
53
54
55
56
57
58
59
60 968 *Biology*, 48(5), 669–678. <https://doi.org/10.1111/jav.01220>

- Grinnell, J. (1913). *Leucosticte tephrocotis dawsoni*: A New Race of Rosy Finch from the Sierra Nevada. *The Condor*, 15(2), 76–79. <https://doi.org/10.2307/1362056>
- Grinnell, J. (1917). Field Tests of Theories Concerning Distributional Control. *The American Naturalist*, 51(602), 115–128. <https://doi.org/10.1086/279591>
- Guo, J., Sun, C., Qu, L., Shen, M., Dou, T., Ma, M., Wang, K., & Yang, N. (2017). Genetic architecture of bone quality variation in layer chickens revealed by a genome-wide association study. *Scientific Reports*, 7(1), 45317. <https://doi.org/10.1038/srep45317>
- He, J., Tu, J., Yu, J., & Jiang, H. (2023). A global assessment of Bergmann's rule in mammals and birds. *Global Change Biology*, 29(18), 5199–5210. <https://doi.org/10.1111/gcb.16860>
- Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., Poss, M. L., Reed, L. K., Storfer, A., & Whitlock, M. C. (2016). Finding the Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. *The American Naturalist*, 188(4), 379–397. <https://doi.org/10.1086/688018>
- Hu, X., Zhang, X., Liu, Z., Li, S., Zheng, X., Nie, Y., Tao, Y., Zhou, X., Wu, W., Yang, G., Zhao, Q., Zhang, Y., Xu, Q., & Mou, C. (2020). Exploration of key regulators driving primary feather follicle induction in goose skin. *Gene*, 731, 144338. <https://doi.org/10.1016/j.gene.2020.144338>
- Johnsson, M., Jonsson, K. B., Andersson, L., Jensen, P., & Wright, D. (2015). Genetic Regulation of Bone Metabolism in the Chicken: Similarities and Differences to Mammalian Systems. *PLOS Genetics*, 11(5), e1005250.

1
2
3
4
5
6
7
8
9
10 991 <https://doi.org/10.1371/journal.pgen.1005250>
11 992 Johnston, R. A., Paxton, K. L., Moore, F. R., Wayne, R. K., & Smith, T. B. (2016).
12 993 Seasonal gene expression in a migratory songbird. *Molecular Ecology*, 25(22), 5680–
13 994 5691. <https://doi.org/10.1111/mec.13879>
14 995 Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology*
15 996 *Letters*, 7(12), 1225–1241. <https://doi.org/10.1111/j.1461-0248.2004.00684.x>
16 997 Kim, H., Kim, H., Seong, P., Arora, D., Shin, D., Park, W., & Park, J.-E. (2021).
17 998 Transcriptomic Response under Heat Stress in Chickens Revealed the Regulation of
18 999 Genes and Alteration of Metabolism to Maintain Homeostasis. *Animals: An Open*
19 1000 *Access Journal from MDPI*, 11(8), 2241. <https://doi.org/10.3390/ani11082241>
20 1001 King, A. S., & McLelland, J. (1984). *Birds: Their structure and function* (2nd ed).
21 1002 Baillière Tindall.
22 1003 Kinsella, C. M., Ruiz-Ruano, F. J., Dion-Côté, A.-M., Charles, A. J., Gossmann, T. I.,
23 1004 Cabrero, J., Kappei, D., Hemmings, N., Simons, M. J. P., Camacho, J. P. M.,
24 1005 Forstmeier, W., & Suh, A. (2019). Programmed DNA elimination of germline
25 1006 development genes in songbirds. *Nature Communications*, 10(1), 5468.
26 1007 <https://doi.org/10.1038/s41467-019-13427-4>
27 1008 Körner, C. (2003). Alpine climate. In C. Körner (Ed.), *Alpine Plant Life: Functional*
28 1009 *Plant Ecology of High Mountain Ecosystems* (pp. 21–30). Springer.
29 1010 https://doi.org/10.1007/978-3-642-18970-8_3
30 1011 Koskenpato, K., Ahola, K., Karstinen, T., & Karell, P. (2016). Is the denser contour
31 1012 feather structure in pale grey than in pheomelanic brown tawny owls *Strix aluco* an

- adaptation to cold environments? *Journal of Avian Biology*, 47(1), 1–6.
- <https://doi.org/10.1111/jav.00746>
- Kuttiyarthu Veetil, N., Henschen, A. E., Hawley, D. M., Melepat, B., Dalloul, R. A., Beneš, V., Adelman, J. S., & Vinkler, M. (2024). Varying conjunctival immune response adaptations of house finch populations to a rapidly evolving bacterial pathogen. *Frontiers in Immunology*, 15.
- <https://doi.org/10.3389/fimmu.2024.1250818>
- Lamichhaney, S., Han, F., Webster, M. T., Andersson, L., Grant, B. R., & Grant, P. R. (2018). Rapid hybrid speciation in Darwin's finches. *Science*, 359(6372), 224–228.
- <https://doi.org/10.1126/science.aao4593>
- Lande, R., & Shannon, S. (1996). The role of genetic variation in adaptation and population persistence in a changing environment. *Evolution*, 50(1), 434–437.
- <https://doi.org/10.1111/j.1558-5646.1996.tb04504.x>
- Lawson, L. P., & Petren, K. (2017). The adaptive genomic landscape of beak morphology in Darwin's finches. *Molecular Ecology*, 26(19), 4978–4989.
- <https://doi.org/10.1111/mec.14166>
- Lei, F.-M., Qu, Y.-H., Gan, Y.-L., Gebauer, A., & Kaiser, M. (2002). The feather microstructure of Passerine sparrows in China. *Journal Für Ornithologie*, 143(2), 205–212. <https://doi.org/10.1046/j.1439-0361.2002.02001.x>
- Lennon, O. (2022). *Do Candidate Genes Distinguish Migratory and Nonmigratory Avian Species?* [Undergraduate Research Scholars Thesis, Texas A&M University]. <https://hdl.handle.net/1969.1/196498>

1
2
3
4
5
6
7
8
9
10 1035 Lenth, R. (2025). *emmeans: Estimated Marginal Means, aka Least-Squares Means*
11 1036 (Version 1.10.7-100002) [R package].
12
13 1037 Lewden, A., Bishop, C. M., & Askew, G. N. (2023). How birds dissipate heat before,
14 during and after flight. *Journal of the Royal Society Interface*, 20(209), 20230442.
15 1038 <https://doi.org/10.1098/rsif.2023.0442>
16
17 1039
18
19 1040 Li, G.-S., Liu, W.-W., Zhang, F., Zhu, F., Yang, F.-X., Hao, J.-P., & Hou, Z.-C. (2020).
20 Genome-wide association study of bone quality and feed efficiency-related traits in
21 1041 Pekin ducks. *Genomics*, 112(6), 5021–5028.
22 1042 <https://doi.org/10.1016/j.ygeno.2020.09.023>
23 1043
24 1044 Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–
25 1045 Wheeler transform. *Bioinformatics*, 25(14), 1754–1760.
26 1046 <https://doi.org/10.1093/bioinformatics/btp324>
27
28 1047 Limber, C., Wagner, G. P., & Prum, R. O. (2024). *Genetic Characterization of the Cell*
29 1048 *Types of in Developing Feathers, and the Evolution of Feather Complexity* (p.
30 1049 2024.11.21.624532). bioRxiv. <https://doi.org/10.1101/2024.11.21.624532>
31 1050 Lockwood, R., Swaddle, J. P., & Rayner, J. M. V. (1998). Avian Wingtip Shape
32 1051 Reconsidered: Wingtip Shape Indices and Morphological Adaptations to Migration.
33 1052 *Journal of Avian Biology*, 29(3), 273–292. <https://doi.org/10.2307/3677110>
34 1053 Loma, J. D., Vicente, M., Tirado, N., Ascui, F., Parada, L. A., Gardon, J., Schlebusch,
35 1054 C., & Broberg, K. (2025). Human adaptation in the Andes Mountains. *Human*
36 1055 *Population Genetics and Genomics*, 5(1).
37 1056 <https://doi.org/10.47248/hpgg2505010002>
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- Lotterhos, K. E., & Whitlock, M. C. (2015). The relative power of genome scans to detect local adaptation depends on sampling design and statistical method. *Molecular Ecology*, 24(5), 1031–1046. <https://doi.org/10.1111/mec.13100>
- Lou, R. N., & Therkildsen, N. O. (2022). Batch effects in population genomic studies with low-coverage whole genome sequencing data: Causes, detection and mitigation. *Molecular Ecology Resources*, 22(5), 1678–1692. <https://doi.org/10.1111/1755-0998.13559>
- Lovell, P. V., Carleton, J. B., & Mello, C. V. (2013). Genomics analysis of potassium channel genes in songbirds reveals molecular specializations of brain circuits for the maintenance and production of learned vocalizations. *BMC Genomics*, 14(1), 470. <https://doi.org/10.1186/1471-2164-14-470>
- Luo, X., Guo, J., Zhang, J., Ma, Z., & Li, H. (2024). Overview of chicken embryo genes related to sex differentiation. *PeerJ*, 12, e17072. <https://doi.org/10.7717/peerj.17072>
- Ma, S., Liu, H., Wang, J., Wang, L., Xi, Y., Liu, Y., Xu, Q., Hu, J., Han, C., Bai, L., Li, L., & Wang, J. (2021). Transcriptome Analysis Reveals Genes Associated With Sexual Dichromatism of Head Feather Color in Mallard. *Frontiers in Genetics*, 12. <https://doi.org/10.3389/fgene.2021.627974>
- MacDougall-Shackleton, S. A., Johnson, R. E., & Hahn, T. P. (2000). Gray-crowned Rosy-Finch (*Leucosticte tephrocotis*). In *The Birds of North America*. Cornell Lab of Ornithology.
- Mamantov, M. A., Gibson-Reinemer, D. K., Linck, E. B., & Sheldon, K. S. (2021).

1
2
3
4
5
6
7
8
9
10 1079 Climate-driven range shifts of montane species vary with elevation. *Global Ecology*
11 and *Biogeography*, 30(4), 784–794. <https://doi.org/10.1111/geb.13246>
12 1080
13 1081 Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., & Zimin, A.
14 (2018). MUMmer4: A fast and versatile genome alignment system. *PLOS*
15 1082 *Computational Biology*, 14(1), e1005944.
16
17 1083 <https://doi.org/10.1371/journal.pcbi.1005944>
18 1084
19 1085 Mascarenhas, R., Meirelles, P. M., & Batalha-Filho, H. (2023). Urbanization drives
20 adaptive evolution in a Neotropical bird. *Current Zoology*, 69(5), 607–619.
21 1086
22 1087 <https://doi.org/10.1093/cz/zoac066>
23
24 1088 Matsuda, Y., & Makino, T. (2024). Comparative genomics reveals convergent signals
25 associated with the high metabolism and longevity in birds and bats. *Proceedings of*
26 1089 *the Royal Society B: Biological Sciences*, 291(2029), 20241068.
27
28 1090 <https://doi.org/10.1098/rspb.2024.1068>
29
30 1091 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernysky, A.,
31 Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The
32 1092 Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation
33 DNA sequencing data. *Genome Research*, 20(9), 1297–1303.
34 1093
35 1094 <https://doi.org/10.1101/gr.107524.110>
36
37 1095 Medina, I., Newton, E., Kearney, M. R., Mulder, R. A., Porter, W. P., & Stuart-Fox, D.
38 (2018). Reflection of near-infrared light confers thermal protection in birds. *Nature*
39 1096 *Communications*, 9(1), 3610. <https://doi.org/10.1038/s41467-018-05898-8>
40
41 1100 Meek, M. H., Beever, E. A., Barbosa, S., Fitzpatrick, S. W., Fletcher, N. K., Mittan-

- Moreau, C. S., Reid, B. N., Campbell-Staton, S. C., Green, N. F., & Hellmann, J. J. (2023). Understanding Local Adaptation to Prepare Populations for Climate Change. *BioScience*, 73(1), 36–47. <https://doi.org/10.1093/biosci/biac101>
- Meiri, S., & Dayan, T. (2003). On the validity of Bergmann's rule. *Journal of Biogeography*, 30(3), 331–351. <https://doi.org/10.1046/j.1365-2699.2003.00837.x>
- Mesner, L. D., Ray, B., Hsu, Y.-H., Manichaikul, A., Lum, E., Bryda, E. C., Rich, S. S., Rosen, C. J., Criqui, M. H., Allison, M., Budoff, M. J., Clemens, T. L., & Farber, C. R. (2014). *Bicc1* is a genetic determinant of osteoblastogenesis and bone mineral density. *The Journal of Clinical Investigation*, 124(6), 2736–2749. <https://doi.org/10.1172/JCI73072>
- Mohammadi, H. (2024a). A Genome-wide Association Study based on Pathway Analysis Related to the Angle Wing in Pekin Ducks using Whole-Genome Sequencing Data. *Research On Animal Production*, 15(2), 32–43. <https://doi.org/10.61186/rap.15.2.32>
- Muto, A., Ikeda, S., Lopez-Burks, M. E., Kikuchi, Y., Calof, A. L., Lander, A. D., & Schilling, T. F. (2014). Nipbl and Mediator Cooperatively Regulate Gene Expression to Control Limb Development. *PLOS Genetics*, 10(9), e1004671. <https://doi.org/10.1371/journal.pgen.1004671>
- Nan, J., Yang, S., Zhang, X., Leng, T., Zhuoma, J., Zhuoma, R., Yuan, J., Pi, J., Sheng, Z., & Li, S. (2023). Identification of candidate genes related to highland adaptation from multiple Chinese local chicken breeds by whole genome sequencing analysis. *Animal Genetics*, 54(1), 55–67. <https://doi.org/10.1111/age.13268>

1
2
3
4
5
6
7
8
9
10 1123 Owen, J. C. (2011). Collecting, processing, and storing avian blood: A review.
11 1124 *Journal of Field Ornithology*, 82(4), 339–354. <https://doi.org/10.1111/j.1557->
12 9263.2011.00338.x
13 1125
14
15 1126 Pap, P. L., Osváth, G., Daubner, T., Nord, A., & Vincze, O. (2020). Down feather
16 morphology reflects adaptation to habitat and thermal conditions across the avian
17 1127 phylogeny. *Evolution*, 74(10), 2365–2376. <https://doi.org/10.1111/evo.14075>
18
19 1128 Pap, P. L., Vincze, O., Wekerle, B., Daubner, T., Vágási, C. I., Nudds, R. L., Dyke, G.
20 J., & Osváth, G. (2017). A phylogenetic comparative analysis reveals correlations
21 1129 between body feather structure and habitat. *Functional Ecology*, 31(6), 1241–1251.
22
23 1130 <https://doi.org/10.1111/1365-2435.12820>
24 1131
25 1132
26 1133 Park, H.-B., Kim, J.-W., & Baek, K.-H. (2020). Regulation of Wnt Signaling through
27 Ubiquitination and Deubiquitination in Cancers. *International Journal of Molecular*
28 1134 *Sciences*, 21(11), Article 11. <https://doi.org/10.3390/ijms21113904>
29
30 1135 Pelletier, F., & Coltman, D. W. (2018). Will human influences on evolutionary
31 dynamics in the wild pervade the Anthropocene? *BMC Biology*, 16(1), 7.
32 1136 <https://doi.org/10.1186/s12915-017-0476-1>
33 1137
34 1138 Poplin, R., Ruano-Rubio, V., DePristo, M. A., Fennell, T. J., Carneiro, M. O., Auwera,
35 G. A. V. der, Kling, D. E., Gauthier, L. D., Levy-Moonshine, A., Roazen, D., Shakir, K.,
36 1139 Thibault, J., Chandran, S., Whelan, C., Lek, M., Gabriel, S., Daly, M. J., Neale, B.,
37 1140 MacArthur, D. G., & Banks, E. (2018). *Scaling accurate genetic variant discovery to*
38 *tens of thousands of samples* (p. 201178). bioRxiv. <https://doi.org/10.1101/201178>
39 1141
40 1142 Pritchard, J. K., & Di Rienzo, A. (2010). Adaptation – not by sweeps alone. *Nature*
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1145 *Reviews Genetics*, 11(10), 665–667. <https://doi.org/10.1038/nrg2880>
- 1146 Prost, S., Armstrong, E. E., Nylander, J., Thomas, G. W. C., Suh, A., Petersen, B.,
- 1147 Dalen, L., Benz, B. W., Blom, M. P. K., Palkopoulou, E., Ericson, P. G. P., & Irestedt,
- 1148 M. (2018). *Genome analyses show strong selection on coloration, morphological*
- 1149 *and behavioral phenotypes in birds-of-paradise* (p. 287086). bioRxiv.
- 1150 <https://doi.org/10.1101/287086>
- 1151 Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D.,
- 1152 Maller, J., Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A
- 1153 Tool Set for Whole-Genome Association and Population-Based Linkage Analyses.
- 1154 *American Journal of Human Genetics*, 81(3), 559–575.
- 1155 Pyle, P. (2022). *Identification Guide to North American Birds, Part I* (Second Edition).
- 1156 Slate Creek Press.
- 1157 Qi, X., Zhang, Q., He, Y., Yang, L., Zhang, X., Shi, P., Yang, L., Liu, Z., Zhang, F., Liu,
- 1158 F., Liu, S., Wu, T., Cui, C., Ouzhuluobu, Bai, C., Baimakangzhuo, Han, J., Zhao, S.,
- 1159 Liang, C., & Su, B. (2019). The Transcriptomic Landscape of Yaks Reveals Molecular
- 1160 Pathways for High Altitude Adaptation. *Genome Biology and Evolution*, 11(1), 72–85.
- 1161 <https://doi.org/10.1093/gbe/evy264>
- 1162 Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for
- 1163 comparing genomic features. *Bioinformatics*, 26(6), 841–842.
- 1164 <https://doi.org/10.1093/bioinformatics/btq033>
- 1165 Ricchetti, I., Taucher, T., Loebick, R., Sin, S. Y. W., Palacios, C., & Lamichhaney, S.
- 1166 (2024). Gene expression variation in geographically diverse populations of two

1
2
3
4
5
6
7
8
9
10 1167 North American songbird species. *Frontiers in Bird Science*, 3.
11 1168 <https://doi.org/10.3389/fbirs.2024.1382657>
12
13 1169 Rodriguez, M. D., Bay, R. A., & Ruegg, K. C. (2025). Telomere Length Differences
14
15 1170 Indicate Climate Change-Induced Stress and Population Decline in a Migratory Bird.
16
17 1171 *Molecular Ecology*, 34(21), e17642. <https://doi.org/10.1111/mec.17642>
18
19 1172 Rogalla, S., Shawkey, M. D., & D’Alba, L. (2022). Thermal effects of plumage
20
21 1173 coloration. *Ibis*, 164(4), 933–948. <https://doi.org/10.1111/ibi.13100>
22
23 1174 Romano, A., Séchaud, R., Hirzel, A. H., & Roulin, A. (2019). Climate-driven
24
25 1175 convergent evolution of plumage colour in a cosmopolitan bird. *Global Ecology and*
26 1176 *Biogeography*, 28(4), 496–507. <https://doi.org/10.1111/geb.12870>
27
28 1177 Romanov, M. N., Abdelmanova, A. S., Fisinin, V. I., Gladyr, E. A., Volkova, N. A.,
29
30 1178 Koshkina, O. A., Rodionov, A. N., Vetokh, A. N., Gusev, I. V., Anshakov, D. V.,
31
32 1179 Stanishevskaya, O. I., Dotsev, A. V., Griffin, D. K., & Zinovieva, N. A. (2023). Selective
33
34 1180 footprints and genes relevant to cold adaptation and other phenotypic traits are
35
36 1181 unscrambled in the genomes of divergently selected chicken breeds. *Journal of*
37 1182 *Animal Science and Biotechnology*, 14(1), 35. [https://doi.org/10.1186/s40104-022-](https://doi.org/10.1186/s40104-022-00813-0)
38
39 1183 00813-0
40
41 1184 Rundel, P. W., Gibson, A. C., & Sharifi, M. R. (2008). The Alpine Flora of the White
42
43 1185 Mountains, California. *Madroño*, 55(3), 202–215. [https://doi.org/10.3120/0024-](https://doi.org/10.3120/0024-9637-55.3.202)
44
45 1186 9637-55.3.202
46
47 1187 Rundel, P. W., & Millar, C. I. (2016). Alpine Ecosystems. In H. Mooney & E. Zavaleta
48
49 1188 (Eds.), *Ecosystems of California* (pp. 613–634). University of California Press.
50
51
52
53
54
55
56
57
58
59
60

- <https://www.degruyterbrill.com/document/doi/10.1525/9780520962170-033/pdf?licenseType=restricted>
- Sadeesh, E. M., Lahamge, M. S., Malik, A., & Ampadi, A. N. (2025). Nuclear Genome-Encoded Mitochondrial OXPHOS Complex I Genes in Female Buffalo Show Tissue-Specific Differences. *Molecular Biotechnology*, 67(6), 2411–2427. <https://doi.org/10.1007/s12033-024-01206-6>
- Sandoval, L., & Barrantes, G. (2019). Is black plumage an adaptation to high elevations in a cosmopolitan bird genus? *Journal of Avian Biology*, 50(7). <https://doi.org/10.1111/jav.02041>
- Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation. *Nature Reviews Genetics*, 14(11), 807–820. <https://doi.org/10.1038/nrg3522>
- Schweizer, R. M., Jones, M. R., Bradburd, G. S., Storz, J. F., Senner, N. R., Wolf, C., & Cheviron, Z. A. (2021a). Broad Concordance in the Spatial Distribution of Adaptive and Neutral Genetic Variation across an Elevational Gradient in Deer Mice. *Molecular Biology and Evolution*, 38(10), 4286–4300. <https://doi.org/10.1093/molbev/msab161>
- Sebastianelli, M., Lukhele, S. M., Secomandi, S., de Souza, S. G., Haase, B., Moysi, M., Nikiforou, C., Hutflus, A., Mountcastle, J., Balacco, J., Pelan, S., Chow, W., Fedrigo, O., Downs, C. T., Monadjem, A., Dingemanse, N. J., Jarvis, E. D., Brelsford, A., vonHoldt, B. M., & Kirschel, A. N. G. (2024). A genomic basis of vocal rhythm in birds. *Nature Communications*, 15(1), 3095. <https://doi.org/10.1038/s41467-024->

1
2
3
4
5
6
7
8
9
10 1211 47305-5
11 1212 Seki, R., Li, C., Fang, Q., Hayashi, S., Egawa, S., Hu, J., Xu, L., Pan, H., Kondo, M.,
12
13 1213 Sato, T., Matsubara, H., Kamiyama, N., Kitajima, K., Saito, D., Liu, Y., Gilbert, M. T.
14
15 1214 P., Zhou, Q., Xu, X., Shiroishi, T., ... Zhang, G. (2017). Functional roles of Aves class-
16
17 1215 specific cis-regulatory elements on macroevolution of bird-specific features.
18
19 1216 *Nature Communications*, 8(1), 14229. <https://doi.org/10.1038/ncomms14229>
20
21 1217 Semenov, G. A., Sonnenberg, B. R., Branch, C. L., Heinen, V. K., Welklin, J. F.,
22
23 1218 Padula, S. R., Patel, A. M., Bridge, E. S., Pravosudov, V. V., & Taylor, S. A. (2024).
24
25 1219 Genes and gene networks underlying spatial cognition in food-caching chickadees.
26
27 1220 *Current Biology*, 34(9), 1930-1939.e4. <https://doi.org/10.1016/j.cub.2024.03.058>
28
29 1221 Seutin, G., White, B. N., & Boag, P. T. (1991). Preservation of avian blood and tissue
30
31 1222 samples for DNA analyses. *Canadian Journal of Zoology*, 69(1), 82–90.
32
33 1223 <https://doi.org/10.1139/z91-013>
34
35 1224 Shafer, A. B. A., Wolf, J. B. W., Alves, P. C., Bergström, L., Bruford, M. W.,
36
37 1225 Brännström, I., Colling, G., Dalén, L., De Meester, L., Ekblom, R., Fawcett, K. D.,
38
39 1226 Fior, S., Hajibabaei, M., Hill, J. A., Hoezel, A. R., Höglund, J., Jensen, E. L., Krause, J.,
40
41 1227 Kristensen, T. N., ... Zieniński, P. (2015). Genomics and the challenging translation
42
43 1228 into conservation practice. *Trends in Ecology & Evolution*, 30(2), 78–87.
44
45 1229 <https://doi.org/10.1016/j.tree.2014.11.009>
46
47 1230 Shakya, S. B. (2020). *Genomics and Population History of Black-Headed Bulbul*
48
49 1231 (*Brachypodius Atriceps*) Color Morphs [Ph.D., Louisiana State University and
50
51
52
53
54
55
56
57
58
59
60 1232 Agricultural & Mechanical College].

- <https://www.proquest.com/docview/2665131551/abstract/2200EDB181A4591PQ/>
- 1
- Sirkiä, P. M., Virolainen, M., & Laaksonen, T. (2010). Melanin coloration has temperature-dependent effects on breeding performance that may maintain phenotypic variation in a passerine bird. *Journal of Evolutionary Biology*, 23(11), 2385–2396. <https://doi.org/10.1111/j.1420-9101.2010.02100.x>
- Stettenheim, P. R. (2000). The Integumentary Morphology of Modern Birds—An Overview. *American Zoologist*, 40(4), 461–477. <https://doi.org/10.1093/icb/40.4.461>
- Steven, S. (2021). *Genome-Wide Analyses of Genomic Variation Regulating Differential Stress Responses in Japanese Quail* [Doctoral dissertation, University of Arkansas]. ProQuest Dissertations & Theses.
- Stinchcombe, J. R., & Hoekstra, H. E. (2008). Combining population genomics and quantitative genetics: Finding the genes underlying ecologically important traits. *Heredity*, 100(2), 158–170. <https://doi.org/10.1038/sj.hdy.6800937>
- Stoutjesdijk, F. (2003). Birds in the cold: The effects of plumage structure and environment on operative temperature, shown by spherical models. *Ornis Svecica*, 13(4), Article 4. <https://doi.org/10.34080/os.v13.22804>
- Struk, A. A., Mugon, J., Huston, A., Scholer, A. A., Stadler, G., Higgins, E. T., Sokolowski, M. B., & Danckert, J. (2019). Self-regulation and the foraging gene (PRKG1) in humans. *Proceedings of the National Academy of Sciences*, 116(10), 4434–4439. <https://doi.org/10.1073/pnas.1809924116>
- Sun, L., Zhou, T., Wan, Q.-H., & Fang, S.-G. (2020). Transcriptome Comparison

1
2
3
4
5
6
7
8
9
10 1255 Reveals Key Components of Nuptial Plumage Coloration in Crested Ibis.
11 1256 *Biomolecules*, 10(6), Article 6. <https://doi.org/10.3390/biom10060905>
12
13 1257 Sun, Y., Li, Y., Jiang, X., Wu, Q., Lin, R., Chen, H., Zhang, M., Zeng, T., Tian, Y., Xu, E.,
14
15 1258 Zhang, Y., & Lu, L. (2024). Genome-wide association study identified candidate
16
17 1259 genes for egg production traits in the Longyan Shan-ma duck. *Poultry Science*,
18
19 1260 103(9), 104032. <https://doi.org/10.1016/j.psj.2024.104032>
20
21 1261 Symonds, M. R. E., & Tattersall, G. J. (2010). Geographical variation in bill size
22
23 1262 across bird species provides evidence for Allen’s rule. *The American Naturalist*,
24
25 1263 176(2), 188–197. <https://doi.org/10.1086/653666>
26 1264 Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable,
27
28 1265 A. L., Fang, T., Doncheva, N. T., Pyysalo, S., Bork, P., Jensen, L. J., & von Mering, C.
29
30 1266 (2023). The STRING database in 2023: Protein-protein association networks and
31
32 1267 functional enrichment analyses for any sequenced genome of interest. *Nucleic*
33
34 1268 *Acids Research*, 51(D1), D638–D646. <https://doi.org/10.1093/nar/gkac1000>
35
36 1269 Tam, V., Patel, N., Turcotte, M., Bossé, Y., Paré, G., & Meyre, D. (2019). Benefits and
37
38 1270 limitations of genome-wide association studies. *Nature Reviews Genetics*, 20(8),
39
40 1271 467–484. <https://doi.org/10.1038/s41576-019-0127-1>
41 1272 Tattersall, G. J., Arnaout, B., & Symonds, M. R. E. (2017). The evolution of the avian
42
43 1273 bill as a thermoregulatory organ. *Biological Reviews*, 92(3), 1630–1656.
44
45 1274 <https://doi.org/10.1111/brv.12299>
46
47 1275 Thomas, P. D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P., & Mi, H.
48
49 1276 (2022). PANTHER: Making genome-scale phylogenetics accessible to all. *Protein*

- 1277 *Science*, 31(1), 8–22. <https://doi.org/10.1002/pro.4218>
- 1278 Tian, S., Zhou, X., Phuntsok, T., Zhao, N., Zhang, D., Ning, C., Li, D., & Zhao, H.
- 1279 (2020). Genomic Analyses Reveal Genetic Adaptations to Tropical Climates in
- 1280 Chickens. *iScience*, 23(11). <https://doi.org/10.1016/j.isci.2020.101644>
- 1281 Twining, H. (1940). Foraging Behavior and Survival in the Sierra Nevada Rosy Finch.
- 1282 *The Condor*, 42(1), 64–72. <https://doi.org/10.2307/1364320>
- 1283 Twumasi, G., Wang, H., Xi, Y., Qi, J., Li, L., Bai, L., & Liu, H. (2024). Genome-Wide
- 1284 Association Studies Reveal Candidate Genes Associated with Pigmentation
- 1285 Patterns of Single Feathers of Tianfu Nonghua Ducks. *Animals*, 14(1), Article 1.
- 1286 <https://doi.org/10.3390/ani14010085>
- 1287 Urban, M. C. (2018). Escalator to extinction. *Proceedings of the National Academy*
- 1288 *of Sciences*, 115(47), 11871–11873. <https://doi.org/10.1073/pnas.1817416115>
- 1289 Videla Rodriguez, E. A., Mitchell, J. B. O., & Smith, V. A. (2022). A Bayesian network
- 1290 structure learning approach to identify genes associated with stress in spleens of
- 1291 chickens. *Scientific Reports*, 12(1), 7482. [https://doi.org/10.1038/s41598-022-](https://doi.org/10.1038/s41598-022-11633-7)
- 1292 [11633-7](https://doi.org/10.1038/s41598-022-11633-7)
- 1293 Wang, C., Li, S., Yu, W., Xin, Q., Li, C., Feng, Y., Peng, X., & Gong, Y. (2011). Cloning
- 1294 and expression profiling of the VLDLR gene associated with egg performance in
- 1295 duck (*Anas platyrhynchos*). *Genetics Selection Evolution*, 43(1), 29.
- 1296 <https://doi.org/10.1186/1297-9686-43-29>
- 1297 Ward, S., Rayner, J. M., Möller, U., Jackson, D. M., Nachtigall, W., & Speakman, J. R.
- 1298 (1999). Heat transfer from starlings *sturnus vulgaris* during flight. *Journal of*

1
2
3
4
5
6
7
8
9
10 1299 *Experimental Biology*, 202 (Pt 12), 1589–1602.
11 1300 <https://doi.org/10.1242/jeb.202.12.1589>
12
13 1301 Weeks, B. C., Harvey, C., Tobias, J. A., Sheard, C., Zhou, Z., & Fouhey, D. F. (2025).
14
15 1302 Longer Wing Bones in Warmer Climates Suggest a Role of Thermoregulation in Bird
16
17 1303 Wing Evolution. *Global Ecology and Biogeography*, 34(4), e70033.
18
19 1304 <https://doi.org/10.1111/geb.70033>
20
21 1305 Weeks, B. C., Willard, D. E., Zimova, M., Ellis, A. A., Witynski, M. L., Hennen, M., &
22
23 1306 Winger, B. M. (2020). Shared morphological consequences of global warming in
24
25 1307 North American migratory birds. *Ecology Letters*, 23(2), 316–325.
26
27 1308 <https://doi.org/10.1111/ele.13434>
28
29 1309 Whitlock, M. C., & Lotterhos, K. E. (2015). Reliable Detection of Loci Responsible for
30 1310 Local Adaptation: Inference of a Null Model through Trimming the Distribution of
31
32 1311 F(ST). *The American Naturalist*, 186 Suppl 1, S24-36.
33
34 1312 <https://doi.org/10.1086/682949>
35
36 1313 Wolf, B. O., & Walsberg, G. E. (2000). The Role of the Plumage in Heat Transfer
37
38 1314 Processes of Birds1. *American Zoologist*, 40(4), 575–584.
39
40 1315 <https://doi.org/10.1093/icb/40.4.575>
41 1316 Wu, Y., Yan, Y., Zhao, Y., Gu, L., Wang, S., & Johnson, D. H. (2021). Genomic bases
42
43 1317 underlying the adaptive radiation of core landbirds. *BMC Ecology and Evolution*,
44
45 1318 27(1), 162. <https://doi.org/10.1186/s12862-021-01888-5>
46
47 1319 Xiao, J., Wu, Y., Chen, R., Lin, Y., Wu, L., Tian, W., & Liu, L. (2008). Expression of
48
49 1320 Pcp4 gene during osteogenic differentiation of bone marrow mesenchymal stem

- cells in vitro. *Molecular and Cellular Biochemistry*, 309(1), 143–150.
- <https://doi.org/10.1007/s11010-007-9652-x>
- Xie, J., Tang, L., Lu, L., Zhang, L., Xi, L., Liu, H.-C., Odle, J., & Luo, X. (2014). Differential Expression of Heat Shock Transcription Factors and Heat Shock Proteins after Acute and Chronic Heat Stress in Laying Chickens (*Gallus gallus*). *PLOS ONE*, 9(7), e102204. <https://doi.org/10.1371/journal.pone.0102204>
- Xiong, D.-H., Liu, X.-G., Guo, Y.-F., Tan, L.-J., Wang, L., Sha, B.-Y., Tang, Z.-H., Pan, F., Yang, T.-L., Chen, X.-D., Lei, S.-F., Yerges, L. M., Zhu, X.-Z., Wheeler, V. W., Patrick, A. L., Bunker, C. H., Guo, Y., Yan, H., Pei, Y.-F., ... Deng, H.-W. (2009). Genome-wide Association and Follow-Up Replication Studies Identified ADAMTS18 and TGFBR3 as Bone Mass Candidate Genes in Different Ethnic Groups. *The American Journal of Human Genetics*, 84(3), 388–398. <https://doi.org/10.1016/j.ajhg.2009.01.025>
- Yan, C., Zhang, Z.-Y., Lv, Y., Wang, Z., Jiang, K., & Li, J.-T. (2022). Genome of *Laudakia sacra* Provides New Insights into High-Altitude Adaptation of Ectotherms. *International Journal of Molecular Sciences*, 23(17), Article 17. <https://doi.org/10.3390/ijms231710081>
- Yeaman, S. (2015). Local Adaptation by Alleles of Small Effect. *The American Naturalist*, 186 Suppl 1, S74–89. <https://doi.org/10.1086/682405>
- Zhang, C., Liu, C., Zhang, J., Zheng, L., Chang, Q., Cui, Z., & Liu, S. (2022). Analysis on the desert adaptability of indigenous sheep in the southern edge of Taklimakan Desert. *Scientific Reports*, 12(1), 12264. <https://doi.org/10.1038/s41598-022->

1
2
3
4
5
6
7
8
9
10 1343 15986-x
11 1344 Zhang, W., Fan, Z., Han, E., Hou, R., Zhang, L., Galaverni, M., Huang, J., Liu, H.,
12
13 1345 Silva, P., Li, P., Pollinger, J. P., Du, L., Zhang, X., Yue, B., Wayne, R. K., & Zhang, Z.
14
15 1346 (2014). Hypoxia Adaptations in the Grey Wolf (*Canis lupus chanco*) from Qinghai-
16
17 1347 Tibet Plateau. *PLOS Genetics*, 10(7), e1004466.
18
19 1348 <https://doi.org/10.1371/journal.pgen.1004466>
20
21 1349 Zhang, X., Yang, F., Zhu, T., Zhao, X., Zhang, J., Wen, J., Zhang, Y., Wang, G., Ren, X.,
22
23 1350 Chen, A., Wang, X., Wang, L., Lv, X., Yang, W., Qu, C., Wang, H., Ning, Z., & Qu, L.
24
25 1351 (2024). Whole genome resequencing reveals genomic regions related to red
26
27 1352 plumage in ducks. *Poultry Science*, 103(6), 103694.
28
29 1353 <https://doi.org/10.1016/j.psj.2024.103694>
30 1354 Zhang, X., Zhu, T., Wang, L., Lv, X., Yang, W., Qu, C., Li, H., Wang, H., Ning, Z., & Qu,
31
32 1355 L. (2023). Genome-Wide Association Study Reveals the Genetic Basis of Duck
33
34 1356 Plumage Colors. *Genes*, 14(4), Article 4. <https://doi.org/10.3390/genes14040856>
35
36 1357 Zheng, S., Ouyang, J., Liu, S., Tang, H., Xiong, Y., Yan, X., & Chen, H. (2023).
37
38 1358 Genomic signatures reveal selection in Lingxian white goose. *Poultry Science*,
39
40 1359 102(1), 102269. <https://doi.org/10.1016/j.psj.2022.102269>
41 1360 Zhou, X., Carbonetto, P., & Stephens, M. (2013). Polygenic Modeling with Bayesian
42
43 1361 Sparse Linear Mixed Models. *PLOS Genetics*, 9(2), e1003264.
44
45 1362 <https://doi.org/10.1371/journal.pgen.1003264>
46
47 1363 Zhuang, Z.-X., Chen, S.-E., Chen, C.-F., Lin, E.-C., & Huang, S.-Y. (2019). Genome-
48
49 1364 wide association study on the body temperature changes of a broiler-type strain
50
51
52
53
54
55
56
57
58
59
60

Taiwan country chickens under acute heat stress. *Journal of Thermal Biology*, 82, 33–42. <https://doi.org/10.1016/j.jtherbio.2019.03.007>

Zimova, M., Weeks, B. C., Willard, D. E., Giery, S. T., Jirinec, V., Burner, R. C., & Winger, B. M. (2023). Body size predicts the rate of contemporary morphological change in birds. *Proceedings of the National Academy of Sciences of the United States of America*, 120(20), e2206971120. <https://doi.org/10.1073/pnas.2206971120>

Conflicts of Interest

The authors declare no conflicts of interest.

Data Accessibility

Raw sequence data have been deposited in the NCBI Sequence Read Archive under BioProject accession PRJNA1297057.

Statistical analysis, GWAS and F_{ST} results are available at <http://datadryad.org/share/DqvYM4bZd7IZHEXy5O9FGkrQ86Q90YoGz3ZOAJoLHmY>

Code and scripts used for processing and analysis of sequence data are available at <https://github.com/erigande/mega-non-model-wgs-snakeflow/>. All other scripts as well as sample metadata are available on GitHub at https://github.com/ecnrobertson/GCRF_Pub.

Benefit-Sharing Statement

Benefits Generated: Benefits from this research accrue from the sharing of our data and results on public databases as described above.


Statement of Contributions

E.C.N.R., T.B., E.Z., K.C.R conceived of the study and E.C.N.R. wrote the majority of the manuscript and oversaw the majority of the analysis. C.M.B. contributed to the bioinformatic analysis. S.D. performed and conceived of the methodology for all feather morphology measurements. T.B. and E.Z. contributed ideas and performed all field sampling. K.C.R., E.Z. and M.B.H. acquired funding. All authors discussed the results and contributed to the final manuscript.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures

A Sampling Scheme



B GWAS Methods



C F_{ST} Methods




Figure 1. A) A map of California highlighting the sampling location and design for two populations of Grey-crowned Rosy-Finch, Piute Pass and White Mountain. [Key environmental variables at each site are noted: elevation is noted for each site along](#)

with an elevation transect (ELEV) and mean temperature of the warmest month (MTWM). B) Overview of methods for GWAS analysis. Morphological measurements were corrected for the effects of sex, age, and body size before being input into BSLMM and LMM models. C) Overview of the methods for the F_{ST} analysis. The output of both analyses were analyzed to identify significant SNPs and related genes.

Body Morphology Results




Figure 2. Violin plots of beak morphology raw measurement: A) beak width, B) nares length, and C) beak depth. Overlaid are the corrected mean trait measurements shown with a crossbar. Significance of the effect of site on the morphological trait is shown for each comparison. Piute Pass is the warm site and White Mountain is the cold site. We found that nares were longer while beak widths and depths were smaller in the warmer environment.

Figure 3. A) Photograph showing a body feather from a Sierra Nevada Gray-crowned Rosy-Finch captured at White Mountain. Highlighted is a microscopic imagesimplification of the pennaceous and plumulaceous barbule structure. B) Violin plots of node density raw measurements, with illustrations above of the structure of plumulaceous barbule nodes. Overlaid on the plots are the corrected mean trait measurements shown with a crossbar. Significance of the effect of site on the morphological trait is shown for each comparison.





Figure 4. Manhattan plot depicting the results of the LMM analysis on beak depth (A), wing chord (B), and pennaceous barbule length (C) with $-\log_{10}(\text{p-value})$ on the y-axis and the mapped Zebra Finch chromosome location on the x-axis. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted. Also plotted are the results of the $p_{\text{Per-site}}$ F_{ST} results (DG), using the 90th quantile threshold of 0.235.

Corresponding genes are also shown and functional categories are highlighted by color



category. See Table 1 for more functional details.

Supplemental Information for:

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch

Robertson, Erica C.N.¹, Brown, Timothy², Deitch, Sophie¹, Bossu, Christine M.¹, Zavaleta, Erika S.², Hooten, Mevin B.³, Ruegg, Kristen C.¹


Supplemental Methodology

Extracting temperature data from AdaptWest, 2022.

The two sampling locations, identified by GPS coordinates collected during fieldwork, were each buffered by 1 km to account for the spatial heterogeneity experienced by the birds. Monthly climate rasters from AdaptWest were filtered to include only June, July, and August, corresponding to the Gray-crowned Rosy-Finch breeding season. For each site, raster layers representing mean, minimum, and maximum monthly temperature were extracted within the 1 km buffer, and pixel values were averaged. The resulting mean, minimum, and maximum temperature values were then summarized for each site. The average values were also plotted across time (Sup

Fig 1).

Sup Fig 1. Average values within a 1km buffer around sample sites for Tave, Tmax, and Tmin of the breeding season plotted against month.



Sexing and Aging of Birds

Aging followed standard guidelines outlined in the *Pyle Identification Guide to North American Birds* (Pyle 1997). Age was determined primarily by examination of skull pneumatization and degree of cranial ossification, supplemented by plumage characteristics (extent of molt limits, feather wear, and coloration). When available, known-age individuals from previous banding or recapture records were also included. All assessments were conducted by trained banders using consistent criteria across field seasons. A full breakdown of individuals by age class and sex is provided in Sup Table 1.

Variation around morphological data collected

Not all individuals were measured for the full suite of morphological traits. In some cases, birds exhibited signs of acute handling stress (e.g., wincing of the eyes, open-beak panting, or rapid breathing), and were released before all measurements could be completed to minimize risk. In other instances, data collection was interrupted by rapidly changing field conditions, such as approaching thunderstorms or hail events, which required immediate cessation of banding activities for personnel safety. As a result, sample sizes differ slightly among traits, reflecting our decision to prioritize bird welfare and field safety over completeness of measurements.

Generating PC axis representing size

To further validate our choice of tarsus as a proxy for body size, we performed a principal component analysis on allometric traits to compare a PC axis representing overall body size with tarsus. We first removed all individuals that were missing information for any of the following traits: wing chord, tarsus, tail length, beak depth, beak width, and beak length. This left us with 136 individuals with complete data. We centered and scaled the variables and performed a principal component analysis on the resulting data. The first axis, PC1, explained 34.3% of the variance while PC2 explained 28.7%. We found the following loadings for PC1 and PC2:

	wing chord	tarsus	tail length	beak depth	beak width	beak length
PC1	0.5900	0.5358	0.5358	0.2094	0.1290	0.0172
PC2	-0.1654	-0.0829	-0.1260	0.6134	0.5789	0.4881

Because of the highly positive loadings on PC1 and the high percent of variance explained, we used this as a proxy for body size. We found that PC1 size was highly correlated with tarsus (0.77) so we proceeded with tarsus as our proxy for body size.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Morphological data and analysis

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Summary of morphological data

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Sup Table 1. Summary of sample sizes, distribution of sexes and ages for each site.

<u>Piute Pass n=72</u>		<u>White Mountains n=98</u>	
<u>Sex</u>	<u>Age</u>	<u>Sex</u>	<u>Age</u>
<u>F:25</u>	<u>SY:30</u>	<u>F:42</u>	<u>SY:55</u>
<u>M:35</u>	<u>ASY:30</u>	<u>M:35</u>	<u>ASY:22</u>
<u>UNK:12</u>	<u>HY:12</u>	<u>UNK:21</u>	<u>HY:21</u>

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Sup Table 2. Summary of trait values for Piute Pass.

<u>Trait</u>	<u>Mean</u>	<u>SD</u>	<u>N</u>	<u>Range</u>
<u>Wing Chord</u>	<u>100.47</u>	<u>3.808</u>	<u>72</u>	<u>92-110</u>
<u>Tarsus</u>	<u>20.001</u>	<u>.729</u>	<u>72</u>	<u>18.1-22.3</u>
<u>Nare Length</u>	<u>2.186</u>	<u>0.4362</u>	<u>57</u>	<u>1.45-3.2</u>
<u>Beak Depth</u>	<u>6.505</u>	<u>0.544</u>	<u>61</u>	<u>4.74-7.79</u>
<u>Beak Width</u>	<u>5.837</u>	<u>0.5310</u>	<u>61</u>	<u>4.5-7.07</u>
<u>Beak Length</u>	<u>11.504</u>	<u>0.5686</u>	<u>61</u>	<u>9.4-12.87</u>
<u>Pen Barbule Length</u>	<u>0.3795</u>	<u>0.0593</u>	<u>54</u>	<u>0.29-0.61</u>
<u>Pen Barbule Density</u>	<u>19.276</u>	<u>3.1449</u>	<u>54</u>	<u>13.25-24.67</u>
<u>Plum Barbule Length</u>	<u>0.8243</u>	<u>0.0902</u>	<u>54</u>	<u>0.64-1.05</u>
<u>Plum Barbule Density</u>	<u>28.353</u>	<u>2.6963</u>	<u>54</u>	<u>21-33.33</u>
<u>Plum Barbule Node Density</u>	<u>6.997</u>	<u>0.5748</u>	<u>54</u>	<u>5.33-8</u>

Sup Table 3. Summary of trait values for White Mountain.

Trait	Mean	SD	N	Range
Wing Chord	98.44	3.435	95	92-106
Tarsus	19.9	.572	95	18.4-21.7
Nare Length	1.87	0.327	93	1.14-2.9
Beak Depth	6.569	0.4757	95	5.56-7.97
Beak Width	6.0202	0.4743	95	5-7
Beak Length	11.2	0.6844	95	9.49-12.88
Pen Barbule Length	0.3631	0.05541	85	0.24-0.52
Pen Barbule Density	19.8313	2.6285	85	13.67-25.33
Plum Barbule Length	0.79822	0.1011	85	0.6-1.11
Plum Barbule Density	28.38039	3.0564	85	20.33-35
Plum Barbule Node Density	7.2745	0.6865	85	5.33-9.33

Morphological Statistical Analysis Model Details

Univariate Model Summary Details

We evaluated a suite of candidate linear models for each morphological trait to assess the effects of body size (tarsus length), age, sex, and site. All models were fit using ordinary least squares in R. The following fixed-effect combinations were tested:

Model ID	Fixed Effects Included
1	tarsus
2	age
3	sex
4	site_code
5	tarsus + age

6	tarsus + sex
7	tarsus + site_code
8	age + sex
9	age + site_code
10	sex + site_code
11	tarsus × age
12	tarsus × sex
13	tarsus × site_code
14	age × sex
15	age × site_code
16	sex × site_code
17	tarsus + age + sex
18	tarsus + age + site_code
19	tarsus + sex + site_code
20	age + sex + site_code
21	tarsus × age × sex
22	tarsus × age × site_code
23	tarsus × sex × site_code
24	age × sex × site_code

Sup Table 4. Top three models tested and resulting AIC information for each of those models for each trait.

Formula	K	AICc	Delta_AICc	AICcWt	Cum.Wt
Beak Depth					
tarsus + age + site_code	6	189.95893	0	0.22197985	0.22197985
tarsus + sex + site_code	6	190.210098	0.25116876	0.19578208	0.41776194
tarsus * sex * site_code	12	190.692707	0.73377728	0.15380691	0.57156885
Beak Width					

tarsus + sex + site_code	6	214.603266	0	0.28176801	0.28176801
		7		8	8
tarsus * sex * site_code	12	214.733299	0.13003272	0.26403132	0.54579934
		4	4	2	
tarsus + age + sex + site_code	7	216.545844	1.94257758	0.10667590	0.65247524
		3	9	4	5
Beak Length					
sex	4	202.858482	0	0.21026054	0.21026054
		2		1	1
age	4	204.054312	1.19583051	0.11563424	0.32589478
		7	2	8	9
tarsus + sex	5	204.275641	1.41715939	0.10352030	0.42941509
		6		1	
Nare Length					
tarsus + sex + site_code	6	124.965750	0	0.35089455	0.35089455
		2		6	6
tarsus + age + sex + site_code	7	125.280982	0.31523265	0.29972621	0.65062077
		9	7	5	1
tarsus + age + site_code	6	127.320330	2.35458055	0.10811500	0.75873577
		8	4	2	3
Wing Chord					
tarsus + age + sex + site_code	7	735.524521	0	0.83590162	0.83590162
		2			
tarsus + sex + site_code	6	738.858897	3.33437659	0.15779909	0.9937007
		8	5		
tarsus + age + sex	6	745.522703	9.99818237	0.00563738	0.99933809
		6	4		
Pennaceous Barbule Length					
sex	4	-	0	0.26538709	0.26538709
		445.925198		9	9
		6			
sex + site_code	5	-445.551068	0.37413064	0.22010928	0.48549638
			3	8	7
age * sex	6	-	1.48309822	0.12642388	0.61192026
		444.442100	2	1	8
		4			
Pennaceous Barbule Density					
tarsus + age	5	616.554421	0	0.24815054	0.24815054
		6		8	8

tarsus + sex	5	616.938818	0.38439685	0.20475971	0.45291026
		4	5	8	6
tarsus + age + site_code	6	617.915651	1.36122961	0.12564001	0.57855028
		2	8	6	3
Plumulaceous Barbule Length					
age + site_code	5	-	0	0.27657032	0.27657032
		267.584944		2	2
		9			
age	4	-	1.04890058	0.16369661	0.44026694
		266.536044	7	8	
		3			
age * site_code	7	-	1.07419116	0.16163966	0.6019066
		266.510753	8		
		7			
Plumulaceous Barbule Density					
tarsus + age	5	672.442234	0	0.30687339	0.30687339
		6			
tarsus + age + sex	6	673.176964	0.73472949	0.21252731	0.51940070
		1	8	1	1
tarsus + sex	5	673.642519	1.20028475	0.16839171	0.68779241
		4		1	2
Plumulaceous Barbule Node Density					
tarsus + age + site_code	6	229.271709	0	0.4507249	0.4507249
		3			
age + site_code	5	231.33502	2.06331062	0.16064578	0.61137068
			5	7	7
tarsus + age + sex + site_code	7	231.48286	2.21115065	0.14919912	0.76056981
			7	9	6

Sup Table 5. Summary of model ran and the model results for the top model for each trait. Asterisk (*) notes which traits are considered significantly different between sites.

Formula	r.squared	adj.r.squared	p.value
Beak Depth *			
tarsus + age + site_code	0.26910403	0.24961347	1.30E-09
Beak Width *			
tarsus + sex + site_code	0.15702313	0.13454374	3.49E-05
Beak Length			
sex	0.52253017	0.51628873	2.75E-25

Nare Length *			
tarsus + sex + site code	0.219043	0.19719804	3.50E-07
Wing Chord *			
tarsus + age + sex + site code	0.6826689	0.6728139	2.35E-38
Pennaceous Barbule Length			
sex	0.31689028	0.30684455	5.56E-12
Pennaceous Barbule Density			
tarsus + age	0.3791724	0.36506268	1.25E-13
Plumulaceous Barbule Length *			
age + site code	0.16133686	0.1426999	2.70E-05
Plumulaceous Barbule Density			
tarsus + age	0.09665926	0.07612878	0.00372514
Plumulaceous Barbule Node Density			
tarsus + age	0.09665926	0.07612878	0.00372514

Sup Table 6.A-J. Detailed summary for each of the trait models run. Sample size (n) is noted for each analysis. Significant site p-values are italicized. When site is present, 95% CI is also shown with the beta coefficient for site.

A. Beak Depth, n=155

term	estimate	std.error	statistic	p.value
(Intercept)	3.37921794	1.20275672	2.80956065	0.00562258
tarsus	0.15757233	0.06065696	2.59776171	0.0103184
age3	-0.0419748	0.08464803	-0.4958742	0.62070894
age4	-0.7813467	0.11173235	-6.9930214	8.29E-11
site codeWMTN	0.23301356	0.0764162	3.04926899	0.00271257
$\beta = 0.46$, 95% CI = 0.16–0.76				

B. Beak Width, n=155

term	estimate	std.error	statistic	p.value
(Intercept)	3.13067772	1.26675195	2.47142127	0.01457617
tarsus	0.13913241	0.06376169	2.18206908	0.03066072
sexM	-0.1290975	0.0855716	-1.5086492	0.13349188
sexUNK	-0.5671672	0.12422359	-4.5656968	1.03E-05
site codeWMTN	0.28128538	0.0818869	3.43504748	0.0007664
$\beta = 0.55$, 95% CI = 0.24–0.87				

C. Beak Length, n=156

term	estimate	std.error	statistic	p.value
(Intercept)	11.5557576	0.05611459	205.931438	6.60E-189
sexM	-0.0851779	0.07849068	-1.0851971	0.27954128
sexUNK	-1.4224242	0.11421569	-12.453843	5.00E-25

D. Nare Length, n=148

term	estimate	std.error	statistic	p.value
(Intercept)	0.00316755	0.92079149	0.00344003	0.99726005
tarsus	0.11403506	0.04632887	2.46142557	0.01502832
sexM	-0.1216435	0.06971025	-1.7449874	0.08313529
sexUNK	-0.2391738	0.07966172	-3.0023678	0.0031627
site_codeWMTN	-0.3084156	0.0611737	-5.0416373	1.38E-06
$\beta = -0.76$, 95% CI = -1.07-(-0.46)				

E. Wing Chord, n=167

term	estimate	std.error	statistic	p.value
(Intercept)	67.5307604	5.49907378	12.2803881	6.99E-25
tarsus	1.47810264	0.27931635	5.29185855	3.91E-07
age3	0.98013348	0.42161033	2.32473783	0.02133471
age4	2.61989277	0.47482379	5.51761055	1.35E-07
sexM	5.18357951	0.39536383	13.1109099	3.49E-27
site_codeWMTN	-1.1884579	0.34055186	-3.4898001	0.0006237
$\beta = -0.32$, 95% CI = -0.50-(-0.14)				

F. Pennaceous Barbule Length, n=139

term	estimate	std.error	statistic	p.value
(Intercept)	0.34640226	0.00621558	55.7312693	1.55E-95
sexM	0.01699774	0.00899133	1.89045812	0.06082431
sexUNK	0.0882362	0.0112384	7.85131195	1.10E-12

G. Pennaceous Barbule Density, n=136

<u>term</u>	<u>estimate</u>	<u>std.error</u>	<u>statistic</u>	<u>p.value</u>
(Intercept)	23.2506487	6.11112099	3.80464545	0.00021631
tarsus	-0.1510297	0.30918622	-0.4884749	0.626024
age3	0.47029958	0.47571193	0.98862262	0.32465654
age4	-4.24769	0.53314041	-7.9673009	6.67E-13

H. Plumulaceous Barbule Length

<u>term</u>	<u>estimate</u>	<u>std.error</u>	<u>statistic</u>	<u>p.value</u>
(Intercept)	0.8237826	0.01620735	50.8277177	6.72E-90
age3	-0.0349157	0.01789566	-1.9510714	0.0531196
age4	0.07217363	0.02106868	3.42563687	0.00081297
site_codeWMTN	-0.0288163	0.01624784	-1.773549	0.07839249
$\beta = -0.30$, 95% CI = -0.62-0.34				

I. Plumulaceous Barbule Density


<u>term</u>	<u>estimate</u>	<u>std.error</u>	<u>statistic</u>	<u>p.value</u>
(Intercept)	25.5415447	7.50508001	3.40323417	0.00088205
tarsus	0.1335034	0.37971223	0.351591	0.72570563
age3	1.16460095	0.58422278	1.99341926	0.048277
age4	-1.3306565	0.65475081	-2.0323098	0.04412831

J. Plumulaceous Barbule Node Density

<u>term</u>	<u>estimate</u>	<u>std.error</u>	<u>statistic</u>	<u>p.value</u>
(Intercept)	9.17335111	1.46944391	6.24273647	5.54E-09
tarsus	-0.1062565	0.07416009	-1.432799	0.15429738
age3	0.27870664	0.11659663	2.39034915	0.01825573
age4	-0.7579252	0.12906466	-5.8724455	3.34E-08
site_codeWMTN	0.26874166	0.0987462	2.72153933	0.00738179
$\beta = 0.41$, 95% CI = 0.11-0.70				


Population Structure Results

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Sup Fig 24. PCA results showing a lack of population structure between the White Mountain (WMNT, blue) and Piute Pass (PIPA, red) populations. There are some outlier individuals but an ADMIXTURE analysis confirmed that 1 population is the most supported.


Extended GWAS Results



Sup Fig 32. Beak length GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.



Sup Fig 43. Beak width GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.



Sup Fig 54. Pennaceous barbule density GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.

GWAS model evaluation


Sup Table 7. λ_{GC} value for each ULMM run, based on p-values.

Trait	λ_{GC}
Wing Chord	1.0287
Nare Length	1.0316
Beak Depth	1.0129
Beak Width	1.0157
Beak Length	0.9684
Pen Barbule Length	1.0273
Pen Barbule Density	1.0232
Plum Barbule Length	0.9773
Plum Barbule Density	1.0009
Plum Barbule Node Density	1.0218

Sup Fig 6. PP plots for all traits.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only








Figure 1. A) A map of California highlighting the sampling location and design for two populations of Grey-crowned Rosy-Finch, Piute Pass and White Mountain. Elevation is noted for each site along with an elevation transect. B) Overview of methods for GWAS analysis. Morphological measurements were corrected for the effects of sex, age, and body size before being input into BSLMM and LMM models. C) Overview of the methods for the F_{ST} analysis. The output of both analyses were analyzed to identify significant SNPs and related genes.

774x989mm (118 x 118 DPI)


Body Morphology Results




A



B




C




A




A1






A2




A3




B Plumulaceous Barbule Node Density




A




B




C



D



Gene Function Categories



Gene	Model	Function	Details	Reference
<i>Wing Chord</i>				
PCP4	BSLMM	Development, Stress Response	Bone development, stress response in quail	Xiao et al 2008; Steven 2021
GRIK2	BSLMM	Reproduction	Egg quality and production in goose and duck	Gao et al 2023; Bhavana et al 2022
AKAP7	BSLMM	Stress Response	Stress response in quail	Stevens 2021
PACRG	BSLMM	Heat Stress	Heat stress response in chickens	Tian et al 2020
HAPLN1	BSLMM	Coloration	Plumage coloration in ducks	Zhang et al 2024
NEGR1	BSLMM	Development, Col- oration	Neural development, plumage coloration in crested ibis	Sun et al 2020
† PGLYRP2	BSLMM/LMM	Immunity, Heat Stress	Immune response in vertebrates, heat stress response in chickens	Hafiz et al 2020; Kim et al 2021
GUCA1C	LMM	Morphology	Visual systems in birds, bone quality in ducks, wing deformity in chickens	Wu et al 2021; Gesemann et al 2023; Li et al 2020; Mohammadi et al 2024
RFX3	LMM	Hypoxia Adaptation	Altitude adaptation in wolves	Zhang et al 2014; Shweizer et al 2014
NIPBL	LMM	Development, DNA Repair	Developmental regulation and DNA repair	Barnes 2019

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

(Continued from previous page)					
Gene	Model	Function	Details	Reference	
CHD1	LMM	Gene Regulation	Modulates gene expression in birds	Fridolfsson et al 2000	
NDUFA11	LMM	Metabolism, Respiration	Respiratory system support in buffalo, mitochondrial activity in songbirds	Clark et al 2024; Sadeesh et al 2025	
Beak Depth	TENM4	Development, Coloration, Reproduction	Neural development in quail, dorsal feather coloration in duck, breeding timing in great tit	Twumasi et al 2024; Steven 2021; Gienapp et al 2017	
	RASL11A	Morphology	Craniofacial development in buteo	Abernathy 2021	
	TINAG	Morphology	Beak development in chicken	Bai et al 2018	
	LAMA4	Migration	Migration timing in Swainson's thrush	Johnston et al 2016	
	LURAP1L	Coloration	Pigmentation in mallards	Ma et al 2021	
	PTPRD	Communication, Coloration	Vocal signaling in flycatchers, pigmentation in mallards	García et al 2023; Ma et al 2021	
	CLRN3	Reproduction, Sex	Sex differentiation in chicken	Lou et al 2024	
	CPEB3	Reproduction	Oocyte meiosis regulator in chicken	Chen et al 2022	

(Continued from previous page)

Gene	Model	Function	Details	Reference
BICC1	BSLMM	Morphology, Body Size	Bone development in chicken and mice, body size in chicken and hermit thrush	Kinsella et al 2019; Johnson et al 2015; Mesner et al 2014
ANK3	BSLMM	Migration	Migration in birds	Lenon 2022
PRKG1	BSLMM	Reproduction, Behavior, Coloration	Ovarian function in duck, foraging behavior in humans, signal transduction and feather pigmentation in ducks	Sun et al 2024; Struk et al 2019; Twumasi et al 2024
OTUD7A	BSLMM	Development	Neural development and adaptation to urban environments in bananaquit	Mascarenhas et al 2023
ERCC8	BSLMM	Hypoxia Adaptation	Altitude adaptation in lizard	Yan et al 2022
MEGF11	BSLMM	Coloration	Dorsal feather coloration in duck	Twumasi et al 2024
DLG1	BSLMM	Development, Regulation	Embryo development in mice, regulatory development in birds	Seki et al 2017
<i>Nare Length</i>				
NRIP1	BSLMM	Reproduction, Adaptation, Metabolism	Female reproduction in goose, nocturnal adaptation in owls, mitochondrial biosynthesis in vertebrates	Zheng et al 2023; SungCho et al 2019; Da Costa Moreno et al 2022

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

4

(Continued from previous page)				
Gene	Model	Function	Details	Reference
ATP1A1	BSLMM	Migration	Migration in birds	Lenon 2022
TENM4	BSLMM	Development, Col- oration, Reproduction	Neural development in quail, dorsal feather coloration in duck, breeding timing in great tit	Twumasi et al 2024; Steven 2021; Gienapp et al 2017
KALRN	BSLMM	Neurodevelopment	Cognitive ability in chickadee	Richetti et al 2024
CNTNAP5	BSLMM	Development, Mem- ory, Hypoxia Adapta- tion	Brain development and signaling in zebra finch, spatial memory in chickadees, hypoxia response in chickens	Gilbert et al 2021; Semenov et al 2024; Li et al 2009
KCTD18	BSLMM	Immunity, Communi- cation	Immune response in vultures, vocal learning in chickens	Chung et al 2015; Lovell et al 2013
PLPP3	BSLMM	Morphology, Metabolism	Culmen length in buteo	Abernathy et al 2021
DLG1	BSLMM	Development, Regula- tion	Embryo development in mice, regulatory de- velopment in birds	Seki et al 2017
ADAMTS18	BSLMM	Morphology, Bone Density	Bone mineral density in humans and chickens	Xiong et al 2009; Guo et al 2017
PSD2	BSLMM	Coloration, Stress Re- sponse	Melanin production in owls, stress response in quail	Ducrest et al 2025; Steven 2021

(Continued from previous page)

Gene	Model	Function	Details	Reference
BRAT1	BSLMM	Stress Response	Stress response in chickens	Rodriguez et al 2022
<i>Pennaceous Barbule Length</i>				
KCNF1	BSLMM	Coastal Adaptation	Coastal adaptation in song sparrow and frogs	Clark et al 2024
CSMD1	BSLMM/LMM	Heat Stress, Hypoxia Adaptation	Heat adaptation in chickens, adaptation to altitude in humans	Hao et al 2024; De Loma et al 2025
NKAIN1	BSLMM	Immunity, Migration	Immune response in house finch, migration tendency in European blackcap	Veetil et al 2024; Delmore et al 2020
KCNJ3	BSLMM	Reproduction	Egg production in duck	Sun et al 2024
AGBL1	BSLMM	Metabolism	Fatty acid composition in chicken	Cho et al 2023
TENM1	BSLMM	Morphology	Craniofacial development in pigeons	Boer et al 2021
<i>Plumulaceous Node Density</i>				
EFNA2	BSLMM	Heat Stress, Morphology, Coloration	Heat stress response in chickens, bone morphogenesis and remodeling in ground tit, color differences in bulbul	Zhuang et al 2019; Cheng et al 2020; Shakya et al 2020
STK10	BSLMM	Morphology	Cranium and bill depth in buteo	Abernathy 2021
<i>Plumulaceous Barbule Density</i>				
BORCS6	BSLMM	Metabolism	Meat quality in chickens	Alsoufi et al 2023
HSF2	BSLMM	Heat Stress	Response to heat stress in chickens	Xie et al 2014

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

9

(Continued from previous page)				
Gene	Model	Function	Details	Reference
PRKN	BSLMM	Domestication	Domestication in birds	Farias-Virgens et al 2025
NRXN1	BSLMM	Neurodevelopment, Communication, Coloration	Neural connections, plumage coloration in crested ibis, vocal rhythm in birds, speech disorders in humans	Sun et al 2020; Sebastianelli et al 2024
ALDH1A1	BSLMM	Hypoxia Adaptation, Metabolism, Vision	Hypoxia adaptation and fat metabolism in rosy finch, eyesight in owl and mice	Bao et al 2016; Funk et al 2023; Borges et al 2019
ADAMTS19	BSLMM	Morphology	Rumpless trait in chickens	Chen et al 2024
HERC4	BSLMM	Coloration, Migration	Plumage color in duck, migration in European blackbirds	Zhang et al 2023; Franchini et al 2017
SMC4	BSLMM	Development	Feather follicle development in goose	Hu et al 2020
SPAG16	BSLMM	Reproduction, Coloration, Longevity	Sexual development in birds-of-paradise, plumage coloration in ducks, longevity in mammals	Prost et a 2018; Zhang et al 2024; Matsuda et al 2024

F_{ST} outliers

Gene	Function	Reference
GDPD5	Lipid Metabolism	Feng et al 2023
NOX4	Heat Stress, Altitude Adaptation	Goel et al 2021; Diebold et al 2010
MAML2	Altitude Adaptation	Schweizer et al 2021
SUPT20H	Altitude Adaptation	Schweizer et al 2021
PPP1R1C	Cold Adaptation	Federova et al 2022
SOX14	Feather Development	Chen et al 2024
MEGF11	Feather Development	Twumasi et al 2024
MSRA	Altitude Adaptation	Jeong et al 2018; Gheyas et al 2021
AKT3	Altitude Adaptation	Buroker et al 2012; Qi et al 2018
DOCK11	Feather Development	Crates et al 2024
SMARCA1	Craniofacial Development	Boer et al 2021
GAB3	Craniofacial Development	Boer et al 2021
ADGRD1	Altitude Adaptation	Nan et al 2021
NPHP4	Vision	Borges et al 2019
ALDH7A1	Beak Development	Bai et al 2014
ADAMTS19	Metabolism	Zhang et al 2022
VLDLR	Lipid Protein	Wang et al 2011
ERCC8	Altitude Adaptation	Yan et al 2022
CTIF	Beak Development	Lawson et al 2017

Supplemental Information for:

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch

Robertson, Erica C.N.¹, Brown, Timothy², Deitch, Sophie¹, Bossu, Christine M.¹, Zavaleta, Erika S.², Hooten, Mevin B.³, Ruegg, Kristen C.¹


Supplemental Methodology

Extracting temperature data from AdaptWest, 2022.

The two sampling locations, identified by GPS coordinates collected during fieldwork, were each buffered by 1 km to account for the spatial heterogeneity experienced by the birds. Monthly climate rasters from AdaptWest were filtered to include only June, July, and August, corresponding to the Gray-crowned Rosy-Finch breeding season. For each site, raster layers representing mean, minimum, and maximum monthly temperature were extracted within the 1 km buffer, and pixel values were averaged. The resulting mean, minimum, and maximum temperature values were then summarized for each site. The average values were also plotted across time (Sup

Fig 1).

Sup Fig 1. Average values within a 1km buffer around sample sites for Tave, Tmax, and Tmin of the breeding season plotted against month.



Sexing and Aging of Birds

Aging followed standard guidelines outlined in the *Pyle Identification Guide to North American Birds* (Pyle 1997). Age was determined primarily by examination of skull pneumatization and degree of cranial ossification, supplemented by plumage characteristics (extent of molt limits, feather wear, and coloration). When available, known-age individuals from previous banding or recapture records were also included. All assessments were conducted by trained banders using consistent criteria across field seasons. A full breakdown of individuals by age class and sex is provided in Sup Table 1.

Variation around morphological data collected

Not all individuals were measured for the full suite of morphological traits. In some cases, birds exhibited signs of acute handling stress (e.g., wincing of the eyes, open-beak panting, or rapid breathing), and were released before all measurements could be completed to minimize risk. In other instances, data collection was interrupted by rapidly changing field conditions, such as approaching thunderstorms or hail events, which required immediate cessation of banding activities for personnel safety. As a result, sample sizes differ slightly among traits, reflecting our decision to prioritize bird welfare and field safety over completeness of measurements.

Generating PC axis representing size

To further validate our choice of tarsus as a proxy for body size, we performed a principal component analysis on allometric traits to compare a PC axis representing overall body size with tarsus. We first removed all individuals that were missing information for any of the following traits: wing chord, tarsus, tail length, beak depth, beak width, and beak length. This left us with 136 individuals with complete data. We centered and scaled the variables and performed a principal component analysis on the resulting data. The first axis, PC1, explained 34.3% of the variance while PC2 explained 28.7%. We found the following loadings for PC1 and PC2:

	wing chord	tarsus	tail length	beak depth	beak width	beak length
PC1	0.5900	0.5358	0.5358	0.2094	0.1290	0.0172
PC2	-0.1654	-0.0829	-0.1260	0.6134	0.5789	0.4881

Because of the highly positive loadings on PC1 and the high percent of variance explained, we used this as a proxy for body size. We found that PC1_size was highly correlated with tarsus (0.77) so we proceeded with tarsus as our proxy for body size.

Morphological data and analysis

Summary of morphological data

Sup Table 1. Summary of sample sizes, distribution of sexes and ages for each site.

Piute Pass n=72		White Mountains n=98	
Sex	Age	Sex	Age
F:25	SY:30	F:42	SY:55
M:35	ASY:30	M:35	ASY:22
UNK:12	HY:12	UNK:21	HY:21

Sup Table 2. Summary of trait values for Piute Pass.

Trait	Mean	SD	N	Range
Wing Chord	100.47	3.808	72	92-110
Tarsus	20.001	.729	72	18.1-22.3
Nare Length	2.186	0.4362	57	1.45-3.2
Beak Depth	6.505	0.544	61	4.74-7.79
Beak Width	5.837	0.5310	61	4.5-7.07
Beak Length	11.504	0.5686	61	9.4-12.87
Pen Barbule Length	0.3795	0.0593	54	0.29-0.61
Pen Barbule Density	19.276	3.1449	54	13.25-24.67
Plum Barbule Length	0.8243	0.0902	54	0.64-1.05
Plum Barbule Density	28.353	2.6963	54	21-33.33
Plum Barbule Node Density	6.997	0.5748	54	5.33-8

Sup Table 3. Summary of trait values for White Mountain.

Trait	Mean	SD	N	Range
Wing Chord	98.44	3.435	95	92-106
Tarsus	19.9	.572	95	18.4-21.7
Nare Length	1.87	0.327	93	1.14-2.9
Beak Depth	6.569	0.4757	95	5.56-7.97
Beak Width	6.0202	0.4743	95	5-7
Beak Length	11.2	0.6844	95	9.49-12.88
Pen Barbule Length	0.3631	0.05541	85	0.24-0.52
Pen Barbule Density	19.8313	2.6285	85	13.67-25.33
Plum Barbule Length	0.79822	0.1011	85	0.6-1.11

Plum Barbule Density	28.38039	3.0564	85	20.33-35
Plum Barbule Node Density	7.2745	0.6865	85	5.33-9.33

Morphological Statistical Analysis Model Details

Univariate Model Summary Details

We evaluated a suite of candidate linear models for each morphological trait to assess the effects of body size (tarsus length), age, sex, and site. All models were fit using ordinary least squares in R. The following fixed-effect combinations were tested:

Model ID	Fixed Effects Included
1	tarsus
2	age
3	sex
4	site_code
5	tarsus + age
6	tarsus + sex
7	tarsus + site_code
8	age + sex
9	age + site_code
10	sex + site_code
11	tarsus × age
12	tarsus × sex
13	tarsus × site_code
14	age × sex
15	age × site_code
16	sex × site_code
17	tarsus + age + sex
18	tarsus + age + site_code
19	tarsus + sex + site_code
20	age + sex + site_code
21	tarsus × age × sex
22	tarsus × age × site_code
23	tarsus × sex × site_code
24	age × sex × site_code

Sup Table 4. Top three models tested and resulting AIC information for each of those models for each trait.

Formula	K	AICc	Delta_AICc	AICcWt	Cum.Wt
Beak Depth					
tarsus + age + site_code	6	189.95893	0	0.2219798 55	0.2219798 55
tarsus + sex + site_code	6	190.210098 8	0.2511687 66	0.1957820 89	0.4177619 44
tarsus * sex * site_code	12	190.692707 3	0.7337772 8	0.1538069 11	0.5715688 55
Beak Width					
tarsus + sex + site_code	6	214.603266 7	0	0.2817680 18	0.2817680 18
tarsus * sex * site_code	12	214.733299 4	0.1300327 24	0.2640313 22	0.5457993 4
tarsus + age + sex + site_code	7	216.545844 3	1.9425775 89	0.1066759 04	0.6524752 45
Beak Length					
sex	4	202.858482 2	0	0.2102605 41	0.2102605 41
age	4	204.054312 7	1.1958305 12	0.1156342 48	0.3258947 89
tarsus + sex	5	204.275641 6	1.4171593 9	0.1035203 01	0.4294150 9
Nare Length					
tarsus + sex + site_code	6	124.965750 2	0	0.3508945 56	0.3508945 56
tarsus + age + sex + site_code	7	125.280982 9	0.3152326 57	0.2997262 15	0.6506207 71
tarsus + age + site_code	6	127.320330 8	2.3545805 54	0.1081150 02	0.7587357 73
Wing Chord					
tarsus + age + sex + site_code	7	735.524521 2	0	0.8359016 2	0.8359016 2
tarsus + sex + site_code	6	738.858897 8	3.3343765 95	0.1577990 9	0.9937007
tarsus + age + sex	6	745.522703 6	9.9981823 74	0.0056373 8	0.9993380 9
Pennaceous Barbule Length					

sex	4	- 445.925198 6	0	0.2653870 99	0.2653870 99
sex + site_code	5	- 445.551068	0.3741306 43	0.2201092 88	0.4854963 87
age * sex	6	- 444.442100 4	1.4830982 22	0.1264238 81	0.6119202 68
Pennaceous Barbule Density					
tarsus + age	5	616.554421 6	0	0.2481505 48	0.2481505 48
tarsus + sex	5	616.938818 4	0.3843968 55	0.2047597 18	0.4529102 66
tarsus + age + site_code	6	617.915651 2	1.3612296 18	0.1256400 16	0.5785502 83
Plumulaceous Barbule Length					
age + site_code	5	- 267.584944 9	0	0.2765703 22	0.2765703 22
age	4	- 266.536044 3	1.0489005 87	0.1636966 18	0.4402669 4
age * site_code	7	- 266.510753 7	1.0741911 68	0.1616396 6	0.6019066
Plumulaceous Barbule Density					
tarsus + age	5	672.442234 6	0	0.3068733 9	0.3068733 9
tarsus + age + sex	6	673.176964 1	0.7347294 98	0.2125273 11	0.5194007 01
tarsus + sex	5	673.642519 4	1.2002847 5	0.1683917 11	0.6877924 12
Plumulaceous Barbule Node Density					
tarsus + age + site_code	6	229.271709 3	0	0.4507249	0.4507249
age + site_code	5	231.33502	2.0633106 25	0.1606457 87	0.6113706 87
tarsus + age + sex + site_code	7	231.48286	2.2111506 57	0.1491991 29	0.7605698 16

Sup Table 5. Summary of model ran and the model results for the top model for each trait. Asterisk (*) notes which traits are considered significantly different between sites.

Formula	r.squared	adj.r.squared	p.value
Beak Depth *			
tarsus + age + site_code	0.26910403	0.24961347	1.30E-09
Beak Width *			
tarsus + sex + site_code	0.15702313	0.13454374	3.49E-05
Beak Length			
sex	0.52253017	0.51628873	2.75E-25
Nare Length *			
tarsus + sex + site_code	0.219043	0.19719804	3.50E-07
Wing Chord *			
tarsus + age + sex + site_code	0.6826689	0.6728139	2.35E-38
Pennaceous Barbule Length			
sex	0.31689028	0.30684455	5.56E-12
Pennaceous Barbule Density			
tarsus + age	0.3791724	0.36506268	1.25E-13
Plumulaceous Barbule Length *			
age + site_code	0.16133686	0.1426999	2.70E-05
Plumulaceous Barbule Density			
tarsus + age	0.09665926	0.07612878	0.00372514
Plumulaceous Barbule Node Density			
tarsus + age	0.09665926	0.07612878	0.00372514

Sup Table 6.A-J. Detailed summary for each of the trait models run. Sample size (n) is noted for each analysis. Significant site p-values are italicized. When site is present, 95% CI is also shown with the beta coefficient for site.

A. Beak Depth, n=155

term	estimate	std.error	statistic	p.value
(Intercept)	3.37921794	1.20275672	2.80956065	0.00562258
tarsus	0.15757233	0.06065696	2.59776171	0.0103184
age3	-0.0419748	0.08464803	-0.4958742	0.62070894
age4	-0.7813467	0.11173235	-6.9930214	8.29E-11
site_codeWMTN	0.23301356	0.0764162	3.04926899	0.00271257
$\beta = 0.46$, 95% CI = 0.16–0.76				

B. Beak Width, n=155

term	estimate	std.error	statistic	p.value
(Intercept)	3.13067772	1.26675195	2.47142127	0.01457617
tarsus	0.13913241	0.06376169	2.18206908	0.03066072
sexM	-0.1290975	0.0855716	-1.5086492	0.13349188
sexUNK	-0.5671672	0.12422359	-4.5656968	1.03E-05
site_codeWMTN	0.28128538	0.0818869	3.43504748	0.0007664
$\beta = 0.55$, 95% CI = 0.24–0.87				

C. Beak Length, n=156

term	estimate	std.error	statistic	p.value
(Intercept)	11.5557576	0.05611459	205.931438	6.60E-189
sexM	-0.0851779	0.07849068	-1.0851971	0.27954128
sexUNK	-1.4224242	0.11421569	-12.453843	5.00E-25

D. Nare Length, n=148

term	estimate	std.error	statistic	p.value
(Intercept)	0.00316755	0.92079149	0.00344003	0.99726005
tarsus	0.11403506	0.04632887	2.46142557	0.01502832
sexM	-0.1216435	0.06971025	-1.7449874	0.08313529
sexUNK	-0.2391738	0.07966172	-3.0023678	0.0031627
site_codeWMTN	-0.3084156	0.0611737	-5.0416373	1.38E-06
$\beta = -0.76$, 95% CI = -1.07–(-0.46)				

E. Wing Chord, n=167

term	estimate	std.error	statistic	p.value
(Intercept)	67.5307604	5.49907378	12.2803881	6.99E-25
tarsus	1.47810264	0.27931635	5.29185855	3.91E-07
age3	0.98013348	0.42161033	2.32473783	0.02133471
age4	2.61989277	0.47482379	5.51761055	1.35E-07
sexM	5.18357951	0.39536383	13.1109099	3.49E-27
site_codeWMTN	-1.1884579	0.34055186	-3.4898001	0.0006237
$\beta = -0.32$, 95% CI = -0.50–(-0.14)				

F. Pennaceous Barbule Length, n=139

term	estimate	std.error	statistic	p.value
(Intercept)	0.34640226	0.00621558	55.7312693	1.55E-95
sexM	0.01699774	0.00899133	1.89045812	0.06082431
sexUNK	0.0882362	0.0112384	7.85131195	1.10E-12

G. Pennaceous Barbule Density, n=136

term	estimate	std.error	statistic	p.value
(Intercept)	23.2506487	6.11112099	3.80464545	0.00021631
tarsus	-0.1510297	0.30918622	-0.4884749	0.626024
age3	0.47029958	0.47571193	0.98862262	0.32465654
age4	-4.24769	0.53314041	-7.9673009	6.67E-13

H. Plumulaceous Barbule Length

term	estimate	std.error	statistic	p.value
(Intercept)	0.8237826	0.01620735	50.8277177	6.72E-90
age3	-0.0349157	0.01789566	-1.9510714	0.0531196
age4	0.07217363	0.02106868	3.42563687	0.00081297
site_codeWMTN	-0.0288163	0.01624784	-1.773549	0.07839249
$\beta = -0.30$, 95% CI = -0.62-0.34				

I. Plumulaceous Barbule Density


term	estimate	std.error	statistic	p.value
(Intercept)	25.5415447	7.50508001	3.40323417	0.00088205
tarsus	0.1335034	0.37971223	0.351591	0.72570563
age3	1.16460095	0.58422278	1.99341926	0.048277
age4	-1.3306565	0.65475081	-2.0323098	0.04412831

J. Plumulaceous Barbule Node Density

term	estimate	std.error	statistic	p.value
(Intercept)	9.17335111	1.46944391	6.24273647	5.54E-09
tarsus	-0.1062565	0.07416009	-1.432799	0.15429738
age3	0.27870664	0.11659663	2.39034915	0.01825573
age4	-0.7579252	0.12906466	-5.8724455	3.34E-08


site_codeWMNTN	0.26874166	0.0987462	2.72153933	0.00738179
	$\beta = 0.41$, 95% CI = 0.11-0.70			

Population Structure Results




Sup Fig 2. PCA results showing a lack of population structure between the White Mountain (WMNT, blue) and Piute Pass (PIPA, red) populations. There are some outlier individuals but an ADMIXTURE analysis confirmed that 1 population is the most supported.


Extended GWAS Results



Sup Fig 3. Beak length GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.



Sup Fig 4. Beak width GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.




Sup Fig 5. Pennaceous barbule density GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.

GWAS model evaluation

Sup Table 7. λ GC value for each ULMM run, based on p-values.

Trait	λ GC
Wing Chord	1.0287
Nare Length	1.0316
Beak Depth	1.0129
Beak Width	1.0157
Beak Length	0.9684
Pen Barbule Length	1.0273
Pen Barbule Density	1.0232
Plum Barbule Length	0.9773
Plum Barbule Density	1.0009
Plum Barbule Node Density	1.0218

Sup Fig 6. PP plots for all traits: A) plumulaceous barbule density, B) plumulaceous barbule node density, C) pennaceous barbule density, D) pennaceous barbule length, E) plumulaceous barbule length, F) beak length, G) beak width, H) beak depth, I) nare length, J) wing chord.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Molecular Ecology

Original Article

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch

Submission ID cc4c246c-6b5d-4ee8-9da6-438449184f85

Submission Version Initial Submission

PDF Generation 25 Jul 2025 14:55:02 EST by Atypon ReX

Authors

Ms. Robertson, Erica C.N.
Corresponding Author
Submitting Author

 [ORCID](https://orcid.org/0000-0002-0869-6300)
<https://orcid.org/0000-0002-0869-6300>

Affiliations
• Department of Biology, Colorado State University, Fort Collins, CO, USA

Mr. Brown, Timothy

Affiliations
• Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA, USA

Ms. Deitch, Sophie

Affiliations
• Department of Biology, Colorado State University, Fort Collins, CO, USA

Dr. Bossu, Christen M.1

Affiliations
• Department of Biology, Colorado State University, Fort Collins, CO, USA

Dr. Zavaleta, Erika S.

Affiliations
• Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA, USA

Dr. Hooten, Mevin B.	
	Affiliations
	• Department of Statistics and Data Sciences, The University of Texas at Austin, Austin, TX, USA
Dr. Ruegg, Kristen C.	
	Affiliations
	• Department of Biology, Colorado State University, Fort Collins, CO, USA
Additional Information	

Files for peer review

All files submitted by the author for peer review are listed below. Files that could not be converted to PDF are indicated; reviewers are able to access them online.

Name	Type of File	Size	Page
GCRF_manuscript_full_draft_figures.docx	Main Document - MS Word	4.5 MB	Page 4
CH1_CF_V2_final.pdf	Figure	1.5 MB	Page 56
GWAS_Fig2_final.pdf	Figure	385.0 KB	Page 57
GWAS_fig3b_final.pdf	Figure	8.6 MB	Page 58
Manhattan_plots_Clean_final.pdf	Figure	1.1 MB	Page 59
FST_Gene_table.pdf	Table	52.1 KB	Page 60
GWAS_Gene_Table.pdf	Table	57.4 KB	Page 61
fst_genes_supplement.csv	Supplementary Material for Review	7.2 KB	<i>Not converted to PDF</i>
GCRF_manuscript_supplement.docx	Supplementary Material for Review	3.2 MB	Page 67
GWAS_genes_supplement.csv	Supplementary Material for Review	5.7 KB	<i>Not converted to PDF</i>

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch

Running Title: Local adaptation in an alpine finch

Authors: Robertson, Erica C.N.¹, Brown, Timothy², Deitch, Sophie¹, Bossu, Christen M.¹, Zavaleta, Erika S.², Hooten, Mevin B.³, Ruegg, Kristen C.¹

Affiliations:

- 1. Department of Biology, Colorado State University, Fort Collins, CO, USA
- 2. Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA, USA
- 3. Department of Statistics and Data Sciences, The University of Texas at Austin, Austin, TX, USA

Corresponding Author: Erica C.N. Robertson

Abstract

Understanding patterns and mechanisms underlying local adaptation is becoming increasingly important for species conservation amid anthropogenically driven environmental change. Alpine systems are experiencing particularly intense pressure from environmental change resulting from increased rates of warming and corresponding loss of snow and ice. We integrate morphological and genetic analyses to identify traits important for local adaptation in one of the highest elevation breeding birds in North America, the Sierra Nevada Gray-crowned Rosy Finch. We performed an in-depth analysis of how traits with known links to thermoregulation in birds such as wing length, bill size, and feather microstructure vary between two populations at sites with contrasting climate and environmental conditions. We identified loci underlying these traits using a genome-wide association study and further examined regions of the

genome related to altitude adaptation and cold tolerance using F_{ST} outlier tests. Together, these results indicate that temperature, food availability, and alpine landscape features may impose multifaceted and potentially conflicting selective pressures on morphological traits important to adaptation in alpine birds. Overall, this work represents one of the first in-depth analyses of the genetic basis of adaptation in an alpine specialist songbird.

Keywords

local adaptation, conservation genomics, genome-wide association study, F_{ST} outlier test, alpine

Introduction

Local adaptation occurs when natural selection favors traits that confer fitness advantages in specific habitats (Endler, 1977; Kawecki & Ebert, 2004; Savolainen et al., 2013). A species' adaptive capacity determines the extent to which it will be able to respond to rapid environmental change (Forester et al., 2022, 2023; Lande & Shannon, 1996; Meek et al., 2023). As a result, identifying genetic and morphological traits involved in local adaptation is a key component of effective species conservation planning (Hoban et al., 2016; Meek et al., 2023). Historically, assessments of the capacity for local adaptation relied on reciprocal transplant experiments that allowed researchers to link variation in putatively ecologically important traits to differences in fitness across environmental gradients (Blanquart et al., 2013; Clausen et al., 1941; Kawecki & Ebert, 2004; Savolainen et al., 2013). However, recent advances in genomic methods have made it possible to detect signatures of selection across the genome and to associate these with environmental variation and ecologically important morphological variation (Capblancq et al., 2020; Hoban et al., 2016; Lotterhos & Whitlock, 2015). We leveraged recent advances in genomic methods to identify

morphological and genetic traits important to local adaptation in an understudied alpine bird system.

Many montane species undergo upward range shifts with climate change (Mamantov et al., 2021). However, for those species already living near the top of a mountain system, there are constraints that may limit the ability to track their niche, a phenomenon termed the “escalator to extinction” (Freeman et al., 2018; Urban, 2018). Alternatively, local adaptation can impact adaptive responses to the environment and allow species to persist despite a restricted capacity for range shifts (Aitken et al., 2008; Capblancq et al., 2020). A first step toward improving our understanding of how high-altitude species will respond to climate change is to identify morphological traits and corresponding regions of the genome involved in adaptation to extreme alpine environments (Blanquart et al., 2013; Pritchard & Di Rienzo, 2010).

Birds living in alpine regions are exposed to extreme environments throughout the year, including variable snow cover, hypoxic conditions, and dramatic temperature changes (Grabherr et al., 2010; Körner, 2003). As a result, traits that contribute directly to thermoregulation are expected to be under strong selective pressure in alpine species. Previous work has shown climate-linked traits often follow Allen’s rule, a hypothesis that posits animals adapted to colder environments will have shorter appendages than animals adapted to warm climates (Allen, 1877). In birds, this rule predicts that beaks and wings will be longer in warmer environments to facilitate greater heat dissipation across a larger surface area, and shorter in colder environments to facilitate heat conservation (Greenberg et al., 2012; Lewden et al., 2023; Symonds & Tattersall, 2010; Tattersall et al., 2017; Ward et al., 1999; Weeks et al., 2020, 2025). For this reason, we predicted that birds in high altitude alpine systems will have shorter beaks and smaller wings.

Although less studied in the context of local adaptation, feather microstructure may also be important to avian species' adaptation to environmental variation in temperature. While substantial evidence suggests feather microstructure plays a key role in determining the thermoregulatory capacity of a bird's plumage (Stettenheim, 2000; Stoutjesdijk, 2003; Wolf & Walsberg, 2000), there is some confusion over how environmental variation may drive feather microstructure differences (D'alba et al., 2017; Koskenpato et al., 2016; Lei et al., 2002; Pap et al., 2017, 2020). In one of the few studies that has been conducted at the population level, Koskenpato et al. (2016) found that tawny owls living in colder environments have significantly denser plumulaceous contour feathers than those living in warmer environments. This suggests that feather density may facilitate heat retention. In contrast, however, Pap et al. (2017) found that European bird species wintering in colder areas had less dense feathers than those wintering in warmer places. Thus, there is a clear need for studies specifically linking environmental variation with population-level differences in feather microstructure.

Recent advances in population genomics have enabled the detection of loci involved in local adaptation, even in non-model and wild species (Faria et al., 2014). Among these, F_{ST} outlier tests are commonly used to identify genomic regions that exhibit elevated population differentiation, which may signal divergent selection (Hoban et al., 2016). However, many adaptive traits are likely polygenic, involving small effects at numerous loci (Pritchard & Di Rienzo, 2010; Yeaman, 2015), and such loci will not be picked up by F_{ST} outlier tests. To address this, genome-wide association studies (GWAS) can be used to link more subtle genetic variation to phenotypic variation in ecologically important traits suspected to be under selection (Stinchcombe & Hoekstra, 2008). Taken together, combining approaches that detect both population-level

divergence and polygenic trait architecture offers a powerful framework for investigating local adaptation.

We investigated the potential for local adaptation in the Gray-crowned Rosy Finch (*Leucosticte tephrocotis dawsoni*), one of the highest-breeding songbirds in North America. Despite long-standing interest (Grinnell, 1913, 1917; Twining, 1940), research on this species has been limited due to the difficulty of accessing its extreme nesting sites located in the rocky cliff faces of the Sierra Nevada and White Mountains of California more than 3,000 meters above sea level. To help fill this knowledge gap we conducted morphological and genomic analysis of two populations breeding at contrasting ends of the elevational and thermal range. Piute Pass, located in the Sierra Nevada, is lower in elevation and warmer than the White Mountains (Fig. 1). To identify traits potentially involved in local adaptation, we quantified variation in beak and feather morphology between the two populations. Based on Allen’s rule, we predicted that birds in the warmer Piute Pass population would have longer wings and larger beaks. Further, we predicted that birds in the colder White Mountains location would exhibit feather traits associated with increased insulation—specifically, longer barbules and higher node density. Lastly, we used genome-wide association studies (GWAS) to identify genes linked to feather and bill morphology and F_{ST} outlier tests to identify other potentially important sources of genetic variation. Overall, our findings provide new insights into the process of local adaptation in an extreme alpine specialist bird and provide key baseline information on the capacity for adaptation in the face of global environmental change.

Methods

Site Selection

Our study compares Rosy Finch populations from two sites in the Sierra Nevada and White Mountains. The Piute Pass location is in the Sierra Nevada at approximately

3500m in elevation and has a warmer average temperature during the warmest month of 11°C. The White Mountain location, near White Mountain Peak, is higher in elevation at 4150m and colder with a mean average temperature during the warmest month of 7.5°C (extracted from AdaptWest Project, 2022) (Fig 1). The White Mountain location is located in the rain shadow of the Sierra Nevada Mountains and receives about a third of the precipitation for the same elevation (Rundel et al., 2008). Between the two sites, Piute Pass receives more precipitation as snow and has a more persistent snowpack than the White Mountain location. Additionally, aquatic environments such as streams, lakes and ephemeral pools are more abundant in the Sierra Nevada than the White Mountains (Rundel et al., 2008). The alpine ecosystems of these mountains can be delineated as communities occurring above tree line (Rundel & Millar, 2016). Environmental stressors such as extreme winter temperatures, short growing seasons, high winds, low partial pressures of O₂, and limited water availability are characteristic of these alpine environments (Grabherr et al., 2010; Rundel & Millar, 2016). Separating the western Sierra Nevada Mountains and the eastern White Mountains is the Owens Valley, a subregion of the Great Basin Desert (*Ecosystems of California*, 2016) (Belnap et al., 2016).

Sample Collection

A total of 171 Gray-crowned Rosy Finches were collected using potter traps White Mountains (n=98) and Piute Pass (n=73) in June and July, the peak of the species' breeding season (MacDougall-Shackleton et al., 2000). Birds showing signs of stress were released, resulting in slight variation in the types of data collected for each individual. Blood samples were collected from 150 of the captured birds using the brachial wing vein and stored in Queen's lysis buffer at room temperature (Owen, 2011; Seutin et al., 1991). At the same time, morphological measurements - tarsus length, wing chord, beak width, beak length, beak depth, and nare length - were also collected

from 154 of the birds using an electronic caliper and 5-10 body feathers were collected from the breast of all individuals. Birds were banded to allow for identification of the individuals and to avoid recapture, then released. All handling and banding of birds was done following the guidelines and protocols of the U.S Geological Survey (USGS) Bird Banding Laboratory (BBL) and comply with permits and permissions from federal and state agencies.

Feather Microstructure Measurements

We measured the microstructure of three feathers for each individual. For each feather, three photos were taken using an Olympus BX51 Microscope: one photo each of an unbroken pennaceous and plumulaceous barb from the center of each region was taken at 4x objective, and one photo of a plumulaceous barbule was taken at 10x objective. Structures were then measured from the photos using the segmented line tool in ImageJ. For the pennaceous and plumulaceous barbule length, we measured one barbule from each region starting from the distal tip to where the barbule meets the barb. For the pennaceous and plumulaceous barbule density, we drew a 0.5mm line along the middle of a barb, counting the number of barbules along that line. And finally, for plumulaceous node density, we drew a 0.2mm line along the center of a barbule, counting the number of nodes along that line (Fig 2A). Each replicate measure for a feather trait was averaged per individual prior to downstream analysis.

Statistical analysis for local adaptation

Body size and sex can confound morphological comparisons among individual birds (D. C. Adams et al., 2020). To account for these effects, we included sex and tarsus length (a proxy for body size) as predictor variables in our models. For each trait, we constructed linear models with tarsus length, sex, age, and site code as predictors and used AIC tests to identify the combination of variables that best explained trait variation. Traits for which site code was a significant predictor were considered

significantly different between the two populations. All p-values for site code predictors were corrected for multiple testing using a Bonferroni correction, and corrected p-values are reported. We used *emmeans* (Lenth, 2025) to calculate average trait values while controlling for predictors other than site code, and these adjusted means were overlaid on the raw data in a violin plot (Fig 2 & 3).

Sequencing and Genotyping

DNA was extracted from blood samples of 150 individuals using Qiagen's DNeasy Blood and Tissue Extraction Kit protocol and quantified with the Qubit dsDNA HS Assay Kit. We used a modified version of Illumina's Nextera Library Preparation protocol to prepare WGS libraries and pooled the libraries by equal mass before sequencing. The resulting libraries were sequenced on 2 lanes of an Illumina NovaSeq 6000 at Novogene Corporation.

To process the raw sequence reads and detect variants, we utilized Snakemake (Molder et al. 2021), a workflow management system that provides efficiency, adaptability, and reproducibility. The pipeline we adapted to our species can be found on Github (<https://github.com/eriqande/mega-non-model-wgs-snakeflow/>). Sequence data were trimmed using fastp (S. Chen et al., 2018) to remove adaptor sequences and polyG tails using a sliding window, and then aligned to a Brown-capped Rosy Finch (*Leucosticte australis*) reference genome (GenBank: GCA_025504685.1) using Burrows-Wheeler Aligner software (H. Li & Durbin, 2009). We marked PCR duplicates using *samtools* (Li et al., 2009) and read groups (sample, lane, library) were added using *picard* (<http://broadinstitute.github.io/picard>). Individual coverage was estimated using *samtools*, and given the range of coverage (7.4-19X), we downsampled bam files to 10x using the *samtools* subsample function (Li et al., 2009). Individual gvcf files were created using GATK's HaplotypeCaller (Poplin et al., 2018) with high base quality score filters (`--min-base-quality-score 33 --minimum-mapping-quality 20`) to remove batch

effects (Lou and Therkildsen 2021). We then parallelized the calling of genotypes across 3 million bp regions across the genome using the GenomicsDBImport and GenotypeGVCFs functions (Auwera & O’Connor, 2020; McKenna et al., 2010). GATK versions above 4.0 call missing data as homozygous reference, which can bias downstream analysis. To avoid this, all loci with a depth of zero were manually marked as missing. To remove systematic errors according to GATK variant quality score recalibration (VQSR) best practices, we hard filtered variants with the following parameters: StrandOddsRatio (>3.0), FisherStrand (>60.0), MappingQuality (>40.0), and Quality by Depth (>2.0). We further filtered Single Nucleotide Polymorphisms (SNPs), removing indels, filtering for missingness (<50%), allele frequencies (>0.5 and <0.95), depth (>4x), quality (>30.0) using *bcftools* (Li, 2011). After filtering, the resulting high-quality SNPs were passed through BEAGLE 4.1 (Browning & Browning, 2016) to impute missing genotypes.

Population genetic structure analysis

Population structure was assessed using a Principal Component Analysis (PCA) followed by ADMIXTURE (Alexander et al., 2009). SNPs identified from sequence data were pruned for linkage disequilibrium in PLINK (Purcell et al., 2007) using a 50kb window, 10 bp window step size, and a R² threshold of 0.1. Eigenvalues for the PCA were then generated based on the pruned SNPs using PLINK and the first two principal components (PC) were plotted in R. To further confirm a lack of population structure ADMIXTURE was used to test k=1-2. The results were plotted using R and the cv values were evaluated.

Genome-Wide Association Study

Two GWAS models were implemented on all traits using GEMMA (Zhou et al., 2013), a Bayesian sparse linear model, BSLMM, and a univariate linear model, ULMM. These GWAS approaches allow us to detect SNPs associated with a given trait while

explicitly accounting for population structure and relatedness through the incorporation of random effects whose correlation structure is characterized by a relatedness matrix. Input files were generated with PLINK (Purcell et al., 2007). We implemented a Bayesian sparse linear mixed model (BSLMM) using 500,000 MCMC burn-in iterations that were discarded and 5 million MCMC iterations that were saved. We then filtered for SNPs that had a mean posterior inclusion probability (PIP) above 0.01 (Fig 4 A & B). A univariate linear mixed model (ULMM) employing the Wald test was also conducted. The Wald test assesses the significance of the SNP genotypes in a linear regression model and is used to test the null hypothesis that the effect size of a genetic variant on the phenotypic trait is zero (i.e., there is no association between the variant and the trait). SNPs were then filtered based on a p-value threshold of 5×10^{-8} , a threshold widely used in GWAS studies (e.g. Chen et al., 2021) (Fig 4 A & B).

Genome Wide F_{ST}

To estimate per site F_{ST} , we used OutFLANK (Whitlock & Lotterhos, 2015). OutFLANK identifies outlier F_{ST} SNPs by modeling the distribution of F_{ST} values across loci using a trimmed likelihood approach to exclude loci under strong selection. It then fits a chi-square distribution to the neutral F_{ST} values and identifies SNPs with unusually high or low F_{ST} as potential outliers, suggesting loci under selection. We used a false discovery rate threshold of 0.005 to determine outlier SNPs. We took the 90th quantile of these F_{ST} values to narrow down the top SNPs for lit review and plotted these on a Manhattan plot (Fig 4 C). We then determined the F_{ST} values for the SNPs previously identified in the GWAS analyses based on the values calculated by OutFLANK.

Identification of associated genes, enrichment analysis and literature review

To identify genes associated with significant SNPs from our GWAS and F_{ST} analysis, we implemented BEDTools (Quinlan & Hall, 2010) closest function with a previously created Brown-capped Rosy Finch gene annotation (Funk et al., 2023) to

identify genes close to the SNP positions. The output from this tool was filtered to find only protein coding genes within 25kb of the given SNP. For plotting purposes, we used the nucmer in MUMmer4.x (Marçais et al., 2018)(Marçais et al. 2018) to align the Brown-capped Rosy Finch genome assembly to the Zebra Finch (*Taeniopygia guttata*) genome assembly (GenBank: GCF_003957565.2). We then filtered the delta alignment output of nucmer to keep only matches that were greater than 400 bps, and used the show-coords to display the coordinates. We then used custom R scripts to convert our scaffold and positions to Zebra Finch chromosome positions for plotting SNPs in Manhattan plots.

Enrichment in gene ontology (GO) terms was performed with Panther 19.0 (Thomas et al., 2022). Enrichment thresholds were set to $p < 0.05$ after Bonferroni correction. Gene network analysis was conducted using STRING (Szklarczyk et al., 2023) with zebra finch as the organismal reference. STRING (Search Tool for the Retrieval of Interacting Genes/Proteins) is a database that predicts and visualizes protein-protein interactions based on known and predicted associations from experiments, literature, and computational methods (Szklarczyk et al., 2023). For F_{ST} genes, all genes associated with the OutFLANK outliers were input into enrichment analyses. Genes were grouped with all beak morphology trait genes run as a single analysis, all feather trait genes run together, and wing chord genes run in another analysis.

We performed a literature search for all of the candidate genes using google scholar and the search terms gene name followed by, bird, trait specific term (when applicable), environmental adaptation, local adaptation (e.g., “XIAP” gene beak, or “DSCAM” gene environmental adaptation). For F_{ST} SNPs, only those genes associated with the 90th quantile of SNPs were evaluated in the literature. Genes were grouped by their key functions or associations as described in the literature (Table 1).

Results

Local adaptation of body and feather morphology

AIC model selection indicated five traits where site was a significant predictor. For wing chord, the best model included tarsus length, age, sex, and site. The model showed high explanatory power (adjusted $R^2=0.673$, $p<0.001$) and showed that birds at the White Mountain location (WMTN) had shorter wings chords compared to Piute Pass (PIPA) ($p=0.003$). The best model for nare length included tarsus, sex, and site and had lower explanatory power (adjusted $R^2=0.197$) but was still highly significant ($p<0.001$). This model found that nare length was shorter in the WMTN population ($p<0.001$). Similarly, beak width had tarsus, age and site as significant predictors (adjusted $R^2=0.135$, $p<0.001$) with WMTN having beaks that are wider than those in PIPA ($p=0.004$). Beak depth had tarsus, age and site as the best predictors (adjusted $R^2=0.250$) and found that WMTN has significantly deeper beaks than PIPA ($p=0.014$). Beak length was not significantly different between the two locations and was best predicted by sex (adjusted $R^2=0.516$, $p<0.001$). For feather microstructure, the only trait that had site as a significant predictor was node density. This model included tarsus and age as well as site (adjusted $R^2=0.32$, $p<0.001$) and showed that node density was higher in the WMTN populations ($p=0.037$). The best model for plumulaceous barbule length based on weighted AIC included site as a predictor; however, the effect size was uncertain, and the effect of site was not statistically significant ($p=0.078$). Pennaceous barbule density had tarsus and age as the best predictors (adjusted $R^2=0.365$, $p<0.001$), pennaceous barbule length had sex as the best predictor (adjusted $R^2=0.307$, $p<0.001$) and plumulaceous barbule density had tarsus and age as the best predictors (adjusted $R^2=0.076$, $p=0.004$).

Evaluation of population structure

Whole-genome sequencing was performed on 150 individuals. Two individuals were removed as their collection location was not available. Variant filtering resulted in 7,739,836 SNPs for subsequent genetic analysis. Linkage pruning removed 5,526,104 SNPs. Visualizing PLINK PCA results revealed little population structure with most individuals clustering together (SupFig 1). ADMIXTURE results confirmed that there is no population structure in these samples as k=1 had the lowest cv error (0.599) compared to k=2 or 3 (0.612 and 0.627).

Genes identified with genome-wide association approach

The univariate linear mixed model identified a total of 64 SNPs for traits wing chord, beak length, beak width, pennaceous barbule length, and pennaceous barbule density. Using a Brown-capped Rosy Finch gene annotation we identified a total of 26 unique genes corresponding to these SNPs (Table1). We filtered out genes that have not been fully characterized (e.g., “LOC100221041”) leaving 14 genes. The Bayesian sparse linear mixed model identified 266 SNPs for wing chord, nare, beak length, depth and width, pennaceous barbule length and density, and plumulaceous barbule length and node density. We identified 146 genes associated with these SNPs (Table 1), 110 of which had been fully characterized. For each trait the global view of p-values for all SNPs and associated genes across both models are represented by Manhattan plots (Fig 4 A-C, SupFig 2-5).

Go term analysis using Panther produced significant results for beak depth and feather traits combined. For beak depth, the top terms were presynaptic membrane assembly ($p=1.44 \times 10^{-5}$), synaptic membrane adhesion (specifically cell adhesion) ($p=1.01 \times 10^{-5}$) and heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules ($p=1.13 \times 10^{-5}$). For the feather traits (plumulaceous barbule and node density combined with pennaceous barbule length), fructosamine catabolic process was a

significant term ($p=1.12\times10^{-6}$). Wing chord had no significant GO terms. String analysis for beak and feather traits as well as wing chord did not find significantly more interactions than expected.

Genes identified through high Fixation index (F_{ST}) between the two populations

Using OutFLANK 831 loci were identified as having significantly high F_{ST} when comparing the White Mountain and Piute Pass populations, corresponding to 265 genes (Fig 4D, Sup Table 2). No GWAS SNPs were flagged as having significant F_{ST} values. The minimum F_{ST} to be considered an outlier was 0.17513. The mean F_{ST} of SNPs not considered outliers was 0.0045. Using a 90th quantile we determined $F_{ST}=0.235$ to be the threshold, narrowing down the SNPs to 105 loci. For the smaller set of loci, bedtools identified 61 related genes (Table 2). We ran STRING and GO term enrichments on both the larger and smaller set of genes. The smaller set did not have a significant number of connections over all or any significantly enriched GO terms. The larger set of 264 genes, however, had an overall significant number of connections ($p<0.05$) and had multiple significant annotated keywords (sourced from UniProt). The most significant was the keyword repeat ($p<0.001$), the next was ubl conjugation pathways ($p<0.05$).

Discussion

Understanding the genetic and morphological basis of local adaptation is critical for predicting how species respond to environmental change. Alpine species face unique conservation challenges due to their extreme environments and thus are a high priority for studies of adaptive capacity. In this study, we investigated morphological and genetic differences between two populations of an alpine specialist songbird located at the extreme ends of its environmental range. Consistent with our predictions, we found wings were shorter and feathers were denser in the colder environment. However, in contrast to our prediction, beak depth and width were larger in the colder environment, suggesting other factors like diet may place contrasting selective pressures on this

ecologically important trait. We also identified numerous genes with known links to limb, facial, and feather development that potentially underlie variation in wing, bill and feather microstructure. Further F_{ST} outlier analyses revealed numerous significant differences between cold and warm populations at SNPs located within genes with clear links to altitudinal adaptation. Together, our results provide important insights into the morphological traits and associated genomic regions associated with microgeographic variation in a high-altitude songbird species.

Trait differences between warm and cold habitats

Ecogeographic rules, such as Allen’s rule, are useful for predicting broad geographic trends in morphology in response to environmental conditions, but observed patterns do not always align with expectations. In our study, we found that wing chord was smaller in the colder environment, suggesting that this morphological trait may play a role in thermoregulation, consistent with Allen’s rule. Heat dissipation from the wings occurs across the vascularized brachial regions, primarily during flight when these areas are exposed to air flow (Lewden et al., 2023; Ward et al., 1999). Wing bone length reflects the extent of vascularized areas on the wings and has been shown to be longer in warmer environments across a broad sample of passerine species (Weeks et al., 2025). However, studies of other alpine bird species have found the opposite pattern: that birds from high-elevation sites have longer wings than those from low-elevation sites (Bears et al., 2008; Ceresa et al., 2024). In these cases, researchers have suggested that the demand for more energetically efficient flight at high elevations may outweigh thermoregulatory selective pressures, demonstrating the often contrasting selective pressures on ecologically important traits. Overall, our morphological analyses support the idea that thermoregulatory demands may play a dominant role in shaping wing morphology in the Gray-crowned Rosy Finch, potentially outweighing the benefits of flight efficiency in high-elevation, cold-adapted species.

Contrary to our predictions based on Allen's rule, beak depth and width were larger in the colder environment. Although temperature is often a key environmental variable exerting selective pressure on alpine taxa, variation in beak morphology between our populations may instead reflect divergent selective pressures associated with foraging ecology—a well-documented driver of beak divergence in birds (Grant et al., 1976; Lamichhaney et al., 2018). In our study system, the Piute Pass population likely has access to more aquatic insects due to the higher availability of aquatic habitats proximate to Rosy Finch breeding sites in the Sierra Nevada (Epanchin et al., 2010; Rundel & Millar, 2016). For example, Epanchin et al. (2010) found that, when abundant, mayflies can comprise up to 38% of the Sierra Nevada Gray-Crowned Rosy Finch diet. In contrast, the White Mountain population may rely more heavily on insects deposited in snowfields, a behavior that could favor a more robust, conical bill for extracting frozen prey from the snow surface. Work is currently underway to investigate the dietary differences between these two populations of Rosy Finch (Tim Brown, personal correspondence), and future work should explore the selective pressures experienced by this species. These results emphasize the complexity of identifying drivers of local adaptation, especially in alpine systems where extreme environments, food scarcity, and variation in migration strategies can create conflicting selection pressures.

Of our feather comparisons, we identified significant differences between cold and warm sites for only one trait - plumulaceous node density. In line with our predictions, plumulaceous node density was higher in the colder, high elevation White Mountain population. Higher node density is known to improve air trapping ability, thus trapping warmth against the body surface (King & McLelland, 1984). Consistent with our results, a study on sparrows found that higher elevation alpine forms had higher node density relative to lowland forms (Lei et al., 2002). A further multi-species comparison

found that species wintering in cold and windy conditions had downy feathers with higher node densities (D'alba et al., 2017, 2017). These interspecific trends are mirrored in our results, suggesting that the White Mountain population has feather adaptations consistent with an increased need for heat retention. The lack of significant results for other feather traits may be explained by the amount of variation in the measurements as well as the sample sizes being small, resulting in an underpowered comparison.

Genome-wide analysis

We used GWAS to identify SNPs within genes involved in traits that differ between warm- and cold-adapted populations described previously – beak width and depth, nares length, wing cord length, and pennaceous barbule density. Because ours is one of the first studies to investigate the genetic basis of feather microstructure variation, we also report on the genes associated with other measured feather traits, even though high variance in these traits prevented our ability to robustly investigate differences between populations occupying cold and warm environments.

Our literature review of GWAS results identified genes with diverse associations. For wing chord, we identified limb and bone development, heat response, and vascularization. Notably, the genes *GUCA1C* and *NIPBL* have been linked to limb development in ducks and zebrafish (G. Li et al., 2020; Mohammadi, 2024; Muto et al., 2014) and may play a similar developmental role in songbirds. We also detected the genes *PGLYRP2* and *PACRG*, which have been shown to be upregulated in chickens experiencing heat stress (Kim et al., 2021; Tian et al., 2020). Together, the limb development and heat response genes we identified as important further support the potential connection between wing length and thermoregulation suggested by our morphological finding of shorter wings in the colder environment. Although GWAS

results do not establish a direct functional link to specific traits (Tam et al., 2019), our findings suggest that future research should investigate the potential role of thermoregulatory selective pressures on genes involved in wing development.

When looking at genes associated with beak morphology, BSLMM found more SNPs than LMM and identified several genes potentially associated with facial structure development and climate adaptation. One gene associated with beak depth was *XIAP*. Duplications of this gene have been found to result in facial dysmorphism in humans, including the underdevelopment of cheekbones and protruding jaws (Di Benedetto et al., 2014). Although a link with beak development in birds has not yet been formally established, this could be an area for further research. Another gene identified for beak width was *MINDY2*, which belongs to a group of deubiquitinating enzymes. These genes are related to the regulation of the Wnt pathway (Park et al., 2020), a pathway that is well established as important for beak development (Nierop 2002). Additional genes such as *RASL11A*, *TINAG*, and *PLPP3* have been directly linked to beak development in avian species (Abernathy, 2021; Bai et al., 2018). Two genes associated with beak depth in our study (*LRP1B* and *UTRN*) were shown to be associated with breeding temperature in a study that used a GWAS approach to identify loci associated with climate variables related to morphology in hermit thrush (N. Adams et al., 2025). Although the beak morphology of our species did not entirely fall in line with predictions based around their thermoregulatory role, the elucidation of genes indicating a relationship between thermoregulation and beak morphology suggests that beaks still play a thermoregulatory role in this species.

Our GWAS on feather traits was, to the best of our knowledge, the first genome-wide association study of feather microstructural traits. Identifying the genetic basis of these fine-scale traits opens new avenues for studying how feather structure evolves in response to environmental pressures and functional demands. One of the genes

associated with pennaceous barbule length in our study, *HBE1*, has also been identified in previous research on feather development. Using RNA sequencing, Limber et al. (2024) found that *HBE1*, a hemoglobin gene (Mao et al., 2023), was highly expressed in feather pulp, the central tissue of the developing feather germ. Additional genes associated with feather traits in our study have been identified as being related to plumage coloration in birds, including *EFNA2*, *NRXN1*, and *HERC4* (Shakya, 2020; L. Sun et al., 2020; X. Zhang et al., 2023). Many other genes identified as being associated with feather traits in our work have not been shown to be associated with feather development or morphology in the literature. Feather morphology remains understudied in the context of local adaptation, and the genetic architecture of these microstructural traits is still poorly understood. Our study offers a foundation for deeper investigation. Expanding this work to include comparative genomic studies across populations or species with more strongly differentiated feather morphology could also help link genotype to phenotype and provide insight into the adaptive significance of feather microstructure.

Our original intent with the F_{ST} outlier tests was to determine if any of the genes identified in the GWAS had variants that diverged significantly between the cold and warm environment populations. However, when comparing our F_{ST} results with SNPs identified in the GWAS, we found no overlap between the two sets. This result is consistent with known limitations of F_{ST} -based outlier scans, which are biased toward detecting loci under strong divergent selection and with large effect sizes (Hoban et al., 2016). F_{ST} measures allele frequency differences between populations and loci with large phenotypic effects are more likely to experience strong selection and exhibit pronounced divergence. In contrast, highly polygenic traits are shaped by small-effect variants that shift subtly in frequency—patterns that often fall below the detection threshold of F_{ST} scans. Our findings underscore the value of combining F_{ST} outlier

analyses with GWAS approaches: while the former is designed to pinpoint loci under strong selection, the latter is better suited for uncovering the complex genetic architecture of polygenic traits. Together, these methods can generate a more complete understanding of the genetic basis for local adaptation.

While we did not find overlap between the F_{ST} and the GWAS results, our F_{ST} analysis highlighted numerous genes involved in hypoxia response, altitude adaptation, and cold adaptation. These results are consistent with our hypothesis that elevation and associated climatic variables are applying divergent selection pressures in this species. Of these genes, *NOX4* was found to be involved in mediating hypoxia-inducible transcription factors, a key component of the cellular response to hypoxia (Diebold et al., 2010). Similarly, *MAML2* and *SUPT20H* were found to be associated with altitude adaptation in deer mice (Schweizer et al., 2021a) and *ADGRD1* was associated with altitude adaptation in Tibetan chickens (Nan et al., 2023). Multiple studies identified *PPP1R1C* as being related to cold adaptation in chickens (Fedorova et al., 2022a; Romanov et al., 2023). These are all traits that would lend themselves to survival in alpine environments, and divergent selection is likely a product of varying selection pressures between the higher and colder White Mountain location and the warmer and lower Piute Pass location.

Our F_{ST} analysis also identified several genes that might be involved with plumage color differences in Rosy Finches. Another gene that showed divergence between these two populations of finch was *PHIP* which has been shown to be related to differentiation in plumage coloration in House Finch (Balenger et al., 2015). Interestingly, when comparing candidate outlier genes from this work with work identifying genes involved with the notable plumage coloration differences across the broader Rosy Finch species complex, there were some overlapping genes. In a GWAS on crown coloration, Funk et al (2023) identified *NDUFAF2*, a gene that was also

significant in this work. For body plumage coloration, our results overlap with those of Funk et al (2023) for the genes *KCNV2*, *PUM3*, and *VLDLR*. Although it was not a component of the current work, other studies have shown that plumage coloration is involved in avian thermoregulation, with darker plumage putatively facilitating heat absorption (Medina et al., 2018; Rogalla et al., 2022), and can show adaptation to local environment (Romano et al., 2019; Sandoval & Barrantes, 2019; SirkiÄ et al., 2010). This provides an important avenue for future work with this species investigating the differences in plumage coloration between populations.

An additional result of our F_{ST} analysis was the identification of a gene, *CTIF*, that has been previously identified as being a climate-adapted gene in a landscape genomic study on the closely related Brown-capped Rosy Finch (DeSaix et al., 2022). Interestingly, this gene was also identified through a contrasting approach using a mixed linear model on beak shape for two species of Darwin’s finches, *Geospiza fortis* and *G. scandens* (Lawson & Petren, 2017). The authors of this paper summarize their results as indicating that inter and intraspecies beak shape variation is a result of a small suite of traits evolving in concert, corresponding to many genes. It is possible that we did not identify *CTIF* genes in our GWAS method because of low power due to smaller variation in the beak traits measured, a result of using continuous measurements between two populations versus categorical comparisons between different species as in Lawson & Petren (2017). Because of the large number of SNPs obtained for our two populations, our F_{ST} analysis likely had higher power and thus detected this gene, in addition to the fact that it had a high F_{ST} of 0.263 (averaged across the 5 SNPs associated with this trait). The overlap of these genes in the two studies described as well as ours makes it a strong candidate as a genetic component of adaptation to environment, potentially through the mechanism of beak morphology.

An important outcome of local adaptation is the maintenance of ecologically important genetic variation (Savolainen et al., 2013). Conserving genetic variation prevents loss of adaptive potential and can underlie rapid adaptive responses to environmental change (Forester et al., 2023). We integrated phenotypic measurements, genome-wide association studies, and F_{ST} analyses, to understand the morphological and genetic basis of local adaptation of an alpine bird species, the Sierra Gray-crowned Rosy Finch. When comparing traits suspected to be involved in thermoregulation for this species, we found that morphological variation across populations only partially aligned with ecogeographic expectations. Wing chord was shorter in the colder environment, consistent with Allen’s rule and likely reflecting thermoregulatory adaptation. However, beak depth and width were larger in colder environments, contrary to predictions, suggesting additional selective pressures such as diet specialization may be playing a role. Comparisons of feather microstructure further showed thermoregulatory adaptation, with higher plumulaceous node density in the high-elevation population likely enhancing insulation. When coupled with the results of our GWAS and F_{ST} outlier scans we gained a clear picture of traits and genes important to adaptation in alpine birds. Importantly, we found that many of the genes underlying wing length, beak depth and feather microstructure had links to both development of similar traits in other organisms as well as thermoregulation. Our genomic and morphological results provide strong evidence that this alpine species is locally adapted to temperature variation. Overall, this work enhances our understanding of the morphological traits and genetic mechanisms underlying local adaptation in an alpine species. The resulting insights will be used inform the delineation of conservation units and support management strategies that seek to preserve the evolutionary processes essential for long-term

persistence of alpine organisms in a changing world (Allendorf et al., 2010; Flanagan et al., 2018; Shafer et al., 2015).

Acknowledgements

We thank Sarah Albright, Alena Arnold, Emma Brown, Carl Brown, Alison Brunschwiler, Vincent Chevreuil, Ryan Cheung, Daisy Cortes, Jessica Gallardo, Karim Hanna, Raquel Lozano, Juan Rebellon, Leticia Santillana, Kevin Silberberg, and Juyung Yoo for their participating in catching, handling, banding, measuring, and sampling birds. We thank Bailey Lubash for contributing to the methodology for measuring feather microstructure characteristics. We thank NovoGene for their assistance with next generation sequencing. We thank Christine Rayne and Amanda Carpenter for library preparation, and the Ruegg and Funk labs at Colorado State University for detailed feedback on early interactions of the project and manuscript. This work utilized the RMACC Summit supercomputer, which is supported by the National Science Foundation (awards ACI-1532235 and ACI-1532236), the University of Colorado Boulder and Colorado State University. The RMACC Summit supercomputer is a joint effort of the University of Colorado Boulder and Colorado State University. Figures were created with [BioRender.com](https://www.biorender.com). Funding for this project was made possible by the National Science Foundation (NSF 2222524, NSF 2222525, and NSF 2222526). Handling, banding, and sampling of Grey-crowned Rosy Finch was performed under Federal Banding Permit 24401 and CFWScientific Collecting Permit (SCP) S-190610001-22081-001-02.

References

<https://doi.org/10.1371/journal.pgen.1003264>

Abernathy, E. (2021). *The Genomic Basis of Population and Adaptive Divergence in Buteo Sister Species Across Multiple Evolutionary and Geographic Scales* [Ph.D., University of California, Davis].

<https://www.proquest.com/docview/2638107596/abstract/BB50D6B5F64067PQ/1>

Adams, D. C., Glynne, E., & Kaliontzopoulou, A. (2020). Interspecific allometry for sexual shape dimorphism: Macroevolution of multivariate sexual phenotypes with application to Rensch’s rule. *Evolution; International Journal of Organic Evolution*, 74(9), 1908–1922. <https://doi.org/10.1111/evo.14049>

Adams, N., Dias, T., Skeen, H. R., Pegan, T., Willard, D. E., Winger, B., Ruegg, K., Weeks, B.

C., & Bay, R. (2025). Genetic and morphological shifts associated with climate change in a migratory bird. *BMC Biology*, 23(1), 3. <https://doi.org/10.1186/s12915-024-02107-5>

AdaptWest Project. (2022). *Gridded current and projected climate data for North America at 1km resolution, generated using the ClimateNA v7.30 software (T. Wang et al., 2022)* [Dataset].

Ahmad, H. I., Asif, A. R., Ahmad, M. J., Jabbar, F., Adnan, M., Ahmed, S., Afzal, G., Saleem, A. H., Li, L., Jiang, H., Simirgiotis, M. J., Rauf, M., & Chen, J. (2020). Adaptive evolution of peptidoglycan recognition protein family regulates the innate signaling against microbial pathogens in vertebrates. *Microbial Pathogenesis*, 147, 104361. <https://doi.org/10.1016/j.micpath.2020.104361>

Aitken, S. N., Yeaman, S., Holliday, J. A., Wang, T., & Curtis-McLane, S. (2008). Adaptation, migration or extirpation: Climate change outcomes for tree populations. *Evolutionary Applications*, 1(1), 95–111. <https://doi.org/10.1111/j.1752-4571.2007.00013.x>

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, 19(9), 1655–1664. <https://doi.org/10.1101/gr.094052.109>

Allen, J. A. (1877). *The Influence of Physical Conditions in the Genesis of Species*.

Allendorf, F. W., Hohenlohe, P. A., & Luikart, G. (2010). Genomics and the future of conservation genetics. *Nature Reviews Genetics*, 11(10), 697–709. <https://doi.org/10.1038/nrg2844>

Alsoufi, M. A., Liu, Y., Cao, C., Zhao, J., Kang, J., Li, M., Wang, K., He, Y., & Ge, C. (2023). Integrated Transcriptomics Profiling in Chahua and Digao Chickens' Breast for Assessment Molecular Mechanism of Meat Quality Traits. *Genes*, 14(1), Article 1.

<https://doi.org/10.3390/genes14010095>

Auwerda, G. V. der, & O'Connor, B. D. (2020). *Genomics in the cloud: Using Docker, GATK, and WDL in Terra* (First edition). O'Reilly.

Bai, H., Sun, Y., Liu, N., Xue, F., Li, Y., Xu, S., Ye, J., Zhang, L., Chen, Y., & Chen, J. (2018). Single SNP- and pathway-based genome-wide association studies for beak deformity in chickens using high-density 600K SNP arrays. *BMC Genomics*, 19(1), 501. <https://doi.org/10.1186/s12864-018-4882-8>

Bai, H., Zhao, N., Li, X., Ding, Y., Guo, Q., Chen, G., & Chang, G. (2024). Whole-genome resequencing identifies candidate genes associated with heat adaptation in chickens. *Poultry Science*, 103(10), 104139. <https://doi.org/10.1016/j.psj.2024.104139>

Bai, H., Zhu, J., Sun, Y., Liu, R., Liu, N., Li, D., Wen, J., & Chen, J. (2014). Identification of Genes Related to Beak Deformity of Chickens Using Digital Gene Expression Profiling. *PLOS ONE*, 9(9), e107050. <https://doi.org/10.1371/journal.pone.0107050>

Balenger, S. L., Bonneaud, C., Sefick, S. A., Edwards, S. V., & Hill, G. E. (2015). Plumage color and pathogen-induced gene expression in a wild bird. *Behavioral Ecology*, 26(4), 1100–1110. <https://doi.org/10.1093/beheco/arv055>

Barnes, N. E., Mendoza, K. M., Strasburg, G. M., Velleman, S. G., & Reed, K. M. (2019). Thermal challenge alters the transcriptional profile of the breast muscle in turkey poults. *Poultry Science*, 98(1), 74–91. <https://doi.org/10.3382/ps/pey401>

Bears, H., C. Drever, M., & Martin, K. (2008). Comparative morphology of dark-eyed juncos *Junco hyemalis* breeding at two elevations: A common aviary experiment. *Journal of Avian Biology*, 39(2), 152–162. <https://doi.org/10.1111/j.2008.0908-8857.04191.x>

Belnap, J., Webb, R. H., Esque, T. C., Brooks, M. L., Defalco, L. A., & Macmahon, J. A. (2016). Deserts. In *Ecosystems of California* (1st ed., pp. 635–668). University of

California Press. <http://www.jstor.org/stable/10.1525/j.ctv1xxzp6.35>

Bhavana, K., Foote, D. J., Srikanth, K., Balakrishnan, C. N., Prabhu, V. R., Sankaralingam, S., Singha, H. S., Gopalakrishnan, A., & Nagarajan, M. (2022). Comparative transcriptome analysis of Indian domestic duck reveals candidate genes associated with egg production. *Scientific Reports*, 12(1), 10943. <https://doi.org/10.1038/s41598-022-15099-5>

Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to measuring local adaptation. *Ecology Letters*, 16(9), 1195–1205. <https://doi.org/10.1111/ele.12150>

Boer, E. F., Maclary, E. T., & Shapiro, M. D. (2021). Complex genetic architecture of three-dimensional craniofacial shape variation in domestic pigeons. *Evolution & Development*, 23(6), 477–495. <https://doi.org/10.1111/ede.12395>

Borges, R., Fonseca, J., Gomes, C., Johnson, W. E., O'Brien, S. J., Zhang, G., Gilbert, M. T. P., Jarvis, E. D., & Antunes, A. (2019). Avian Binocularity and Adaptation to Nocturnal Environments: Genomic Insights from a Highly Derived Visual Phenotype. *Genome Biology and Evolution*, 11(8), 2244–2255. <https://doi.org/10.1093/gbe/evz111>

Browning, B. L., & Browning, S. R. (2016). Genotype Imputation with Millions of Reference Samples. *American Journal of Human Genetics*, 98(1), 116–126. <https://doi.org/10.1016/j.ajhg.2015.11.020>

Buroker, N. E., Ning, X.-H., Zhou, Z.-N., Li, K., Cen, W.-J., Wu, X.-F., Zhu, W.-Z., Scott, C. R., & Chen, S.-H. (2012). AKT3, ANGPTL4, eNOS3, and VEGFA associations with high altitude sickness in Han and Tibetan Chinese at the Qinghai-Tibetan Plateau. *International Journal of Hematology*, 96(2), 200–213. <https://doi.org/10.1007/s12185-012-1117-7>

Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Exposito-Alonso, M., & Keller, S. R. (2020). Genomic Prediction of (Mal)Adaptation Across Current and Future Climatic Landscapes. *Annual Review of Ecology, Evolution, and Systematics*, 51(Volume 51, 2020), 245–269. <https://doi.org/10.1146/annurev-ecolsys-020720-042553>

Ceresa, F., Brambilla, M., Kvist, L., Vitulano, S., Pes, M., Tomasi, L., Pedrini, P., Bettega, C., Anderle, M., Hilpold, A., & Kranebitter, P. (2024). Wing morphology changes with habitat availability and elevation in an alpine-specialist bird. *Global Ecology and Conservation*, 56, e03276. <https://doi.org/10.1016/j.gecco.2024.e03276>

Chen, C.-K., Chang, Y.-M., Jiang, T.-X., Yue, Z., Liu, T.-Y., Lu, J., Yu, Z., Lin, J.-J., Vu, T.-D., Huang, T.-Y., Harn, H. I.-C., Ng, C. S., Wu, P., Chuong, C.-M., & Li, W.-H. (2024). Conserved regulatory switches for the transition from natal down to juvenile feather in birds. *Nature Communications*, 15(1), 4174. <https://doi.org/10.1038/s41467-024-48303-3>

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, 34(17), i884–i890. <https://doi.org/10.1093/bioinformatics/bty560>

Chen, X., Bai, X., Liu, H., Zhao, B., Yan, Z., Hou, Y., & Chu, Q. (2022). Population Genomic Sequencing Delineates Global Landscape of Copy Number Variations that Drive Domestication and Breed Formation of in Chicken. *Frontiers in Genetics*, 13. <https://doi.org/10.3389/fgene.2022.830393>

Chen, Z., Boehnke, M., Wen, X., & Mukherjee, B. (2021). Revisiting the genome-wide significance threshold for common variant GWAS. *G3 Genes|Genomes|Genetics*, 11(2), jkaa056. <https://doi.org/10.1093/g3journal/jkaa056>

Cheng, Y., Miller, M. J., Zhang, D., Song, G., Jia, C., Qu, Y., & Lei, F. (2020). Comparative Genomics Reveals Evolution of a Beak Morphology Locus in a High-Altitude

Songbird. *Molecular Biology and Evolution*, 37(10), 2983–2988.

<https://doi.org/10.1093/molbev/msaa157>

Cho, E., Kim, M., Cho, S., So, H.-J., Lee, K.-T., Cha, J., Jin, D., & Lee, J. H. (2023). A genome-wide association study for the fatty acid composition of breast meat in an F2 crossbred chicken population. *Journal of Animal Science and Technology*, 65(4), 735–747. <https://doi.org/10.5187/jast.2023.e1>

Cho, Y. S., Jun, J. H., Kim, J. A., Kim, H.-M., Chung, O., Kang, S.-G., Park, J.-Y., Kim, H.-J., Kim, S., Kim, H.-J., Jang, J., Na, K.-J., Kim, J., Park, S. G., Lee, H.-Y., Manica, A., Mindell, D. P., Fuchs, J., Edwards, J. S., ... Bhak, J. (2019). Raptor genomes reveal evolutionary signatures of predatory and nocturnal lifestyles. *Genome Biology*, 20(1), 181. <https://doi.org/10.1186/s13059-019-1793-1>

Chung, O., Jin, S., Cho, Y. S., Lim, J., Kim, H., Jho, S., Kim, H.-M., Jun, J., Lee, H., Chon, A., Ko, J., Edwards, J., Weber, J. A., Han, K., O'Brien, S. J., Manica, A., Bhak, J., & Paek, W. K. (2015). The first whole genome and transcriptome of the cinereous vulture reveals adaptation in the gastric and immune defense systems and possible convergent evolution between the Old and New World vultures. *Genome Biology*, 16(1), 215. <https://doi.org/10.1186/s13059-015-0780-4>

Clark, J., Lim, H. C., Olsen, B., & Kovach, A. (2024). *Whole-genome data uncover the basis of local adaptation with gene flow in a threatened coastal songbird*. <https://doi.org/DOI: 10.22541/au.172114945.59127531/v1>

Clausen, J., Keck, D. D., & Hiesey, W. M. (1941). Regional Differentiation in Plant Species. *The American Naturalist*, 75(758), 231–250. <https://doi.org/10.1086/280955>

Crates, R., von Takach, B., Young, C. M., Stojanovic, D., Neaves, L. E., Murphy, L., Gautschi, D., Hogg, C. J., Heinsohn, R., Bell, P., & Farquharson, K. A. (2024). Genomic insights into the critically endangered King Island scrubtit. *Journal of*

Heredity, 115(5), 552–564. <https://doi.org/10.1093/jhered/esae029>

da Costa Moreno, H. (2022). *Molecular Evolution of Vertebrate Genes Involved in Mitochondrial Biogenesis: Unravelling the Origins of Birds’ Endothermy* [Masters Thesis, Universidade do Porto]. ProQuest Dissertations & Theses.

D’alba, L., Carlsen, T. H., Ásgeirsson, Á., Shawkey, M. D., & Jónsson, J. E. (2017). Contributions of feather microstructure to eider down insulation properties. *Journal of Avian Biology*, 48(8), 1150–1157. <https://doi.org/10.1111/jav.01294>

Delmore, K., Illera, J. C., Pérez-Tris, J., Segelbacher, G., Lugo Ramos, J. S., Durieux, G., Ishigohoka, J., & Liedvogel, M. (2020). The evolutionary history and genomics of European blackcap migration. *eLife*, 9, e54462. <https://doi.org/10.7554/eLife.54462>

DeSaix, M., George, T., Seglund, A., Spellman, G., Zavaleta, E., & Ruegg, K. (2022). Forecasting climate change response in an alpine specialist songbird reveals the importance of considering novel climate. *Diversity and Distributions*, 28. <https://doi.org/10.1111/ddi.13628>

Di Benedetto, D., Musumeci, S. A., Avola, E., Alberti, A., Buono, S., Scuderi, C., Grillo, L., Galesi, O., Spalletta, A., Giudice, M. L., Luciano, D., Vinci, M., Bianca, S., Romano, C., & Fichera, M. (2014). Definition of minimal duplicated region encompassing the XIAP and STAG2 genes in the Xq25 microduplication syndrome. *American Journal of Medical Genetics. Part A*, 164A(8), 1923–1930. <https://doi.org/10.1002/ajmg.a.36570>

Diebold, I., Petry, A., Hess, J., & Görlach, A. (2010). The NADPH Oxidase Subunit NOX4 Is a New Target Gene of the Hypoxia-inducible Factor-1. *Molecular Biology of the Cell*, 21(12), 2087–2096. <https://doi.org/10.1091/mbc.e09-12-1003>

Ducrest, A.-L., San-Jose, L. M., Neuenschwander, S., Schmid-Siegert, E., Simon, C., Pagni, M., Iseli, C., Richter, H., Guex, N., Cumer, T., Beaudoin, E., Dupasquier, M.,

- Charruau, P., Ducouret, P., Xenarios, I., Goudet, J., & Roulin, A. (2025). Melanin and Neurotransmitter Signalling Genes Are Differentially Co-Expressed in Growing Feathers of White and Rufous Barn Owls. *Pigment Cell & Melanoma Research*, 38(2), e70001. <https://doi.org/10.1111/pcmr.70001>
- Endler, J. A. (1977). *Geographic Variation, Speciation and Clines. (MPB-10), Volume 10*. Princeton University Press. <https://doi.org/10.2307/j.ctvx5wbdg>
- Epanchin, P. N., Knapp, R. A., & Lawler, S. P. (2010). Nonnative trout impact an alpine-nesting bird by altering aquatic-insect subsidies. *Ecology*, 91(8), 2406–2415. <https://doi.org/10.1890/09-1974.1>
- Faria, R., Renaut, S., Galindo, J., Pinho, C., Melo-Ferreira, J., Melo, M., Jones, F., Salzburger, W., Schluter, D., & Butlin, R. (2014). Advances in Ecological Speciation: An integrative approach. *Molecular Ecology*, 23(3), 513–521. <https://doi.org/10.1111/mec.12616>
- Farias-Virgens, M., Peede, D., Deacon, T., Okanoya, K., White, S. A., & Huerta-Sanchez, E. (2025). The genomics of the domestication syndrome in a songbird model species. *Communications Biology*, 8(1), 853. <https://doi.org/10.1038/s42003-025-08235-0>
- Fedorova, E. S., Dementieva, N. V., Shcherbakov, Y. S., & Stanishevskaya, O. I. (2022a). Identification of Key Candidate Genes in Runs of Homozygosity of the Genome of Two Chicken Breeds, Associated with Cold Adaptation. *Biology*, 11(4), Article 4. <https://doi.org/10.3390/biology11040547>
- Feng, J., Zhu, W., Shi, H., Peng, D., Zang, L., Wang, Y., ZhaXi, L., BaiMa, J., Amevor, F. K., Wang, X., Ma, X., & Zhao, X. (2023). Analysis of the Selection Signal of the Tibetan Black Chicken Genome Based on Whole-Genome Sequencing. *Genes*, 14(9), Article 9. <https://doi.org/10.3390/genes14091672>
- Flanagan, S. P., Forester, B. R., Latch, E. K., Aitken, S. N., & Hoban, S. (2018). Guidelines

for planning genomic assessment and monitoring of locally adaptive variation to inform species conservation. *Evolutionary Applications*, 11(7), 1035–1052.
<https://doi.org/10.1111/eva.12569>

Forester, B. R., Beever, E. A., Darst, C., Szymanski, J., & Funk, W. C. (2022). Linking evolutionary potential to extinction risk: Applications and future directions. *Frontiers in Ecology and the Environment*, 20(9), 507–515.
<https://doi.org/10.1002/fee.2552>

Forester, B. R., Day, C. C., Ruegg, K., & Landguth, E. L. (2023). Evolutionary potential mitigates extinction risk under climate change in the endangered southwestern willow flycatcher. *Journal of Heredity*, 114(4), 341–353.
<https://doi.org/10.1093/jhered/esac067>

Franchini, P., Irisarri, I., Fudickar, A., Schmidt, A., Meyer, A., Wikelski, M., & Partecke, J. (2017). Animal tracking meets migration genomics: Transcriptomic analysis of a partially migratory bird species. *Molecular Ecology*, 26(12), 3204–3216.
<https://doi.org/10.1111/mec.14108>

Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018). Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the National Academy of Sciences*, 115(47), 11982–11987. <https://doi.org/10.1073/pnas.1804224115>

Fridolfsson, A.-K., & Ellegren, H. (2000). Molecular Evolution of the Avian CHD1 Genes on the Z and W Sex Chromosomes. *Genetics*, 155(4), 1903–1912.
<https://doi.org/10.1093/genetics/155.4.1903>

Funk, E. R., Spellman, G. M., Winker, K., Withrow, J. J., Ruegg, K. C., & Taylor, S. A. (2023). The genetic basis of plumage coloration and elevation adaptation in a clade of recently diverged alpine and arctic songbirds. *Evolution; International Journal of*

- Organic Evolution*, 77(3), 705–717. <https://doi.org/10.1093/evolut/qpac064>
- Gao, G., Zhang, K., Zhao, X., Xu, G., Xie, Y., Zhou, L., Zhang, C., & Wang, Q. (2023). Identification of Molecular Markers Associated with Goose Egg Quality Through Genome-Wide Association Analysis. *Scientia Agricultura Sinica*, 56(19), 3894–3904. <https://doi.org/10.3864/j.issn.0578-1752.2023.19.015>
- García, N. C., Campagna, L., Rush, A. C., Bowie, R. C. K., & Lovette, I. J. (2023). Comparative genomics of two *Empidonax* flycatchers reveal candidate genes for bird song production. *Evolution*, 77(8), 1818–1828. <https://doi.org/10.1093/evolut/qpac096>
- Gesemann, M., & Neuhauss, S. C. F. (2023). Evolution of visual guanylyl cyclases and their activating proteins with respect to clade and species-specific visual system adaptation. *Frontiers in Molecular Neuroscience*, 16. <https://doi.org/10.3389/fnmol.2023.1131093>
- Gheyas, A. A., Vallejo-Trujillo, A., Kebede, A., Lozano-Jaramillo, M., Dessie, T., Smith, J., & Hanotte, O. (2021). Integrated Environmental and Genomic Analysis Reveals the Drivers of Local Adaptation in African Indigenous Chickens. *Molecular Biology and Evolution*, 38(10), 4268–4285. <https://doi.org/10.1093/molbev/msab156>
- Gienapp, P., Laine, V. N., Mateman, A. C., van Oers, K., & Visser, M. E. (2017). Environment-Dependent Genotype-Phenotype Associations in Avian Breeding Time. *Frontiers in Genetics*, 8. <https://doi.org/10.3389/fgene.2017.00102>
- Grabherr, G., Gottfried, M., & Pauli, H. (2010). Climate Change Impacts in Alpine Environments. *Geography Compass*, 4(8), 1133–1153. <https://doi.org/10.1111/j.1749-8198.2010.00356.x>
- Grant, P. R., Grant, B. R., Smith, J. N., Abbott, I. J., & Abbott, L. K. (1976). Darwin's finches: Population variation and natural selection. *Proceedings of the National Academy of*

Sciences, 73(1), 257–261. <https://doi.org/10.1073/pnas.73.1.257>

Greenberg, R., Cadena, V., Danner, R. M., & Tattersall, G. (2012). Heat Loss May Explain Bill Size Differences between Birds Occupying Different Habitats. *PLOS ONE*, 7(7), e40933. <https://doi.org/10.1371/journal.pone.0040933>

Grinnell, J. (1913). *Leucosticte tephrocotis dawsoni*: A New Race of Rosy Finch from the Sierra Nevada. *The Condor*, 15(2), 76–79. <https://doi.org/10.2307/1362056>

Grinnell, J. (1917). Field Tests of Theories Concerning Distributional Control. *The American Naturalist*, 51(602), 115–128. <https://doi.org/10.1086/279591>

Guo, J., Sun, C., Qu, L., Shen, M., Dou, T., Ma, M., Wang, K., & Yang, N. (2017). Genetic architecture of bone quality variation in layer chickens revealed by a genome-wide association study. *Scientific Reports*, 7(1), 45317. <https://doi.org/10.1038/srep45317>

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., Poss, M. L., Reed, L. K., Storfer, A., & Whitlock, M. C. (2016). Finding the Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. *The American Naturalist*, 188(4), 379–397. <https://doi.org/10.1086/688018>

Hu, X., Zhang, X., Liu, Z., Li, S., Zheng, X., Nie, Y., Tao, Y., Zhou, X., Wu, W., Yang, G., Zhao, Q., Zhang, Y., Xu, Q., & Mou, C. (2020). Exploration of key regulators driving primary feather follicle induction in goose skin. *Gene*, 731, 144338. <https://doi.org/10.1016/j.gene.2020.144338>

Johnsson, M., Jonsson, K. B., Andersson, L., Jensen, P., & Wright, D. (2015). Genetic Regulation of Bone Metabolism in the Chicken: Similarities and Differences to Mammalian Systems. *PLOS Genetics*, 11(5), e1005250. <https://doi.org/10.1371/journal.pgen.1005250>

Johnston, R. A., Paxton, K. L., Moore, F. R., Wayne, R. K., & Smith, T. B. (2016). Seasonal

gene expression in a migratory songbird. *Molecular Ecology*, 25(22), 5680–5691.

<https://doi.org/10.1111/mec.13879>

Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, 7(12), 1225–1241. <https://doi.org/10.1111/j.1461-0248.2004.00684.x>

Kim, H., Kim, H., Seong, P., Arora, D., Shin, D., Park, W., & Park, J.-E. (2021).

Transcriptomic Response under Heat Stress in Chickens Revealed the Regulation of

Genes and Alteration of Metabolism to Maintain Homeostasis. *Animals: An Open*

Access Journal from MDPI, 11(8), 2241. <https://doi.org/10.3390/ani11082241>

King, A. S., & McLelland, J. (1984). *Birds: Their structure and function* (2nd ed). Baillière Tindall.

Kinsella, C. M., Ruiz-Ruano, F. J., Dion-Côté, A.-M., Charles, A. J., Gossmann, T. I.,

Cabrero, J., Kappei, D., Hemmings, N., Simons, M. J. P., Camacho, J. P. M.,

Forstmeier, W., & Suh, A. (2019). Programmed DNA elimination of germline

development genes in songbirds. *Nature Communications*, 10(1), 5468.

<https://doi.org/10.1038/s41467-019-13427-4>

Körner, C. (2003). Alpine climate. In C. Körner (Ed.), *Alpine Plant Life: Functional Plant*

Ecology of High Mountain Ecosystems (pp. 21–30). Springer.

https://doi.org/10.1007/978-3-642-18970-8_3

Koskenpato, K., Ahola, K., Karstinen, T., & Karell, P. (2016). Is the denser contour feather

structure in pale grey than in pheomelanic brown tawny owls *Strix aluco* an

adaptation to cold environments? *Journal of Avian Biology*, 47(1), 1–6.

<https://doi.org/10.1111/jav.00746>

Kuttiyarthu Veetil, N., Henschen, A. E., Hawley, D. M., Melepat, B., Dalloul, R. A., Beneš, V.,

Adelman, J. S., & Vinkler, M. (2024). Varying conjunctival immune response

adaptations of house finch populations to a rapidly evolving bacterial pathogen.

Frontiers in Immunology, 15. <https://doi.org/10.3389/fimmu.2024.1250818>

Lamichhaney, S., Han, F., Webster, M. T., Andersson, L., Grant, B. R., & Grant, P. R. (2018). Rapid hybrid speciation in Darwin’s finches. *Science*, 359(6372), 224–228. <https://doi.org/10.1126/science.aao4593>

Lande, R., & Shannon, S. (1996). THE ROLE OF GENETIC VARIATION IN ADAPTATION AND POPULATION PERSISTENCE IN A CHANGING ENVIRONMENT. *Evolution*, 50(1), 434–437. <https://doi.org/10.1111/j.1558-5646.1996.tb04504.x>

Lawson, L. P., & Petren, K. (2017). The adaptive genomic landscape of beak morphology in Darwin’s finches. *Molecular Ecology*, 26(19), 4978–4989. <https://doi.org/10.1111/mec.14166>

Lei, F.-M., Qu, Y.-H., Gan, Y.-L., Gebauer, A., & Kaiser, M. (2002). The feather microstructure of Passerine sparrows in China. *Journal Für Ornithologie*, 143(2), 205–212. <https://doi.org/10.1046/j.1439-0361.2002.02001.x>

Lennon, O. (2022). *Do Candidate Genes Distinguish Migratory and Nonmigratory Avian Species?* [Undergraduate Research Scholars Thesis, Texas A&M University]. <https://hdl.handle.net/1969.1/196498>

Lenth, R. (2025). *emmeans: Estimated Marginal Means, aka Least-Squares Means* (Version 1.10.7-100002) [R package].

Lewden, A., Bishop, C. M., & Askew, G. N. (2023). How birds dissipate heat before, during and after flight. *Journal of The Royal Society Interface*, 20(209), 20230442. <https://doi.org/10.1098/rsif.2023.0442>

Li, G.-S., Liu, W.-W., Zhang, F., Zhu, F., Yang, F.-X., Hao, J.-P., & Hou, Z.-C. (2020). Genome-wide association study of bone quality and feed efficiency-related traits in Pekin ducks. *Genomics*, 112(6), 5021–5028. <https://doi.org/10.1016/j.ygeno.2020.09.023>

- Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*, 25(14), 1754–1760.
<https://doi.org/10.1093/bioinformatics/btp324>
- Limber, C., Wagner, G. P., & Prum, R. O. (2024). *Genetic Characterization of the Cell Types of in Developing Feathers, and the Evolution of Feather Complexity* (p. 2024.11.21.624532). bioRxiv. <https://doi.org/10.1101/2024.11.21.624532>
- Loma, J. D., Vicente, M., Tirado, N., Ascui, F., Parada, L. A., Gardon, J., Schlebusch, C., & Broberg, K. (2025). Human adaptation in the Andes Mountains. *Human Population Genetics and Genomics*, 5(1). <https://doi.org/10.47248/hpgg2505010002>
- Lotterhos, K. E., & Whitlock, M. C. (2015). The relative power of genome scans to detect local adaptation depends on sampling design and statistical method. *Molecular Ecology*, 24(5), 1031–1046. <https://doi.org/10.1111/mec.13100>
- Lovell, P. V., Carleton, J. B., & Mello, C. V. (2013). Genomics analysis of potassium channel genes in songbirds reveals molecular specializations of brain circuits for the maintenance and production of learned vocalizations. *BMC Genomics*, 14(1), 470. <https://doi.org/10.1186/1471-2164-14-470>
- Luo, X., Guo, J., Zhang, J., Ma, Z., & Li, H. (2024). Overview of chicken embryo genes related to sex differentiation. *PeerJ*, 12, e17072. <https://doi.org/10.7717/peerj.17072>
- Ma, S., Liu, H., Wang, J., Wang, L., Xi, Y., Liu, Y., Xu, Q., Hu, J., Han, C., Bai, L., Li, L., & Wang, J. (2021). Transcriptome Analysis Reveals Genes Associated With Sexual Dichromatism of Head Feather Color in Mallard. *Frontiers in Genetics*, 12. <https://doi.org/10.3389/fgene.2021.627974>
- MacDougall-Shackleton, S. A., Johnson, R. E., & Hahn, T. P. (2000). Gray-crowned Rosy-Finch (*Leucosticte tephrocotis*). In *The Birds of North America*. Cornell Lab of Ornithology.

Mamantov, M. A., Gibson-Reinemer, D. K., Linck, E. B., & Sheldon, K. S. (2021). Climate-driven range shifts of montane species vary with elevation. *Global Ecology and Biogeography*, 30(4), 784–794. <https://doi.org/10.1111/geb.13246>

Marçais, G., Delcher, A. L., Phillippy, A. M., Coston, R., Salzberg, S. L., & Zimin, A. (2018). MUMmer4: A fast and versatile genome alignment system. *PLOS Computational Biology*, 14(1), e1005944. <https://doi.org/10.1371/journal.pcbi.1005944>

Mascarenhas, R., Meirelles, P. M., & Batalha-Filho, H. (2023). Urbanization drives adaptive evolution in a Neotropical bird. *Current Zoology*, 69(5), 607–619. <https://doi.org/10.1093/cz/zoac066>

Matsuda, Y., & Makino, T. (2024). Comparative genomics reveals convergent signals associated with the high metabolism and longevity in birds and bats. *Proceedings of the Royal Society B: Biological Sciences*, 291(2029), 20241068. <https://doi.org/10.1098/rspb.2024.1068>

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, 20(9), 1297–1303. <https://doi.org/10.1101/gr.107524.110>

Medina, I., Newton, E., Kearney, M. R., Mulder, R. A., Porter, W. P., & Stuart-Fox, D. (2018). Reflection of near-infrared light confers thermal protection in birds. *Nature Communications*, 9(1), 3610. <https://doi.org/10.1038/s41467-018-05898-8>

Meek, M. H., Beever, E. A., Barbosa, S., Fitzpatrick, S. W., Fletcher, N. K., Mittan-Moreau, C. S., Reid, B. N., Campbell-Staton, S. C., Green, N. F., & Hellmann, J. J. (2023). Understanding Local Adaptation to Prepare Populations for Climate Change. *BioScience*, 73(1), 36–47. <https://doi.org/10.1093/biosci/biac101>

- Mesner, L. D., Ray, B., Hsu, Y.-H., Manichaikul, A., Lum, E., Bryda, E. C., Rich, S. S., Rosen, C. J., Criqui, M. H., Allison, M., Budoff, M. J., Clemens, T. L., & Farber, C. R. (2014). *Bicc1* is a genetic determinant of osteoblastogenesis and bone mineral density. *The Journal of Clinical Investigation*, 124(6), 2736–2749. <https://doi.org/10.1172/JCI73072>
- Mohammadi, H. (2024a). A Genome-wide Association Study based on Pathway Analysis Related to the Angle Wing in Pekin Ducks using Whole-Genome Sequencing Data. *Research On Animal Production*, 15(2), 32–43. <https://doi.org/10.61186/rap.15.2.32>
- Mooney, H., Zavaleta, E., & Chapin, M. C. (Eds.). (2016). *Ecosystems of California* (1st ed.). University of California Press. <https://www.jstor.org/stable/10.1525/j.ctv1xxzp6>
- Muto, A., Ikeda, S., Lopez-Burks, M. E., Kikuchi, Y., Calof, A. L., Lander, A. D., & Schilling, T. F. (2014). Nipbl and Mediator Cooperatively Regulate Gene Expression to Control Limb Development. *PLOS Genetics*, 10(9), e1004671. <https://doi.org/10.1371/journal.pgen.1004671>
- Nan, J., Yang, S., Zhang, X., Leng, T., Zhuoma, J., Zhuoma, R., Yuan, J., Pi, J., Sheng, Z., & Li, S. (2023). Identification of candidate genes related to highland adaptation from multiple Chinese local chicken breeds by whole genome sequencing analysis. *Animal Genetics*, 54(1), 55–67. <https://doi.org/10.1111/age.13268>
- Owen, J. C. (2011). Collecting, processing, and storing avian blood: A review. *Journal of Field Ornithology*, 82(4), 339–354. <https://doi.org/10.1111/j.1557-9263.2011.00338.x>
- Pap, P. L., Osváth, G., Daubner, T., Nord, A., & Vincze, O. (2020). Down feather morphology reflects adaptation to habitat and thermal conditions across the avian phylogeny. *Evolution*, 74(10), 2365–2376. <https://doi.org/10.1111/evo.14075>
- Pap, P. L., Vincze, O., Wekerle, B., Daubner, T., Vágási, C. I., Nudds, R. L., Dyke, G. J., &

Osváth, G. (2017). A phylogenetic comparative analysis reveals correlations between body feather structure and habitat. *Functional Ecology*, 31(6), 1241–1251. <https://doi.org/10.1111/1365-2435.12820>

Park, H.-B., Kim, J.-W., & Baek, K.-H. (2020). Regulation of Wnt Signaling through Ubiquitination and Deubiquitination in Cancers. *International Journal of Molecular Sciences*, 21(11), Article 11. <https://doi.org/10.3390/ijms21113904>

Poplin, R., Ruano-Rubio, V., DePristo, M. A., Fennell, T. J., Carneiro, M. O., Auwera, G. A. V. der, Kling, D. E., Gauthier, L. D., Levy-Moonshine, A., Roazen, D., Shakir, K., Thibault, J., Chandran, S., Whelan, C., Lek, M., Gabriel, S., Daly, M. J., Neale, B., MacArthur, D. G., & Banks, E. (2018). *Scaling accurate genetic variant discovery to tens of thousands of samples* (p. 201178). bioRxiv. <https://doi.org/10.1101/201178>

Pritchard, J. K., & Di Rienzo, A. (2010). Adaptation – not by sweeps alone. *Nature Reviews Genetics*, 11(10), 665–667. <https://doi.org/10.1038/nrg2880>

Prost, S., Armstrong, E. E., Nylander, J., Thomas, G. W. C., Suh, A., Petersen, B., Dalen, L., Benz, B. W., Blom, M. P. K., Palkopoulou, E., Ericson, P. G. P., & Irestedt, M. (2018). *Genome analyses show strong selection on coloration, morphological and behavioral phenotypes in birds-of-paradise* (p. 287086). bioRxiv. <https://doi.org/10.1101/287086>

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *American Journal of Human Genetics*, 81(3), 559–575.

Qi, X., Zhang, Q., He, Y., Yang, L., Zhang, X., Shi, P., Yang, L., Liu, Z., Zhang, F., Liu, F., Liu, S., Wu, T., Cui, C., Ouzhuluobu, Bai, C., Baimakangzhuo, Han, J., Zhao, S., Liang, C., & Su, B. (2019). The Transcriptomic Landscape of Yaks Reveals Molecular Pathways

for High Altitude Adaptation. *Genome Biology and Evolution*, 11(1), 72–85.

<https://doi.org/10.1093/gbe/evy264>

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*, 26(6), 841–842.

<https://doi.org/10.1093/bioinformatics/btq033>

Ricchetti, I., Taucher, T., Loebick, R., Sin, S. Y. W., Palacios, C., & Lamichhaney, S. (2024).

Gene expression variation in geographically diverse populations of two North American songbird species. *Frontiers in Bird Science*, 3.

<https://doi.org/10.3389/fbirs.2024.1382657>

Rogalla, S., Shawkey, M. D., & D’Alba, L. (2022). Thermal effects of plumage coloration.

Ibis, 164(4), 933–948. <https://doi.org/10.1111/ibi.13100>

Romano, A., Séchaud, R., Hirzel, A. H., & Roulin, A. (2019). Climate-driven convergent evolution of plumage colour in a cosmopolitan bird. *Global Ecology and*

Biogeography, 28(4), 496–507. <https://doi.org/10.1111/geb.12870>

Romanov, M. N., Abdelmanova, A. S., Fisinin, V. I., Gladyr, E. A., Volkova, N. A., Koshkina,

O. A., Rodionov, A. N., Vetokh, A. N., Gusev, I. V., Anshakov, D. V., Stanishevskaya,

O. I., Dotsev, A. V., Griffin, D. K., & Zinovieva, N. A. (2023). Selective footprints and

genes relevant to cold adaptation and other phenotypic traits are unscrambled in

the genomes of divergently selected chicken breeds. *Journal of Animal Science and*

Biotechnology, 14(1), 35. <https://doi.org/10.1186/s40104-022-00813-0>

Rundel, P. W., Gibson, A. C., & Sharifi, M. R. (2008). THE ALPINE FLORA OF THE WHITE

MOUNTAINS, CALIFORNIA. *Madroño*, 55(3), 202–215. <https://doi.org/10.3120/0024-9637-55.3.202>

Rundel, P. W., & Millar, C. I. (2016). TWENTY-NINE. Alpine Ecosystems. In H. Mooney & E.

Zavaleta (Eds.), *Ecosystems of California* (pp. 613–634). University of California

Press. <https://www.degruyterbrill.com/document/doi/10.1525/9780520962170-033/pdf?licenseType=restricted>

Sadeesh, E. M., Lahamge, M. S., Malik, A., & Ampadi, A. N. (2025). Nuclear Genome-Encoded Mitochondrial OXPHOS Complex I Genes in Female Buffalo Show Tissue-Specific Differences. *Molecular Biotechnology*, 67(6), 2411–2427. <https://doi.org/10.1007/s12033-024-01206-6>

Sandoval, L., & Barrantes, G. (2019). Is black plumage an adaptation to high elevations in a cosmopolitan bird genus? *Journal of Avian Biology*, 50(7). <https://doi.org/10.1111/jav.02041>

Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation. *Nature Reviews Genetics*, 14(11), 807–820. <https://doi.org/10.1038/nrg3522>

Schweizer, R. M., Jones, M. R., Bradburd, G. S., Storz, J. F., Senner, N. R., Wolf, C., & Cheviron, Z. A. (2021a). Broad Concordance in the Spatial Distribution of Adaptive and Neutral Genetic Variation across an Elevational Gradient in Deer Mice. *Molecular Biology and Evolution*, 38(10), 4286–4300. <https://doi.org/10.1093/molbev/msab161>

Sebastianelli, M., Lukhele, S. M., Secomandi, S., de Souza, S. G., Haase, B., Moysi, M., Nikiforou, C., Hutfluss, A., Mountcastle, J., Balacco, J., Pelan, S., Chow, W., Fedrigo, O., Downs, C. T., Monadjem, A., Dingemanse, N. J., Jarvis, E. D., Brelsford, A., vonHoldt, B. M., & Kirschel, A. N. G. (2024). A genomic basis of vocal rhythm in birds. *Nature Communications*, 15(1), 3095. <https://doi.org/10.1038/s41467-024-47305-5>

Seki, R., Li, C., Fang, Q., Hayashi, S., Egawa, S., Hu, J., Xu, L., Pan, H., Kondo, M., Sato, T., Matsubara, H., Kamiyama, N., Kitajima, K., Saito, D., Liu, Y., Gilbert, M. T. P., Zhou, Q., Xu, X., Shiroishi, T., ... Zhang, G. (2017). Functional roles of Aves class-specific

cis-regulatory elements on macroevolution of bird-specific features. *Nature Communications*, 8(1), 14229. <https://doi.org/10.1038/ncomms14229>

Semenov, G. A., Sonnenberg, B. R., Branch, C. L., Heinen, V. K., Welklin, J. F., Padula, S. R., Patel, A. M., Bridge, E. S., Pravosudov, V. V., & Taylor, S. A. (2024). Genes and gene networks underlying spatial cognition in food-caching chickadees. *Current Biology*, 34(9), 1930–1939.e4. <https://doi.org/10.1016/j.cub.2024.03.058>

Seutin, G., White, B. N., & Boag, P. T. (1991). Preservation of avian blood and tissue samples for DNA analyses. *Canadian Journal of Zoology*, 69(1), 82–90. <https://doi.org/10.1139/z91-013>

Shafer, A. B. A., Wolf, J. B. W., Alves, P. C., Bergström, L., Bruford, M. W., Brännström, I., Colling, G., Dalén, L., De Meester, L., Ekblom, R., Fawcett, K. D., Fior, S., Hajibabaei, M., Hill, J. A., Hoezel, A. R., Höglund, J., Jensen, E. L., Krause, J., Kristensen, T. N., ... Zieliński, P. (2015). Genomics and the challenging translation into conservation practice. *Trends in Ecology & Evolution*, 30(2), 78–87. <https://doi.org/10.1016/j.tree.2014.11.009>

Shakya, S. B. (2020). *Genomics and Population History of Black-Headed Bulbul (Brachypodius Atriceps) Color Morphs* [Ph.D., Louisiana State University and Agricultural & Mechanical College]. <https://www.proquest.com/docview/2665131551/abstract/2200EDB181A4591PQ/>

1

Sirkiä, P. M., Virolainen, M., & Laaksonen, T. (2010). Melanin coloration has temperature-dependent effects on breeding performance that may maintain phenotypic variation in a passerine bird. *Journal of Evolutionary Biology*, 23(11), 2385–2396. <https://doi.org/10.1111/j.1420-9101.2010.02100.x>

Stettenheim, P. R. (2000). The Integumentary Morphology of Modern Birds—An Overview1.

American Zoologist, 40(4), 461–477. <https://doi.org/10.1093/icb/40.4.461>

Steven, S. (2021). *Genome-Wide Analyses of Genomic Variation Regulating Differential Stress Responses in Japanese Quail* [Doctoral dissertation, University of Arkansas]. ProQuest Dissertations & Theses.

Stinchcombe, J. R., & Hoekstra, H. E. (2008). Combining population genomics and quantitative genetics: Finding the genes underlying ecologically important traits. *Heredity*, 100(2), 158–170. <https://doi.org/10.1038/sj.hdy.6800937>

Stoutjesdijk, F. (2003). Birds in the cold: The effects of plumage structure and environment on operative temperature, shown by spherical models. *Ornis Svecica*, 13(4), Article 4. <https://doi.org/10.34080/os.v13.22804>

Struk, A. A., Mugon, J., Huston, A., Scholer, A. A., Stadler, G., Higgins, E. T., Sokolowski, M. B., & Danckert, J. (2019). Self-regulation and the foraging gene (PRKG1) in humans. *Proceedings of the National Academy of Sciences*, 116(10), 4434–4439. <https://doi.org/10.1073/pnas.1809924116>

Sun, L., Zhou, T., Wan, Q.-H., & Fang, S.-G. (2020). Transcriptome Comparison Reveals Key Components of Nuptial Plumage Coloration in Crested Ibis. *Biomolecules*, 10(6), Article 6. <https://doi.org/10.3390/biom10060905>

Sun, Y., Li, Y., Jiang, X., Wu, Q., Lin, R., Chen, H., Zhang, M., Zeng, T., Tian, Y., Xu, E., Zhang, Y., & Lu, L. (2024). Genome-wide association study identified candidate genes for egg production traits in the Longyan Shan-ma duck. *Poultry Science*, 103(9), 104032. <https://doi.org/10.1016/j.psj.2024.104032>

Symonds, M. R. E., & Tattersall, G. J. (2010). Geographical variation in bill size across bird species provides evidence for Allen’s rule. *The American Naturalist*, 176(2), 188–197. <https://doi.org/10.1086/653666>

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A. L.,

- Fang, T., Doncheva, N. T., Pyysalo, S., Bork, P., Jensen, L. J., & von Mering, C. (2023). The STRING database in 2023: Protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Research*, 51(D1), D638–D646. <https://doi.org/10.1093/nar/gkac1000>
- Tam, V., Patel, N., Turcotte, M., Bossé, Y., Paré, G., & Meyre, D. (2019). Benefits and limitations of genome-wide association studies. *Nature Reviews Genetics*, 20(8), 467–484. <https://doi.org/10.1038/s41576-019-0127-1>
- Tattersall, G. J., Arnaout, B., & Symonds, M. R. E. (2017). The evolution of the avian bill as a thermoregulatory organ. *Biological Reviews*, 92(3), 1630–1656. <https://doi.org/10.1111/brv.12299>
- Thomas, P. D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P., & Mi, H. (2022). PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Science*, 31(1), 8–22. <https://doi.org/10.1002/pro.4218>
- Tian, S., Zhou, X., Phuntsok, T., Zhao, N., Zhang, D., Ning, C., Li, D., & Zhao, H. (2020). Genomic Analyses Reveal Genetic Adaptations to Tropical Climates in Chickens. *iScience*, 23(11). <https://doi.org/10.1016/j.isci.2020.101644>
- Twining, H. (1940). Foraging Behavior and Survival in the Sierra Nevada Rosy Finch. *The Condor*, 42(1), 64–72. <https://doi.org/10.2307/1364320>
- Twumasi, G., Wang, H., Xi, Y., Qi, J., Li, L., Bai, L., & Liu, H. (2024). Genome-Wide Association Studies Reveal Candidate Genes Associated with Pigmentation Patterns of Single Feathers of Tianfu Nonghua Ducks. *Animals*, 14(1), Article 1. <https://doi.org/10.3390/ani14010085>
- Urban, M. C. (2018). Escalator to extinction. *Proceedings of the National Academy of Sciences*, 115(47), 11871–11873. <https://doi.org/10.1073/pnas.1817416115>
- Videla Rodriguez, E. A., Mitchell, J. B. O., & Smith, V. A. (2022). A Bayesian network

structure learning approach to identify genes associated with stress in spleens of chickens. *Scientific Reports*, 12(1), 7482. <https://doi.org/10.1038/s41598-022-11633-7>

Wang, C., Li, S., Yu, W., Xin, Q., Li, C., Feng, Y., Peng, X., & Gong, Y. (2011). Cloning and expression profiling of the VLDLR gene associated with egg performance in duck (*Anas platyrhynchos*). *Genetics Selection Evolution*, 43(1), 29. <https://doi.org/10.1186/1297-9686-43-29>

Ward, S., Rayner, J. M., MOLLer, U., Jackson, D. M., Nachtigall, W., & Speakman, J. R. (1999). Heat transfer from starlings *sturnus vulgaris* during flight. *The Journal of Experimental Biology*, 202 (Pt 12), 1589–1602. <https://doi.org/10.1242/jeb.202.12.1589>

Weeks, B. C., Harvey, C., Tobias, J. A., Sheard, C., Zhou, Z., & Fouhey, D. F. (2025). Longer Wing Bones in Warmer Climates Suggest a Role of Thermoregulation in Bird Wing Evolution. *Global Ecology and Biogeography*, 34(4), e70033. <https://doi.org/10.1111/geb.70033>

Weeks, B. C., Willard, D. E., Zimova, M., Ellis, A. A., Witynski, M. L., Hennen, M., & Winger, B. M. (2020). Shared morphological consequences of global warming in North American migratory birds. *Ecology Letters*, 23(2), 316–325. <https://doi.org/10.1111/ele.13434>

Whitlock, M. C., & Lotterhos, K. E. (2015). Reliable Detection of Loci Responsible for Local Adaptation: Inference of a Null Model through Trimming the Distribution of F_{ST} . *The American Naturalist*, 186 Suppl 1, S24–36. <https://doi.org/10.1086/682949>

Wolf, B. O., & Walsberg, G. E. (2000). The Role of the Plumage in Heat Transfer Processes of Birds1. *American Zoologist*, 40(4), 575–584. <https://doi.org/10.1093/icb/40.4.575>

Wu, Y., Yan, Y., Zhao, Y., Gu, L., Wang, S., & Johnson, D. H. (2021). Genomic bases

- underlying the adaptive radiation of core landbirds. *BMC Ecology and Evolution*, 21(1), 162. <https://doi.org/10.1186/s12862-021-01888-5>
- Xiao, J., Wu, Y., Chen, R., Lin, Y., Wu, L., Tian, W., & Liu, L. (2008). Expression of Pcp4 gene during osteogenic differentiation of bone marrow mesenchymal stem cells in vitro. *Molecular and Cellular Biochemistry*, 309(1), 143–150. <https://doi.org/10.1007/s11010-007-9652-x>
- Xie, J., Tang, L., Lu, L., Zhang, L., Xi, L., Liu, H.-C., Odle, J., & Luo, X. (2014). Differential Expression of Heat Shock Transcription Factors and Heat Shock Proteins after Acute and Chronic Heat Stress in Laying Chickens (*Gallus gallus*). *PLOS ONE*, 9(7), e102204. <https://doi.org/10.1371/journal.pone.0102204>
- Xiong, D.-H., Liu, X.-G., Guo, Y.-F., Tan, L.-J., Wang, L., Sha, B.-Y., Tang, Z.-H., Pan, F., Yang, T.-L., Chen, X.-D., Lei, S.-F., Yerges, L. M., Zhu, X.-Z., Wheeler, V. W., Patrick, A. L., Bunker, C. H., Guo, Y., Yan, H., Pei, Y.-F., ... Deng, H.-W. (2009). Genome-wide Association and Follow-Up Replication Studies Identified ADAMTS18 and TGFBR3 as Bone Mass Candidate Genes in Different Ethnic Groups. *The American Journal of Human Genetics*, 84(3), 388–398. <https://doi.org/10.1016/j.ajhg.2009.01.025>
- Yan, C., Zhang, Z.-Y., Lv, Y., Wang, Z., Jiang, K., & Li, J.-T. (2022). Genome of *Laudakia sacra* Provides New Insights into High-Altitude Adaptation of Ectotherms. *International Journal of Molecular Sciences*, 23(17), Article 17. <https://doi.org/10.3390/ijms231710081>
- Yeaman, S. (2015). Local Adaptation by Alleles of Small Effect. *The American Naturalist*, 186 Suppl 1, S74-89. <https://doi.org/10.1086/682405>
- Zhang, C., Liu, C., Zhang, J., Zheng, L., Chang, Q., Cui, Z., & Liu, S. (2022). Analysis on the desert adaptability of indigenous sheep in the southern edge of Taklimakan Desert. *Scientific Reports*, 12(1), 12264. <https://doi.org/10.1038/s41598-022-15986-x>

Zhang, W., Fan, Z., Han, E., Hou, R., Zhang, L., Galaverni, M., Huang, J., Liu, H., Silva, P., Li, P., Pollinger, J. P., Du, L., Zhang, X., Yue, B., Wayne, R. K., & Zhang, Z. (2014). Hypoxia Adaptations in the Grey Wolf (*Canis lupus chanco*) from Qinghai-Tibet Plateau. *PLOS Genetics*, 10(7), e1004466. <https://doi.org/10.1371/journal.pgen.1004466>

Zhang, X., Yang, F., Zhu, T., Zhao, X., Zhang, J., Wen, J., Zhang, Y., Wang, G., Ren, X., Chen, A., Wang, X., Wang, L., Lv, X., Yang, W., Qu, C., Wang, H., Ning, Z., & Qu, L. (2024). Whole genome resequencing reveals genomic regions related to red plumage in ducks. *Poultry Science*, 103(6), 103694. <https://doi.org/10.1016/j.psj.2024.103694>

Zhang, X., Zhu, T., Wang, L., Lv, X., Yang, W., Qu, C., Li, H., Wang, H., Ning, Z., & Qu, L. (2023). Genome-Wide Association Study Reveals the Genetic Basis of Duck Plumage Colors. *Genes*, 14(4), Article 4. <https://doi.org/10.3390/genes14040856>

Zheng, S., Ouyang, J., Liu, S., Tang, H., Xiong, Y., Yan, X., & Chen, H. (2023). Genomic signatures reveal selection in Lingxian white goose. *Poultry Science*, 102(1), 102269. <https://doi.org/10.1016/j.psj.2022.102269>

Zhou, X., Carbonetto, P., & Stephens, M. (2013). Polygenic Modeling with Bayesian Sparse Linear Mixed Models. *PLOS Genetics*, 9(2), e1003264. <https://doi.org/10.1371/journal.pgen.1003264>

Zhuang, Z.-X., Chen, S.-E., Chen, C.-F., Lin, E.-C., & Huang, S.-Y. (2019). Genome-wide association study on the body temperature changes of a broiler-type strain Taiwan country chickens under acute heat stress. *Journal of Thermal Biology*, 82, 33–42. <https://doi.org/10.1016/j.jtherbio.2019.03.007>

Data Accessibility

Raw sequence data have been deposited in the NCBI Sequence Read Archive under BioProject accession PRJNA1297057.

Statistical analysis, GWAS and F_{ST} results are available at <http://datadryad.org/share/DqvYM4bZd7lZHExY5O9FGkrQ86Q90YoGz3ZOAJoLHmY>

Code and scripts used for processing and analysis as well as sample metadata are available on GitHub at https://github.com/ecnrobertson/GCRF_Pub

Benefit-Sharing Statement

Benefits Generated: Benefits from this research accrue from the sharing of our data and results on public databases as described above.

Statement of Contributions

E.C.N.R, T.B., E.Z., K.C.R conceived of the study and E.C.N.R. wrote the majority of the manuscript and oversaw the majority of the analysis. C.M.B. contributed to the bioinformatic analysis. S.D. performed and conceived of the methodology for all feather morphology measurements. T.B. and E.Z. contributed ideas and performed all field sampling. K.C.R., E.Z. and M.B.H. acquired funding. All authors discussed the results and contributed to the final manuscript.

Figures




Figure 1. A) A map of California

highlighting the sampling location and design for two populations of Grey-crowned Rosy-Finch, Piute Pass and White Mountain. Key environmental variables at each site are noted: elevation (ELEV) and mean temperature of the warmest month (MTWM). B) Overview of methods for GWAS analysis. Morphological measurements were corrected for the effects of sex, age, and body size before being input into BSLMM and LMM models. The output was analyzed to identify significant SNPs and related genes.

Body Morphology Results




Figure 2. Violin plots of beak morphology raw measurement: A) beak width, B) nare length, and C) beak depth. Overlaid are the corrected mean trait measurements shown with a crossbar. Significance of the effect of site on the morphological trait is shown for each comparison. Piute Pass is the warm site and White Mountain is the cold site. We found that nares were longer while beak widths and depths were smaller in the warmer environment.






Figure 3. A) Photograph showing a body feather from a Sierra Nevada Grey-crowned Rosy-Finch captured at White Mountain. Highlighted is a simplification of the plumulaceous barbule structure. B) Violin plots of node density raw measurements, with illustrations above of the structure of plumulaceous barbule nodes. Overlaid on the plots are the corrected mean trait measurements shown with a crossbar. Significance of the effect of site on the morphological trait is shown for each comparison.

Figure 4. Manhattan plot depicting the results of the LMM analysis on beak depth (A), wing chord (B), and pennaceous barbule length with $-\log_{10}(\text{p-value})$ on the y-axis and the mapped Zebra Finch chromosome location on the x-axis. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted. Also plotted are the results of the Per-site F_{st} results (C), using the 90th quantile threshold of 0.235. Corresponding genes are also shown and functional categories are highlighted by color category. See Table 1 for more




functional details.





B GWAS Methods




C F_{ST} Methods




Body Morphology Results






1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

B Plumulaceous Barbule Node Density




A

Wing Chord GWAS




B

Beak Depth GWAS




C

Pennaceous Barbule Length GWAS



D

Per-Site F_{ST}



Gene Function Categories



Table 1: Summary of F_{ST} Outlier Results. Show are the genes for which function information were available, from the narrower (90th quantile) F_{ST} results. Gene functions are color coded following the key in Figure 4.

<i>F_{ST} outliers</i>		
Gene	Function	Reference
GDPD5	Lipid Metabolism	Feng et al 2023
NOX4	Heat Stress, Altitude Adaptation	Goel et al 2021; Diebold et al 2010
MAML2	Altitude Adaptation	Schweizer et al 2021
SUPT20H	Altitude Adaptation	Schweizer et al 2021
PPP1R1C	Cold Adaptation	Federova et al 2022
SOX14	Feather Development	Chen et al 2024
MEGF11	Feather Development	Twumasi et al 2024
MSRA	Altitude Adaptation	Jeong et al 2018; Gheyas et al 2021
AKT3	Altitude Adaptation	Buroker et al 2012; Qi et al 2018
DOCK11	Feather Development	Crates et al 2024
SMARCA1	Craniofacial Development	Boer et al 2021
GAB3	Craniofacial Development	Boer et al 2021
ADGRD1	Altitude Adaptation	Nan et al 2021
NPHP4	Vision	Borges et al 2019
ALDH7A1	Beak Development	Bai et al 2014
ADAMTS19	Metabolism	Zhang et al 2022
VLDLR	Lipid Protein	Wang et al 2011
ERCC8	Altitude Adaptation	Yan et al 2022
CTIF	Beak Development	Lawson et al 2017

Table 1: Summary of GWAS Results by Trait. Show are the genes for which function information is available. Gene functions are color coded following the key in Figure 4.

Gene	Model	Function	Details	Reference
<i>Wing Chord</i>				
PCP4	BSLMM	Development, Stress Response	Bone development, stress response in quail	Xiao et al. 2022
GRIK2	BSLMM	Reproduction	Egg quality and production in goose and duck	Gao et al. 2022
AKAP7	BSLMM	Stress Response	Stress response in quail	Stevenson et al. 2022
PACRG	BSLMM	Heat Stress	Heat stress response in chickens	Tian et al. 2022
HAPLN1	BSLMM	Coloration	Plumage coloration in ducks	Zhang et al. 2022
NEGR1	BSLMM	Development, Coloration	Neural development, plumage coloration in crested ibis	Sun et al. 2022
PGLYRP2	BSLMM/LMM	Immunity, Heat Stress	Immune response in vertebrates, heat stress response in chickens	Hafizi et al. 2022
GUCA1C	LMM	Morphology	Visual systems in birds, bone quality in ducks, wing deformity in chickens	Kim et al. 2022
RFX3	LMM	Hypoxia Adaptation	Altitude adaptation in wolves	Wu et al. 2022

(Continued from previous page)					
Gene	Model	Function	Details	Ref.	
NIPBL	LMM	Development, DNA Repair	Developmental regulation and DNA repair	Ba	
CHD1	LMM	Gene Regulation	Modulates gene expression in birds	Fri 200	
NDUFA11	LMM	Metabolism, Respiration	Respiratory system support in buffalo, mitochondrial activity in songbirds	Clas Sae 202	
<i>Beak Depth</i>					
TENM4	BSLMM	Development, Coloration, Reproduction	Neural development in quail, dorsal feather coloration in duck, breeding timing in great tit	Tw 202 202 et al	
RASL11A	BSLMM	Morphology	Craniofacial development in buteo	Ab	
TINAG	BSLMM	Morphology	Beak development in chicken	Ba	
LAMA4	BSLMM	Migration	Migration timing in Swainson's thrush	Job 201	
LURAP1L	BSLMM	Coloration	Pigmentation in mallards	Ma	
PTPRD	BSLMM	Communication, Coloration	Vocal signaling in flycatchers, pigmentation in mallards	Ga 202 202	
CLRN3	BSLMM	Reproduction, Sex Differentiation	Sex differentiation in chicken	Lo	
CPEB3	BSLMM	Reproduction	Oocyte meiosis regulator in chicken	Ch	

(Continued from previous page)					
Gene	Model	Function	Details	Ref	
BICC1	BSLMM	Morphology, Body Size	Bone development in chicken and mice, body size in chicken and hermit thrush	Kin 2018	
ANK3	BSLMM	Migration	Migration in birds	son 2018	
PRKG1	BSLMM	Reproduction, Behavior, Coloration	Ovarian function in duck, foraging behavior in humans, signal transduction and feather pigmentation in ducks	Me 2018	
OTUD7A	BSLMM	Development	Neural development and adaptation to urban environments in bananaquit	Len 2018	
ERCC8	BSLMM	Hypoxia Adaptation	Altitude adaptation in lizard	Sur 2018	
MEGF11	BSLMM	Coloration	Dorsal feather coloration in duck	Str 2018	
DLG1	BSLMM	Development, Regulation	Embryo development in mice, regulatory development in birds	Tw 2018	
<i>Nare Length</i>					
NRIP1	BSLMM	Reproduction, Adaptation, Metabolism	Female reproduction in goose, nocturnal adaptation in owls, mitochondrial biosynthesis in vertebrates	Sek 2018	
<i>Zhang et al 2018</i>					

4

(Continued from previous page)				
Gene	Model	Function	Details	Ref.
ATP1A1	BSLMM	Migration	Migration in birds	Ler...
TENM4	BSLMM	Development, Col- oration, Reproduction	Neural development in quail, dorsal feather coloration in duck, breeding timing in great tit	Tw... 202... 202... et ...
KALRN	BSLMM	Neurodevelopment	Cognitive ability in chickadee	Ric... 202...
CNTNAP5	BSLMM	Development, Mem- ory, Hypoxia Adapta- tion	Brain development and signaling in zebra finch, spatial memory in chickadees, hypoxia response in chickens	Gil... 202... et ... al ...
KCTD18	BSLMM	Immunity, Communi- cation	Immune response in vultures, vocal learning in chickens	Ch... 201... al ...
PLPP3	BSLMM	Morphology, Metabolism	Culmen length in buteo, formation of goose fatty liver	Ab... 202... 201...
DLG1	BSLMM	Development, Regula- tion	Embryo development in mice, regulatory de- velopment in birds	Sek...
ADAMTS18	BSLMM	Morphology, Bone Density	Bone mineral density in humans and chickens	Xie... Gu...
PSD2	BSLMM	Coloration, Stress Re- sponse	Melanin production in owls, stress response in quail	Du... 202... 202...

(Continued from previous page)				
Gene	Model	Function	Details	Ref.
BRAT1	BSLMM	Stress Response	Stress response in chickens	Ro 202
<i>Pennaceous Barbule Length</i>				
KCNF1	BSLMM	Coastal Adaptation	Coastal adaptation in song sparrow and frogs	Cl
CSMD1	BSLMM/LMM	Heat Stress, Hypoxia Adaptation	Heat adaptation in chickens, adaptation to altitude in humans	Ha De 202
NKAIN1	BSLMM	Immunity, Migration	Immune response in house finch, migration tendency in European blackcap	Ve De 202
KCNJ3	BSLMM	Reproduction	Egg production in duck	Su
AGBL1	BSLMM	Metabolism	Fatty acid composition in chicken	Ch
TENM1	BSLMM	Morphology	Craniofacial development in pigeons	Bo
<i>Plumulaceous Node Density</i>				
EFNA2	BSLMM	Heat Stress, Morphology, Coloration	Heat stress response in chickens, bone morphogenesis and remodeling in ground tit, color differences in bulbul	Zh 201 al et a
STK10	BSLMM	Morphology	Cranium and bill depth in buteo	Ab
<i>Plumulaceous Barbule Density</i>				
BORCS6	BSLMM	Metabolism	Meat quality in chickens	Als 202
HSF2	BSLMM	Heat Stress	Response to heat stress in chickens	Xie


(Continued from previous page)				
Gene	Model	Function	Details	Ref.
PRKN	BSLMM	Domestication	Domestication in birds	Fan et al. 2019
NRXN1	BSLMM	Neurodevelopment, Communication, Coloration	Neural connections, plumage coloration in crested ibis, vocal rhythm in birds, speech disorders in humans	Sun et al. 2020
ALDH1A1	BSLMM	Hypoxia Adaptation, Metabolism, Vision	Hypoxia adaptation and fat metabolism in rosy finch, eyesight in owl and mice	Baig et al. 2020
ADAMTS19	BSLMM	Morphology	Rumpless trait in chickens	Chen et al. 2020
HERC4	BSLMM	Coloration, Migration	Plumage color in duck, migration in European blackbirds	Zhang et al. 2020
SMC4	BSLMM	Development	Feather follicle development in goose	Hu et al. 2020
SPAG16	BSLMM	Reproduction, Coloration, Longevity	Sexual development in birds-of-paradise, plumage coloration in ducks, longevity in mammals	Probst et al. 2020

Supplemental Information for:

The genetic and morphological basis of local adaptation to elevational extremes in an alpine finch


Robertson, Erica C.N.¹, Brown, Timothy², Deitch, Sophie¹, Bossu, Christen M.¹,
Zavaleta, Erika S.², Hooten, Mevin B.³, Ruegg, Kristen C.¹

Population Structure Results




Sup Fig 1. PCA results showing a lack of population structure between the White Mountain (WMNT, blue) and Piute Pass (PIPA, red) populations. There are some outlier individuals but an ADMIXTURE analysis confirmed that 1 population is the most supported.


Extended GWAS Results




Sup Fig 2. Beak length GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.



Sup Fig 3. Beak width GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.



Sup Fig 4. Pennaceous barbule density GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.



Sup Fig 5. Tail length GWAS results. Significant SNPs from the BSLMM analysis are overlaid onto the LMM results, plotted with their corresponding p-values and corresponding genes are noted if available.