

Key Points:

- The interaction between the Tethyan domain and the Paleo-Pacific domain began in the Latest Triassic
- An inward dipping double subduction model can explain the interaction between the Tethyan and Paleo-Pacific domains
- Inward-dipping double subduction systems are more likely to produce juvenile crust compared to one-sided subduction systems

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

Y. Yan,
yanyi@gig.ac.cn

Citation:

Zhao, Q., Yan, Y., Carter, A., Clift, P. D., Wu, J., Asis, J. B., et al. (2025). Interaction between the Tethyan and Paleo-Pacific tectonic domains in Southeast Asia: Late Triassic initiation of an inward-dipping double subduction system. *Tectonics*, *44*, e2024TC008762. <https://doi.org/10.1029/2024TC008762>

Received 3 DEC 2024
Accepted 22 JUL 2025

Author Contributions:








Investigation: Qi Zhao, Junaidi Bin Asis, Shihao Gou

Writing – original draft: Qi Zhao

Writing – review & editing:

Andrew Carter, Peter D. Clift, Jonny Wu, Zuofei Zhu, Yang Zhou

Interaction Between the Tethyan and Paleo-Pacific Tectonic Domains in Southeast Asia: Late Triassic Initiation of an Inward-Dipping Double Subduction System

Qi Zhao¹ , Yi Yan¹ , Andrew Carter² , Peter D. Clift^{3,4} , Jonny Wu⁵ , Junaidi Bin Asis⁶ , Zuofei Zhu⁷, Shihao Gou¹, and Yang Zhou⁸ 

¹State Key Laboratory of Deep Earth Processes and Resources, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, China, ²School of Natural Sciences, Birkbeck, University of London, London, UK, ³Department of Earth Sciences, University College London, London, UK, ⁴Institute of Marine and Environmental Sciences, University of Szczecin, Szczecin, Poland, ⁵Department of Geosciences, University of Arizona, Tucson, AZ, USA, ⁶Faculty of Science and Natural Resources, Kota Kinabalu, Malaysia, ⁷Sanya Institute of South China Sea Geology, Guangzhou Marine Geological Survey, Sanya, China, ⁸School of National Safety and Emergency Management, Beijing Normal University, Zhuhai, China

Abstract Interaction between Tethys and the Paleo-Pacific subduction zones in Southeast Asia during the Mesozoic remains poorly understood. Using new and published zircon U-Pb and Hf data sets from Borneo (Paleo-Pacific domain) and Sumatra (Tethyan domain), we propose that isotopically juvenile magmatism was active on both sides of Sundaland due to the initiation of inward-dipping double subduction during the latest Triassic when Indochina collided with Sibumasu, as evidenced by a pronounced positive shift in zircon $\epsilon\text{Hf}(t)$ values from both Cenozoic sedimentary successions and Mesozoic magmatic rocks in Sumatra and Borneo. From the latest Triassic to Cretaceous, the contrasting positive $\epsilon\text{Hf}(t)$ values ranges between Borneo and Sumatra, with Borneo showing a broad range and Sumatra a narrower variability, imply that the inward-dipping double subduction system evolved asymmetrically due to differences in slab dip angles between the subducting Meso-Tethys and Paleo-Pacific oceanic lithosphere. After 80 Ma, this asymmetric double subduction system was disrupted, marked by the complete cessation of arc magmatism in Borneo while isotopically juvenile magmatism continued on Sumatra. Our findings emphasize that, when compared to the contemporary single-sided subduction system of the western Meso-Tethyan domain and the northern Paleo-Pacific domain, SE Asia developed more juvenile crust due to large-scale upper plate extension driven by inward-dipping double subduction.

Plain Language Summary The interaction between the Tethyan and Paleo-Pacific domains has shaped the formation of the present megacontinent Eurasia. However, how the two subduction zones (Tethyan and Paleo-Pacific domains) interacted and evolved, and how they impacted on the geology of Southeast Asia is not yet clear. Using radiometric ages and hafnium isotopes from magmatic and detrital zircon, we found that the interaction between the Tethyan and Paleo-Pacific domains in Southeast Asia can be explained by an inward-dipping double subduction model that began in the Late Triassic. Furthermore, our results highlight that the strengthened juvenile crustal signatures in Southeast Asia because of the large-scale extension of upper plate in an inward-dipping double subduction system.

1. Introduction

The last supercontinent cycle was characterized by the breakup of Pangea, a process that began in the Triassic (De Min et al., 2020; Golonka, 2007; Olsen, 1997), leading to the formation of the present megacontinent Eurasia, regarded as a potential precursor to the future supercontinent Amasia (Wang et al., 2020). The assembly of Eurasia was driven by broadly E-W-trending Tethyan-style collisional orogens (Tethyan tectonic domain) and the approximately N-S-trending circum-Pacific-style accretionary orogens (Paleo-Pacific tectonic domain). These trends converge in Southeast Asia, influencing both mantle convection patterns and climate (Li et al., 2019; Metcalfe, 2011; Nance et al., 2014; Wang et al., 2020, 2024).

The post Triassic evolution of western Tethyan and the northern Paleo-Pacific tectonic domains have been extensively studied. The western Tethyan tectonic domains are characterized by closure of the Devonian-Triassic Paleo-Tethys in the Late Triassic (Fan et al., 2024; Hu et al., 2014), as well as the subduction and demise of the




Figure 1. (a) Simplified tectonic map illustrating the Tethys tectonic domains and the Pacific tectonic domains, along with the temporal evolution of the ocean basins (refer to Metcalfe, 2021; Zhou et al., 2023 and reference therein). (b) Location of compiled Mesozoic magmatic and detrital zircons, as well as the new detrital zircons in the interaction zone. Further details are provided in Table S3 of Supporting Information S1.

Permian–Cretaceous Meso-Tethys and Jurassic–Cretaceous Ceno-Tethys, between the Triassic and Paleogene (Hall, 2012; Metcalfe, 2013). These developments were accompanied by the accretion to Eurasia of continental blocks derived from the northern margin of Gondwana (e.g., the South Qiangtang and Lhasa terranes; Figure 1a; Metcalfe, 2017, 2021). In the Phanerozoic the Paleo-Pacific plate has been subducting along the East Asian continental margin in various stages, although the timing of its initial subduction remains debated. For South China, proposed subduction initiation of Paleo-Pacific plate spans from 500 Ma to the Permian or Triassic (Figure 1a; Isozaki et al., 2010; Li & Li, 2007; Pastor-Galán et al., 2021; Zhu et al., 2013; Zhou et al., 2023), while the North China Craton records Early Jurassic onset of Paleo-Pacific plate subduction (Qiu, Kong et al., 2022; Qiu, Li et al., 2022; Wu et al., 2019; Zhu & Xu, 2019).

However, the interaction and evolution of these two subduction zones (Tethyan and Paleo-Pacific domains), as well as their impact on the geology of Southeast Asia, remain unclear. Present-day West Sumatra and West and Southwest Borneo, which are separated from the Malay Peninsula by the Sunda Shelf, and the Bangka and

Belitung islands, represent the easternmost extent of the Tethyan domain (Figure 1b; Li et al., 2020; Metcalfe, 2017, 2021; Zhang et al., 2019) and the southernmost extent of the Paleo-Pacific domain (Breitfeld, Davies, et al., 2020; Breitfeld, Hennig-Breitfeld et al., 2020; Hennig et al., 2017; Metcalfe, 1996; Wang, Liu, et al., 2022; Wang et al., 2023, 2024). Two principal models have been proposed regarding the origin of Southwest Borneo: (a) It represents an allochthonous terrane that accreted to SE Asia during the Late Jurassic or Early Cretaceous (Breitfeld, Davies, et al., 2020; Breitfeld, Hennig-Breitfeld et al., 2020; Hall, 2012; Hennig et al., 2017; Metcalfe, 2009), or (b) it was autochthonous, undergoing accretion linked to Paleo-Pacific plate subduction since the Earliest Jurassic or possibly earlier (Wang, Liu, et al., 2022; Wang et al., 2023, 2024). Nevertheless, Southwest Borneo—or at least its northwestern portion (the Northwest Schwaner zone, termed West Borneo in Hennig et al., 2017)—was situated along the southeastern margin of the Sundaland continent since the Late Triassic. Tectonic reconstructions constrain accretion of the West Sumatra block onto the Sundaland margin to the Late Triassic, after which a Mesozoic subduction-related arc developed within a narrow belt throughout West Sumatra (Barber & Crow, 2009; Barber et al., 2005; Cobbing, 2005; Li et al., 2020; Metcalfe, 2021). Thus, the Mesozoic magmatic rocks in West Sumatra and West and/or Southwest Borneo record, respectively, continental accretion and growth related to the northeastward subduction of the Tethyan lithosphere (e.g., Li et al., 2020) and the westward subduction of the Paleo-Pacific plate (Breitfeld et al., 2017; Breitfeld, Davies et al., 2020; Hennig et al., 2017; Wang et al., 2024). These rocks provide an important record for understanding how these two subduction systems interacted.

In this study, we examined compiled regional magmatic and detrital zircon U-Pb data in the West and Southwest Borneo and West Sumatra, and integrated these results with new zircon U-Pb and Hf isotopic data obtained from samples collected across northern Borneo (Sabah). Zircon data from Mesozoic igneous rocks were used to track the spatial distribution of magmatism through time and Mesozoic detrital zircon ages from Upper Oligocene to Neogene sedimentary rocks capture temporal variations in magmatism across the sediment source regions. Collectively, these data sets capture regional temporal and spatial changes in Mesozoic magmatism providing fresh new insights into the tectono-magmatic evolution and underlying geodynamic processes across the interaction zone (Figure 1a) between the Tethyan and Pacific-Pacific domains.

2. Geological Background

Southeast Asia was formed by the successive assembly of different continental fragments that rifted from Gondwana since the Paleozoic (Metcalfe, 1988, 2017). During the Triassic, major continental blocks, such as Sibumasu, Indochina, West and SW Borneo were positioned between the Tethyan and Paleo-Pacific domains (Figure 1a; e.g., Metcalfe, 2017), herein referred to as the interaction zone of both. The origin of Southwest Borneo remains debated. It may represent either a continental terrane rifted from northwestern Australia (e.g., Breitfeld, Davies, et al., 2020; Hall, 2012; Hennig et al., 2017; Metcalfe, 2009), or a Paleo-Pacific active continental margin similar or attached to West Borneo (e.g., Wang, Liu, et al., 2022; Wang et al., 2023, 2024). In contrast, West Borneo (containing the Northwest Schwaner Zone) was unambiguously situated at the western and southernmost part of the Paleo-Pacific subduction zone in the Triassic, where most of the subduction-related Mesozoic igneous and metamorphic rocks were developed (Figure 1b; Breitfeld, Davies, et al., 2020; Haile et al., 1977; Hennig et al., 2017; Williams et al., 1988). Upper Mesozoic to Paleogene fluvio-deltaic sedimentary rocks and deep-marine turbidites are mainly developed in Sarawak and Sabah, respectively. Sabah, situated at the northern tip of Borneo, is thereby dominated mainly by Paleogene-Neogene clastic sedimentary rocks deposited on the Mesozoic Chert-Spilitic Formation/ophiolite basement (Hutchison, 2005). Cenozoic rifting of the Sunda Shelf caused by the propagating South China Sea, along with associated tectonic events in Borneo resulted in erosion and deposition of large volumes of Lower Oligocene to Neogene sediments in Sabah (Breitfeld, Hall et al., 2023; Breitfeld, Hennig-Breitfeld et al., 2023; Hutchison, 1996, 2005; Hall, 2013; Hall & Breitfeld, 2017). Although Early Cenozoic drainage reorganization in Borneo triggered dynamic sediment provenance fluctuations between the Malay-Thai Peninsula and West and Southwest Borneo (e.g., Breitfeld, Hennig-Breitfeld, et al., 2020; Galin et al., 2017; Hennig-Breitfeld et al., 2019), most Upper Oligocene to Neogene sedimentary rocks in Sabah are dominated by Cretaceous zircons and were mainly derived by multi-recycling from West and Southwest Borneo with minor inputs from inland Sundaland (e.g., Malay Peninsula) or mainland Asia (Breitfeld et al., 2017; Breitfeld, Hennig-Breitfeld, et al., 2020, 2023; Breitfeld, Hall et al., 2023; Galin et al., 2017; Hennig-Breitfeld et al., 2019; Quek, Lee, et al., 2021; van Hattum et al., 2006, 2013). As a result, these Cretaceous

zircons-dominant sedimentary successions represent an important archive of continental growth history and magmatism related to subduction of the southernmost Paleo-Pacific oceanic slab beneath Borneo in the Mesozoic.

By contrast, Sumatra lies at the easternmost Tethyan domain and was partly formed from accretion of the Woyla Arc, and the West and East Sumatra blocks in the Cretaceous (Advokaat et al., 2018; Figure 1b). Rocks exposed in Sumatra record subduction of Mesozoic Tethyan lithosphere following the closure of the Paleo-Tethyan Ocean in the Middle to Late Triassic (Barber & Crow, 2009; Metcalfe, 1996). The Woyla Arc is mainly characterized by a Lower Cretaceous volcanic-sedimentary succession, comprising basaltic to andesitic volcanic rocks interbedded with clastic sedimentary units. It is interpreted as an intra-oceanic subduction-related arc that accreted to Sundaland in the Middle-Late Cretaceous due to the absence of continental rocks (Barber, 2000; Barber et al., 2005; Metcalfe, 2017, 2021). West and East Sumatra are separated by the Medial Sumatra Tectonic Zone, characterized as a major crustal shear zone/strike-slip fault system rather than a suture zone, given the complete absence of exposed ophiolitic components (e.g., Barber & Crow, 2009; Metcalfe, 2013). West Sumatra was probably emplaced by the Medial Sumatra Tectonic Zone in the Late Triassic (Barber & Crow, 2009). Triassic-Jurassic and Cretaceous granitoids that represent a subduction-related arc magmatic suite are mainly distributed within a narrow belt in West Sumatra, representing the southeastern extension of the Meso-Tethyan arc system (Cobbing, 2005; Gasparon & Varne, 1995; Li et al., 2020; Zhang et al., 2019). Thus, detrital and magmatic zircon data in West Sumatra record the onset of crustal growth related to subduction of the easternmost Tethyan lithosphere after the Late Triassic (Li et al., 2020; Zhang et al., 2019).

3. Methods

3.1. Sampling and Analytical Methods

A total of five representative sandstones were collected from strata in Sabah (northern Borneo), which include the Cretaceous turbidite slice within the ophiolitic basement (TB-10B), Labang Formation (Late Oligocene, SK-06), Garinono Mélange (Early Miocene, SK-02), Kumaunt Formation (Early Miocene, K-01) and Tanjong Formation (Early-Middle Miocene, K-03). Zircon U-Pb-(Hf) isotopic analyses were conducted on these samples to constrain temporal variations in magmatism along the southernmost segment of the Paleo-Pacific subduction zone.

Zircon grains were separated using standard density and magnetic techniques after Mange and Maurer (1992). U-Pb dating was performed by laser ablation-inductively coupled plasma-mass spectroscopy (LA-ICP-MS) at Wuhan SampleSolution Analytical Technology Co., Ltd., Wuhan, China using the Agilent 7900 ICP-MS coupled with a GeolasPro laser ablation system. The diameter of the analytical spot is 32 μm . Zircon 91500 (1,065 Ma; Wiedenbeck et al., 1995) and glass NIST610 (Pearce et al., 1997) served as external standards for U-Pb dating and trace element calibration. Data is provided in Table S1 of Supporting Information S1.

From the 506 dated grains, 117 concordant zircons with ages younger than 500 Ma were selected for Lu-Hf isotopic analysis by LA-MC (multicollector) ICP-MS to track the Mesozoic tectono-magmatic evolution of Borneo. In-situ Lu-Hf isotopic measurements were performed at Wuhan SampleSolution Analytical Technology Co., Ltd employing a Neptune Plus MC-ICP-MS (Thermo Fisher Scientific, Germany) coupled with a Geolas HD excimer ArF laser ablation system (Coherent, Göttingen). The diameter of the ablation spot is 44 μm . Analyses followed established Hu et al. (2012) operational frameworks. Plešovice, 91500 and GJ-1 are analyzed simultaneously with our samples to ensure the reliability of our data (Zhang et al., 2020). Their external precision (2SD) is better than 0.000020, with measured values showing concordance with published reference values within analytical uncertainty. ICPMSDataCal was used to perform off-line selection and integration of analyzed signals and mass bias calibration (Liu et al., 2010). Data is provided in Table S2 of Supporting Information S1.

3.2. Data Compilation

To determine the interaction and evolution of the two subduction zones (Paleo-Pacific and Tethyan domains), we compiled Neogene detrital zircon U-Pb-(Hf) data from Sabah (Breitfeld, Hall, et al., 2023; Zhang et al., 2023) and Sumatra (Hsu, 2016; Zhang et al., 2019). Zircon U-Pb-(Hf) data of Mesozoic magmatic rocks from Sumatra (Li et al., 2020), West Borneo and Southwest Borneo (Breitfeld et al., 2017; Breitfeld, Davies et al., 2020; Gan et al., 2022; Hennig et al., 2017; Qian et al., 2022; Setiawan et al., 2013; van Hattum et al., 2013; Wang, Qian et al., 2021; Wang, Zhang et al., 2021; Wang, Liu et al., 2022; Wang, Wu et al., 2022), Indochina (Hieu et al., 2015; Hou et al., 2019; Minh et al., 2020; Nguyen et al., 2019; Shellnutt et al., 2013; Shi et al., 2015; Tran




Figure 2. Cathodoluminescent (CL) images of zircons from the Cretaceous turbidite slice and Upper Oligocene-Neogene sandstones sampled in this study.

et al., 2014; Wang et al., 2016), Malay Peninsula (Cao et al., 2020; Ghani et al., 2013; Gillespie et al., 2019; Hazad et al., 2019; Jamil et al., 2016; Liew & Page, 1985; Liu et al., 2020; Ng et al., 2015; Oliver et al., 2014; Qian et al., 2023; Quek, Lai, et al., 2021; Searle et al., 2012; Wang, Qian, et al., 2021; Yu et al., 2022), Bangka and Belitung Island (Ng et al., 2017; Wang, Qian, et al., 2021) were also compiled.

To ensure data reliability and comprehensiveness, all data were compiled directly from original studies, retaining the authors' own processing approaches—including outlier removal and concordance threshold selection. Compiled data from the Mesozoic magmatic rocks and the Neogene detrital samples comprising 7486 zircon U-Pb ages and 4224 Hf isotopic measurements. The locations of the new study samples and compiled samples are shown in Figure 1b. Compiled zircon age data and Hf isotopic data are shown in Table S3 and S4 of Supporting Information S1, respectively.

4. Results

All new zircon grains from the Cretaceous turbidite slice and Upper Oligocene-Neogene sandstones in Sabah display lengths of 60–200 μm, and axial ratios spanning 1:1 to 3:1 (Figure 2). Most of these detrital zircon data yield ages younger than 500 Ma (Table S1 in Supporting Information S1), comprising a total of 114 Paleozoic, 232 Mesozoic and 11 Cenozoic age. Grains exhibit oscillatory zoning (Figure 2), and have Th/U ratios >0.1, consistent with a magmatic origin (Corfu et al., 2003). The Triassic and Cretaceous zircons display subrounded to euhedral morphologies, suggesting first cycle to moderately recycled input (Figure 2).

The five samples exhibit comparable zircon age distributions, while the Cretaceous turbidite slice within the ophiolitic basement (TB-10B) contains a higher abundance of Permo-Triassic and Devonian-Carboniferous zircons (Figure 3). A limited population of Oligocene to Early Miocene zircons exists in K-03A of Tanjong




Figure 3. Kernel density estimation (KDE) spectra of detrital zircons from the Cretaceous turbidite slice and Upper Oligocene-Neogene sandstones sampled in this study.

Formation, documenting potential magmatism associated with Proto-South China Sea subduction during this interval (e.g., Breitfeld, Hall, et al., 2023). When integrated with detrital zircons from published Neogene sedimentary rock samples in Sabah (e.g., Breitfeld, Hall, et al., 2023; Zhang et al., 2023), these zircons maintained a distribution pattern dominated by Late Cretaceous (~100 Ma) ages, while the Kernel Density Estimate (KDE) plots show minor age peaks in the Permo-Triassic (~210–260 Ma) and Devonian-Carboniferous (~330–380 Ma; Figure 4). The $eHf(t)$ values of these detrital zircons from five samples began to increase in the Latest Triassic, and became overwhelmingly positive during the Jurassic and Cretaceous ($eHf(t) = -5.2 \sim +14.9$; Figure 5).

Figure 5 shows magmatic activity across West Sumatra and West and Southwest Borneo from the Permian, through the Triassic when Sibumasu collided with Indochina, to Early Miocene. Prior to the latest Triassic, magmatic zircons are scarce in both Borneo and West Sumatra. However, their detrital zircons exhibit scattered $eHf(t)$ values, with detrital zircons from West Sumatra showing more positive $eHf(t)$ values relative to those from




Figure 4. A Kernel density estimation (KDE) spectra of published zircon U-Pb ages of Mesozoic magmatic rocks around Sundaland. New and compiled detrital zircons U-Pb ages from the Upper Oligocene-Neogene sedimentary rocks in Sabah and Sumatra are also shown for comparison. For the references pertaining to the compiled data set, please refer to the Methods section. Table S1 and S3 in Supporting Information S1 present the new and compiled zircon age data, respectively.

West and Southwest Borneo. After the latest Triassic, on the western side (West Sumatra), magmatic zircons exhibit a narrow range in positive ϵ_{Hf} values whereas detrital zircons show a progressive trend toward increasingly positive ϵ_{Hf} values. This is not seen in the West and Southwest Borneo data, which instead shows a wider scattering of positive ϵ_{Hf} values since the latest Triassic. Subduction-related magmatism in the West and Southwest Borneo appears to have ceased for a considerable period following 80 Ma. By contrast detrital zircon data from West Sumatra imply ongoing magmatism and a continuation of the trend of increasing positive ϵ_{Hf} values.

5. Discussion

5.1. Upper Oligocene-Neogene Sediments in Sabah: A Record of Magmatism in the Southern Pacific-Pacific Domain

Previous provenance analyses indicates that sources of Cenozoic sedimentary rocks preserved in Sarawak and Sabah are mainly from West Borneo, Southwest Borneo and the Malay Peninsula with some grains may have been recycled from older sedimentary rocks (e.g., Breitfeld, Hennig-Breitfeld, et al., 2020, 2023; Breitfeld, Hall et al., 2023; Galin et al., 2017; Hennig-Breitfeld et al., 2019; van Hattum et al., 2013). The Malay Peninsula mainly provided Permian-Triassic zircons, while West Borneo yielded Triassic and Cretaceous zircons, and Southwest Borneo dominantly supplied Cretaceous age grains (Figure 4). New detrital zircon data from Upper Oligocene-Neogene sandstones are also dominated by Cretaceous ages (Figure 3), suggesting that sediment was mainly sourced from West Borneo and Southwest Borneo via the proto-Rajang River drainage system by recycling or directly from the arc, rather than from the Malay Peninsula (e.g., Breitfeld, Hennig-Breitfeld, et al., 2020). The subrounded to euhedral morphological signatures of Triassic zircons indicate that they may directly originate from the Triassic magmatic rocks of West Borneo (Figure 2), while sedimentary recycling of the Kuching-Sibu zone incorporating Malay-Thai and West Borneo provenance components remains a plausible mechanism (e.g., Breitfeld, Hennig-Breitfeld, et al., 2020, 2023; Breitfeld, Hall et al., 2023). Similar to the KDE age spectra, an MDS plot (Figure 6; Vermeesch et al., 2016) also shows zircon age distributions from Sabah samples plotting closest to zircon ages from magmatic rocks of West Borneo and Southwest Borneo, suggesting that this was the primary source area.

Magmatism in West Borneo and Southwest Borneo has been recognized as divided into Triassic and Jurassic subduction-related magmatic episodes and a Cretaceous magmatic arc associated with the Paleo-Pacific plate subduction (Breitfeld et al., 2017; Breitfeld, Davies et al., 2020; Hennig et al., 2017; Wang, Qian, et al., 2021; Wang, Zhang et al., 2021; Wang, Liu et al., 2022; Wang et al., 2024; Williams et al., 1988), while some Early Jurassic rift-related within-plate magmatism also developed in Southwest Borneo (Breitfeld, Davies, et al., 2020). Zircon $\epsilon_{\text{Hf}}(t)$ values serve as a powerful tracer for sediment provenance (e.g., Zhang et al., 2023), with their capacity to distinguish between mantle-derived material (more positive) and older crustal components (characterized by strongly negative $\epsilon_{\text{Hf}}(t)$ values) providing critical insights into the relative contributions of these endmembers to magmatic systems (Sundell & Macdonald, 2022). Most detrital zircons younger than 200 Ma have $\epsilon_{\text{Hf}}(t)$ values similar to those of West Borneo and Southwest Borneo magmatic rocks (Figure 5), and consistent with the implications of the KDEs and MDS plots that suggest a dominant supply from West Borneo and Southwest Borneo. A scarcity of detrital zircons postdating ~ 80 Ma aligns with the waning and cessation of subduction-related magmatic activity in West Borneo and Southwest Borneo (Breitfeld et al., 2017; Breitfeld, Davies et al., 2020; Hennig et al., 2017; Wang et al., 2024; Williams et al., 1988). A small number of




Figure 5. Zircon ages versus zircon $\epsilon\text{Hf}(t)$ values of the Mesozoic magmatic rocks and Neogene sedimentary rocks in West Sumatra and Borneo. Magmatic zircons compiled from Mesozoic igneous rocks in the Malay Peninsula are also shown for comparison. For the references pertaining to the compiled data set, please refer to the Methods section. Table S2 and S4 in Supporting Information S1 present the new and compiled zircon Hf isotope data, respectively.




Figure 6. A Multi-Dimensional Scalar (MDS; Vermeesch et al., 2016) plot for compiled zircon U-Pb ages from Mesozoic magmatic rocks around Sundaland, (a) with or (b) without the new and compiled Neogene detrital zircon data in Borneo and Sumatra. For the references pertaining to the compiled data set, please refer to the Methods section.

zircons younger than 200 Ma fall within the $\epsilon\text{Hf}(t)$ range of granites in the Malay Peninsula, indicating limited input from that area (e.g., Breitfeld, Hennig-Breitfeld, et al., 2020). The $\epsilon\text{Hf}(t)$ values of our detrital zircons from Sabah, that represent an archive of magmatism across West Borneo and Southwest Borneo, began to increase in the Latest Triassic, and became overwhelmingly positive during the Jurassic and Cretaceous (Figure 5). The compiled magmatic zircons from West Borneo and Southwest Borneo also have heterogeneous $\epsilon\text{Hf}(t)$ values (varying by up to ~ 20 ϵ units) in the Latest Triassic, followed by fully positive $\epsilon\text{Hf}(t)$ values with juvenile features (Figure 5). Although the detrital zircons in the Upper Oligocene-Neogene strata was interpreted as multi-recycling products from Rajang Group and/or Crocker Formation (Breitfeld, Hennig-Breitfeld, et al., 2020, 2023; Breitfeld, Hall et al., 2023; Hennig-Breitfeld et al., 2019), the close similarity of results between Sabah and West and Southwest Borneo supports the view that the Neogene sediments in Sabah effectively document the temporal changes in magmatic activity in West Borneo and Southwest Borneo since the Mesozoic, thus serving as a record of the magmatic evolution of the southern Paleo-Pacific domain.

5.2. Spatiotemporal Evolution of Magmatism in Each Subduction System

Published whole-rock elemental and Sr-Nd-Pb-(Hf-O) isotopic results indicate that the Mesozoic magmatic rocks in West Borneo were the products of subduction of the Paleo-Pacific plate under the East Asia active continental margin (Wang, Qian, et al., 2021; Wang, Zhang et al., 2021; Wang, Liu et al., 2022; Wang et al., 2024), while some Jurassic magmatic rocks in Southwest Borneo are interpreted as rift-related within-plate granites (Breitfeld, Davies, et al., 2020). Mesozoic granitoids in Sumatra represent the products of subduction magmatism in the easternmost Tethys (Li et al., 2020). Temporally, Mesozoic magmatism in West Borneo and Southwest Borneo exhibits some similarities in age with Sumatra, as shown by the magmatic zircon U-Pb ages on the KDE and MDS plots (Figures 4 and 6). During the Early Permian to Middle Triassic, magmatism associated with Paleo-Tethyan subduction predominantly occurred in East Malaysia, exhibiting distinct negative $\epsilon\text{Hf}(t)$ signatures (Figure 5 e.g., Gillespie et al., 2019; Liu et al., 2020; Ng et al., 2015; Wang, Qian, et al., 2021). The positive $\epsilon\text{Hf}(t)$ values of detrital zircons from Sumatra likely preserve information predating its accretion to Sundaland, given that Sumatra may represent the southern extension of the West Burma block (Barber & Crow, 2009; Metcalfe, 2013). In contrast, westward subduction of the Paleo-Pacific plate remained limited or inactive until the Late Triassic (e.g., Zhou et al., 2023). Therefore, only sporadic magmatism from this period is preserved in West Sumatra and West Borneo (Figure 5). Since latest Triassic, the temporal trends in Hf isotope variations are broadly similar in magmatic and detrital zircons from both Sumatra and Borneo, although $\epsilon\text{Hf}(t)$ values of magmatic and detrital zircons in Sumatra are consistently higher and more uniform than those in Borneo from the same period (Figure 5). This suggests that evolution of the Mesozoic magmatic arc was temporally synchronous across the two active continental margins (Sumatra and Borneo). The notable variation (~ 20 ϵ units) in $\epsilon\text{Hf}(t)$ values seen in latest Triassic magmatic and detrital zircons (220–200 Ma) from both Sumatra and Borneo, representing the western and eastern boundaries of Sundaland, suggest a mix of evolved and juvenile crust in each subduction system immediately after Triassic collision between Indochina and Sibumasu. A trend to positive $\epsilon\text{Hf}(t)$ values on Borneo and Sumatra was established during the Jurassic and Cretaceous, indicating that juvenile crustal growth is prevalent on both sides of the interaction zone (Figure 5). In the central part of the overriding plate (Malay Peninsula), magmatic rocks remain isotopically evolved, implying that far from the trench the continental mantle lithosphere is thick (e.g., Chapman et al., 2017). Detrital data show a magmatic lull after ~ 80 Ma in West and Southwest Borneo, indicating that subduction-related arc magmatism in West and Southwest Borneo had largely ceased by that time (Breitfeld, Davies, et al., 2020; Hennig et al., 2017), unlike Sumatra where magmatism continued throughout the Cenozoic (Figures 4 and 5).

Spatially, the post-Latest Triassic magmatic rocks of West Sumatra are distributed along a narrow belt near the trench with high and uniformly positive $\epsilon\text{Hf}(t)$ values, while those in West and Southwest Borneo are relatively more widely distributed with lower positive $\epsilon\text{Hf}(t)$ values away from the trench (Figures 1b and 7a), potentially due to differences in the angle, age, and rate of the subducting plates. Although plate kinematics are not well-constrained due to a lack of preserved oceanic lithosphere, reconstructions of global plate motion show that from the Jurassic to the Cretaceous, there is no consistent pattern in reconstructed convergence rates between the two margins of Sundaland (Figure 7b; Müller et al., 2016), this suggests that convergence rate is not the primary factor influencing the spatial distribution patterns of Mesozoic magmatic rocks. However, global plate reconstructions predict the eastern margin of interaction zone was mainly influenced by subduction of older Paleo-Pacific oceanic lithosphere that would be consistent with accreted ocean plate stratigraphy ages in SW Hokkaido,




Figure 7. (a) Spatial distribution of zircon $\epsilon Hf(t)$ values Mesozoic magmatic rocks (200–66 Ma) in West Sumatra, Malay Peninsula, and West and Southwest Borneo. Filled circles represent the mean $\epsilon Hf(t)$ values of magmatic zircons. (b) Convergence rate and oceanic plate age along the two subduction zones of Sundaland (orange line = Paleo-Pacific; blue = Tethyan) based on global plate reconstructions of Müller et al. (2016, 2019).

Japan (Ishiga & Ishiyama, 1987). By contrast, West Sumatra initially experienced subduction of relatively younger oceanic lithosphere that separating it from the Woyla Arc (Advokaat et al., 2018). Subduction of more steeply dipping young oceanic lithosphere would tend to form a narrow arc near the trench due to their weaker ability to carry volatiles to the lower crust, compared to more gently dipping subduction of a colder and older serpentinized Paleo-Pacific oceanic lithosphere (e.g., Bastias-Silva et al., 2024).




Figure 8. Input of juvenile material into Sumatra and Borneo, initiated in the Latest Triassic and continued to develop from the Jurassic to the Cretaceous due to the extension of the inward-dipping double subduction.

5.3. Inward Dipping Double Subduction Initiated in Latest Triassic

Spatiotemporal variations in radiogenic isotope compositions can shed light on the evolution of subduction zone dynamics. In Sumatra, a fundamental restructuring of the arc system in the Latest Triassic is identified based on the marked increase in $\epsilon\text{Hf}(t)$ values observed in zircons from Mesozoic granitoids, as well as detrital zircons from Cenozoic strata and modern river sediments (Figure 5; Li et al., 2020; Zhang et al., 2019). An extensional fringing arc related to NW-SE trending extensional basins in Sumatra (Barber & Crow, 2009) or slab retreat or rollback that occurred shortly after the Middle-Late Triassic collision of Sibumasu with Indochina and East Malaya, have been considered as indicative of the initial input of juvenile material in Sumatra (Li et al., 2020).

For Borneo, the presence of Mesozoic supra-subduction zone ophiolites and Mesozoic-Cenozoic basin indicates that Borneo has been subjected to a protracted extensional setting since the Late Triassic (Burton-Johnson et al., 2020). The development of the Triassic Sadong and Kuching Formations and Cretaceous Pedawan Formation—characterized by forearc basin successions in West Borneo—provides critical evidence for syn-depositional extensional tectonics (Breitfeld et al., 2017; Breitfeld, Burley et al., 2023; Mazumder et al., 2021; Pieters et al., 1993), likely associated with episodic slab rollback during subduction of the Paleo-Pacific plate (Zhou et al., 2023). Significant extensional collapse around 215 Ma also characterizes the Sibumasu terrane, particularly in northern Thailand and potentially other regions (Morley, 2018). Therefore, the notable variation (220–200 Ma, ~ 20 ϵ units) in magmatic and detrital zircon $\epsilon\text{Hf}(t)$ values from both Sumatra and Borneo suggest the addition of juvenile material in an extensional setting in each subduction system.

The initiation of double subduction is usually associated with collision, such as the Solomon subduction zone (Sun et al., 2021). Numerical modeling indicates that during the early stages of a double subduction system, the stress on the overriding plate either remained in an extensional state (Lyu et al., 2019) or transitioned from compression to extension, typically within the first few million years (Zhang et al., 2024). Therefore, we consider that an inward-dipping double subduction system underneath Sundaland (the interaction zone) began to be established immediately after latest Triassic collision between Indochina and Sibumasu (Figure 8).




Figure 9. (a) Palaeogeographic reconstructions for the Pale-Pacific and Eastern Tethys in late Triassic (after Metcalfe, 2017). Notes the enlarged view of the interaction zone. (b) Density curves of $\epsilon Hf(t)$ from latest Triassic to early Late Cretaceous (200–80 Ma) for batholith in the Tethys and Pale-Pacific domains. Data of batholith in the South Qiangtang and South China are from Gong et al. (2024), Zhang et al. (2023) and references therein.

During the Jurassic and Cretaceous, the positive $\epsilon Hf(t)$ values on Borneo and Sumatra indicate that the melt source region of the two subduction zones is predominantly depleted asthenospheric mantle but, mixed with varying proportions of isotopically evolved continental lithospheric mantle. The contrasting ranges of positive $\epsilon Hf(t)$ values imply the double subduction system evolved asymmetrically. The narrow spatial distribution observed in the arc magmatism of Sumatra and their uniformly positive $\epsilon Hf(t)$ isotopic values is thus attributed to the absence of continental lithospheric mantle near the trench during the high-angle subduction of the Meso-Tethys slab (e.g., Chapman et al., 2017). In contrast, the wide arc magma belt and broader Hf isotopic values in Borneo represent varying degrees of involvement of continental mantle lithosphere during more shallow dipping subduction of the Paleo-Pacific oceanic crust in that time (Zhou et al., 2023; Figure 8). Around 80 Ma, this stable asymmetric double subduction system was disrupted, marked by the complete cessation of subduction-related arc magmatism in Borneo while magmatism continued in West Sumatra. In SE Vietnam (north of West Borneo) subduction-related magmatism had also ceased by c. 80 Ma (Hennig-Breitfeld et al., 2021), implying the waning of the Paleo-Pacific subduction zone at that time. The magmatic arc in West Sumatra has maintained isotopically juvenile signals even after 80 Ma (Figure 5). This aligns with numerical models suggesting that large-scale extension, along with extensive and vigorous mantle upwellings, persists even after one of the two subductions ceases (Li et al., 2024).

5.4. Juvenile Crust Production in SE Asia Linked to Double Subduction

Our data show a significant isotopic shift from isotopically evolved to isotopically juvenile in the continental arc of the Sundaland margin around 200 Ma, triggered by collision between Sibumasu and Indochina and the initiation of an inward-dipping double subduction regime (Figure 5). During the latest Triassic to Jurassic, Southeast Asia served as an interaction zone connecting the Qiangtang block of western Meso-Tethys and South China of the northern Paleo-Pacific, where single subduction developed (Figure 9a). However, isotopic data for contemporary batholith from South Qiangtang in the Meso-Tethys domain and South China in the Paleo-Pacific domain exhibit $\epsilon Hf(t)$ values lower than those measured from West and Southwest Borneo and West Sumatra in the interaction zone (Figure 9b). This raises the possibility that, compared to single subduction in the Meso-

Tethys or Paleo-Pacific domains, an inward-dipping double subduction system is more conducive to the formation of juvenile crust.

Numerical simulations indicate that inward-dipping double subduction results in larger-scale upper plate extension and greater volumes of upwelling mantle compared to single subduction (Li et al., 2024; Lyu et al., 2019; Zhang et al., 2024). This fits with evidence of widespread extension across Sumatra and Borneo (Barber & Crow, 2009; Breifeld et al., 2017; Breifeld, Burley et al., 2023; Burton-Johnson et al., 2020; Mazumder et al., 2021; Pieters & Noya, 1993; Zhou et al., 2023). Isotopically juvenile signature in Sumatra persist to the present day (Figure 5) and may be a residual feature inherited from the time of double subduction (e.g., Li et al., 2024). Our study emphasizes that the role of inward-dipping double subduction in Southeast Asia, including its influence on mantle structure and temperature.

6. Conclusions

Records of magmatic and detrital zircons from both Borneo and Sumatra show an inward-dipping double subduction system beneath Sundaland initiated in the latest Triassic, coinciding with the collision of Indochina and Sibumasu. Magmatism persisted on both West and Southwest Borneo and West Sumatra from the latest Triassic to the Cretaceous; however, it evolved asymmetrically due to differences in the subducting oceanic lithospheres—specifically, the younger, steeply dipping Meso-Tethys and the older, more shallowly dipping Paleo-Pacific oceanic lithosphere. After 80 Ma, subduction-related magmatism on the Borneo side ceased following the termination of subduction of Paleo-Pacific oceanic crust, while on the Sumatra side, magmatism with isotopically juvenile compositions continued. Furthermore, our study establishes a possible causal link between the strengthened juvenile crustal signatures in Southeast Asia and the large-scale extension of the overriding lithospheric plate in an inward-dipping double subduction system.

Data Availability Statement

The Supporting Information and Table S1–S4 in Supporting Information S1 is available at <https://doi.org/10.17632/8g669bt2bt.1> (Zhao, 2025).

References

- Advokaat, E. L., Bongers, M. L. M., Rudyawan, A., BouDagher-Fadel, M. K., Langereis, C. G., & van Hinsbergen, D. J. J. (2018). Early Cretaceous origin of the Woyla Arc (Sumatra, Indonesia) on the Australian plate. *Earth and Planetary Science Letters*, 498, 348–361. <https://doi.org/10.1016/j.epsl.2018.07.001>
- Barber, A. J. (2000). The origin of the Woyla Terranes in Sumatra and the late Mesozoic evolution of the Sundaland margin. *Journal of Asian Earth Sciences*, 18(6), 713–738. [https://doi.org/10.1016/S1367-9120\(00\)00024-9](https://doi.org/10.1016/S1367-9120(00)00024-9)
- Barber, A. J., & Crow, M. J. (2009). Structure of Sumatra and its implications for the tectonic assembly of Southeast Asia and the destruction of paleotethys. *Island Arc*, 18(1), 3–20. <https://doi.org/10.1111/j.1440-1738.2008.00631.x>
- Barber, A. J., Crow, M. J., & Milsom, J. S. (2005). *Sumatra: Geology, resources and tectonic evolution* (Vol. 31, p. 290). Geological Society [London] Memoir.
- Bastias-Silva, J., Burton-Johnson, A., Chew, D., Riley, T., Jara, W., & Chiaradia, M. (2024). A temporal control on the isotopic compositions of the Antarctic Peninsula arc. *Communications Earth and Environment*, 5(1), 157. <https://doi.org/10.1038/s43247-024-01301-1>
- Breifeld, H. T., Hall, R., Suggate, S. M., van Hattum, M. W. A., Hennig-Breifeld, J., BouDagher-Fadel, M. K., et al. (2023). Neogene sedimentary successions in northern and central Sabah: Provenance and tectonic implications. In *Stratigraphy of Geo- and biodynamic processes* (pp. 71–119). <https://doi.org/10.1016/bs.sats.2023.08.009>
- Breifeld, H. T., Burley, S. D., Galin, T., Hennig-Breifeld, J., & Rajali, R. (2023). The Kuching formation: A deep marine equivalent of the Sadong formation, and its implications for the early Mesozoic tectonic evolution of Western and southern Borneo. *Bulletin Geological Society of Malaysia*, 76, 101–129. <https://doi.org/10.7186/BGSM76202308>
- Breifeld, H. T., Davies, L., Hall, R., Armstrong, R., Forster, M., Lister, G., et al. (2020). Mesozoic paleo-pacific subduction beneath SW Borneo: U–pb geochronology of the Schwaneer Granitoids and the Pinoh Metamorphic group. *Frontiers in Earth Science*, 8, 568715. <https://doi.org/10.3389/feart.2020.568715>
- Breifeld, H. T., Hall, R., Galin, T., Forster, M. A., & BouDagher-Fadel, M. K. (2017). A Triassic to Cretaceous Sundaland–Pacific subduction margin in West Sarawak, Borneo. *Tectonophysics*, 694, 35–56. <https://doi.org/10.1016/j.tecto.2016.11.034>
- Breifeld, H. T., Hennig-Breifeld, J., Anthony, G., BouDagher-Fadel, M., Vermeesch, P., Lünsdorf, K., et al. (2023). The onshore West Baram Delta deposits: Provenance and drainage in the middle Miocene to Pliocene in NW Borneo and comparison to the Champion Delta. *Marine and Petroleum Geology*, 158, 106537. <https://doi.org/10.1016/j.marpetgeo.2023.106537>
- Breifeld, H. T., Hennig-Breifeld, J., BouDagher-Fadel, M. K., Hall, R., & Galin, T. (2020). Oligocene-Miocene drainage evolution of NW Borneo: Stratigraphy, sedimentology and provenance of Tatau-Nyalau province sediments. *Journal of Asian Earth Sciences*, 195, 104331. <https://doi.org/10.1016/j.jseaes.2020.104331>
- Burton-Johnson, A., Macpherson, C. G., Millar, I. L., Whitehouse, M. J., Ottley, C. J., & Nowell, G. M. (2020). A Triassic to Jurassic arc in north Borneo: Geochronology, geochemistry, and genesis of the segama valley felsic intrusions and the Sabah ophiolite. *Gondwana Research*, 84, 229–244. <https://doi.org/10.1016/j.gr.2020.03.006>

Acknowledgments

This work was funded by the National Key Research and Development Program of China (2023YFF0803403), the Strategic Priority Research Program (B) of the Chinese Academy of Sciences (XDB0840200), the National Natural Science Foundation (42202248), the Guangdong Basic and Applied Basic Research Foundation (2023A1515011786) of China and the Director General's Scientific Research Fund of Guangzhou Marine Geological Survey, China (2023GMGSJZJ00007). We are grateful to the Editor Margaret Rusmore, the Associate Editor, an anonymous reviewer and H. Tim Breifeld for their careful and constructive reviews on our paper.

- Cao, J., Yang, X., Du, G., & Li, H. (2020). Genesis and tectonic setting of the Malaysian Waterfall granites and tin deposit: Constraints from LA-ICP (MC)-MS zircon U-Pb and cassiterite dating and Sr-Nd-Hf isotopes. *Ore Geology Reviews*, *118*, 103336. <https://doi.org/10.1016/j.oregeorev.2020.103336>
- Chapman, J. B., Ducea, M. N., Kapp, P., Gehrels, G. E., & DeCelles, P. G. (2017). Spatial and temporal radiogenic isotopic trends of magmatism in Cordilleran orogens. *Gondwana Research*, *48*, 189–204. <https://doi.org/10.1016/j.gr.2017.04.019>
- Cobbing, E. J. (2005). Granites. In A. J. Barber, et al., (eds.), *Sumatra: Geology, Resources and Tectonic Evolution* (Vol. 31(1)), pp. 54–62. Geological Society [London] Memoir. <https://doi.org/10.1144/gsl.mem.2005.031.01.05>
- Corfu, F., Hanchar, J. M., Hoskin, P. W., & Kinny, P. (2003). Atlas of zircon textures. *Reviews in Mineralogy and Geochemistry*, *53*(1), 469–500. <https://doi.org/10.2113/0530469>
- De Min, A., Velicogna, M., Ziberna, L., Chiaradia, M., Alberti, A., & Marzoli, A. (2020). Triassic magmatism in the European Southern Alps as an early phase of Pangea break-up. *Geological Magazine*, *157*(11), 1800–1822. <https://doi.org/10.1017/S0016756820000084>
- Fan, J.-J., Zhang, B.-C., Zhou, J.-B., Niu, Y., Sun, S.-L., Lv, J.-P., et al. (2024). The Meso-Tethys Ocean: The nature, extension and spatial-temporal evolution. *Earth-Science Reviews*, *255*, 104839. <https://doi.org/10.1016/j.earscirev.2024.104839>
- Galín, T., Breitfeld, H. T., Hall, R., & Sevastjanova, I. (2017). Provenance of the Cretaceous–Eocene Rajang Group submarine fan, Sarawak, Malaysia from light and heavy mineral assemblages and U-Pb zircon geochronology. *Gondwana Research*, *51*, 209–233. <https://doi.org/10.1016/j.gr.2017.07.016>
- Gan, C., Qian, X., Wang, Y., Feng, Q., Zhang, Y., Asis, J. B., & Ao, S. (2022). Late cretaceous granitoids along the Northern Kuching Zone: Implications for the paleo-pacific subduction in Borneo. *Lithosphere*, *2022*(1), 3310613. <https://doi.org/10.2113/2022/3310613>
- Gasparon, M., & Varne, R. (1995). Sumatran granitoids and their relationship to Southeast Asian terranes. *Tectonophysics*, *251*(1), 277–299. [https://doi.org/10.1016/0040-1951\(95\)00083-6](https://doi.org/10.1016/0040-1951(95)00083-6)
- Ghani, A. A., Lo, C.-H., & Chung, S.-L. (2013). Basaltic dykes of the Eastern Belt of Peninsular Malaysia: The effects of the difference in crustal thickness of Sibumasu and Indochina. *Journal of Asian Earth Sciences*, *77*, 127–139. <https://doi.org/10.1016/j.jseas.2013.08.004>
- Gillespie, M. R., Kendall, R. S., Leslie, A. G., Millar, I. L., Dodd, T. J. H., Kearsey, T. I., et al. (2019). The igneous rocks of Singapore: New insights to Palaeozoic and Mesozoic assembly of the Sukhothai Arc. *Journal of Asian Earth Sciences*, *183*, 103940. <https://doi.org/10.1016/j.jseas.2019.103940>
- Golonka, J. (2007). Late Triassic and early Jurassic palaeogeography of the world. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *244*(1–4), 297–307. <https://doi.org/10.1016/j.palaeo.2006.06.041>
- Gong, N., Zhang, S.-Q., Qi, H., Yuan, G.-L., Li, J., Wang, G.-H., et al. (2024). Two stage Mesozoic Oceanic subduction and related mantle metasomatism beneath the South Qiangtang terrane with implications for post-collisional magmatism. *Gondwana Research*, *136*, 219–235. <https://doi.org/10.1016/j.gr.2024.09.001>
- Haile, N. S., McElhinny, M. W., & McDougall, I. (1977). Palaeomagnetic data and radiometric ages from the Cretaceous of West Kalimantan (Borneo), and their significance in interpreting regional structure. *Journal of the Geological Society*, *133*(2), 133–144. <https://doi.org/10.1144/gsjgs.133.2.0133>
- Hall, R. (2012). Late Jurassic–Cenozoic reconstructions of the Indonesian region and the Indian Ocean. *Tectonophysics*, *570–571*, 1–41. <https://doi.org/10.1016/j.tecto.2012.04.021>
- Hall, R. (2013). Contraction and extension in northern Borneo driven by subduction rollback. *Journal of Asian Earth Sciences*, *76*, 399–411. <https://doi.org/10.1016/j.jseas.2013.04.010>
- Hall, R., & Breitfeld, T. (2017). Nature and demise of the Proto-South China Sea. *Bulletin Geological Society of Malaysia*, *63*, 61–76. <https://doi.org/10.7186/bgsm63201703>
- Hazad, F. I., Ghani, A. A., & Lo, C. H. (2019). Arc related dioritic–granodioritic magmatism from southeastern Peninsular Malaysia and its tectonic implication. *Cretaceous Research*, *95*, 208–224. <https://doi.org/10.1016/j.cretres.2018.10.016>
- Hennig, J., Breitfeld, H. T., Hall, R., & Nugraha, A. M. S. (2017). The Mesozoic tectono-magmatic evolution at the Paleo-Pacific subduction zone in West Borneo. *Gondwana Research*, *48*, 292–310. <https://doi.org/10.1016/j.gr.2017.05.001>
- Hennig-Breitfeld, J., Breitfeld, H. T., Hall, R., BouDagher-Fadel, M., & Thirlwall, M. (2019). A new upper Paleogene to Neogene stratigraphy for Sarawak and Labuan in northwestern Borneo: Paleogeography of the eastern Sundaland margin. *Earth-Science Reviews*, *190*, 1–32. <https://doi.org/10.1016/j.earscirev.2018.12.006>
- Hennig-Breitfeld, J., Breitfeld, H. T., Sang, D. Q., Vinh, M. K., Long, T. V., Thirlwall, M., & Cuong, T. X. (2021). Ages and character of igneous rocks of the Da Lat Zone in SE Vietnam and adjacent offshore regions (Cuu Long and Nam Con Son basins). *Journal of Asian Earth Sciences*, *218*, 104878. <https://doi.org/10.1016/j.jseas.2021.104878>
- Hieu, P. T., Yang, Y. Z., Binh, D. Q., Nguyen, T. B. T., Dung, L. T., & Chen, F. (2015). Late Permian to Early Triassic crustal evolution of the Kontum massif, central Vietnam: Zircon U-Pb ages and geochemical and Nd-Hf isotopic composition of the Hai Van granitoid complex. *International Geology Review*, *57*(15), 1877–1888. <https://doi.org/10.1080/00206814.2015.1031194>
- Hou, L., Xiong, F., Wang, W., Guo, L., Peng, H., Ni, S., & Zhang, Q. (2019). Carboniferous-Triassic felsic igneous rocks and typical mineral deposits in the Truong Son orogenic belt, SE Asia: Implications for Paleo-Tethyan tectonic evolution and metallogeny. *Ore Geology Reviews*, *112*, 103036. <https://doi.org/10.1016/j.oregeorev.2019.103036>
- Hsu, C. C. (2016). *Detrital zircon U-Pb and Hf isotopic study in northwestern Sumatra, Indonesia*. [Master Thesis] (pp. 1–158). National Taiwan University.
- Hu, P. Y., Li, C., Li, J., Wang, M., Xie, C. M., & Wu, Y. W. (2014). Zircon U–Pb–Hf isotopes and whole-rock geochemistry of gneissic granites from the Jitang complex in Leiwuqi area, eastern Tibet, China: Record of the closure of the Paleo-Tethys Ocean. *Tectonophysics*, *623*, 83–99. <https://doi.org/10.1016/j.tecto.2014.03.018>
- Hu, Z. C., Liu, Y. S., Gao, S., Liu, W., Yang, L., Zhang, W., et al. (2012). Improved in situ Hf isotope ratio analysis of zircon using newly designed X skimmer cone and jet sample cone in combination with the addition of nitrogen by laser ablation multiple collector ICP-MS. *Journal of Analytical Atomic Spectrometry*, *27*(9), 1391–1399. <https://doi.org/10.1039/C2JA30078H>
- Hutchison, C. S. (1996). The “Rajang Accretionary Prism” and “Lupar Line” problem of Borneo. In *Tectonic Evolution of SE Asia*. R. Hall, & D. J. Blundell (Eds.), Geological Society of London Special Publication. (Vol. 106(1)), pp. 247–261. <https://doi.org/10.1144/gsl.sp.1996.106.01.16>
- Hutchison, C. S. (2005). *Geology of North-West Borneo: Sarawak, Brunei and Sabah* (p. 421). Elsevier.
- Ishiga, H., & Ishiyama, D. (1987). Jurassic accretionary complex in Kaminokuni Terrane, southwestern Hokkaido, Japan. *Mining Geology*, *37*, 381–394.
- Isozaki, Y., Aoki, K., Nakama, T., & Yanai, S. (2010). New insight into a subduction-related orogen: A reappraisal of the geotectonic framework and evolution of the Japanese Islands. *Gondwana Research*, *18*(1), 82–105. <https://doi.org/10.1016/j.gr.2010.02.015>
- Jamil, A., Ghani, A. A., Zaw, K., Osman, S., & Quek, L. X. (2016). Origin and tectonic implications of the ~200 Ma, collision-related Jerai pluton of the Western Granite Belt, Peninsular Malaysia. *Journal of Asian Earth Sciences*, *127*, 32–46. <https://doi.org/10.1016/j.jseas.2016.06.004>

- Li, J., Sha, J., McLoughlin, S., & Wang, X. (2019). Mesozoic and Cenozoic palaeogeography, palaeoclimate and palaeoecology in the eastern Tethys. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 515, 1–5. <https://doi.org/10.1016/j.palaeo.2018.11.014>
- Li, L.-S., Capitanio, F. A., Cawood, P. A., Wu, B.-J., Zhai, M.-G., & Wang, X.-L. (2024). Double subduction controls on long-lived continental tectonics and subcontinental mantle temperatures. *Geology*, 52(11), 836–840. <https://doi.org/10.1130/g52232.1>
- Li, S., Chung, S.-L., Lai, Y.-M., Ghani, A. A., Lee, H.-Y., & Murtadha, S. (2020). Mesozoic juvenile crustal formation in the easternmost Tethys: Zircon Hf isotopic evidence from Sumatran Granitoids, Indonesia. *Geology*, 48(10), 1002–1005. <https://doi.org/10.1130/g47304.1>
- Li, Z. X., & Li, X. H. (2007). Formation of the 1300-km-wide intracontinental orogen and postorogenic magmatic province in Mesozoic South China: A flat-slab subduction model. *Geology*, 35(2), 179–182. <https://doi.org/10.1130/g23193a.1>
- Liew, T. C., & Page, R. W. (1985). U-Pb zircon dating of granitoid plutons from the West Coast Province of Peninsular Malaysia. *Journal of the Geological Society*, 142(3), 515–526. <https://doi.org/10.1144/gsjgs.142.3.0515>
- Liu, L., Hu, R.-Z., Zhong, H., Yang, J.-H., Kang, L.-F., Zhang, X.-C., et al. (2020). Petrogenesis of multistage S-type granites from the Malay Peninsula in the Southeast Asian tin belt and their relationship to Tethyan evolution. *Gondwana Research*, 84, 20–37. <https://doi.org/10.1016/j.gr.2020.02.013>
- Liu, Y. S., Gao, S., Hu, Z. C., Gao, C. G., Zong, K. Q., & Wang, D. B. (2010). Continental and oceanic crust recycling-induced melt-peridotite interactions in the Trans-North China Orogen: U-pb dating, Hf isotopes and trace elements in zircons of mantle xenoliths. *Journal of Petrology*, 51(1–2), 537–571. <https://doi.org/10.1093/petrology/egp082>
- Lyu, T., Zhu, Z., & Wu, B. (2019). Subducting slab morphology and mantle transition zone upwelling in double-slab subduction models with inward-dipping directions. *Geophysical Journal International*, 218(3), 2089–2105. <https://doi.org/10.1093/gji/ggz268>
- Mange, M. A., & Maurer, H. F. W. (1992). *Heavy minerals in colour*. Chapman and Hall. <https://doi.org/10.1007/978-94-011-2308-2>
- Martinez, F., & Taylor, B. (2002). Mantle wedge control on back-arc crustal accretion. *Nature*, 416(6879), 417–420. <https://doi.org/10.1038/416417a>
- Mazumder, R., Anthony, F. B. M., Say, B. T. S., Roy, S., Hajri, A. A., Ohta, T., et al. (2021). Cretaceous Forearc sedimentation and contemporary basin tectonics in northwestern Borneo: New sedimentological insights from Pedawan formation, Kuching zone, east Malaysia. *The Journal of Geology*, 129(4), 391–415. <https://doi.org/10.1086/715790>
- Metcalfe, I. (1988). Origin and assembly of Southeast Asian continental terranes. In *Gondwana and Tethys* M. G. Audley-Charles, & A. Hallam (Eds.), Geological Society of London Special Publication (Vol. 37(1), pp. 101–118). <https://doi.org/10.1144/gsl.sp.1988.037.01.08>
- Metcalfe, I. (1996). Pre-Cretaceous evolution of SE Asian terrane, In R. Hall, & D. Blundell, (eds.), *Tectonic Evolution of Southeast Asia* (Vol. 106(1), pp. 97–122). Geological Society [London] Special Publication. <https://doi.org/10.1144/GSL.SP.1996.106.01.09>
- Metcalfe, I. (2009). Late Palaeozoic and Mesozoic tectonic and palaeogeographical evolution of SE Asia. In E. Buffetaut, G. Cuny, J. Le Loeuff, & V. Suteethorn (Eds.), *Late Palaeozoic and Mesozoic Ecosystems in SE Asia* (Vol. 315(1), pp. 7–23). Geological Society of London Special Publication. <https://doi.org/10.1144/sp315.2>
- Metcalfe, I. (2011). Palaeozoic–Mesozoic history of SE Asia. In R. Hall, M. A. Cottam, & M. E. J. Wilson (Eds.), *The SE Asian gateway: History and tectonics of the Australia-Asia collision*. Geological Society of London. <https://doi.org/10.1144/sp355.2>
- Metcalfe, I. (2013). Gondwana dispersion and Asian accretion: Tectonic and palaeogeographic evolution of eastern Tethys. *Journal of Asian Earth Sciences*, 66, 1–33. <https://doi.org/10.1016/j.jseae.2012.12.020>
- Metcalfe, I. (2017). Tectonic evolution of Sundaland. *Bulletin Geological Society of Malaysia*, 63, 27–60. <https://doi.org/10.7186/bgsm63201702>
- Metcalfe, I. (2021). Multiple Tethyan ocean basins and orogenic belts in Asia. *Gondwana Research*, 100, 87–130. <https://doi.org/10.1016/j.gr.2021.01.012>
- Minh, N. T., Dung, N. T., Hung, D. D., Minh, P., Yu, Y., & Hieu, P. T. (2020). Zircon U-Pb ages, geochemistry and isotopic characteristics of the Chu Lai granitic pluton in the Kontum massif, central Vietnam. *Mineralogy and Petrology*, 114(4), 289–303. <https://doi.org/10.1007/s00710-020-00707-x>
- Morley, C. K. (2018). Understanding Sibumasu in the context of ribbon continents. *Gondwana Research*, 64, 184–215. <https://doi.org/10.1016/j.gr.2018.07.006>
- Müller, R. D., Seton, M., Zahirovic, S., Williams, S. E., Matthews, K. J., Wright, N. M., et al. (2016). Ocean Basin evolution and global-scale plate reorganization events since Pangea breakup. *Annual Review of Earth and Planetary Sciences*, 44(1), 107–138. <https://doi.org/10.1146/annurev-earth-060115-012211>
- Müller, R. D., Zahirovic, S., Williams, S. E., Cannon, J., Seton, M., Bower, D. J., et al. (2019). A global plate model including lithospheric deformation along major rifts and orogens since the Triassic. *Tectonics*, 38(6), 1884–1907. <https://doi.org/10.1029/2018tc005462>
- Nance, R. D., Murphy, J. B., & Santosh, M. (2014). The supercontinent cycle: A retrospective essay. *Gondwana Research*, 25(1), 4–29. <https://doi.org/10.1016/j.gr.2012.12.026>
- Ng, S. W.-P., Whitehouse, M. J., Roselee, M. H., Teschner, C., Murtadha, S., Oliver, G. J. H., et al. (2017). Late Triassic granites from Bangka, Indonesia: A continuation of the Main Range granite province of the South-East Asian Tin Belt. *Journal of Asian Earth Sciences*, 138, 548–561. <https://doi.org/10.1016/j.jseae.2017.03.002>
- Ng, S. W.-P., Whitehouse, M. J., Searle, M. P., Robb, L. J., Ghani, A. A., Chung, S.-L., et al. (2015). Petrogenesis of Malaysian granitoids in the Southeast Asian tin belt: Part 2. U-Pb zircon geochronology and tectonic model. *Geological Society of America Bulletin*, 127(9–10), 1238–1258. <https://doi.org/10.1130/b31214.1>
- Nguyen, H. H. (2019). *The Cenozoic evolution of the South Vietnam margin of the South China Sea and the origin of coastal placer deposits*. [Ph. D. thesis] (p. 177). University of London.
- Oliver, G., Zaw, K., Hotson, M., Meffre, S., & Manka, T. (2014). U-Pb zircon geochronology of early Permian to Late Triassic rocks from Singapore and Johor: A plate tectonic reinterpretation. *Gondwana Research*, 26(1), 132–143. <https://doi.org/10.1016/j.gr.2013.03.019>
- Olsen, P. E. (1997). Stratigraphic record of the early Mesozoic breakup of Pangea in the Laurasia-Gondwana rift system. *Annual Review of Earth and Planetary Sciences*, 25(1), 337–401. <https://doi.org/10.1146/annurev.earth.25.1.337>
- Pastor-Galán, D., Spencer, C. J., Furukawa, T., & Tsujimori, T. (2021). Evidence for crustal removal, tectonic erosion and flare-ups from the Japanese evolving forearc sediment provenance. *Earth and Planetary Science Letters*, 564, 116893. <https://doi.org/10.1016/j.epsl.2021.116893>
- Pearce, N. J. G., Perkins, W. T., Westgate, J. A., Gorton, M. P., Jackson, S. E., Neal, C. R., & Chenerly, S. P. (1997). A compilation of new and published major and trace element data for NIST SRM 610 and NIST SRM 612 Glass Reference materials. *Geostandards Newsletter*, 21(1), 115–144. <https://doi.org/10.1111/j.1751-908X.1997.tb00538.x>
- Pieters, P. E., & Surono Noya, Y. (1993). *Geology of the Putussibau Sheet area, Kalimantan* (Vol. 1). Geological Survey of Indonesia, Directorate of Mineral Resources, Geological Research and Development Centre, Bandung, Quadrangle 1616.250000.
- Qian, X., Bai, T., Yu, Y., Mustapha, K. A., Sheldrick, T. C., Gan, C., & Wang, Y. (2023). Multiple stages of continental rifting in Eastern Peninsular Malaysia: New insights from Jurassic–Cretaceous Granitoids. *Journal of the Geological Society*, 180(5), jgs2023–jgs2025. <https://doi.org/10.1144/jgs2023-025>

- Qian, X., Yu, Y., Wang, Y., Gan, C., Zhang, Y., Asis, J. B., & Ao, S. (2022). Late Cretaceous Nature of SW Borneo and paleo-pacific subduction: New insights from the granitoids in the schwaner Mountains. *Lithosphere*, 2022(1), 8483732. <https://doi.org/10.2113/2022/8483732>
- Qiu, L., Kong, R., Yan, D.-P., Mu, H.-X., Sun, W., Sun, S., et al. (2022). Paleo-Pacific plate subduction on the eastern Asian margin: Insights from the Jurassic foreland system of the overriding plate. *Geological Society of America Bulletin*, 134(9–10), 2305–2320. <https://doi.org/10.1130/b36118.1>
- Qiu, L., Li, X., Li, X., Yan, D.-P., Ren, M., Zhang, L., & Cheng, G. (2022). Petrogenesis of early cretaceous intermediate to felsic rocks in Shanghai, South China: Magmatic response to Paleo-Pacific plate subduction. *Tectonophysics*, 838, 229469. <https://doi.org/10.1016/j.tecto.2022.229469>
- Quek, L., Lai, Y.-M., Ghani, A. A., Roselee, M. H., Lee, H.-Y., Iizuka, Y., et al. (2021). Peninsular Malaysia transitional geodynamic process from Gondwana to Pangaea: New constraints from 500 to 200 Ma magmatic zircon U-Pb ages and Hf isotopic compositions. *Gondwana Research*, 94, 56–72. <https://doi.org/10.1016/j.gr.2021.03.001>
- Quek, L. X., Lee, T.-Y., Ghani, A. A., Lai, Y.-M., Roselee, M. H., Lee, H.-Y., et al. (2021). Tracing detrital signature from Indochina in Peninsular Malaysia fluvial sediment: Possible detrital zircon recycling into West Borneo Cenozoic sediments. *Journal of Asian Earth Sciences*, 218, 104876. <https://doi.org/10.1016/j.jseae.2021.104876>
- Searle, M. P., Whitehouse, M. J., Robb, L. J., Ghani, A. A., Hutchison, C. S., Sone, M., et al. (2012). Tectonic evolution of the Sibumasu-Indochina terrane collision zone in Thailand and Malaysia: Constraints from new U-Pb zircon chronology of SE Asian tin granitoids. *Journal of the Geological Society*, 169(4), 489–500. <https://doi.org/10.1144/0016-76492011-107>
- Setiawan, N. I., Osanai, Y., Nakano, N., Adachi, T., Setiadj, L. D., & Wahyudiono, J. (2013). Late Triassic metatonalite from the Schwaner Mountains in West Kalimantan and its contribution to sedimentary provenance in the Sundaland. *Berita Sedimentologi*, 12, 4–12.
- Shellnutt, J. G., Lan, C. Y., Van Long, T., Usuki, T., Yang, H. J., Mertzman, S. A., et al. (2013). Formation of Cretaceous Cordilleran and post-orogenic granites and their microgranular enclaves from the Dalat zone, southern Vietnam: Tectonic implications for the evolution of Southeast Asia. *Lithos*, 182–183, 229–241. <https://doi.org/10.1016/j.lithos.2013.09.016>
- Shi, M. F., Lin, F. C., Fan, W. Y., Deng, Q., Cong, F., Tran, M. D., et al. (2015). Zircon U-Pb ages and geochemistry of granitoids in the Truong Son terrane, Vietnam: Tectonic and metallogenic implications. *Journal of Asian Earth Sciences*, 101, 101–120. <https://doi.org/10.1016/j.jseae.2015.02.001>
- Sun, B., Kaus, B. J. P., Yang, J., Lu, G., Wang, X., Wang, K., & Zhao, L. (2021). Subduction polarity reversal triggered by Oceanic Plateau accretion: Implications for induced subduction initiation. *Geophysical Research Letters*, 48(24), e2021GL095299. <https://doi.org/10.1029/2021gl095299>
- Sundell, K. E., & Macdonald, F. A. (2022). The tectonic context of hafnium isotopes in zircon. *Earth and Planetary Science Letters*, 584, 117426. <https://doi.org/10.1016/j.epsl.2022.117426>
- Tran, H. T., Zaw, K., Halpin, J. A., Manaka, T., Meffre, S., Lai, C. K., et al. (2014). The tam Ky-Phuoc Son Shear Zone in central Vietnam: Tectonic and metallogenic implications. *Gondwana Research*, 26(1), 144–164. <https://doi.org/10.1016/j.gr.2013.04.008>
- van Hattum, M. W. A., Hall, R., Pickard, A. L., & Nichols, G. J. (2006). Southeast Asian sediments not from Asia: Provenance and geochronology of north Borneo sandstones. *Geology*, 34(7), 589–592. <https://doi.org/10.1130/g21939.1>
- van Hattum, M. W. A., Hall, R., Pickard, A. L., & Nichols, G. J. (2013). Provenance and geochronology of Cenozoic sandstones of northern Borneo. *Journal of Asian Earth Sciences*, 76, 266–282. <https://doi.org/10.1016/j.jseae.2013.02.033>
- Vermeesch, P., Resentini, A., & Garzanti, E. (2016). An R package for statistical provenance analysis. *Sedimentary Geology*, 336, 14–25. <https://doi.org/10.1016/j.sedgeo.2016.01.009>
- Wang, C., Mitchell, R. N., Murphy, J. B., Peng, P., & Spencer, C. J. (2020). The role of megacontinents in the supercontinent cycle. *Geology*, 49(4), 402–406. <https://doi.org/10.1130/g47988.1>
- Wang, S., Mo, Y., Wang, C., & Ye, P. (2016). Paleotethyan evolution of the Indochina block as deduced from granites in northern Laos. *Gondwana Research*, 38, 183–196. <https://doi.org/10.1016/j.gr.2015.11.011>
- Wang, Y., Liu, Z., Murtadha, S., Cawood, P. A., Qian, X., Ghani, A., et al. (2022). Jurassic subduction of the Paleo-Pacific plate in Southeast Asia: New insights from the igneous and sedimentary rocks in West Borneo. *Journal of Asian Earth Sciences*, 232, 105111. <https://doi.org/10.1016/j.jseae.2022.105111>
- Wang, Y., Qian, X., Bin Asis, J., Cawood, P. A., Wu, S., Zhang, Y., et al. (2023). “Where, when and why” for the arc-trench gap from Mesozoic Paleo-Pacific subduction zone: Sabah Triassic-Cretaceous igneous records in East Borneo. *Gondwana Research*, 117, 117–138. <https://doi.org/10.1016/j.gr.2023.01.008>
- Wang, Y., Qian, X., Cawood, P. A., Gan, C., Zhang, Y., Zhang, F., et al. (2024). Temporal-spatial patterns of Mesozoic Paleo-Pacific and Tethyan supra-subduction systems in SE Asia: Key observations and controversies in Borneo and its surroundings. *Earth-Science Reviews*, 252, 104762. <https://doi.org/10.1016/j.earscirev.2024.104762>
- Wang, Y., Qian, X., Zhang, Y., Gan, C., Zhang, A., Zhang, F., et al. (2021). Southern extension of the Paleotethyan zone in SE Asia: Evidence from the Permo-Triassic granitoids in Malaysia and West Indonesia. *Lithos*, 398–399, 398–399. <https://doi.org/10.1016/j.lithos.2021.106336>
- Wang, Y., Wu, S., Qian, X., Cawood, P. A., Lu, X., Gan, C., et al. (2022b). Early Cretaceous subduction in NW Kalimantan: Geochronological and geochemical constraints from the Raya and Mensibau igneous rocks. *Gondwana Research*, 101, 243–256. <https://doi.org/10.1016/j.gr.2021.08.006>
- Wang, Y., Zhang, A., Qian, X., Asis, J. B., Feng, Q., Gan, C., et al. (2021). Cretaceous Kuching accretionary orogenesis in Malaysia Sarawak: Geochronological and geochemical constraints from mafic and sedimentary rocks. *Lithos*, 400–401, 400–401. <https://doi.org/10.1016/j.lithos.2021.106425>
- Wiedenbeck, M., Allé, P., Corfu, F., Griffin, W. L., Meier, M., Oberli, F., et al. (1995). Three natural zircon standards for U-Th-Pb, Lu-Hf, trace element and REE analyses. *Geostandards Newsletter*, 19, 1–23. <https://doi.org/10.1111/j.1751-908X.1995.tb00147.x>
- Williams, P. R., Johnston, C. R., Almond, R. A., & Simamora, W. H. (1988). Late Cretaceous to early tertiary structural elements of West Kalimantan. *Tectonophysics*, 148(3–4), 279–297. [https://doi.org/10.1016/0040-1951\(88\)90135-7](https://doi.org/10.1016/0040-1951(88)90135-7)
- Wu, F. Y., Yang, J. H., Xu, Y. G., Wilde, S. A., & Richard, J. W. (2019). Destruction of the north China craton in the Mesozoic. *Annual Review of Earth and Planetary Sciences*, 47(1), 173–195. <https://doi.org/10.1146/annurev-earth-053018-060342>
- Yu, Y., Qian, X., Azlan Mustapha, K., Sheldrick, T. C., Gan, C., Zhang, Y., & Wang, Y. (2022). Late Paleozoic–Early Mesozoic granitic rocks in Eastern Peninsular Malaysia: New insights for the subduction and evolution of the Paleo-Tethys. *Journal of Asian Earth Sciences*, 239, 105427. <https://doi.org/10.1016/j.jseae.2022.105427>
- Zhang, K., Liao, J., & Gerya, T. (2024). Onset of double subduction controls plate motion reorganisation. *Nature Communications*, 15(1), 1513. <https://doi.org/10.1038/s41467-024-44764-8>
- Zhang, W., Hu, Z., & Spectroscopy, A. (2020). Estimation of isotopic reference values for pure materials and geological reference materials. *Atomic Spectroscopy*, 41(3), 93–102. <https://doi.org/10.46770/AS.2020.03.001>

- Zhang, X., Chung, S.-L., Lai, Y.-M., Ghani, A. A., Murtadha, S., Lee, H.-Y., & Hsu, C.-C. (2019). A 6000-km-long Neo-Tethyan arc system with coherent magmatic flare-ups and lulls in South Asia. *Geology*, *47*(6), 573–576. <https://doi.org/10.1130/g46172.1>
- Zhang, Z.-Y., Hou, Z.-Q., Lü, Q.-T., Zhang, X.-W., Pan, X.-F., Fan, X.-K., et al. (2023). Crustal architectural controls on critical metal ore systems in South China based on Hf isotopic mapping. *Geology*, *51*(8), 738–742. <https://doi.org/10.1130/g51203.1>
- Zhao, Q. (2025). AGU Tectonics Supporting Information for: Interaction between the Tethyan and Paleo-Pacific tectonic domains in Southeast Asia: Late Triassic initiation of an inward-dipping double subduction system [Dataset]. *Mendeley Data*, *V1*. <https://doi.org/10.17632/8g669bt2bt.1>
- Zhou, Y., Carter, A., Wu, J., Yao, Y., Zhu, R., Liu, H., et al. (2023). Nature of the Paleo-Pacific subduction along the East Asian Continental Margin in the Mesozoic: Insights from the sedimentary record of West Sarawak, Borneo. *Geophysical Research Letters*, *50*(8), e2022GL102370. <https://doi.org/10.1029/2022gl102370>
- Zhu, K. Y., Li, Z. X., Xu, X. S., & Wilde, S. A. (2013). Late Triassic melting of a thickened crust in southeastern China: Evidence for flat-slab subduction of the Paleo-Pacific plate. *Journal of Asian Earth Sciences*, *74*, 265–279. <https://doi.org/10.1016/j.jseaes.2013.01.010>
- Zhu, R. X., & Xu, Y. G. (2019). The subduction of the West Pacific Plate and the destruction of the North China Craton. *Science China Earth Sciences*, *62*(9), 1340–1350. <https://doi.org/10.1007/s11430-018-9356-y>